



ELSEVIER



LOV-based reporters for fluorescence imaging

Anthony M Buckley¹, Jan Petersen², Andrew J Roe¹,
Gillian R Douce¹ and John M Christie²

Chromophore-binding domains from plant and bacterial photoreceptor proteins have recently gathered increasing attention as new sources of genetically encoded fluorescent proteins (FPs). In particular, FPs based on the flavin-binding LOV (light, oxygen, or voltage sensing) domain offer advantages over green fluorescent protein (GFP) owing to their smaller size, pH and thermal stability, utility under anaerobic conditions and their ability to generate reactive oxygen species. This review focuses on the potential applications of this emerging class of fluorescent reporters, discusses the advantages and limitations of LOV-based FPs, whilst offering insights regarding the further development of this technology for bioimaging and photodynamic therapy.

Addresses

¹Institute of Infection, Immunology & Inflammation, College of Medical, Veterinary & Life Sciences, University of Glasgow, Glasgow G12 8QQ, UK

²Institute of Molecular Cell and Systems Biology, College of Medical, Veterinary & Life Sciences, University of Glasgow, Glasgow G12 8QQ, UK

Corresponding author: Christie, John M (John.Christie@glasgow.ac.uk)

Current Opinion in Chemical Biology 2015, 27:39–45

This review comes from a themed issue on **Molecular imaging**

Edited by **Samie Jaffrey** and **Atsushi Miyawaki**

<http://dx.doi.org/10.1016/j.cbpa.2015.05.011>

1367-5931/© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Green fluorescent protein (GFP) from the jellyfish *Aequorea victoria* is widely used throughout cell and developmental biology to study protein dynamics within living cells [1]. Since its introduction, an extensive range of variants and related FPs have been generated through molecular evolution exhibiting improved fluorescence properties, as well as different emission colors that span the entire visible spectrum [2]. However, despite its wide range of applications, GFP and related FPs are reported to have several limitations. For instance, GFP requires molecular oxygen to catalyze the posttranslational cyclization of its fluorophore [3], which can restrict its utility under low-oxygen conditions such as tumor growth [4]. Its relatively large size (~25 kDa) can also lead to steric constraints associated with dysfunctional GFP fusion

proteins [5]. Slow maturation of the GFP fluorophore, as well as temperature and pH stability are other factors to consider. Consequently, considerable interest has been directed towards developing alternative genetically encoded fluorescent reporters to overcome these limitations.


LOV-domain function and fluorescence

LOV domains form a subset of the Per-ARNT-Sim (PAS) domain superfamily and were first identified in proteins regulated either by light, oxygen or voltage, hence the acronym LOV [6]. LOV domains specifically function as photosensory modules and typically bind flavin mononucleotide (FMN) as an ultraviolet (UV)/blue light-absorbing chromophore within the protein that comprises five antiparallel β -sheets and several α -helices (Figure 1a). These relatively small light-sensing moieties (~12–19 kDa; Table 1) are typically linked to a diverse range of effector domains, which together function as blue-light receptors in plants, fungi and bacteria [7]. In this context, the LOV domain serves as a modular photoswitch, placing protein activities, such as kinases, phosphodiesterases and DNA-binding under light control. This modularity of regulation has lately been exploited to create artificial photoswitches for the optical control of biological processes in an emerging field known as optogenetics [8].

The FMN chromophore provides the LOV domain with a yellow color, as well as an intrinsic green fluorescence when excited with UV/blue light [9]. Upon photoexcitation, LOV domains undergo a reversible photocycle involving formation of a covalent adduct between the FMN chromophore and a conserved cysteine residue within the protein (Figure 1b). FMN-cysteinyl adduct formation is accompanied by a loss of fluorescence that subsequently recovers when the protein is incubated in darkness coinciding with photoadduct decay [10,11]. Replacement of this cysteine within the LOV domain abolishes its photochemical activity, improving the fluorescent properties of the protein [12,13]. Mutation of this invariant cysteine is therefore a prerequisite for the generation LOV-based FPs [14*].

Several LOV-based or flavin-based FPs (FbFPs) have now been isolated and are available from a variety of sources (Table 1). These include BsFbFP and its *Escherichia coli* codon-optimized counterpart EcFbFP derived from the bacterial photoreceptor YtvA from *Bacillus subtilis* and PpFbFP, which was developed from a sensory box protein (SB2) from *Pseudomonas putida* [15*]. These

Figure 1



Structure and fluorescence properties of the LOV-based FP iLOV. **(a)** Structure of iLOV (PDB entry 4EES) visualized using PyMOL. The FMN fluorophore is shown as spheres. **(b)** Natural photocycle of the LOV domain. The FMN chromophore in the LOV domain emits green fluorescence upon photoexcitation. Blue light (BL) irradiation induces the formation of a covalent adduct between the FMN chromophore and a conserved cysteine residue that is no longer fluorescent. The photoadduct decays back to the fluorescing state in darkness or in response to near UV irradiation. **(c)** Excitation and emission spectra of iLOV fluorescence. The fluorescence excitation spectrum (dashed blue) was obtained using an emission wavelength of 495 nm, whereas the fluorescence emission spectrum (green) was measured using an excitation wavelength of 450 nm.

LOV-based FPs exhibit a broad wavelength excitation range between 300 and 500 nm with an excitation and emission maximum \sim 450 nm and \sim 495 nm, respectively (Figure 1c). An excitation peak is also visible \sim 380 nm, consistent with the absorbance properties of flavin-based photoreceptors [6]. The requirement for FMN does not restrict its utility as a fluorescent reporter, as this appears to be in plentiful supply in bacterial, plant and human cells [16^{••}] nor limit their targeting to subcellular compartments [16^{••},17]. Moreover, the fast maturation of LOV fluorescence (with minutes) constitutes one advantage over GFP as a noninvasive real-time reporter of transcriptional activities [18]. However, it is worth noting that LOV-based FPs engineered thus far exhibit fluorescence quantum yields (Q_F) that are significantly lower (\sim 0.2–0.4; Table 1) to that of GFP (0.6) [2], which could impact their utility for certain reporter applications.

Brightness and photostability

Several improvements in the brightness and photostability of LOV-based FPs have been made since they were first introduced in 2007 [15[•]]. This includes mutation of the invariant cysteine to facilitate reporter function as well as molecular evolution approaches. In addition, targeted mutagenesis of F37 in PpFbFP has produced variants with 2-fold increases in Q_F [19], whilst additional improvements have been made through mutations at Y112 and Q116 [20[•]]. Whether these mutations can be combined (F37, Y112 and Q116) to enhance the fluorescence of PpFbFP remains to be investigated.

Mutagenesis and DNA shuffling of LOV-coding sequences derived from *Arabidopsis thaliana*, together with fluorescence screening in living plant cells, led to the isolation of an improved LOV-based FP known as iLOV [16^{••}]. Subsequent molecular evolution of iLOV produced variants that are more readily detectable by confocal imaging in living cells due to order-of-magnitude increases in photostability [21] (Figure 2a). However, the Q_F of these photostable iLOV (phiLOV) derivatives appears to be reduced compared those of iLOV and other LOV-based FPs [20[•]], suggesting that there could be trade-offs to consider when engineering for photostability enhancements. In addition, there are likely to be limits in improving the overall brightness of LOV-based FPs given that their fluorescence is dictated by the flavin cofactor. While small spectral blue shifts in fluorescence emission (\sim 10 nm) have been reported [20[•],21], extending these properties beyond the green region of the spectrum will not be possible without replacing the FMN with a different fluorophore.

Natural diversity

In addition to mutagenesis, recent genome searches have uncovered a wide range of genetically diverse LOV-coding sequences that can be further exploited for fluorescence applications. For example, DsFbFP, derived from a LOV-coding sequence isolated from the marine α -proteobacterium *Dinoroseobacter shibae*, exhibits improved fluorescence over PpFbFP [20[•]], whilst CreiLOV, obtained from a photoreceptor protein from the fresh water alga *Chlamydomonas reinhardtii*, shows enhancements in brightness, photostability and pH tolerance over LOV-based FPs such as iLOV [22]. Thermal stability enhancements in reporter activity have also been reported for CreiLOV, through targeted mutagenesis of EcFbFP [23] and from genome mining for LOV-coding sequences from thermotolerant organisms [8]. Disappointingly, LOV-coding sequences appear to be absent from extremophiles [24], thus limiting the opportunity to exploit this source of genetic diversity.

Utility in low-oxygen environments

One hallmark of LOV-based FPs is the capacity of these proteins to fluoresce under low-oxygen environments

Table 1

Properties of LOV-based FPs

| Organism | Protein | FP | Molecular mass (kDa) | Oligomeric state | Quantum yield (Q_F) | Photostability $t_{50\%}$ (min) [20*] | Comments | Refs |
|----------------------------------|---------|----------------|----------------------|------------------|-------------------------|---------------------------------------|---|-----------------------------------|
| <i>Bacillus subtilis</i> | YtvA | EcFbFP | 15.1 | Dimer | 0.34–0.44 | 2.8 | <i>E. coli</i> codon optimized version of BsFbFP | [15*,18,20*] |
| <i>Pseudomonas putida</i> | SB1 | Pp1FbFP | 16.3 | Dimer | 0.27 | 6.8 | Increased brightness over Pp2FbFP | [20*] |
| | SB2 | Pp2FbFP | 16.3 | Dimer | 0.17–0.22 | 2.7 | Increased brightness over Pp2FbFP | [15*,18,20*] |
| | | Pp2FbFP F37T/S | 16.3 | Dimer | 0.24–0.30 | | | |
| | | | Pp2FbFP Y112L | 16.3 | Dimer | 0.30 | Increased brightness over Pp2FbFP | [20*] |
| | | | Pp2FbFP Q116V | 16.3 | Dimer | 0.26 | 1.5 | Increased brightness over Pp2FbFP |
| | | Pp2FbFP L30M | 16.3 | Dimer | 0.25 | 1.2 | Enhanced O ₂ photosensitization over miniSOG | [52] |
| <i>Arabidopsis thaliana</i> | Phot2 | iLOV | 12.1 | Monomer | 0.34–0.44 | | Smallest LOV reporter | [16**,18,22] |
| | | phiLOV2.1 | 12.1 | Monomer | 0.20 | 13 | Improved <i>in vivo</i> photostability over iLOV | [20*,21] |
| | | miniSOG | 12.1 | Monomer | 0.37–0.41 | 2.9 | ¹ O ₂ photosensitization | [20*,41**] |
| <i>Chlamydomonas reinhardtii</i> | Phot | CreiLOV | 13.0 | Monomer | 0.51 | | Enhanced brightness, photostability, thermal and pH tolerance <i>in vitro</i> | [22] |
| <i>Vaucheria frigida</i> | Aureo1 | VafLOV | 19.4 | Dimer | 0.23 | | | [22] |
| <i>Dinoroseobacter shibae</i> | Unknown | DsFbFP | 15.2 | Dimer | 0.35 | 0.4 | | [20*] |

[15*]. This property makes the application of this technology very attractive for the study of anaerobic cellular processes. For example, studies have shown that EcFbFP could surpass yellow fluorescent protein (YFP) as a fluorescent reporter in anaerobically grown cultures of *E. coli* and *Rhodobacter capsulatus* [15*,25,26], as well as yeast [27] and a range of mammalian cells [28]. In addition, LOV-based FPs have been used to track host cell infections by anaerobic pathogens under physiologically relevant conditions [29–31]. Similarly, our recent work has evaluated the utility of phiLOV2.1 as a translational fusion in the obligate anaerobe *Clostridium difficile*. Genetically fusing phiLOV2.1 to the bacterial cytokinetic protein FtsZ enables efficient monitoring of its localization to the septum during binary fission, where it becomes polymerized to form a characteristic Z-ring at the mid-cell (Figure 2b).

When size matters

Another benefit of LOV-based FPs over GFP is its smaller size. While some LOV-based FPs form dimers *in vitro*, iLOV, its derivatives and the newly isolated CreiLOV are monomeric (Table 1) and relatively small in size (110–118 amino acids). These latter properties confer advantages in creating translation fusions where inclusion of larger FPs could impact on protein function due to steric constraints [32]. For example, whilst fusion of GFP is known to compromise the functionality of some viral movement proteins, iLOV was found to be less


disruptive when fused to the movement protein of tobacco mosaic virus (Figure 2c) and the coat protein of potato mop-top virus [16**]. Furthermore, the coding sequence of iLOV (~300 nt) is considerably smaller than that of GFP (~700 nt) and provides a reduced genetic load with respect to viral packaging constraints. Hence, iLOV is reported to outperform GFP as a real-time reporter of viral infection dynamics and has successfully extended the range of plant and animal viruses that can be fluorescently tagged *in vivo* [16**,33,34].

Biosensor applications

In addition, LOV-based FPs have also been used in the development of several biosensor applications. For instance, iLOV fluorescence is quenched in the presence of copper at physiological pH and is reversible in the presence of the metal ion chelator EDTA [35]. Two asparagine residues within the vicinity of the FMN fluorophore at positions 401 and 425 (based on the amino acid positions in *Arabidopsis* phototropin 2 from which iLOV was derived) are proposed to coordinate copper binding. While mutation of these residues is required to confirm their involvement, these studies clearly demonstrate the potential for LOV-based FPs to serve as intracellular metal sensors.

The flavin-binding properties of LOV-based FPs also make them attractive candidates for developing

Figure 2



Utility of LOV-based FPs for bioimaging. **(a)** Photobleaching properties of iLOV and phiLOV expressed in liquid droplets of *E. coli*. Images were taken before photobleaching, immediately after bleaching and post-recovery after 5 min. Adapted from Christie *et al.* [21]. **(b)** Protein localization in the obligate anaerobic bacterium *Clostridium difficile*. The LOV-based FP, phiLOV2.1 was fused to the cytokinetic protein FtsZ. Green fluorescence from FtsZ-phiLOV2.1 is localized at the mid-cell during bacterial cell division (left). Accumulation of red fluorescence from the lipophilic membrane dye FM4-64 showing increased membrane biogenesis at the mid-cell (centre). Merge of green and red fluorescence overlapping at the mid-cell (right). Arrow denotes the point of cellular division. Adapted from Buckley *et al.* [58]. **(c)** Tobacco mosaic virus (TMV)-based expression of iLOV driven from a viral subgenomic promoter. White arrows show systemic movement of TMV expressing the iLOV reporter in the upper leaves of tobacco at 4 days post inoculation. Adapted from Chapman *et al.* [16**].

fluorescent sensors of photo-induced electron transport (PET), which can be used to monitor pH changes within living cells. For this purpose, iLOV was reengineered to incorporate unnatural tyrosine analogs at position 486 that exhibit a reduced pK_a [36**]. The phenolate anion of this side chain quenches iLOV fluorescence at physiological pH by PET. However, protonation of the tyrosine analog enhances fluorescence by uncoupling the PET pathway. Consequently, *E. coli* cells expressing the variant iLOVU3 show increased fluorescence when the pH of the medium was lowered 7–5. This ‘acid turn-on’ property of iLOVU3 therefore holds great promise for monitoring cellular pH changes and has been used effectively to track bacterial phagocytosis into macrophages [36**].

Oxidation of the tyrosine analog in iLOVU3 also blocks the PET quenching pathway, opening up new possibilities for using LOV-based FPs as redox sensors [36**]. Moreover, fusion of EcFbFP to YFP has proved successful in generating a Förster resonance energy transfer (FRET)-based biosensor named FluBO for visualizing temporal changes in oxygen levels in *E. coli* [37]. In this case, efficient FRET from the EcFbFP donor to the YFP acceptor is only observed in the presence of cellular oxygen.

Singlet oxygen generation

Aside from their intrinsic fluorescence, flavins offer great potential as photosensitizers for singlet oxygen generation (SOG). Consequently, LOV-based FPs have been developed as phototoxic modules capable of triggering cell death [38*,39*,40]. MiniSOG was the first LOV-based FP engineered for singlet oxygen (1O_2) photosensitization [41**]. In response to light, miniSOG generates a sufficient amount of 1O_2 to induce diaminobenzidine photooxidation and polymerization, which is discernible by correlative light and electron microscopy [41**]. Exquisite protein localization and complex protein structures within persevered subcellular organelles have been obtained with high resolution (~ 8 – 10 nm) using miniSOG as a genetically encoded tag [42–46]. This photosensitizing ability of miniSOG has therefore created new possibilities for electron microscopy. Additional applications for miniSOG include new methods for optical immunodetection [47], the development of new methodologies for imaging protein–protein interactions in intact cells [48**] and chromophore-assisted light inactivation, a powerful technique where 1O_2 photosensitization can be used to selectively interfere with the function of proteins in close proximity [49]. Detailed photochemical characterization indicates that miniSOG initiates other photo-induced electron transfer processes in addition to 1O_2 production [50,51]. Further photosensitizer improvements could thus be obtained through protein engineering strategies. Indeed, a Pp2FbFP derivative has recently been reported with a 1O_2 quantum yield that is ~ 3 -fold higher than that of miniSOG [52]. Establishing the 1O_2 photosensitizing capabilities of existing LOV-based FPs (Table 1) will also be important to minimize any potential phototoxic side effects during live cell imaging.

Super-resolution microscopy

Photoswitchable FPs are of considerable interest for super-resolution microscopy applications. The natural photocycle of the LOV domain has recently been exploited for such purposes. Irradiation with near UV light is reported to accelerate photoadduct decay within the LOV domain [53] (Figure 1b). Similarly, LOV-containing proteins, including YtvA, can photoswitch between their fluorescing, dark state and their non-fluorescing, active state upon exposure to blue light and near UV light, respectively [54]. This photoswitchable property can facilitate the

distinction between neighboring fluorophore molecules by separating their fluorescence emission in time thereby conferring resolution beyond the diffraction limit of light microscopes [55]. Using Fluorescence photoactivation localization microscopy (FPALM), the intracellular location of YtvA molecules within *E. coli* cell was resolved with an average precision of 35 nm [54]. These photochromic properties of the LOV domain will contribute to extending the range of genetically encoded tools available for super-resolution microscopy.

Conclusions

Since the first identification of the flavin-binding property of the LOV domain over 15 years ago [9], much progress has been made in characterizing the biophysical and structural properties of these photosensory modules. Rational and random protein-engineering approaches have proved successful in harnessing the fluorescence potential of these, and other photosensor domains, for specific applications in cell biology and biotechnology [56]. These photosensors are prevalent in bacteria, fungi, plant and algae thereby enhancing the diversity that can be used for these design principles. Further development is now required to identify additional LOV variants with improved fluorescence, photostability and photosensitizing properties. Whether recent computational predictions can be used to incorporate mutations within the fluorophore-binding pocket in order to shift the fluorescence emission maximum to longer wavelengths should also be investigated [57]. With the advantages of this emerging class of FPs now firmly established, a bright future lies ahead in maximizing their utility for a wide range of applications from fluorescence imaging to cancer therapy.

Acknowledgements

We apologize to colleagues whose work could not be cited due to space constraints. JMC acknowledges project support from the UK Biotechnology and Biological Sciences Research Council (BB/J016047/1; BB/K008129/1; BB/M002128/1), the Gatsby Charitable Foundation and the New Phytologist Trust. GRD gratefully acknowledges support from the Wellcome Trust (086418).

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Tsien RY: **Constructing and exploiting the fluorescent protein paintbox (Nobel Lecture)**. *Angew Chem Int Ed Engl* 2009, **48**:5612-5626.
2. Kremers GJ, Gilbert SG, Cranfill PJ, Davidson MW, Piston DW: **Fluorescent proteins at a glance**. *J Cell Sci* 2011, **124**:157-160.
3. Heim R, Prasher DC, Tsien RY: **Wavelength mutations and posttranslational autooxidation of green fluorescent protein**. *Proc Natl Acad Sci U S A* 1994, **91**:12501-12504.
4. Coralli C, Cemazar M, Kanthou C, Tozer GM, Dachs GU: **Limitations of the reporter green fluorescent protein under simulated tumor conditions**. *Cancer Res* 2001, **61**:4784-4790.
5. Thomas CL, Maule AJ: **Limitations on the use of fused green fluorescent protein to investigate structure-function relationships for the cauliflower mosaic virus movement protein**. *J Gen Virol* 2000, **81**:1851-1855.
6. Christie JM, Blackwood L, Petersen J, Sullivan S: **Plant flavoprotein photoreceptors**. *Plant Cell Physiol* 2015, **56**:401-413.
7. Losi A, Gartner W: **The evolution of flavin-binding photoreceptors: an ancient chromophore serving trendy blue-light sensors**. *Annu Rev Plant Biol* 2012, **63**:49-72.
8. Christie JM, Gawthorne J, Young G, Fraser NJ, Roe AJ: **LOV to BLUF: flavoprotein contributions to the optogenetic toolkit**. *Mol Plant* 2012, **5**:533-544.
9. Christie JM, Salomon M, Nozue K, Wada M, Briggs WR: **LOV (light, oxygen, or voltage) domains of the blue-light photoreceptor phototropin (nph1): binding sites for the chromophore flavin mononucleotide**. *Proc Natl Acad Sci U S A* 1999, **96**:8779-8783.
10. Kasahara M, Swartz TE, Olney MA, Onodera A, Mochizuki N, Fukuzawa H, Asamizu E, Tabata S, Kanegae H, Takano M et al.: **Photochemical properties of the flavin mononucleotide-binding domains of the phototropins from *Arabidopsis*, rice, and *Chlamydomonas reinhardtii***. *Plant Physiol* 2002, **129**:762-773.
11. Christie JM, Corchnoy SB, Swartz TE, Hokenson M, Han IS, Briggs WR, Bogomolni RA: **Steric interactions stabilize the signaling state of the LOV2 domain of phototropin 1**. *Biochemistry* 2007, **46**:9310-9319.
12. Salomon M, Christie JM, Knieb E, Lempert U, Briggs WR: **Photochemical and mutational analysis of the FMN-binding domains of the plant blue light receptor, phototropin**. *Biochemistry* 2000, **39**:9401-9410.
13. Swartz TE, Corchnoy SB, Christie JM, Lewis JW, Szundi I, Briggs WR, Bogomolni RA: **The photocycle of a flavin-binding domain of the blue light photoreceptor phototropin**. *J Biol Chem* 2001, **276**:36493-36500.
14. Mukherjee A, Schroeder CM: **Flavin-based fluorescent proteins: emerging paradigms in biological imaging**. *Curr Opin Biotechnol* 2015, **31C**:16-23.
- Comprehensive review providing detailed coverage on the utility of LOV-based FPs.
15. Drepper T, Eggert T, Circolone F, Heck A, Krauss U, Guterl JK, Wendorff M, Losi A, Gartner W, Jaeger KE: **Reporter proteins for in vivo fluorescence without oxygen**. *Nat Biotechnol* 2007, **25**:443-445.
- First evidence demonstrating the utility of LOV domains as *in vivo* fluorescent reporters under anaerobic conditions.
16. Chapman S, Faulkner C, Kaiserli E, Garcia-Mata C, Savenkov EI, Roberts AG, Oparka KJ, Christie JM: **The photoreversible fluorescent protein iLOV outperforms GFP as a reporter of plant virus infection**. *Proc Natl Acad Sci U S A* 2008, **105**:20038-20043.
- First study showing that LOV-based FPs offer greater utility as real-time reporters of virus infection.
17. Zhang LC, Risoul V, Latifi A, Christie JM, Zhang CC: **Exploring the size limit of protein diffusion through the periplasm in cyanobacterium *Anabaena* sp. PCC 7120 using the 13 kDa iLOV fluorescent protein**. *Res Microbiol* 2013, **164**:710-717.
18. Mukherjee A, Walker J, Weyant KB, Schroeder CM: **Characterization of flavin-based fluorescent proteins: an emerging class of fluorescent reporters**. *PLoS ONE* 2013, **8**:e64753.
19. Mukherjee A, Weyant KB, Walker J, Schroeder CM: **Directed evolution of bright mutants of an oxygen-independent flavin-binding fluorescent protein from *Pseudomonas putida***. *J Biol Eng* 2012, **6**:20.
20. Wingen M, Potzkei J, Endres S, Casini G, Rupprecht C, Fahlke C, Krauss U, Jaeger KE, Drepper T, Gensch T: **The photophysics of LOV-based fluorescent proteins – new tools for cell biology**. *Photochem Photobiol Sci* 2014, **13**:875-883.
- Up-to-date comparison of the fluorescent properties of different LOV-based FPs.

21. Christie JM, Hitomi K, Arvai AS, Hartfield KA, Mettlen M, Pratt AJ, Tainer JA, Getzoff ED: **Structural tuning of the fluorescent protein iLOV for improved photostability.** *J Biol Chem* 2012, **287**:22295-22304.
22. Mukherjee A, Weyant KB, Agrawal U, Walker J, Cann IKO, Schroeder CM: **Engineering and characterization of new LOV-based fluorescent proteins from *Chlamydomonas reinhardtii* and *Vaucheria frigida*.** *ACS Synth Biol* 2015, **4**:371-377.
23. Song X, Wang Y, Shu Z, Hong J, Li T, Yao L: **Engineering a more thermostable blue light photo receptor *Bacillus subtilis* YtvA LOV domain by a computer aided rational design method.** *PLoS Comput Biol* 2013, **9**:e1003129.
24. Krauss U, Minh BQ, Losi A, Gartner W, Eggert T, von Haeseler A, Jaeger KE: **Distribution and phylogeny of light–oxygen–voltage–blue-light-signaling proteins in the three kingdoms of life.** *J Bacteriol* 2009, **191**:7234-7242.
25. Drepper T, Huber R, Heck A, Circolone F, Hillmer AK, Buchs J, Jaeger KE: **Flavin mononucleotide-based fluorescent reporter proteins outperform green fluorescent protein-like proteins as quantitative in vivo real-time reporters.** *Appl Environ Microbiol* 2010, **76**:5990-5994.
26. Krol JE, Rogers LM, Krone SM, Top EM: **Dual reporter system for in situ detection of plasmid transfer under aerobic and anaerobic conditions.** *Appl Environ Microbiol* 2010, **76**:4553-4556.
27. Tielker D, Eichhof I, Jaeger KE, Ernst JF: **Flavin mononucleotide-based fluorescent protein as an oxygen-independent reporter in *Candida albicans* and *Saccharomyces cerevisiae*.** *Eukaryot Cell* 2009, **8**:913-915.
28. Walter J, Hausmann S, Drepper T, Puls M, Eggert T, Dihne M: **Flavin mononucleotide-based fluorescent proteins function in mammalian cells without oxygen requirement.** *PLoS ONE* 2012, **7**:e43921.
29. Landete JM, Peiroten A, Rodriguez E, Margolles A, Medina M, Arques JL: **Anaerobic green fluorescent protein as a marker of *Bifidobacterium* strains.** *Int J Food Microbiol* 2014, **175**:6-13.
30. Lobo LA, Smith CJ, Rocha ER: **Flavin mononucleotide (FMN)-based fluorescent protein (FbFP) as reporter for gene expression in the anaerobe *Bacteroides fragilis*.** *FEMS Microbiol Lett* 2011, **317**:67-74.
31. Choi CH, DeGuzman JV, Lamont RJ, Yilmaz O: **Genetic transformation of an obligate anaerobe, *P. gingivalis* for FMN-green fluorescent protein expression in studying host–microbe interaction.** *PLoS ONE* 2011, **6**:e18499.
32. Gawthorne JA, Reddick LE, Akpunarlieva SN, Beckham KS, Christie JM, Alto NM, Gabrielsen M, Roe AJ: **Express your LOV: an engineered flavoprotein as a reporter for protein expression and purification.** *PLoS ONE* 2012, **7**:e52962.
33. Seago J, Juleff N, Moffat K, Berryman S, Christie JM, Charleston B, Jackson T: **An infectious recombinant foot-and-mouth disease virus expressing a fluorescent marker protein.** *J Gen Virol* 2013, **94**:1517-1527.
34. van den Wollenberg DJ, Dautzenberg IJ, Ros W, Lipinska AD, van den Hengel SK, Hoeben RC: **Replicating reoviruses with a transgene replacing the codons for the head domain of the viral spike.** *Gene Ther* 2015, **22**:51-63.
35. Ravikumar Y, Nadarajan SP, Lee CS, Rhee JK, Yun H: **A new generation fluorescent based metal sensor – iLOV protein.** *J Microbiol Biotechnol* 2014, **25**:503-510.
36. Liu X, Jiang L, Li J, Wang L, Yu Y, Zhou Q, Lv X, Gong W, Lu Y, Wang J: **Significant expansion of fluorescent protein sensing ability through the genetic incorporation of superior photo-induced electron-transfer quenchers.** *J Am Chem Soc* 2014, **136**:13094-13097.
- Ingenious study engineering iLOV as a fluorescent sensor of photo-induced electron transport to monitor pH and redox changes.
37. Potzkei J, Kunze M, Drepper T, Gensch T, Jaeger KE, Buchs J: **Real-time determination of intracellular oxygen in bacteria using a genetically encoded FRET-based biosensor.** *BMC Biol* 2012, **10**:28.
38. Mironova KE, Proshkina GM, Ryabova AV, Stremovskiy OA, Lukyanov SA, Petrov RV, Deyev SM: **Genetically encoded immunophotosensitizer 4D5scFv-miniSOG is a highly selective agent for targeted photokilling of tumor cells in vitro.** *Theranostics* 2013, **3**:831-840.
- Insightful paper describing how singlet oxygen generation by miniSOG can be used to target cell death as a potential therapy for breast cancer.
39. Qi YB, Garren EJ, Shu X, Tsien RY, Jin Y: **Photo-inducible cell ablation in *Caenorhabditis elegans* using the genetically encoded singlet oxygen generating protein miniSOG.** *Proc Natl Acad Sci U S A* 2012, **109**:7499-7504.
- First paper to show the photokilling potential of miniSOG.
40. Ryumina AP, Serebrovskaya EO, Shirmanova MV, Snopova LB, Kuznetsova MM, Turchin IV, Ignatova NI, Klementieva NV, Fradkov AF, Shakhov BE et al.: **Flavoprotein miniSOG as a genetically encoded photosensitizer for cancer cells.** *Biochim Biophys Acta* 2013, **1830**:5059-5067.
41. Shu X, Lev-Ram V, Deerinck TJ, Qi Y, Ramko EB, Davidson MW, Jin Y, Ellisman MH, Tsien RY: **A genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms.** *PLoS Biol* 2011, **9**:e1001041.
- First demonstration of the utility of LOV-based FPs as singlet oxygen generators opening new possibilities for correlative light and electron microscopy.
42. Burgers PP, Ma Y, Margarucci L, Mackey M, van der Heyden MA, Ellisman M, Scholten A, Taylor SS, Heck AJ: **A small novel A-kinase anchoring protein (AKAP) that localizes specifically protein kinase A-regulatory subunit I (PKA-RI) to the plasma membrane.** *J Biol Chem* 2012, **287**:43789-43797.
43. Ludwig A, Howard G, Mendoza-Topaz C, Deerinck T, Mackey M, Sandin S, Ellisman MH, Nichols BJ: **Molecular composition and ultrastructure of the caveolar coat complex.** *PLoS Biol* 2013, **11**:e1001640.
44. Cleyrat C, Darehshouri A, Steinkamp MP, Vilaine M, Boassa D, Ellisman MH, Hermouet S, Wilson BS: **Mpl traffics to the cell surface through conventional and unconventional routes.** *Traffic* 2014, **15**:961-982.
45. Boassa D, Berlanga ML, Yang MA, Terada M, Hu J, Bushong EA, Hwang M, Masliah E, George JM, Ellisman MH: **Mapping the subcellular distribution of alpha-synuclein in neurons using genetically encoded probes for correlated light and electron microscopy: implications for Parkinson's disease pathogenesis.** *J Neurosci* 2013, **33**:2605-2615.
46. Boassa D, Nguyen P, Hu J, Ellisman MH, Sosinsky GE: **Pannexin2 oligomers localize in the membranes of endosomal vesicles in mammalian cells while Pannexin1 channels traffic to the plasma membrane.** *Front Cell Neurosci* 2014, **8**:468.
47. Wurtzler EM, Wendell D: **A streptavidin-SOG chimera for all-optical immunoassays.** *Biomacromolecules* 2014, **15**:228-233.
48. To TL, Fadul MJ, Shu X: **Singlet oxygen triplet energy transfer-based imaging technology for mapping protein–protein proximity in intact cells.** *Nat Commun* 2014, **5**:4072.
- Key study exploiting singlet oxygen generation by miniSOG to develop a new imaging technology for mapping protein proximity in large macromolecular complexes.
49. Lin JY, Sann SB, Zhou K, Nabavi S, Proulx CD, Malinow R, Jin Y, Tsien RY: **Optogenetic inhibition of synaptic release with chromophore-assisted light inactivation (CALI).** *Neuron* 2013, **79**:241-253.
50. Pimenta FM, Jensen RL, Breitenbach T, Etzerodt M, Ogilby PR: **Oxygen-dependent photochemistry and photophysics of 'miniSOG' a protein-encased flavin.** *Photochem Photobiol* 2013, **89**:1116-1126.
51. Ruiz-Gonzalez R, Cortajarena AL, Mejias SH, Agut M, Nonell S, Flors C: **Singlet oxygen generation by the genetically encoded tag miniSOG.** *J Am Chem Soc* 2013, **135**:9564-9567.
52. Torra J, Burgos-Caminal A, Endres S, Wingen M, Drepper T, Gensch T, Ruiz-Gonzalez R, Nonell S: **Singlet oxygen photosensitisation by the fluorescent protein Pp2FbFP L30M, a novel derivative of *Pseudomonas putida* flavin-binding Pp2FbFP.** *Photochem Photobiol Sci* 2015, **14**:280-287.

53. Kennis JT, van Stokkum IH, Crosson S, Gauden M, Moffat K, van Grondelle R: **The LOV2 domain of phototropin: a reversible photochromic switch.** *J Am Chem Soc* 2004, **126**:4512-4513.
54. Losi A, Gartner W, Raffelberg S, Cella Zanacchi F, Bianchini P, Diaspro A, Mandalari C, Abbruzzetti S, Viappiani C: **A photochromic bacterial photoreceptor with potential for super-resolution microscopy.** *Photochem Photobiol Sci* 2013, **12**:231-235.
55. Betzig E, Patterson GH, Sougrat R, Lindwasser OW, Olenych S, Bonifacino JS, Davidson MW, Lippincott-Schwartz J, Hess HF: **Imaging intracellular fluorescent proteins at nanometer resolution.** *Science* 2006, **313**:1642-1645.
56. Shcherbakova DM, Shemetov AA, Kaberniuk AA, Verkhusha VV: **Natural photoreceptors as a source of fluorescent proteins, biosensors, and optogenetic tools.** *Annu Rev Biochem* 2015, **84**:519-550.
57. Khrenova MG, Nemukhin AV, Domratcheva T: **Theoretical characterization of the flavin-based fluorescent protein iLOV and its Q489K mutant.** *J Phys Chem* 2015, **119**: 5176-5183.
58. Buckley A, Spencer J, Jukes C, Candish D, Irvine J, Fagan RP, Fairweather N, Roe AJ, Christie JM, Douce G: **Lighting up *Clostridium difficile*: reporting gene expression using fluorescent LOV domains.** *Sci Rep* 2015. (in preparation).