Larsen, N. K., Funder, S., Linge, H., Möller, P., Schomacker, A., Fabel, D., Xu, S., and Kjær, K. H. (2016) A Younger Dryas 1 re-advance of local glaciers in North Greenland. *Quaternary Science Reviews*, 147, pp. 47-58.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/111461/>

Deposited on: 23 October 2015

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

A Younger Dryas re-advance of local glaciers in North Greenland

Nicolaj K. Larsen^{a,b,*}, Svend Funder^b, Henriette Linge^c, Per Möller^d, Anders Schomacker^b, Derek Fabel^e, Sheng Xu^e, Kurt H. Kjær^b

^a*Department of Geoscience, Aarhus University, Høegh Guldbergs Gade 2, 8000 Aarhus C, Denmark*

^b*Centre for GeoGenetics, Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5-7, 1350 Copenhagen K, Denmark*

^c*Department of Earth Science, University of Bergen, and Bjerknes Centre for Climate Research, Postbox 7803, N-5020 Bergen, Norway*

^d*Department of Geology, Lund University, Sweden*

^e*AMS Laboratory, Scottish Universities Environmental Research Centre, East Kilbride G75 0QF, Scotland, UK*

Abstract

The Younger Dryas (YD) is a well-constrained cold event from 12,900 to 11,700 years ago but it remains unclear how the cooling and subsequent abrupt warming recorded in ice cores was translated into ice margin fluctuations in Greenland. Here we present ¹⁰Be surface exposure ages from three moraines in front of local glaciers on a 50 km stretch along the north coast of Greenland, facing the Arctic Ocean. Ten ages range from 11.6 ± 0.5 to 27.2 ± 0.9 ka with a mean age of 12.5 ± 0.7 ka after exclusion of two outliers. We consider this to be a minimum age for the abandonment of the moraines. The ages of the moraines are furthermore constrained using Optically Stimulated Luminescence (OSL) dating of epishelf sediments, which were deposited prior to the ice advance that formed the moraines, yielding a maximum age of 12.4 ± 0.6 ka, and bracketing the formation and subsequent abandonment of the moraines to within the interval 11.8-13.0 ka ago. This is the first time a synchronous YD glacier advance and subsequent retreat has been recorded for several independent glaciers in Greenland. In most other areas, there is no evidence for re-advance and glaciers were retreating during YD. We explain the different behaviour of the glaciers in northernmost Greenland as a function of their remoteness from the Atlantic Meridional Overturning Circulation (AMOC), which in other areas has been held responsible for modifying the YD drop in temperatures.

1. Introduction

The Younger Dryas (YD) marks a cold period at the end of the last glaciation, from 12.9 to 11.7 ka ago (Rasmussen et al., 2006). In spite of rising insolation during the YD, temperatures dropped, probably because the Atlantic Meridional Overturning Circulation (AMOC) was reduced or even shut down by large amounts of meltwater running off the melting ice sheets (Broecker et al., 1989; Denton et al., 2005; Carlson, 2013). The YD is especially pronounced in areas bordering the North Atlantic, and nowhere is it seen more clearly than in ice cores from the top of the Greenland ice sheet where the oscillation began with a c. 6°C drop in temperatures over a period of 200 years, and ended with an abrupt rise of c. 10°C over a period of only 60 years (Steffensen et al., 2008; Buizert et al., 2014). Knowledge about how these abrupt climate changes are translated into ice marginal behaviour along the Greenland ice sheet is of importance for evaluating the impact of future warming on the ice margin.

Here we present new evidence of a YD advance and subsequent retreat by three outlet glaciers from the North Cap, the local ice cap over the North Greenland mountain range (Fig. 1). The maximum positions of this advance in respective valley are marked by prominent terminal moraines, which ages are determined by new and previously published surface exposure ages, combined with previously published Optically Stimulated Luminescence (OSL) ages. We furthermore discuss the implications of our findings in the view of previous studies on the YD ice margin behaviour in other parts of Greenland.

2. Study area

The north coast of Greenland represents the land area closest to the North Pole. It comprises a 10 to 15 km wide coastal plain, bordered to the south by 500-1500 m high mountains. The coastal plain and connected valleys impinging into the mountain range southwards, host landforms and sediments recording glacial and marine events since the Last Glacial Maximum (LGM) in Greenland (Larsen et al., 2010). Based on the distribution of erratic boulders it was established that a local ice cap, the North Cap, developed over the mountain range in Peary Land during LGM, which to the south and southeast merged with the Greenland ice sheet (Koch, 1923; Dawes, 1986; Funder, 1989). On the coastal plain, erratics, till fabric measurements, and striations show that the outlet glaciers from the local ice caps merged with the Greenland ice sheet to form shelf-based ice in the Arctic Ocean that was deflected eastwards along the coast (Funder and Larsen, 1982; Dawes, 1986; Larsen et al.,

2010; Jakobsson et al., 2014). The deflection of the glaciers was most likely a result of buttressing by thick multiyear (palaeocrystic) sea ice in the Arctic Ocean (Bradley and England, 2008), which forced the local glaciers from Peary Land to flow along the coast (Larsen et al., 2010; Jakobsson et al., 2014). During the initial deglaciation large epishelf lakes were formed between the shelf-based ice and the mountains (Larsen et al., 2010; Möller et al., 2010). In these lakes, thick successions of laminated glaciolacustrine sediments were deposited on the coastal plain and in Sifs valley up to an elevation of 110 m a.s.l. (Figs. 2-3). OSL ages of the epishelf lake sediments ranged from 135 to 12.4 ka and the large spread suggests that they were differently affected by incomplete bleaching, with the youngest age serving as a maximum age for the deposition of the glaciolacustrine sediments (Larsen et al., 2010; Möller et al., 2010). During the final break-up of the shelf-based ice at ~10.1 cal. ka BP the coastal plain and the major valleys were inundated and marine sediments were deposited up to 40 m a.s.l. (Möller et al., 2010; Funder et al., 2011a). The isostatic uplift of the coastal plain following the deglaciation left a succession of beach ridges with driftwood that was used to constrain the Holocene sea ice history in the Arctic Ocean (Funder et al., 2011a). Following the deglaciation the local glaciers re-advanced through all major valleys along the coast and formed major terminal moraines on the coastal plain. Previously one of these moraines (Constable Bugt) was dated to between 9.6 and 6.3 cal. ka BP and there was further evidence of a second re-advance in Sifs Valley between 5.5-5.0 cal. ka BP (Möller et al., 2010). The two Holocene re-advances were linked to increased precipitation caused by more open water conditions in the Arctic Ocean during the warm Holocene Thermal Maximum. Our new results indicate that the moraine originated earlier, during the YD.

3. Methods

Ten samples from three terminal moraines on the coastal plain at Henson Bugt, Constable Bugt and Bliss Bugt were collected for surface exposure dating in this study (Fig. 2). Eight boulders were sampled using hammer and chisel, as well as a rock drill. The sampled boulder lithologies are meta-sandstone and quartz (Table 1). Two amalgamated pebble samples were collected from the sediment surface; rounded milky quartz pebbles were collected in sample bags in the field and split into uniform size categories (uniform a, b, c axis) before crushing. Sample locations and elevations were recorded in the field with a hand-held GPS. Clinometer measurements were taken for each sample for determination of the site-specific topographic shielding. Sample thickness was noted in the field (checked before crushing), as well as observations with regard to potential weathering loss (e.g. surface roughness, ventifacts,

differential weathering, striae) and snow shielding (e.g. geomorphology, snow patch distribution, lichen distribution and type).

3.1 ^{10}Be sample preparation and ^{10}Be age calculation

All rock samples were crushed at the Swedish Museum of Natural History (Stockholm), and processed at the School of Geographical and Earth Sciences (University of Glasgow). The 0.25-0.5 mm fraction from crushing and pulverizing was enriched with respect to quartz by treatment with HCl/HNO₃, magnetic separation, and H₃PO₄. Purification of the quartz fraction was done by successive HF/HNO₃ leaches, and purity (i.e. [Al] < 100 µg/g) was assessed by measuring the Al concentration using flame atomic absorption spectrometry (AAS) and inductively coupled plasma optical emission spectrometry (ICP-OES).

Preparation of samples for Be measurement was done at the Glasgow University – Scottish Universities Environmental Research Centre (GU – SUERC) Cosmogenic Isotope Laboratory, following procedures modified from Child et al. (2000). After carrier addition (see Table 1 for amounts and concentrations), clean quartz (11-27 g) was dissolved in 40 % HF. Be was extracted from solution by anion and cation exchange, precipitated as Be(OH)₂, dried and oxidized to BeO. AMS targets were made by mixing BeO and Nb (1:6) and pressing this into cathodes. The $^{10}\text{Be}/^9\text{Be}$ of these targets were measured at the SUERC AMS facility (Xu et al., 2010), and normalised to a nominal value of NIST SRM 4325 of 3.06×10^{-11} . Samples were prepared and measured between 2007 and 2013. Corresponding process blanks gave $^{10}\text{Be}/^9\text{Be}$ ratios between 1.7 and 6.1×10^{-15} (see Table 1 for sample/blank ratios and uncertainties). The calculated ^{10}Be concentrations have 1σ analytical errors between 3.4 and 6.0 %.

The CRONUS-Earth online calculator was used to estimate surface exposure ages from the calculated ^{10}Be concentrations (Balco et al., 2008). We report exposure ages ($\pm 1\sigma$, Table 1) calculated according to the St scaling scheme using the Arctic ^{10}Be production rate from Young et al. (2013b) (spallation-induced production rate of 3.93 ± 0.15 atoms $\text{g}^{-1} \text{a}^{-1}$, CRONUS-Earth; Alternate calibration data sets, Wrapper script 2.2, Main calculator 2.1, constants 2.2.1, muons 1.1) (Balco et al., 2008). The Arctic production rate is appropriate because of its geographic coverage and because the calibration data set extends back to ca. 16 ka. The analytical uncertainties are used when comparing results internally (i.e. ^{10}Be data only), whereas the systematic uncertainty (parentheses, last column, Table 1) should be used

when comparing the results with ages obtained using different cosmogenic nuclides or independent dating methods.

No correction for erosion, snow cover or landform degradation has been applied, but this does not mean that these factors are irrelevant – just hard to constrain. Erosion is very lithology specific, ranging from close to zero (amorphous quartz pebbles), granular weathering (meta-sandstones), to random loss of pieces (quartz veins/lenses). Snow cover can potentially be present most of the year, but no data on duration, thickness and density is available. Landform degradation encompasses several processes, such as wind deflation (evident from pebble lags) and freeze-thaw activity (polygonal cracking patterns, ice-wedge polygons and patterned ground), transforming the ridges from their initial unknown geometry to the present-day gently sloping features.

4. Results

4.1. Henson Bugt

In Henson Bugt, a large terminal moraine is located ~10 km from the present ice margin of a local glacier draining a small ice cap (Fig. 2). The ~30-40 m high and ~600 m wide moraine displays occasional large boulders of which three were sampled for surface exposure dating (Fig. 4). The small diameter of lichens on the boulder surfaces suggests an age of only 100 years (E.S. Hansen, pers.com.). However, the ‘lichen-kill’ effect of semi-permanent snowbeds in the recent past (Lévesque and Svoboda, 1999; Matthews, 2005) would hamper the applicability of lichenometric dating. Three boulder samples (PEA 0719, 0720, 0721) yield ^{10}Be surface exposure ages of 12.5 ± 0.6 , 13.5 ± 0.7 , and 12.4 ± 0.6 ka (Table 1, Fig. 3), revealing statistically indistinguishable ages, allowing a mean arithmetic age of 12.8 ± 0.8 ka to be calculated.

4.2. Constable Bugt

In Constable Bugt, a ~60 to 100 m high and ~1 km wide terminal moraine blocks the entrance to Sifs Valley (Figs. 2, 5). The moraine, located ~17 km from the present valley glacier margin, has a core composed of diamict and, in addition to this, glaciotectonically up-thrusted fine-grained glaciolacustrine sediments on its proximal part (Möller et al., 2010). Six processed samples (PEA 0611, 0613, 0618, 0621, 0645, 0647) range from 11.6 ± 0.5 to 27.2 ± 0.9 ka (Table 1, Fig. 3). From its stable setting, cobble on bedrock, sample PEA 0647 is expected to provide the most reliable ^{10}Be surface exposure age (assuming no inheritance),

and this cobble gives 12.6 ± 0.7 ka. Samples PEA 0613, 0618, 0645 and 0647 yield statistically indistinguishable ages, resulting in an arithmetic mean age of 12.1 ± 0.6 ka for the moraine ridge. The two older ^{10}Be surface exposure ages can likely be explained by inheritance. Occasional boulders showing inheritance are to be expected, however, the amalgamated pebble sample (PEA 0611) yields the oldest age, suggesting that the re-worked material comprising the moraine has a complex exposure history prior to moraine ridge formation.

4.3. Bliss Bugt

In Bliss Bugt a ~10-30 m high moraine is located ~8 km from the present ice margin of Moore Glacier (Fig. 2). The surface consists of sorted coarse-grained sediments with few scattered boulders (Fig. 6). One amalgamated sample of surface pebbles from the sediment ridge gives a ^{10}Be surface exposure age of 12.8 ± 0.7 ka (Table 1, Fig. 3). This sample was included in the study by Möller et al. (2010), but because its ^{10}Be surface exposure age was considered to be an outlier (significantly older than the data set from Bliss Bugt at that time), no conclusions were drawn. However, since the age overlaps with the new ^{10}Be data from the moraine ridges at Constable Bugt and Henson Bugt we find it to be reliable.

5. Discussion

5.1. Younger Dryas moraines in North Greenland

In total, we dated eight boulders and two amalgamated pebble samples from the moraines on the north coast of Greenland. Two samples are significantly older than the rest and considered as outliers, suffering from inheritance. However, the consistent ages of the majority of samples suggest that inheritance is not a significant problem; otherwise a more scattered age distribution would have been expected, such as noted for erratics on moraines in Scoresby Sund (Kelly et al., 2008). We attempted to minimize the post-depositional effects by sampling the largest boulders on the flat surface of the moraine crests at Henson Bugt and Constable Bugt. Long-term moraine degradation also plays an insignificant role in West Greenland where the Fjord Stade moraines were precisely dated to 9.3 and 8.2 ka by ^{10}Be measurements, controlled by threshold lake records (Young et al., 2011; Young et al., 2013a). These uncertainties, in addition to other processes such as shielding by long lasting snow cover, moraine degradation where boulders are slowly exhumed until the moraine stabilizes, or because of dead-ice melting (Heyman et al., 2011; Houmark-Nielsen et al., 2012), have the potentials of making the ages too young. We therefore treat all ages as minimum ages.

Accordingly, we calculate an arithmetic mean minimum age of 12.5 ± 0.7 ka (n=8), excluding the two outliers mentioned above (Fig. 7).

Clearly, our new ^{10}Be results yielding a YD age of the moraines seems incompatible with our previous interpretation which suggests an age between 9.6 and 6.3 cal. ka BP (Möller et al., 2010). However, the two interpretations are not necessarily mutually exclusive. The consistency of ^{10}Be ages on three independent moraines along 50 km of coast is very compelling. So is also the radiocarbon age (~ 10.3 cal. ka BP) of the glaciotectonically up-thrusted sediments on the proximal side of the moraine in Constable Bugt (site CB0705; Fig. 5 in Möller et al., 2010), suggesting a younger age for this part of the moraine. With the new ^{10}Be ages along the major part of the Constable Bugt moraine, we suggest that its core was formed during the YD advance, while the up-thrusted sediments on its proximal side are related to a younger re-advance between 9.6 and 6.3 cal. ka BP (Fig. 8). Given that the coastal plain was inundated 10.1 cal. ka BP, following the break-up of the ice on the shelf, marine sediments were deposited on the distal part of the moraine up to 40 m a.s.l. in Constable Bugt. Thus, the surficial diamict with molluscs (~ 10.1 and 9.6 cal. ka BP) located below the marine limit (site CB0706; Fig. 5 in Möller et al., 2010) is re-interpreted. We thus suggest it represents a marine drop till or, more likely, a young solifluction lobe with diamictized marine sediment on the distal part of the moraine, rather than being glacially remoulded marine sediment from the interior of Sifs valley at terminal moraine formation, as suggested by Möller et al. (2010).

We can furthermore constrain the age of moraines by re-evaluating the retrieved OSL ages of the epishelf sediments, deposited during the break-up of the LGM shelf-based ice and before the moraines were formed (Larsen et al., 2010; Möller et al., 2010). The large spread of older OSL ages, from 135 to as young as 12.4 ka, with the youngest ages likely a maximum age interval for the formation of the moraines as discussed above (Figs. 3, 7). Besides giving a lower age constrain for the moraine, the presence of epishelf sediments in Sifs valley furthermore demonstrates that the glacier advanced to the moraine from up-valley, *i.e.* the moraine was not formed by still-stand during stepwise deglaciation.

In summary, the age of the moraines are bracketed by the minimum limiting ^{10}Be surface exposure ages of 12.5 ± 0.7 and the maximum limiting OSL age of 12.4 ± 0.6 ka, implying that the moraines were formed and abandoned within an age interval of 11.8-13.0 ka. The ^{10}Be surface exposure ages are derived from three terminal moraines formed by independent glaciers over a distance of 50 km. This indicates that the glaciers advanced as a response to external climate forcing rather than local topography or surging, and we conclude

that the moraines formed during an advance in response to the YD cooling. However, the large uncertainties both in surface exposure and OSL ages preclude an assessment of the duration of moraine formation, and its more precise age within the YD.

5.2. Younger Dryas ice margin fluctuations in Greenland

In other parts of Greenland the ice margin response to the YD cooling is more ambiguous. This is seen from recent ^{10}Be surface exposure dating of moraines and detailed seismic studies on the shelf, accompanied by AMS ^{14}C -dated marine cores (Fig. 9). Below is a brief review of the evidence and the climatic background given by the authors.

5.2.1. East Greenland

In the East Greenland Fjord Zone, between 70° and 77°N, a belt of moraines - the Milne Land moraines - were formed by outlet glaciers from the ice sheet at the mouths of all major fjords (Fig. 9). The moraines occur in swarms forming a generally 5-10 km but up to 25 km wide belt. Distinct weathering limits between the outer moraines and the terrain distal to them and also their impressive dimensions – probably the highest and longest moraine ridges in Greenland – suggest that the moraine formation began with a re-advance after a period when the glaciers had been farther back in the fjords (Hjort, 1981; Funder, 1989; Landvik, 1994). In support of this, marine cores from the outer fjord area of Scoresby Sund may indicate that, this part of the fjord was deglaciated already during the Allerød (Marienfeld, 1992; Dowdeswell et al., 1994; Funder et al., 1998). Cessation of ice rafted debris (IRD) deposition on the shelf along the fjord zone during the Bølling (c. 13.8 ka cal. BP), suggests that, by this time, the major outlet glaciers had retreated from the inner shelf and terminated in the fjords where a large part of their IRD was trapped (Stein et al., 1993; Funder et al., 1998). Concordantly, this evidence suggests that the outer Milne Land moraines were formed during re-advance or a long lasting still stand, and not as a step during deglaciation.

Results from radiocarbon dating of marine shells associated with the Milne Land moraines indicate that, in the entire region, the belt of moraines was abandoned and rapid deglaciation begun by the end of the Preboreal Oscillation (11.3-11.15 ka cal. BP) (Björck et al., 1997). Both in Scoresby Sund and on Hochstetter Forland, 500 km to the north, the moraine formation was accompanied by a fall of 30-40 m in relative sea level from the outer to the inner moraine, suggesting that they were formed during a protracted period of time (Funder, 1978; Hjort and Björck, 1984; Hall et al., 2010). This has later been confirmed by an

extensive dataset comprising both ^{10}Be exposure and AMS ^{14}C ages from the moraines in Scoresby Sund. These data indicate that the advance of the region's largest outlet glacier most likely culminated in late Allerød times (>12.9 ka cal. BP), and that moraine formation ceased at 11-11.2 ka cal. BP (Hall et al., 2008). ^{10}Be surface exposure dating of erratic boulders on moraines from a local ice cap, correlated with the Milne Land moraines, gives a slightly older age for the period, from 13 to 11.6 ka (Kelly et al., 2008).

Unfortunately the time resolution of the dating prevents conclusions about the character of the retreat from the oldest moraines - whether it was gradual or punctuated by significant re-advance, or possibly even consisted of only two distinct re-advances – one in late Allerød times and the other in the late Preboreal, because no moraines have been directly dated to YD, neither in Scoresby Sund, nor in more northerly areas. Hjort and Björck (1984) suggested that the moraines on Hochstetter Forland were separated by an ice free interval, followed by a re-advance, and Denton et al. (2005) and Hall et al. (2010) suggested that the Preboreal re-advance could have obliterated the missing moraines from late YD. Still, even though no moraines have been dated to the YD, it is a reasonable assumption that the glacier fronts were located within the zone of the Milne Land moraines during the YD, and could be represented by undated moraines between the outer and inner Milne Land moraines. This would imply that the YD here was a period of oscillatory retreat within a narrow a zone.

Denton et al. (2005) noticed a mismatch between the YD temperatures deduced from moraine altitudes and those derived from borehole temperatures in nearby ice cores. It was suggested that while the borehole temperatures record mean annual temperature, the equilibrium-line altitudes deduced from moraines, mainly reflect summer temperatures. This implies that the YD temperature drop was mainly a winter phenomenon with little effect on the ice margin. This increase in seasonality was seen as a result of a reduction in the AMOC, which led to increased sea ice and reduced inflow of advective heat to the North Atlantic (Denton et al., 2005).

In contrast to this, in the Kangerlussuaq area to the south (c. 67°N), both shelf and fjord experienced large-scale retreat of the ice sheet margin in the YD (Fig. 9). This was shown by the occurrence of light isotopes in foraminifer shells from marine sediments on the shelf, as well as onset of IRD deposition during YD, which indicate a large supply of meltwater and break-up of the ice sheet margin (Jennings et al., 2006). A “major construction feature” c. 50 km off the coast could be a YD terminal moraine (Andrews et al., 1997), and is the only possible indication of re-advance during or before YD. It was suggested that the massive retreat was caused by advection of warm Atlantic intermediate water through the

Irminger Current, which overruled the climatic cooling. This should also have persisted when the ice margin retreated into the long and deep fjord, which was almost completely deglaciated during late YD, as shown recently by ^{10}Be surface exposure ages (Dyke et al., 2014).

5.2.2. Southeast Greenland

Farther to the south, at Sermilik (66°N) and Bernstorffs Fjord (64°N) (Fig. 9), ^{10}Be surface exposure ages and ^{14}C ages lake sediments show that large outlet glaciers here remained on the inner shelf until c. 11 ka (Long et al., 2008; Roberts et al., 2008; Dyke et al., 2014; Larsen et al., 2015). Apparently the warm Irminger Current water did not reach this coast and ice margin, in spite of its appearance at the shelf break outside Bernstorffs Fjord, as shown by sea-bed and core data (Kuijpers et al., 2003). Here, the warm Irminger Current water appeared already at 14.5 cal. ka BP, continuing through the YD with no apparent increase in sea ice or reduction in AMOC (Kuijpers et al., 2003; Knutz et al., 2011). It is difficult to reconcile the land and marine records from this area, especially since the shelf here has a width of only 50 km, but the ice sheet margin apparently stayed on the inner shelf during the YD.

5.2.3. South Greenland

In a marine sediment core from the Eirik Drift to the south of Greenland's southern tip (c. 58°N) Carlson et al. (2008) found an increase in Greenland detritus during YD, implying increased runoff from the ice sheet (Fig. 9). This is in agreement with evidence from isolation lakes in the Nanortalik district in southernmost Greenland, indicating rapid uplift, i.e. ice thinning retreat, especially during the late YD (Sparrenbom et al., 2006). In a lake core near the southern tip of Greenland (c. 60°N) - the only lake sediment record in Greenland reaching beyond YD - the Allerød/YD transition is characterized by an increase in lake productivity and occurrence of a warmth demanding diatom, indicating that YD was warmer than Allerød (Björck et al., 2002). This "anomalous" warming was explained as a result of reduced AMOC and increased sea ice and seasonality (Björck et al., 2002). This is the same mechanism that was proposed to afford the glacier advance in Scoresby Sund, but here in the south the increased seasonality resulted in summer temperatures that were high enough to promote melting. It is notable that the beginning of the Holocene in this lake is marked by a sharp transition from minerogenic to organic sediment – akin to the boundary

seen in NW European lakes, and so far the only evidence of the abrupt warming at the end of the YD found in ice-free Greenland.

5.2.4. West Greenland

In West Greenland, outlet glaciers in all major fjords coalesced on the shelf. Here, two large moraine systems, the Hellefiske and Fiskebanke moraines were traced for c. 500 km during oil-prospecting in the 1970's (Kelly, 1985; Funder et al., 2011b) (Fig. 9). The Hellefiske moraine follows the shelf edge, while the Fiskebanke moraine reflects a lobate ice sheet margin on the inner shelf and in the shelf troughs. These moraines have alternatively been referred to as Saalian/LGM or LGM/YD in their formation (Funder et al. 2011). Recent results from Disko Bugt show that a large outlet glacier reached the shelf edge, more than 200 km off the coast, at the LGM (see below), suggesting that the ice sheet margin also reached the much narrower shelf edge farther south, and that the oldest of the moraines were not older than the LGM. However, as none of the moraines have yet been dated, and as no YD moraines have so far been observed elsewhere in West Greenland, it remains an open question if either the Hellefiske or Fiskebanke moraines were formed during the YD, but recent ^{10}Be surface exposure ages from coastal areas show that the ice sheet margin remained on the shelf in this part of Greenland up until shortly before 11 or even 10 ka ago (Roberts et al., 2009; Larsen et al., 2014).

In Disko Bugt, the largest drainage trough in western Greenland (c. 70°N), ^{14}C ages of shells from marine cores show that the large ice stream, which occupied the trough outside Disko Bugt, may have begun its retreat at 13.8 cal. ka BP (Fig. 9) (O'Cofaigh et al. 2012). However, by c. 12.2 cal. ka BP, in mid-YD times, the retreat was interrupted by a short-lived re-advance almost to the shelf edge, followed by "instantaneous" collapse (O'Cofaigh et al., 2012; Kelley et al., 2013; Jennings et al., 2014; Rinterknecht et al., 2014). The re-advance amounted to almost 100 km along a c. 50 km wide front. This is seen from ^{14}C ages of shells from marine cores, both incorporated in till and *in situ* in the overlying marine mud on the inner shelf (O'Cofaigh et al., 2012; Rinterknecht et al., 2014). The collapse brought the ice margin back from the shelf-edge c. 200 km to the mouth of Disko Bugt (Fig. 9). The short duration of this re-advance/retreat suggests that it could represent a surge of a thin glacier lodged in the trough and controlled by subglacial topography and ice thickness, and not directly by the YD cooling (O'Cofaigh et al., 2012). Both before and especially after the re-advance, the Disko Bugt glacier retreated rapidly over the shelf through large-scale calving (Jennings et al., 2014). ^{10}Be surface exposure ages from the coastal area at the mouth of

Disko Bugt show that deglaciation of the bay began between 12.2 and 11.6 ka ago (Kelley et al., 2013; Rinterknecht et al., 2014). According to Rinterknecht et al. (2014) the collapse of the shelf-bound part of the Disko Bugt ice stream was triggered by the arrival of warm subsurface water as a result of a reduced AMOC.

5.2.5. Northwest Greenland

Farther north, in the Uummannaq Trough (c. 71°N), ¹⁴C dated marine cores show that deglaciation from the outer shelf began c. 14.8 cal. ka BP, and ¹⁰Be surface exposure ages from coastal mountains show that by 12.4 ka ago the outer and mid shelf, as well as the mountains, were cleared of ice (Fig. 9). After this, the recession picked up speed, and in the following millennium 100 km of the fjords were deglaciated (Roberts et al., 2013).

Accordingly, the YD, and especially its mid and late parts, appears to be a period of retreat.

Along the northwest coast of Greenland, the YD ice sheet margin was also located on the shelf. A marine core from the entrance to Nares Strait between Canada and Greenland (c. 77°N) shows that the cored site was deglaciated before 12.5 cal. ka BP and that meltwater sediments dominated for some centuries (Knudsen et al., 2008).

5.3. Synthesis of the Greenland ice sheet's response to the YD cooling

As noted above, the circumstance that three major glaciers over a distance of 50 km from east to west advanced and retreated synchronously imply that this advance was not caused by local forcing, such as the short-lived re-advance on the shelf off Disko Bugt, but by external, climatic factors. Surprisingly, the survey of ice margin behavior in other parts of Greenland shows that no other moraines have been dated to the YD, although some moraines are under discussion (Fig. 9). The Milne Land moraines, the only other evidence in Greenland for significant late glacial ice margin re-advance, apparently culminated in late Allerød times, but smaller re-advances of the outlet glaciers may have occurred during YD, represented by undated moraines in the area. In more southerly areas, wherever a record is available, the ice margins seem to have retreated from the shelf and in the fjords as shown by the records from Kangerlussuaq, Nanortalik, Uumannaq Fjord and especially Disko Bugt, where the retreat in late YD amounted to large scale collapse (Jennings et al., 2006; Sparrenbom et al., 2006; Carlson et al., 2008; O'Cofaigh et al., 2012; Roberts et al., 2013; Dyke et al., 2014; Jennings et al., 2014; Rinterknecht et al., 2014).

There is therefore a striking mismatch between the dramatic cooling and later abrupt warming seen in ice cores, and the retreat of the ice margins at most sites where a record is

available. This has generally been explained as a result of a reduction in AMOC and accompanying increase of seasonality (Björck et al., 2002, Denton et al., 2005, Carlson et al., 2008, Hall et al., 2008, Kelly et al., 2008, Rinterknecht et al., 2014). The reduced AMOC is thought to have been caused by a large supply of meltwater to the North Atlantic from melting ice sheets, which freshened the sea surface and reduced the advection of heat into the North Atlantic. This in turn increased sea ice extent, which sealed the ocean from exchange with the atmosphere, resulting in higher seasonality in Greenland. In addition, reduction in the AMOC may also have increased production of warm subsurface water at low latitudes, which advected northwards and promoted melting of shelf-bound ice margins (Marcott et al., 2011), as noted for the shelf-bound ice margins at Kangerlussuaq and Disko Bugt (Jennings et al., 2006, Rinterknecht et al., 2014).

These mechanisms may explain why the cold oscillation seen in ice cores did not lead to significant re-advance, and all processes are tied up to oceanographic perturbations originating in the North Atlantic. It is therefore noteworthy that areas with significant YD ice margin retreat in Greenland all are located in areas which today lie alongside the warm Irminger and Greenland Currents, which transport warm Atlantic water to the coasts of Southeast and West Greenland, while the areas with advance or sluggish retreat – Peary Land and the East Greenland Fjord Zone – are outside this realm (Fig. 9). This underlines Carlson et al.'s (2008) contention that ice sheets responded differently to cooling, depending on their geographic distribution relative to heat transport. In concert with this, Buizert et al. (2014) from analyses of ice cores showed that YD seasonality, a measure of the AMOC intensity, decreases from south to north, away from the influence of warm Atlantic water. We suggest that the reason for the YD glacier advance/retreat in Peary Land, at a time when ice margins generally retreated, may lie in the respective distances of these areas from North Atlantic advective heat, with Greenland's north coast being the farthest away. However, the coupling between the YD cooling and the subsequent glacier response remains enigmatic, in part because there are very few well-dated sites from this period in Greenland.

6. Conclusion

We have constrained the age of three terminal moraines in front of local glaciers along a 50 km stretch on the north coast of Greenland, using OSL and surface exposure dating techniques. The minimum limiting ^{10}Be surface exposure ages on erratics on the moraines show a mean age of 12.5 ± 0.7 ka and the maximum limiting OSL age for epishelf-lake sediments immediately preceding moraine formation give an age of 12.4 ± 0.6 ka. These ages

bracket the formation and subsequent abandonment of the moraines to within the interval 11.8-13.0 ka. The synchronous advance of independent glaciers suggests that they responded to external climate forcing rather than being controlled by topography or being the result of glacier surging. Occurrence of older glaciolacustrine sediments up valley from one of the moraines shows that this valley was ice free before the advance, and this is the first record of YD glacier re-advance/retreat in Greenland. In all other studied areas of Greenland the ice margins were retreating at this time, probably sluggishly in northern East Greenland, but rapidly in more southerly areas. The difference in ice margin response to the YD cooling may be explained by distance to the AMOC, with East and North Greenland being the farthest away. Where data are available, the retreat began before YD, and there is so far none, or little, evidence of a direct ice margin response, neither to the early YD cooling nor to the subsequent rapid Preboreal warming. This may to some extent owe to the limits of the available dating techniques and inaccessibility of critical sites, but in the light of the ongoing warming and its dramatic effect on the Greenland ice sheet, the problem deserves more attention.

Acknowledgements

The Danish National Research Council (FNU), Carlsbergfondet and the Swedish Research Council (VR) supported the project. NKL acknowledges support from Villum Fonden. Laboratory analyses benefited from internal funding from the Bjerknes Centre for Climate Research, HL thanks the Swedish Museum of Natural History for access to their crushing facilities, and Maria Miguens-Rodriguez, Delia Gheorghiu, Hannah Mathers, Fotini Karakitsos and Alessa Geiger for assistance during laboratory processing of the samples. Eric Steen Hansen and Eske Willerslev provided field assistance. We thank Guest editor Jason Briner, Nicolas Young and an anonymous reviewer for their constructive comments, which helped improve the manuscript.

References

- Andrews, J.T., Smith, L.M., Preston, R., Cooper, T., Jennings, A.E., 1997. Spatial and temporal patterns of iceberg rafting (IRD) along the East Greenland margin, ca 68 degrees N, over the last 14 cal ka. *Journal of Quaternary Science* 12, 1-13.
- Balco, G., Stone, J.O., Lifton, N.A., Dunai, T.J., 2008. A complete and easily accessible means of calculating surface exposure ages or erosion rates from Be-10 and Al-26 measurements. *Quat Geochronol* 3, 174-195.

473 Björck, S., Bennike, O., Rosen, P., Andresen, C.S., Bohncke, S., Kaas, E., Conley, D., 2002.
 474 Anomalously mild Younger Dryas summer conditions in southern Greenland. *Geology*
 475 30, 427-430.

476 Björck, S., Rundgren, M., Ingolfsson, O., Funder, S., 1997. The Preboreal oscillation around
 477 the Nordic Seas: terrestrial and lacustrine responses. *Journal of Quaternary Science* 12,
 478 455-465.

479 Bradley, R.S., England, J.H., 2008. The Younger Dryas and the Sea of ancient ice.
 480 *Quaternary Res* 70, 1-10.

481 Broecker, W.S., Kennett, J.P., Flower, B.P., Teller, J.T., Trumbore, S., Bonani, G., Wolfli,
 482 W., 1989. Routing of Meltwater from the Laurentide Ice-Sheet during the Younger
 483 Dryas Cold Episode. *Nature* 341, 318-321.

484 Buizert, C., Gkinis, V., Severinghaus, J.P., He, F., Lecavalier, B.S., Kindler, P., Leuenberger,
 485 M., Carlson, A.E., Vinther, B., Masson-Delmotte, V., White, J.W.C., Liu, Z.Y., Otto-
 486 Bliesner, B., Brook, E.J., 2014. Greenland temperature response to climate forcing
 487 during the last deglaciation. *Science* 345, 1177-1180.

488 Carlson, A., Stoner, J.S., Donnelly, J.P., Hillaire-Marcel, C., 2008. Response of the southern
 489 Greenland Ice Sheet during the last two deglaciations. *Geology* 36, 359-362.

490 Carlson, A.E., 2013. The Younger Dryas Climate Event. *Encyclopedia of Quaternary Science*
 491 3, 126-134.

492 Child, D., Elliott, G., Mifsud, C., Smith, A.M., Fink, D., 2000. Sample processing for earth
 493 science studies at ANTARES. *Nuclear Instruments and Methods in Physics Research*
 494 Section B172, 856-860.

495 Dawes, P.R., 1986. Glacial erratics on the Arctic Ocean margin of North Greenland:
 496 implications for an extensive ice-shelf. *Bulletin of the Geological Society of Denmark*
 497 35, 59-69.

498 Denton, G.H., Alley, R.B., Comer, G.C., Broecker, W.S., 2005. The role of seasonality in
 499 abrupt climate change. *Quaternary Science Reviews* 24, 1159-1182.

500 Dowdeswell, J.A., Uenzelmannneben, G., Whittington, R.J., Marienfeld, P., 1994. The Late
 501 Quaternary Sedimentary Record in Scoresby Sund, East Greenland. *Boreas* 23, 294-
 502 310.

503 Dunne, J., Elmore, D., Muzikar, P., 1999. Scaling factors for the rates of production of
 504 cosmogenic nuclides for geometric shielding and attenuation at depth on sloped
 505 surfaces. *Geomorphology* 27, 3-11.

506 Dyke, L.M., Hughes, A.L.C., Murray, T., Hiemstra, J.F., Andresen, C.S., Rodes, A., 2014.
507 Evidence for the asynchronous retreat of large outlet glaciers in southeast Greenland at
508 the end of the last glaciation. *Quaternary Science Reviews* 99, 244-259.

509 Funder, S., 1978. Holocene stratigraphy and vegetation history in the Scoresby Sund area,
510 East Greenland. *Bulletin Grønlands Geologiske Undersøgelse* 129, 1-66.

511 Funder, S., 1989. Quaternary geology of the ice-free areas and adjacent shelves og
512 Greenland, In: Fulton, R.J. (Ed.), *Quaternary geology of Canada and Greenland*.
513 Geological Survey of Canada, *Geology of Canada*, pp. 743-792.

514 Funder, S., Goosse, H., Jepsen, H., Kaas, E., Kjær, K.H., Korsgaard, N.J., Larsen, N.K.,
515 Linderson, H., Lysa, A., Möller, P., Olsen, J., Willerslev, E., 2011a. A 10,000-Year
516 Record of Arctic Ocean Sea-Ice Variability-View from the Beach. *Science* 333, 747-
517 750.

518 Funder, S., Hjort, C., Landvik, J.Y., Nam, S.I., Reeh, N., Stein, R., 1998. History of a stable
519 ice margin East Greenland during the Middle and Upper Pleistocene. *Quaternary*
520 *Science Reviews* 17, 77-123.

521 Funder, S., Kjeldsen, K.K., Kjær, K.H., Ó Cofaigh, C., 2011b. The Greenland ice sheet
522 during the last 300.000 years: a review. *Developments in Quaternary Science* 15, 699-
523 713.

524 Funder, S., Larsen, O., 1982. Implications of volcanic erratics in Quaternary deposits of
525 North Greenland. *Bulletin of the Geological Society of Denmark* 31, 57-61.

526 Gosse, J.C., Phillips, F.M., 2001. Terrestrial in situ cosmogenic nuclides: theory and
527 application. *Quaternary Science Reviews* 20, 1475-1560.

528 Hall, B.L., Baroni, C., Denton, G.H., 2008. The most extensive Holocene advance in the
529 Stauning Alper, East Greenland, occurred in the Little Ice Age *Polar Res* 27, 128-134.

530 Hall, B.L., Baroni, C., Denton, G.H., 2010. Relative sea-level changes, Schuchert Dal, East
531 Greenland, with implications for ice extent in late-glacial and Holocene times.
532 *Quaternary Science Reviews* 29, 3370-3378.

533 Heyman, J., Stroeve, A.P., Harbor, J.M., Caffee, M.W., 2011. Too young or too old:
534 Evaluating cosmogenic exposure dating based on an analysis of compiled boulder
535 exposure ages. *Earth and Planetary Science Letters* 302, 71-80.

536 Hjort, C., 1981. A Glacial Chronology for Northern East Greenland. *Boreas* 10, 259-274.

537 Hjort, C., Björck, S., 1984. A re-evaluation of the glacial chronology in northern East
538 Greenland. *Geologiska Foreningens i Stockholm Förhandlingar* 105, 235-243.

539 Houmark-Nielsen, M., Linge, H., Fabel, D., Schnabel, C., Xu, S., Wilcken, K.M., Binnie, S.,
 540 2012. Cosmogenic surface exposure dating the last deglaciation in Denmark:
 541 Discrepancies with independent age constraints suggest delayed periglacial landform
 542 stabilisation. *Quat Geochronol* 13, 1-17.

543 Jakobsson, M., Andreassen, K., Bjarnadottir, L.R., Dove, D., Dowdeswell, J.A., England,
 544 J.H., Funder, S., Hogan, K., Ingólfsson, O., Jennings, A., Larsen, N.K., Kirchner, N.,
 545 Landvik, J.Y., Mayer, L., Mikkelsen, N., Møller, P., Niessen, F., Nilsson, J., O'Regan,
 546 M., Polyak, L., Norgaard-Pedersen, N., Stein, R., 2014. Arctic Ocean glacial history.
 547 *Quaternary Science Reviews* 92, 40-67.

548 Jennings, A.E., Hald, M., Smith, M., Andrews, J.T., 2006. Freshwater forcing from the
 549 Greenland Ice Sheet during the Younger Dryas: evidence from southeastern Greenland
 550 shelf cores. *Quaternary Science Reviews* 25, 282-298.

551 Jennings, A.E., Walton, M.E., Cofaigh, C.O., Kilfeather, A., Andrews, J.T., Ortiz, J.D., De
 552 Vernal, A., Dowdeswell, J.A., 2014. Paleoenvironments during Younger Dryas-Early
 553 Holocene retreat of the Greenland Ice Sheet from outer Disko Trough, central west
 554 Greenland. *Journal of Quaternary Science* 29, 27-40.

555 Kelley, S.E., Briner, J.P., Young, N.E., 2013. Rapid ice retreat in Disko Bugt supported by
 556 Be-10 dating of the last recession of the western Greenland Ice Sheet. *Quaternary
 557 Science Reviews* 82, 13-22.

558 Kelly, M., 1985. A review of the Quaternary geology of western Greenland, In: Andrews,
 559 J.T. (Ed.), *Quaternary Environments in Eastern Canadian Arctic, Baffin Bay and
 560 Western Greenland*. Allen and Unwin, Boston, pp. 461–501.

561 Kelly, M.A., Lowell, T.V., Hall, B.L., Schaefer, J.M., Finkel, R.C., Goehring, B.M., Alley,
 562 R.B., Denton, G.H., 2008. A Be-10 chronology of lateglacial and Holocene mountain
 563 glaciation in the Scoresby Sund region, east Greenland: implications for seasonality
 564 during lateglacial time. *Quaternary Science Reviews* 27, 2273-2282.

565 Knudsen, K.L., Stabell, B., Seidenkrantz, M.S., Eiriksson, J., Blake, W., 2008. Deglacial and
 566 Holocene conditions in northernmost Baffin Bay: sediments, foraminifera, diatoms and
 567 stable isotopes. *Boreas* 37, 346-376.

568 Knutz, P.C., Sicre, M.A., Ebbesen, H., Christiansen, S., Kuijpers, A., 2011. Multiple-stage
 569 deglacial retreat of the southern Greenland Ice Sheet linked with Irminger Current
 570 warm water transport. *Paleoceanography* 26, Doi 10.1029/2010pa002053.

571 Koch, L., 1923. Preliminary report upon the geology of Peary Land, Arctic Greenland *Am J
 572 Sci* 5, 190-199.

573 Kuijpers, A., Troelstra, S.R., Prins, M.A., Linthout, K., Akhmetzhanov, A., Bouryak, S.,
 574 Bachmann, M.F., Lassen, S., Rasmussen, S., Jensen, J.B., 2003. Late Quaternary
 575 sedimentary processes and ocean circulation changes at the Southeast Greenland
 576 margin. *Marine Geology* 195, 109-129.

577 Landvik, J.Y., 1994. The Last Glaciation of Germania-Land and Adjacent Areas, Northeast
 578 Greenland. *Journal of Quaternary Science* 9, 81-92.

579 Larsen, N.K., Funder, S., Kjær, K.H., Kjeldsen, K.K., Knudsen, M.F., Linge, H., 2014. Rapid
 580 early Holocene ice retreat in West Greenland. *Quaternary Science Reviews* 92, 310-
 581 323.

582 Larsen, N.K., Kjær, K.H., Funder, S., Möller, P., van der Meer, J.J.M., Schomacker, A.,
 583 Linge, H., Darby, D.A., 2010. Late Quaternary glaciation history of northernmost
 584 Greenland - Evidence of shelf-based ice. *Quaternary Science Reviews* 29, 3399-3414.

585 Larsen, N.K., Kjær, K.H., Lecavalier, B., Bjørk, A.A., Colding, S., Huybrechts, P., Jakobsen,
 586 K.E., Kjeldsen, K.K., Knudsen, K.L., Odgaard, B., Olsen, J., 2015. The response of the
 587 southern Greenland ice sheet to the Holocene thermal maximum. *Geology* 43, 291-294.

588 Lévesque, E., Svoboda, J., 1999. Vegetation re-establishment in polar "lichen-kill"
 589 landscapes: a case study of the Little Ice Age impact. *Polar Res* 18, 221-228.

590 Lohne, O.S., Mangerud, J., Birks, H.H., 2013. Precise C-14 ages of the Vedde and
 591 Saksunarvatn ashes and the Younger Dryas boundaries from western Norway and their
 592 comparison with the Greenland Ice Core (GICC05) chronology. *Journal of Quaternary*
 593 *Science* 28, 490-500.

594 Long, A.J., Roberts, D.H., Simpson, M.J.R., Dawson, S., Milne, G.A., Huybrechts, P., 2008.
 595 Late Weichselian relative sea-level changes and ice sheet history in southeast
 596 Greenland. *Earth and Planetary Science Letters* 272, 8-18.

597 Marcott, S.A., Clark, P.U., Padman, L., Klinkhammer, G.P., Springer, S.R., Liu, Z.Y., Otto-
 598 Bliesner, B.L., Carlson, A.E., Ungerer, A., Padman, J., He, F., Cheng, J., Schmittner,
 599 A., 2011. Ice-shelf collapse from subsurface warming as a trigger for Heinrich events.
 600 *Proceedings of the National Academy of Sciences of the United States of America* 108,
 601 13415-13419.

602 Marienfeld, P., 1992. Recent Sedimentary Processes in Scoresby Sund, East Greenland.
 603 *Boreas* 21, 169-186.

604 Matthews, J.A., 2005. 'Little Ice Age' glacier variations in Jotunheimen, southern Norway: a
 605 study in regionally controlled lichenometric dating of recessional moraines with
 606 implications for climate and lichen growth rates. *Holocene* 15, 1-19.

607 Möller, P., Larsen, N.K., Kjær, K.H., Funder, S., Schomacker, A., Linge, H., Fabel, D., 2010.
 608 Early to middle Holocene valley glaciations on northernmost Greenland. *Quaternary*
 609 *Science Reviews* 29, 3379-3398.

610 Muscheler, R., Adolphi, F., Knudsen, M.F., 2014. Assessing the differences between the
 611 IntCal and Greenland ice-core time scales for the last 14,000 years via the common
 612 cosmogenic radionuclide variations. *Quaternary Science Reviews* 106, 81-87.

613 O'Cofaigh, C., Dowdeswell, J.A., Jennings, A.E., Hogan, K.A., Kilfeather, A., Hiemstra, J.F.,
 614 Noormets, R., Evans, J., McCarthy, D.J., Andrews, J.T., Lloyd, J.M., Moros, M., 2012.
 615 An extensive and dynamic ice sheet on the West Greenland shelf during the last glacial
 616 cycle. *Geology* 41, 219-222.

617 Rasmussen, S.O., Andersen, K.K., Svensson, A.M., Steffensen, J.P., Vinther, B.M., Clausen,
 618 H.B., Siggaard-Andersen, M.L., Johnsen, S.J., Larsen, L.B., Dahl-Jensen, D., Bigler,
 619 M., Rothlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M.E., Ruth, U., 2006. A
 620 new Greenland ice core chronology for the last glacial termination. *Journal of*
 621 *Geophysical Research-Atmospheres* 111.

622 Rinterknecht, V., Jomelli, V., Brunstein, D., Favier, V., Masson-Delmotte, V., Bourles, D.,
 623 Leanni, L., Schlappy, R., 2014. Unstable ice stream in Greenland during the Younger
 624 Dryas cold event. *Geology* 42, 759-762.

625 Roberts, D.H., Long, A.J., Schnabel, C., Davies, B.J., Xu, S., Simpson, M.J.R., Huybrechts,
 626 P., 2009. Ice sheet extent and early deglacial history of the southwestern sector of the
 627 Greenland Ice Sheet. *Quaternary Science Reviews* 28, 2760-2773.

628 Roberts, D.H., Long, A.J., Schnabel, C., Freeman, S., Simpson, M.J.R., 2008. The deglacial
 629 history of southeast sector of the Greenland Ice Sheet during the Last Glacial
 630 Maximum. *Quaternary Science Reviews* 27, 1505-1516.

631 Roberts, D.H., Rea, B.R., Lane, T.P., Schnabel, C., Rodes, A., 2013. New constraints on
 632 Greenland ice sheet dynamics during the last glacial cycle: Evidence from the
 633 Uummannaq ice stream system. *J Geophys Res-Earth* 118, 519-541.

634 Sparrenbom, C.J., Bennike, O., Björck, S., Lambeck, K., 2006. Relative sea-level changes
 635 since 15 000 cal. yr BP in the Nanortalik area, southern Greenland. *Journal Of*
 636 *Quaternary Science* 21, 29-48.

637 Steffensen, J.P., Andersen, K.K., Bigler, M., Clausen, H.B., Dahl-Jensen, D., Fischer, H.,
 638 Goto-Azuma, K., Hansson, M., Johnsen, S.J., Jouzel, J., Masson-Delmotte, V., Popp,
 639 T., Rasmussen, S.O., Rothlisberger, R., Ruth, U., Stauffer, B., Siggaard-Andersen,
 640 M.L., Sveinbjornsdottir, A.E., Svensson, A., White, J.W.C., 2008. High-resolution

Greenland Ice Core data show abrupt climate change happens in few years. *Science* 321, 680-684.

Stein, R., Grobe, H., Hubberten, H., Marienfeld, P., Nam, S., 1993. Latest Pleistocene to Holocene Changes in Glaciomarine Sedimentation in Scoresby Sund and Along the Adjacent East Greenland Continental-Margin - Preliminary-Results. *Geo-Mar Lett* 13, 9-16.

Xu, S., Dougans, A.B., Freeman, S.P.H.T., Schnabel, C., Wilcken, K.M., 2010. Improved Be-10 and Al-26-AMS with a 5 MV spectrometer. *Nucl Instrum Meth B* 268, 736-738.

Young, N., Briner, J., Axford, Y., Csatho, B., Babonis, G.S., Rood, D., Finkel, R., 2011. Response of a marine-terminating Greenland outlet glacier to abrupt cooling 8200 and 9300 years ago. *Geophysical Research Letters* 38, L24701, doi:24710.21029/22011GL049639.

Young, N.E., Briner, J.P., Rood, D.H., Finkel, R.C., Corbett, L.B., Bierman, P.R., 2013a. Age of the Fjord Stade moraines in the Disko Bugt region, western Greenland, and the 9.3 and 8.2 ka cooling events. *Quaternary Science Reviews* 60, 76-90.

Young, N.E., Schaefer, J.M., Briner, J.P., Goehring, B.M., 2013b. A Be-10 production-rate calibration for the Arctic. *Journal of Quaternary Science* 28, 515-526.

Figure captions

Figure 1: Overview of North Greenland. Dashed line defines the boundary between the Greenland ice sheet and the independent North Cap (Koch, 1923).

Figure 2: North coast of Greenland with location names mentioned in the text. The inset geomorphological maps show the ^{10}Be sample sites on the moraines at A) Henson Bugt, B) Constable Bugt and C) Bliss Bugt (modified from Larsen et al., 2010).

Figure 3: (A) OSL ages of glaciolacustrine sediments (orange) deposited in late glacial epishelf lakes between the shelf-based ice (hatched) and the mountains (Larsen et al., 2010; Möller et al., 2010). Note incomplete bleaching causes the large spread of ages and the youngest age is likely the most reliable. (B) ^{10}Be surface exposure ages of boulders on terminal moraines in Henson Bugt, Constable Bugt and Bliss Bugt (red). Blue stippled line marks the highest marine level on the north coast, c. 40 m a.s.l. (Möller et al., 2010).

Figure 4: (A) View from the west of the terminal moraine in Henson Bugt. Dashed yellow line indicates the location of the sampled boulders. (B) Setting of boulder and (C) surface of boulder from which sample PEA0719 was collected. (D) Setting of boulder and (E) surface of boulder from which sample PEA0720 was collected. (F) Setting of boulder and (G) surface of boulder from which sample PEA0721 was collected.

Figure 5: (A) View from the east of the terminal moraine at Constable Bugt. Dashed yellow line follows the crest of main ridge west and east of Sifs River. (B) Boulder on the proximal slope of the western part of the moraine ridge from which sample PEA0618 was collected. Rock drill to the left of the boulder is for scale. Approximate locations of samples PEA0611 and PEA0613 are indicated. (C) Close-up of the location of the low and flat boulder surface from which sample PEA0613 was collected. (D) Wide, flat part of the terminal moraine with a small quartz boulder from which sample PEA0621 was collected. (E) Sample PEA0645 was collected from a boulder protruding from the till on the eastern part of the moraine ridge, rock drill for scale. (F) Flat cobble (6 cm thick) on exposed bedrock on the eastern part of the moraine ridge, collected and processed as sample PEA0647.

Figure 6: (A) View from the southeast of the western part of the terminal moraine ridge at outer Bliss Bugt. Dashed black line follows the crest of the ridge. (B) View to the south along the ridge. (C) View to the north along the ridge. The dashed line outlines the same part of the ridge as (A). (D) Sample PEA0663 consisted of 20 quartz pebbles collected around the hammer.

Figure 7: Comparison of the moraine ages with Greenland ice core climate records. The age of the moraines are bracketed by the (A) maximum limiting OSL ages and (B) minimum limiting ^{10}Be ages. Note that the three oldest OSL have not been plotted. (C) $\delta^{15}\text{N}$ based temperature reconstruction (average of NEEM, GISP2 and NGRIP reconstructions) (Buizert et al., 2014) and (D) $\delta^{18}\text{O}$ record from NGRIP showing an abrupt onset and termination of the Younger Dryas (Steffensen et al., 2008).

Figure 8: The terminal moraine in Constable Bugt and the associated reworked organic remains in the up-thrusted glaciolacustrine silt (orange) on the proximal side and marine molluscs in diamict on the distal side of the moraine, used to constrain the age of the moraine. On top of the moraine are the new cosmogenic exposure ages from Constable Bugt.

709 In the present interpretation the moraine was formed during the Younger Dryas and the up-
710 thrust sediment on the distal side are related to a younger re-advance 9.6 to 6.3 cal. ka BP.
711 The marine molluscs in the surficial diamict are interpreted as, by solifluction processes,
712 redeposited marine sediments (see text for further discussion).
713
714 Figure 9: Compilation of observed and inferred Younger Dryas ice marginal positions in
715 Greenland (see main text for data source). Underlined place names refer to the marine coring
716 sites mentioned in the text.

Table 1

Table 1. Summary of ¹⁰Be sample information for rock samples from northernmost Greenland.

Sample ^a	Elevation (m asl)	Sample type	Lithology	Latitude (°N)	Longitude (°W)	Shielding factor ^b	Thickness ^c (cm) (factor)	Quartz ^d (g)	Be carrier ^e (g)	¹⁰ Be/ ⁹ Be ^{f,g} (x10 ⁻¹⁵)	¹⁰ Be/ ⁹ Be ^{blanks, h} (x10 ⁻¹⁵)	¹⁰ Be conc. ⁱ at g ⁻¹ SiO ₂) (10 ⁴)	¹⁰ Be age ^{j, k} (ka)
<i>Constable Bugt - terminal moraine ridge</i>													
PEA 0611	130	pebbles (n=20)	quartz	83,5877	32,2603	0,9963	2.0 (0.9836)	23,303	0.1264 ^{*1}	231.16±6.16	1.71±0.70	13.84±0.47	27.16 ± 0.93 (1.40)
PEA 0613	123	boulder	meta-sandstone	83,5879	32,2360	0,9975	3.0 (0.9756)	17,200	0.2256 ^{*2}	73.90±2.30	6.10±0.84	5.83±0.27	11.56 ± 0.53 (0.69)
PEA 0618 [#]	86	boulder	meta-sandstone	83,5880	32,1679	0,9986	3.8 (0.9696)	16,499	0.8722 ^{*3}	60.51±2.14	4.50±0.79	5.94±0.30	12.35 ± 0.63 (0.79)
PEA 0621	82	boulder	quartz	83,5890	32,1396	0,9993	3.5 (0.9716)	27,288	0.1273 ^{*1}	133.65±4.77	1.71±0.70	6.84±0.29	14.27 ± 0.61 (0.82)
PEA 0645 [#]	112	boulder	meta-sandstone	83,5945	31,8538	0,9996	3.5 (0.9716)	22,843	0.8262 ^{*3}	86.35±4.13	4.50±0.79	5.93±0.34	11.94 ± 0.69 (0.83)
PEA 0647 [#]	112	cobble on bedrock	meta-sandstone	83,5944	31,8555	0,9996	6.0 (0.9519)	10,762	0.8250 ^{*3}	44.45±1.70	4.50±0.79	6.14±0.37	12.60 ± 0.76 (0.90)
<i>arithmetic mean (n=4):</i>													12.11 ± 0.64
<i>Henson Bugt - terminal moraine ridge</i>													
PEA 0719	34	boulder	quartz	83,6142	32,9454	0,9989	5.0 (0.9597)	22,517	0.1272 ^{*1}	90.61±3.69	1.71±0.70	5.58±0.27	12.47 ± 0.61 (0.77)
PEA 0720	34	boulder	quartz	83,6136	32,9428	0,9988	2.0 (0.9836)	23,239	0.1272 ^{*1}	103.40±4.40	1.71±0.70	6.19±0.31	13.50 ± 0.67 (0.85)
PEA 0721	37	boulder	quartz	83,6135	32,9113	0,9986	4.0 (0.9676)	22,227	0.1273 ^{*1}	90.12±3.74	1.71±0.70	5.63±0.28	12.43 ± 0.62 (0.78)
<i>arithmetic mean (n=3):</i>													12.80 ± 0.77
<i>Bliss Bugt - terminal moraine ridge</i>													
PEA 0663 [#]	23	pebbles (n=20)	quartz	83,5617	29,4832	0,9999	1.0 (0.9918)	18,655	0.8309 ^{*4}	70.40±3.03	4.66±0.90	5.85±0.33	12.80 ± 0.72 (0.87)
<i>All moraines</i>													<i>arithmetic mean (n=8):</i> 12.46 ± 0.73

[#] Samples included in Möller et al. (2010) and recalculated in this study.

^a All samples processed and measured at SUERC. ^b Geometric shielding correction was computed after Dunne et al. (1999). ^c Sample thickness measured from the surface, correction factor calculated assuming an exponential reduction in ¹⁰Be production rate with depth (Gosse & Phillips 2000; Balco et al. 2008). ^d All samples use a density value of 2.65 g cm⁻³. ^e Carrier concentrations: *1(1664.175 mg g⁻¹), *2 (980.39 mg g⁻¹), *3 (300.0 mg g⁻¹), *4 (299.0 mg g⁻¹). ^f Isotope ratios normalised to the NIST SRM 4325 ¹⁰Be standard with a nominal value of ¹⁰Be/⁹Be=3.06x10⁻¹¹. ^g Uncertainties are reported at the 1σ confidence level. ^h Procedural blank values used to correct for background. ⁱ Propagated uncertainties include error in the blank and counting statistics. ^j ¹⁰Be surface exposure ages and corresponding analytical uncertainties calculated using the CRONUS-Earth online calculator (Balco et al. 2008) version 2.2, assuming no atmospheric pressure anomalies (std model), no significant denudation during exposure (e=0 mm a⁻¹), no prior exposure, and no temporal shielding (snow, sediment, vegetation). ^k Ages calculated using the Arctic ¹⁰Be production rate of Young et al. (2013). Propagated errors in the calculated ages include uncertainties of the ¹⁰Be production rate and of the ¹⁰Be decay constant. Errors in paranthesis are the systematic uncertainties.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8