

Editorial

Secondary analysis of data on comorbidity/multimorbidity: a call for papers

Marjan van den Akker¹, Jane Gunn², Stewart W. Mercer³, Martin Fortin⁴, Susan M. Smith⁵

¹Department of Family Medicine, School of Public Health and Primary Care (CAPHRI), Maastricht University, Maastricht, The Netherlands; and Department of General Practice, KU Leuven, Leuven, Belgium; ²Primary Care Research Unit, Department of General Practice, University of Melbourne, Melbourne, Victoria, Australia; ³General Practice and Primary Care, Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK; ⁴Department of Family Medicine and Emergency Medicine, Université de Sherbrooke, Sherbrooke, Quebec, Canada; and Centre Intégré Universitaire de santé et de services sociaux du Saguenay Lac St-Jean, Quebec, Canada; ⁵HRB Centre for Primary Care Research, Department of General Practice, RCSI Medical School, Dublin, Ireland

Journal of Comorbidity 2015;5(1):120–121

Keywords: secondary data analysis, open access, comorbidity, multimorbidity, multiple chronic conditions, chronic diseases

Introduction

Despite the high proportion and growing number of people with comorbidity/multimorbidity, clinical trials often exclude this group, leading to a limited evidence base to guide policy and practice for these individuals [1–5]. This evidence gap can potentially be addressed by secondary analysis of studies that were not originally designed to specifically examine comorbidity/multimorbidity, but have collected information from participants on co-occurring conditions. For example, secondary data analysis from randomized controlled trials may shed light on whether there is a differential impact of interventions on people with comorbidity/multimorbidity. Furthermore, data regarding comorbidity/multimorbidity can often be obtained from registration networks or administrative data sets. These types of data sets can address a range of epidemiological research questions, such as:

- What is the prevalence and incidence of comorbidity/multimorbidity?
 - For example, the prevalence and incidence of comorbidity of cancer taken from a general practice-based registration [6]
- What are the patterns of multimorbidity?
- In the case of different data sets, cross-national comparisons of disease patterns can be performed, such as the recent analysis of primary care electronic records in Spain and the Netherlands [7]
- What is the impact of multimorbidity or specific comorbidities on patient outcomes?
 - An example includes the analysis of data from the Veterans Administration in the USA on the role of comorbidities in the prognosis of heart failure [8]. Another example is the impact of comorbid chronic conditions on the quality of diabetes care [9]
 - The Belgium BELFRAIL study, which comprised subjects aged 80 years and older, included medical histories reported by general practitioners in their database and also reported on multimorbidity as a predictor of adverse events [10]
- What is the association between multimorbidity and related phenomena?
 - For example, the Swedish study that examined the association of disability in relation to multimorbidity and disease clustering in older adults [11]

Correspondence: Marjan van den Akker, Department of Family Medicine, School for Public Health and Primary Care (CAPHRI), Maastricht University, Maastricht, The Netherlands.
Tel.: +31 43 388 2321; fax: +31 43 361 9344;
E-mail: marjan.vandenakker@maastrichtuniversity.nl

Received: Sep 6, 2015; Accepted: Sep 6, 2015; Published: Sep 14, 2015

- Another example is an Australian study examining patterns of multimorbidity in relation to functional decline [12].

Many databases include information on (chronic) diseases and are therefore available for further analysis of interesting and relevant issues related to comorbidity/multimorbidity.

Why secondary data?

The purpose of using secondary data is to exploit the already available data and interpret them in new ways – beyond the research hypotheses for which the studies were designed. This enables researchers to explore specific questions that have not been adequately addressed and increases outputs and knowledge translation from funded research.

Preparation and submission of secondary data analyses

The *Journal of Comorbidity* is calling for submissions of original articles based on the analysis of secondary data (i.e. existing data resources). We are interested

in a broad range of research related to comorbidity/multimorbidity, and welcome secondary analyses using data from various sources, such as randomized trials and administrative, cross-sectional, and longitudinal data. Data sources may be small, such as regionally or locally based data sets, or large, nationally representative data sets.

We welcome original research of secondary analyses and invite submissions that specifically address issues relevant to comorbidity or multimorbidity. Authors should consult the journal's website (www.jcomorbidity.com) for guidelines on preparing and submitting their paper. The journal's usual peer review, editorial processes, and standards will be followed.

Conflicts of interest

The authors declare no conflicts of interest.

Funding

Martin Fortin receives funding from the Canadian Institutes of Health Research (CIHR). Stewart W. Mercer receives funding from the Scottish School of Primary Care. Susan M. Smith receives funding from the Health Research Board Centre for Primary Care Research Centre in Ireland.

References

- 1 Fortin M, Soubhi H, Hudon C, Bayliss EA, van den Akker M. Multimorbidity's many challenges. *Br Med J* 2007;334:1016–7. View Item
- 2 Fortin M, Lapointe L, Hudon C, Vanasse A. Multimorbidity is common to family practice. Is it commonly researched? *Can Fam Physician* 2005;51:244–51.
- 3 Caughey GE, Roughhead EE. Multimorbidity research challenges: where do we go from here? *J Comorbidity* 2011;1:8–10. View Item
- 4 Mercer SW, Gunn J, Wyke S. Improving the health of people with multimorbidity: the need for prospective cohort studies. *J Comorbidity* 2011;1:4–7. View Item
- 5 Smith MS, Soubhi H, Fortin M, Hudon C, O'Dowd T. Interventions for improving outcomes in patients with multimorbidity in primary care and community settings. *Cochrane Database Syst Rev* 2012;4:CD006560. View Item
- 6 Deckx L, van den Akker M, Metsemakers J, Knottnerus A, Schellevis F, Buntinx F. Chronic diseases among older cancer survivors. *J Cancer Epidemiol* 2012;2012:206414. View Item
- 7 Poblador-Plou B, van den Akker M, Vos R, Calderón-Larrañaga A, Metsemakers J, Prados-Torres A. Similar multimorbidity patterns in primary care patients from two European regions: results of a factor analysis. *PLoS One* 2014;9:e100375. View Item
- 8 Kheirbek RE, Alemi F, Fletcher R. Heart failure prognosis: comorbidities matter. *J Palliat Med* 2015;18:447–52. View Item
- 9 Magnan EM, Palta M, Mahoney JE, Pandhi N, Bolt DM, Fink J, et al. The relationship of individual comorbid chronic conditions to diabetes care quality. *BMJ Open Diab Res Care* 2015;3:e000080. View Item
- 10 Boeckxstaens P, Vaes B, Van Pottelbergh G, De Sutter A, Legrand D, Adriaensen W, et al. Multimorbidity measures were poor predictors of adverse events in patients aged ≥80 years: a prospective cohort study. *J Clin Epidemiol* 2015;68:220–7. View Item
- 11 Marengoni A, Angleman S, Fratiglioni L. Prevalence of disability according to multimorbidity and disease clustering: a population-based study. *J Comorbidity* 2011;1:11–8. View Item
- 12 Jackson CA, Jones M, Tooth L, Mishra GD, Byles J, Dobson A. Multimorbidity patterns are differentially associated with functional ability and decline in a longitudinal cohort of older women. *Age Ageing* 2015;44:810–6. View Item