

Sebastiano, M. et al. (2016) High levels of mercury and low levels of persistent organic pollutants in a tropical seabird in French Guiana, the Magnificent frigatebird, Fregata magnificens. *Environmental Pollution*, 214, pp. 384-393. (doi:10.1016/j.envpol.2016.03.070)

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

http://eprints.gla.ac.uk/121333/

Deposited on: 29 July 2016

Enlighten – Research publications by members of the University of Glasgow http://eprints.gla.ac.uk

1	High levels of mercury and low levels of persistent organic pollutants in a
2	tropical seabird in French Guiana, the Magnificent frigatebird, Fregata
3	magnificens
4	
5	Manrico Sebastiano <sup>1*</sup> , Paco Bustamante <sup>2</sup> , David Costantini <sup>1</sup> , Igor Eulaers <sup>1,3</sup> , Govindan
6	Malarvannan <sup>4</sup> , Paula Mendez-Fernandez <sup>2</sup> , Carine Churlaud <sup>2</sup> , Pierre Blévin <sup>5</sup> , Antoine
7	Hauselmann <sup>6</sup> , Giacomo Dell'Omo <sup>7</sup> , Adrian Covaci <sup>4</sup> , Marcel Eens <sup>1</sup> , Olivier Chastel <sup>5</sup>
8	
9	1. Behavioural Ecology & Ecophysiology group, Department of Biology, University of
10	Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
11	2. Littoral Environnement et Sociétés, UMR 7266 CNRS-Université La Rochelle, La
12	Rochelle, France
13	3. Department of Bioscience, Aarhus University, Frederiksborgsvej 399, PO Box 358, 4000,
14	Roskilde, Denmark
15	4. Toxicological Centre, Department of Pharmaceutical Sciences, University of Antwerp,
16	Universiteitsplein 1, 2610 Wilrijk, Belgium
17	5. Centre d'Etudes Biologiques de Chizé (CEBC), UMR7372- CNRS/Univ. La Rochelle, F-
18	79360, France
19	6. Association GEPOG, 15 Av Louis Pasteur, 97300 Cayenne, French Guiana
20	7. Ornis italica, Piazza Crati 15, I-00199 Roma, Italy
21	
22	
23	* Corresponding author: <u>Manrico.Sebastiano@uantwerpen.be</u> Tel. 0032/32652285
24	

#### 25 Abstract

In the present study, trace elements and persistent organic pollutants (POPs) were quantified 26 27 from Magnificent frigatebirds (Fregata magnificens) breeding at a southern Atlantic island. Stable isotope ratio of carbon ( $\delta^{13}$ C) and nitrogen ( $\delta^{15}$ N) were also measured to infer the role 28 29 of foraging habitat on the contamination. For another group from the same colony, GPS tracks 30 were recorded to identify potential foraging areas where the birds may get contaminated. 31 Fourteen trace elements were targeted as well as a total of 40 individual POPs, including 32 organochlorine pesticides (OCPs), polychlorinated biphenyls (PCBs) and polybrominated 33 diphenyl ethers (PBDEs). The concentration of Hg in the blood was up to 6 times higher in adults (5.81  $\pm$  1.27 µg g<sup>-1</sup> dw.) than in nestlings (0.99  $\pm$  0.23 µg g<sup>-1</sup> dw.). A similar pattern 34 was found for POPs.  $\Sigma$ PCBs was the prevalent group both in adults (median 673, range 336 – 35 2801 pg  $g^{-1}$  ww.) and nestlings (median 41, range 19 – 232 pg  $g^{-1}$  ww.), followed by the sum 36 37 of dichlorodiphenyltrichloroethanes and metabolities ( $\Sigma$ DDTs), showing a median value of 220 (range 75 – 2342 pg  $g^{-1}$  ww.) in adults and 25 (range 13 – 206 pg  $g^{-1}$  ww.) in nestlings. 38 39 The isotope data suggested that the accumulation of trace elements and POPs between adults 40 and nestlings could be due to parental foraging in two different areas during incubation and 41 chick rearing, respectively, or due to a shift in the feeding strategies along the breeding 42 season. In conclusion, our work showed high Hg concentration in frigatebirds compared to 43 non-contaminated seabird populations, while other trace elements showed lower values within 44 the expected range in other seabird species. Finally, POP exposure was found generally lower 45 than that previously measured in other seabird species.

- 47
- 48
- 49

### 50 **Capsule abstract**

- 51 In the present study we found high levels of mercury and low levels of persistent organic
- 52 pollutants in a tropical seabird breeding in a protected area.

- 54 Keywords: trace elements, persistent organic pollutants, seabirds, contaminants, French
- 55 Guiana.

#### 56 Introduction

57 Since the last few decades, there has been a significant increase of trace element 58 contamination of the environment and, among those trace elements, mercury (Hg) is a highly 59 toxic non-essential metal. Overall, Hg derives from both natural and anthropogenic sources, 60 but human activities have increased the global amount of circulating Hg. Once deposited in 61 aquatic ecosystems, inorganic Hg is subject to biotic reactions (e.g. methylation) resulting in 62 the production of methylmercury (Me-Hg). Me-Hg is the highly toxic form of Hg in 63 organisms that assimilated it via food intake. Once incorporated in organisms, Me-Hg 64 biomagnifies within food webs from lower to higher trophic levels. Hg has neurological and endocrinological effects and impacts reproduction, behaviour, development, and ultimately 65 66 demography in humans and wildlife (Wolfe et al. 1998; Tan et al. 2009), especially in those 67 species which occupy a high trophic level (e.g. seabirds), and are therefore potentially 68 exposed to high contaminant loads (Frederick and Jayasena 2010; Tartu et al 2013; Goutte et 69 al. 2014a). Birds are also vulnerable to other trace metals, particularly to non-essential trace 70 elements such as silver (Ag), cadmium (Cd) and lead (Pb), which although much less studied 71 (Burger 2008), have the potential to negatively affect reproduction, survival and growth 72 (Scheuhammer 1987; Larison et al. 2000). High exposure to essential trace elements has been 73 sometime associated with negative effects in birds (Sánchez-Virosta et al. 2015), but since 74 wild birds are often exposed to a mixture of trace elements, it is generally difficult to 75 demonstrate a causal link between environmental levels of specific compounds and health 76 impairments (Burger 2008, Sánchez-Virosta et al. 2015). Similarly, several persistent organic 77 pollutants (POPs), have been associated with many physiological, immune, endocrine, fitness 78 and demographic consequences and with a decrease in the reproductive success (Bustnes et al. 79 2006; Verreault et al. 2010, Erikstad et al. 2013; Costantini et al. 2014). Although a long term 80 study on the spatial and temporal trends of POPs revealed that these compounds are expected 81 to decline in the Northern Hemisphere (Braune et al. 2005), they appear to still represent a 82 potential threat to adult survival and thus for population dynamics (Goutte et al. 2015). 83 Several studies on seabirds have focused their attention on the contamination in the polar 84 regions (Goutte et al. 2014b; Goutte et al. 2015; Tartu et al. 2015; Bustnes et al. 2015), which 85 are indeed considered a sink for Hg and organic pollutants (Gabrielsen and Henriksen 2001). 86 Most of these contaminants, including Hg from coal burning sources and pesticides used in 87 agriculture are primarily released from the industrialised areas, and their transport to the 88 Arctic region occurs mainly via the atmosphere but also through large rivers and oceanic 89 currents (Gabrielsen and Henriksen 2001). Compared to polar breeding sites, the level of knowledge is much less about contaminant exposure of seabirds in tropical regions. 90 91 Moreover, since individual detection probabilities of seabirds at breeding colonies are 92 generally high because of high overall site fidelity (Gauthier et al. 2012), and since long lived 93 apex predators should be particularly exposed to persistent and biomagnifying contaminants 94 (Rowe 2008), many seabirds species are ideal models to assess the physiological and 95 behavioural effects of environmental pollution.

96 The main goal of this study was to investigate the presence of trace elements and 97 POPs in a long-lived seabird, Magnificent frigatebirds (Fregata magnificens, hereafter 98 frigatebirds) breeding at Grand Connétable Island, a small island of the coasts of French 99 Guiana, which offers a unique situation to study contaminants in a multiple stressor 100 framework. The assessment of POPs and toxic trace elements in tropical regions, which are 101 well known for their complex ecosystem structure and their high biodiversity, is a significant 102 environmental pollution issue. Information on POPs and trace elements is missing in high 103 trophic level species in this region, and an assessment of contaminant exposure has been 104 previously focussed only on Hg accumulation in humans and fish (Fréry et al. 2001; Fujimura 105 et al. 2012). Moreover, since stable carbon and nitrogen isotope measurements have been successfully used to describe the trophodynamics of trace elements and POPs in marine ecosystems (Bearhop et al. 2000; Eulaers et al. 2014), stable isotopes were analysed to study the role of dietary contaminant pathway. Additionally, Global Positioning System (GPS) tracking was conducted on adult frigatebirds of this colony to identify the foraging areas and then the possible sources of the contamination during reproduction.

#### 112 Material and methods

#### 113 **2.1 Sample collection**

114 The field sampling was carried out in 2013 on Grand Connétable island, a protected area located off the Atlantic coast of South America (French Guiana, 4°49'30N; 51°56'00W). This 115 116 island hosts a unique colony of Magnificent frigatebird that is considered one of the most 117 important in South America, and represents the only breeding site for this seabird species in 118 French Guiana (Dujardin and Tostain 1990). Breeding adults (n = 20, 11 females and 9 males 119 during the incubation/early brooding stage) and 30 days old nestlings (n = 20) were captured by hand or with a nose at the end of a fishing rod (Chastel et al. 2005) on May 27<sup>th</sup> - 28<sup>th</sup> and 120 June 25<sup>th</sup>, respectively. Adults and nestlings were not related to each other. Within few 121 122 minutes after capture, 2 mL of blood were collected from the brachial vein using a 123 heparinized syringe and a 25G needle. Samples were immediately put on ice and centrifuged 124 in the field within less than 1 hour to separate plasma (to be used for POPs) and red blood 125 cells (to be used for trace elements and stable isotopes). After centrifugation, both plasma and 126 red blood cells were kept in dry ice until the end of the field work and, when at the laboratory, were kept in a -20 °C freezer until laboratory analysis. 127

128

#### 129 **2.2 Stable isotope analysis**

130 The isotopic niche of frigatebirds was used as a proxy of their ecological niche, with  $\delta^{13}$ C 131 values of seabirds indicating foraging habitats and  $\delta^{15}$ N values indicating trophic level 132 (Newsome et al. 2007). The stable isotopic method is based on time-integrated assimilated 133 food, with different tissues recording trophic information over different time scales. In the 134 present study,  $\delta^{13}$ C and  $\delta^{15}$ N values were measured in red blood cells, which provide trophic 135 information on a few weeks before sampling (Hobson & Clark 1993). Analyses were 136 performed on lyophilised red blood cells of which 0.30 ± 0.05 mg subsamples were weighed 137 in tin cups for stable isotope analyses. Isotopic analyses were performed at the Littoral 138 Environnement et Sociétés (LIENSs) laboratory at the University of La Rochelle (France) 139 with a Thermo Scientific Delta V Advantage mass spectrometer coupled to a Thermo 140 Scientific Flash EA1112 elemental analyser. The results are expressed in the usual  $\delta$  (‰) 141 notation relative to the deviation from international reference standards (Pee Dee Belemnite for  $\delta^{13}$ C and atmospheric nitrogen for  $\delta^{15}$ N). Based on replicate measurements of internal 142 laboratory standards, the experimental precision did not exceed  $\pm 0.15$  and  $\pm 0.20\%$  for  $\delta^{13}$ C 143 and  $\delta^{15}$ N, respectively. 144

145

#### 146 **2.3 Contaminant analysis**

147 2.3.1 Trace elements

148 The analysis of trace element concentrations was carried out by the Littoral Environnement et 149 Sociétés (LIENSs) laboratory at the University of La Rochelle (France). Fourteen trace 150 elements were analysed on lyophilized red blood cells. Total Hg was quantified with an Altec 151 Advanced Mercury Analyzer AMA 254 spectrophotometer. Prior and after freeze-drying, 152 blood samples were weighed to determine the percentage of water in blood, and aliquots 153 ranging from 5 to 10 mg were analysed for quality assessment, as described in Bustamante et 154 al. (2008). Arsenic (As), chromium (Cr), copper (Cu), iron (Fe), manganese (Mn), selenium 155 (Se), and zinc (Zn) were analyzed using a Varian Vista-Pro ICP-OES and silver (Ag), 156 cadmium (Cd), cobalt (Co), nickel (Ni), lead (Pb), and vanadium (V) using a Series II Thermo 157 Fisher Scientific ICP-MS (aliquots mass: 50 – 200 mg dw.) as described in Bustamante et al. 158 (2008). These elements were selected on 2 bases: a first set of non-essential elements (Ag, Cd, 159 Hg and Pb) and a second set of essential trace elements whose metabolism is disrupted by the 160 non-essential ones (Bustamante et al. 2008). Certified Reference Materials (CRM; dogfish 161 liver DOLT-3, NRCC, and lobster hepatopancreas TORT-2, NRCC) were treated and analysed in the same way as the samples. Results were in good agreement with the certified values, and the standard deviations were low, proving good repeatability of the method. The results for CRMs displayed recoveries of the elements ranging from 88% to 116% (n = 10). All the results for trace elements are presented in absolute concentrations in  $\mu g g^{-1}$  dry weight (dw.).

167

168 2.3.2 POPs

169 The analysis of POPs was performed at the Toxicological Centre of the University of 170 Antwerp (Belgium). The analytical protocol was based on the methods described earlier by Eulaers et al. (2011) and consisted in the processing of 1 mL of plasma by solid-phase 171 172 extraction and clean-up on silica acidified with sulfuric acid (44% w/w). The protocol allowed 173 for the analysis for 26 PCB congeners (CB 28, 49, 52, 74, 99, 101, 105, 118, 128, 138, 146, 174 153, 156, 170, 171, 174, 177, 180, 183, 187, 194, 196, 199, 203, 206, and 209), organochlorine pesticides (OCPs), amongst which dichlorodiphenyltrichloroethane (p,p')-175 176 DDT) and its metabolite dichlorodiphenyldichloroethylene (p, p'-DDE), hexachlorobenzene 177 (HCB), cis-nonachlor (CN), trans-nonachlor (TN), oxychlordane (OxC),  $\beta$ - and y-178 hexachlorocyclohexanes (HCHs), and 7 polybrominated diphenyl ethers (PBDEs: BDE 28, 179 47, 99, 100, 153, 154, and 183). Internal standards (CB 143, ε-HCH and BDE 77) were used 180 to quantify the targeted compounds using gas chromatography (Agilent GC 6890, Palo Alto, 181 CA, USA) coupled to mass spectrometry (Agilent MS 5973). Most PCB congeners, as well as 182 *p*,*p*'-DDT and *p*,*p*'-DDE were separated using a HT-8 capillary column (30 m\*0.22 mm\*0.25 183 µm; SGE Analytical Science, Zulte, Belgium), with the mass spectrometer operated in 184 electron impact ionization mode. The remaining PCB congeners, as well as HCB, CHLs 185 (Chlordanes), HCHs, and PBDEs were separated using a DB-5 capillary column (30 m\*0.25 mm\*0.25 µm; J&W Scientific, Folsom, CA, USA) and the mass spectrometer was operated in 186

electron capture negative ionization mode. Mean  $\pm$  SD recoveries of the internal standards CB 143 and BDE 77 were 86  $\pm$  6 % and 93  $\pm$  10 %, respectively. Procedural blanks were analysed every 12<sup>th</sup> plasma sample, and plasma concentrations were corrected for average procedural blank values. The limit of quantification (LOQ) was compound-specifically set at 3\*SD of the procedural blank concentration or, for compounds not detected in blanks, set at a 10:1 signal to noise ratio.

193

#### 194 2.4 Global Positioning System (GPS) transmitters

During the breeding season of 2011, from July 5<sup>th</sup> to 7<sup>th</sup>, 12 brooding adults were equipped with GPS data loggers (Gipsy, Technosmart, Rome, Italy), which recorded GPS locations per second for 32 up to 85 h. GPS data loggers were taped to the back or tail feathers using Tesa© tape, and weighted ~20g, which represented <2% of the bird weight. Since birds needed to be recaptured to recover the GPS, data were recovered from 7 GPS units only.

200

#### 201 **2.5 Statistical analysis**

202 A principal component analysis (PCA) based on correlation matrix with a direct oblimin 203 factor rotation (i.e., oblique) solution was used to reduce the number of variables into a few 204 representative variables explaining variability in metal accumulation. This approach was 205 preferred instead of examining each metal or POP separately, because (i) concentrations are 206 usually correlated with each other and (ii) this enabled us to reduce the number of statistical 207 models because running many models may increase the chance for type II error. Trace 208 elements and POPs with concentrations below the LOQ were replaced with a value equal to 209 <sup>1</sup>/<sub>2</sub>\*LOQ. Ag, Cd, Co, Cr, Ni, and V had a concentration below the LOQ in all individuals and 210 therefore were not included in the PCA on trace elements, while the other trace elements were 211 quantified in all individuals. Among POPs, PBDEs group was not included in the PCA, since 212 concentrations were below the LOQ for each individual. Then, compounds from the same 213 class were grouped (PCBs, DDTs, and CHLs), and the PCA was applied. In addition, the 214 suitability of the use of PCA to reduce data was tested through the Kaiser-Mayer-Olkin measure of sampling adequacy (K-M-O = 0.70 for trace elements and K-M-O = 0.60 for 215 216 POPs) and the Bartlett's test of sphericity (p < 0.01 for both PCAs), showing the appropriate 217 power of the PCA. After examination of the scree plot, the number of significant principal 218 components was selected on the basis of the Kaiser criterion with eigenvalue higher than 1 219 (Kaiser 1960). According to Frontier (1976), eigenvalues are considered interpretable if they 220 exceed eigenvalues generated by the broken-stick model, so the Broken Stick model 221 performed with the PAST software (3.08 version) was utilized to underline which axes 222 significantly explained variance in our data-set. To compare differences among adults and 223 nestlings in the content of POP and trace elements, a parametric test was used when data were 224 normally distributed, and non-parametric test were utilized when data were not normally 225 distributed. The Spearman's rho test was used to test correlations among different POPs. Data 226 on POPs have been reported as median value since they showed a wide range among samples, 227 hence the mean value would have been an overestimation. Finally, the correlation among 228 trace elements and stable isotope values and among POP groups and stable isotope values 229 were also estimated, respectively, using the Spearman's rho correlation. Since in order to 230 decrease Hg toxicity there should be an amount of Se available equal or higher than that of Hg so that the molar ratio of Se:Hg is greater than 1 (Raymond and Ralston 2009), the molar ratio 231 Se:Hg was calculated using the formula "molar concentration (mol  $g^{-1}$  of dw.) = concentration 232 ( $\mu$ g g<sup>-1</sup> dw.) \* 1000 / atomic weight (g mol<sup>-1</sup>)". All statistical analyses were performed using 233 234 SPSS (22.0.0 version).

236 **Results** 

#### 237 **3.1 Trace elements**

238 Of the fourteen trace elements analysed, six had a concentration below the LOQ (Ag, Cd, Co, Cr, Ni and V) both in adults and nestlings while the remaining eight were quantifiable in all 239 240 individuals, including both essential (As, Cu, Fe, Mn, Se, and Zn) and non-essential (Hg and 241 Pb) elements (Table 1). Fe and Zn reported the highest concentrations among essential 242 elements (2413  $\pm$  68 in adults and 2330  $\pm$  80 in nestlings for Fe, and 19.44  $\pm$  0.90 in adults and  $26.93 \pm 2.95$  in nestlings for Zn expressed as  $\mu g g^{-1}$  dw.). Notably, Hg had a quantifiable 243 244 concentration in all individuals and showed the highest concentration among non-essential elements (5.81  $\pm$  1.27 in adults and 0.99  $\pm$  0.23 µg g<sup>-1</sup> dw. in nestlings; Table 1 and S1). 245 Blood concentrations of As, Fe, Pb and Se were significantly higher in adults than in nestlings 246 247 (p < 0.05), while Mn and Zn were significantly higher in nestlings (p < 0.01), and Cu was 248 similar between adults and nestlings (p = 0.06, Table 1). In particular, Hg showed 249 significantly higher concentrations among the two groups, with adults showing a mean 250 concentration of six times higher than the one for nestlings (p < 0.01). The Se:Hg molar ratio 251 in adults (3.9) was almost 4 times lower than the one in nestlings (15). PCA reduced the 252 targeted eight trace elements to three components (explaining 46.22%, 18.15% and 13.58% of 253 the total variance, respectively), while the Broken Stick model suggested to focus on PC1 254 only. As, Hg, Fe, Mn, and Zn were associated with the first axis (Figure 1), and according to 255 the t-test, the age of the individuals (nestlings or adults) was a significant variable explaining 256 the variation of trace elements along PC1 (t = -10.70, p < 0.01; Figure 2). In this scenario, 257 46.22 % of the total variance was explained by the differences in trace element concentrations 258 between adults and nestlings. Adult females and males differed only for Mn (higher in 259 females, p = 0.03) and Se (higher in males, p = 0.03) concentration.

Of the 40 POP compounds targeted, 15 were not detected in both adults and nestlings, and 262 263 some congeners were below the LOQ for one group only (either adults or nestlings, Table 2). On average,  $\Sigma PCBs$ ,  $\Sigma CHLs$ ,  $\Sigma DDTs$  were higher in adults than in nestlings (p < 0.01), while 264 265 HCBs (p = 0.383) and  $\Sigma$ HCHs (p = 0.718) were similar between adults and nestlings. PBDEs were not detected in any sample (Table 2). **SPCBs** was the most important group based in 266 terms of concentration, showing a median (range) of 673 pg  $g^{-1}$  ww. (336 - 2801 pg  $g^{-1}$  ww.) 267 in adults and 41 pg g<sup>-1</sup> ww. (19 - 232 pg g<sup>-1</sup> ww.) in nestlings, followed by  $\sum$ DDTs at 220 pg 268  $g^{-1}$  ww. (75 - 2342 pg  $g^{-1}$  ww.) in adults and 25 pg  $g^{-1}$  ww. (13 - 206 pg  $g^{-1}$  ww.) in nestlings. 269 Among adults, the congeners CB 153, 268 pg  $g^{-1}$  ww. (114 - 869 pg  $g^{-1}$  ww.) and CB 180, 165 270 pg g<sup>-1</sup> ww. (78 - 879 pg g<sup>-1</sup> ww.), contributed most to the  $\Sigma PCBs$  (34% and 24%, 271 respectively), while among nestlings, CB 153, 17 pg g<sup>-1</sup> ww. (< 1.0 - 84 pg g<sup>-1</sup> ww.) and CB 272 180, 8 pg g<sup>-1</sup> ww. (3.0 - 40 pg g<sup>-1</sup> ww.) contributed most to  $\Sigma$ PCBs (22 % and 11 %, 273 274 respectively) (Table 2). Finally, there was a prevalence of heptaCBs ( $40 \pm 9$  % in adults and 275  $25 \pm 4$  % in nestlings) and hexaCBs ( $46 \pm 14$  % in adults and  $34 \pm 9$  % in nestlings; Figure 276 S1). DDE was the only congener to differ significantly between adult males and females, with 277 higher values in males (p < 0.01).

278 Spearman's rho correlation coefficients among POP groups were positive between 279 PCBs and CHLs (r = 0.91, p < 0.01), PCBs and DDTs (r = 0.90, p < 0.01), CHLs and DDTs (r280 = 0.88, p < 0.01), HCB and HCHs (r = 0.44, p < 0.01). Finally, the PCA reduced the targeted 281 POPs to a number of two components (explaining 45.72% and 24.76% of the total variance, 282 respectively), while Broken Stick model suggested PC1 as the only significant axis. The 283 results of the PCA for the POP profiles are presented in Figure 3. The POPs profile 284 significantly differed between adults and nestlings along the PC1 (t = -11.13, p < 0.01; Figure 285 4).

#### 287 **3.3 Stable isotopes and GPS**

Differences between adults and nestlings for stable isotope values were significant for  $\delta^{13}$ C (adults = -15.01 ± 0.11, nestlings = -15.19 ± 0.09; p < 0.01), while they were not different for  $\delta^{15}$ N (adults = 13.37 ± 0.20, nestlings = 13.41 ± 0.28; p = 0.99; Figure 5).

Hg was significantly positively correlated to  $\delta^{15}$ N in both adults (r = 0.84, p < 0.01) and nestlings (r = 0.52, p = 0.02). Moreover, in adults only, there was a significant positive correlation between  $\delta^{15}$ N and As (r = 0.66, p = 0.01), and significant negative correlations between  $\delta^{15}$ N and other trace elements were limited to Pb (r = -0.52, p = 0.02) and Zn (r = -0.66, p < 0.02). Adults also showed a positive correlation between  $\delta^{13}$ C and Hg (r = 0.51, p = 0.02), while there were no significant correlations among stable isotope values and POPs both in adults and in nestlings.

Finally, GPS tracks showed that 6 out of 7 adults alternated long trips toward Brazilian coasts, south of the Grand Connétable colony, with short trips near the island (Figure 6). They showed a wide variance in the trips and, overall, covered an average distance per foraging roundtrip (one way and return) of 219.3 km with a standard deviation of 173.1 km, with the longest trip being 513.9 km and the shortest 5 km.

303

#### 304 **Discussion**

#### **305 4.1 Trace elements**

Our results showed the presence of a high blood level of Hg in Magnificent frigatebirds breeding in French Guiana. In 2009, the National Forestry Office estimated that in French Guiana 1,333 km of watercourses and 12,000 hectares of tropical forest were directly affected by gold mining (Mansillon et al. 2009), and that the number of illegal mining sites was recently estimated between 500 and 900 (Tudesque et al. 2012). In addition, the changing 311 geomorphology of the Amazon soil is an additional source of Hg (de Oliveira 2001), so that 312 Hg has become a primary pollutant in the Amazonian basin (Roulet et al. 1999) and is a 313 matter of great concern in French Guiana (Fujimura et al. 2012). Even so, an evaluation of its 314 impact on local wildlife is, however, still missing. A previous study has shown how 315 frigatebirds may move up to 1.400km away from the breeding colony outside the breeding 316 season (Weimerskirch et al. 2006), and may therefore be contaminated far from French 317 Guiana. However, the high Hg levels found in both adult and nestling frigatebirds suggest a 318 contamination in the lower trophic levels from the coasts of French Guiana up to the upper 319 Brazilian coasts, which includes the foraging areas of our study population during the 320 breeding season (Figure 6). Since Hg biomagnifies within food webs (Lavoie et al. 2013), 321 adults usually show higher concentrations than nestlings (Carravieri et al. 2014). Consistently, 322 Hg was around six times higher in adults than nestlings (Table 1), and our results showed 323 French Guiana frigatebirds to have values of Hg similar to highly Hg-contaminated species 324 (e.g., Diomedea exulans, Stercorarius skua) (see Table S1). Such blood Hg concentrations 325 have been associated with both a reduction of parental commitment (Tartu et al. 2016) and of 326 the breeding success (Goutte et al. 2014a). In addition, similar Hg concentrations have been 327 shown to interfere with several endocrine mechanisms (Tartu et al. 2013, 2014) and to 328 increase oxidative stress (Costantini et al. 2014), a condition that may decrease reproductive 329 success (Costantini 2014) and facilitate herpes infection (Sebastiano et al. 2016). In this 330 scenario, it is important to take into consideration that Hg in the nestlings' red blood cells 331 reflects Hg exposure since hatching as well as maternal Hg transfer through the eggs (Lewis 332 et al. 1993), while adult Hg concentrations reflect the exposure since the last moult (Dauwe et 333 al. 2003). So, the Hg content found in the blood of adults might be lower than actually is, 334 since birds are able to excrete Hg in feathers (Dauwe et al. 2003).

The PCA showed that the high Hg concentration is coupled with high levels of As, Fe, and Se and low levels of Mn and Zn, while Cu and Pb did not show a related pattern (Figure 1). However, for some trace elements such as Cu, Fe, Mn, and Pb, concentrations were very low as compared to literature values in other seabird species (Summers et al. 2014; Carravieri et al. 2014). Interestingly, among non-essential trace elements, Ag and Cd were below the LOQ for every sample, Pb concentrations were very low, while Hg was the only non-essential trace element with high concentrations.

342 A previous study has underlined that As concentrations varied widely among different 343 tissues, being higher in liver and muscle tissues, and varied with the age of the organism, 344 geographic location, and proximity to anthropogenic activities (Eisler 1988). In birds, 345 inorganic As is considered highly toxic in comparison with organic compounds of this 346 element and may disrupt reproduction, and trigger sub-lethal effects or even induce 347 individual's death (Eisler 1994; Kunito et al. 2008). However, marine animals have only a 348 limited ability to bioaccumulate inorganic arsenic from solution (Neff 1997), so As 349 concentrations in living organisms are generally low (Braune and Noble 2009), and 350 concentrations of As in frigatebirds are much lower than the threshold levels of other seabirds, 351 and therefore should not represent a threat for this population (Eisler 1994).

352 In contrast to non-essential elements, Zn is an essential micronutrient and its 353 deficiency has been associated to an increase in oxidative stress and DNA damage, and a 354 decrease in antioxidant defences (Song et al. 2009). Zn is also one of the main component of 355 metallothioneins, a group of proteins which play an essential role in heavy metal 356 detoxification (Siscar et al. 2013). A comparison among tissues and different species is 357 difficult to interpret, but Zn content showed concentrations similar to other seabird species 358 (Carvalho et al. 2013; Fromant et al. 2016). However, the PCA has underlined a strong 359 lowering of the Zn content in the individuals with higher levels of Hg. As a result, since Zn

has a stimulatory action on the immune response, further studies are warranted in order to clarify if the decrease in Zn content with the increase in Hg might reduce the immune competence of this seabird.

363 In a different way, Se, besides being an essential constituent of selenoproteins utilised 364 as a cofactor for reduction of glutathione peroxidases, (Beckett and Arthur 2005) is also 365 important for the detoxification of Hg exposure. In fact, previous studies have emphasized the 366 "protective effect" of Se on Hg toxicity (Raymond and Ralston 2009). Its protective effect 367 was initially presumed to involve Se sequestration of Hg, thereby preventing its harmful 368 effects. However, as more has become understood about Se physiology, the mechanism of 369 MeHg/Hg toxicity and the mechanism of Se protective effect have also become clear. The 370 high affinity between Hg and Se results in Hg binding to Se (Ralston and Raymond 2010), 371 with the consequent generation of mercuric selenide (HgSe), which is well known to be a 372 non-toxic form in marine mammals and birds (Nigro and Leonzion 1996; Ikemoto et al. 373 2004). In order to be able to decrease Hg toxicity, there should be an amount of Se available 374 higher than that of Hg so that the molar ratio of Se:Hg is greater than 1 (Raymond and 375 Ralston 2009). Since the molar ratio was 3.9 for adults and 15 for nestlings, and since the 376 PCA has shown that individuals with high levels of Hg tend to have higher levels of Se, it is 377 likely that Se is contributing to the detoxification of Hg (Sørmo et al. 2011), preventing from 378 Hg toxic effects more in nestlings than in adults. However, Se in blood, which was higher in 379 males, was lower compared to other seabirds (Fromant et al. 2016), and further investigation 380 is needed to understand if the amount of this essential trace element is adequate to contribute 381 to the organisms' physiological functions.

382

383 **4.2 POPs** 

384 To the best of our knowledge, no studies have previously described plasma POP levels in Magnificent frigatebirds or more generally in French Guiana seabirds. In Mexico, a study of 385 386 Magnificent frigatebirds eggs detected low levels of OCPs and PCBs (Trefry et al. 2013). In 387 French Guiana only one study has investigated whole blood levels of OCPs and PCBs in the 388 eggs of leatherback turtles *Dermochelys coriacea*, and found low levels (Girlet et al. 2010). 389 As mentioned earlier, most seabird POP studies have been conducted on polar and especially 390 Arctic species. In the present study, POP concentrations were generally much lower than what 391 has been found previously in polar seabirds (e.g. Tartu et al. 2015a). The contamination with 392 organochlorine compounds and their metabolites can lead to lethal as well as sub-lethal 393 effects in wildlife (Beyer et al. 1996). In particular, DDTs are highly relevant for apex 394 seabirds, since they are associated with eggshell thinning and thus reduced reproductive 395 success (Beyer et al. 1996). Our results pointed out that the p,p'-DDT content was below the LOQ, while p,p'-DDE showed a median value of 220 pg g<sup>-1</sup> ww. in adults (being higher in 396 males), and 25 pg  $g^{-1}$  ww. in nestlings, respectively, which are much lower than other top 397 398 predator seabirds (Bustnes et al. 2006). POPs have never been measured in frigatebird plasma 399 and a reliable comparison with other tissues cannot be made since different tissues show 400 different toxicodynamics.

401 Comparisons with other species showed PCB 153 and other PCBs, HCB and p,p'-402 DDE to be similar to low contaminated populations of common eider *Somateria mollissima* in 403 the sub-Arctic and high Arctic regions (Bustnes et al. 2012; Fenstad et al. 2014), but much 404 lower than in moderately POPs-contaminated Antarctic seabirds, like the snow petrel 405 (*Pagodroma nivea*; Tartu et al. 2015a). Although  $\Sigma$ PCBs show higher concentrations than the other chemical classes, with a median of 673 pg  $g^{-1}$  ww. in adults and 41 pg  $g^{-1}$  ww. in 406 407 nestlings, these concentrations are much lower than those reported in the blood of 7 polar 408 seabirds among which the extremely contaminated Glaucous gull Larus hyperboreus (Tartu et 409 al. 2015b). In polar seabirds specifically, high PCB contamination is associated with 410 concentration from dozens (mean 47,000 pg  $g^{-1}$  ww., Tartu et al. 2015a) up to hundreds of 411 times higher (mean 448,700 pg  $g^{-1}$  ww., Bustnes et al. 2006) than those we found in 412 frigatebirds from French Guiana.

413

#### 414 **4.3 Stable isotopes and GPS**

415 Differences and similarities in trace elements and POPs between nestlings and adults may be explained by trophic ecology. For example, nestlings may differ from adults in  $\delta^{15}$ N if 416 417 they are fed on a different diet or a different trophic level (Overman and Parrish 2001). This 418 explanation is supported by several studies on seabirds (Hobson 1993; Schmutz and Hobson 419 1998), which found that adults provide to their offspring a food different from that they feed 420 on. One way to increase energy gain per unit time of nestlings would be to increase the size of 421 the fish caught for the nestlings, a strategy that has been recorded in other seabirds (Bugge et 422 al. 2011). However, our results do not support this hypothesis. The similar stable nitrogen 423 isotope values between nestlings and adults suggest that they feed on similar trophic level 424 prey.

425 On the other hand, the stable carbon isotope values in the present study showed adults to have significantly higher  $\delta^{13}$ C values than nestlings (Figure 5). A latitudinal decline in  $\delta^{13}$ C 426 427 values has been documented in marine mammals and seabirds (Kelly 2000), and studies have shown patterns which might suggest a decreasing  $\delta^{13}$ C from the coast to the open sea (Eulaers 428 429 et al. 2014), but information of such stratification in French Guiana is not available. Since at 430 this stage of development, frigatebird nestlings are not able to fly, the stable isotope values in 431 nestlings reflect the prey provided by the adults. Hence, the different carbon stable isotope 432 values between adults and nestlings might be explained in two different ways: (a) adults may 433 get their food in a different feeding area than where they forage for their nestlings (GPS tracks 434 of the breeding season 2011 showed how most adults alternated short trips, mostly to the 435 north, with more long trips in the direction of the Brazilian coasts) (Figure 6); (b) adults may 436 have changed their feeding strategies between the incubation stage and the chick rearing period. In fact, since  $\delta^{13}$ C in seabird red blood cells reflects up to three-four weeks before the 437 blood sampling (Hobson and Clark 1992),  $\delta^{13}$ C in adults might have reflected the foraging 438 habitat during the incubation period. In addition,  $\delta^{13}$ C in nestlings might have reflected the 439 440 foraging habitat during the beginning of the chick rearing, since nestlings were around 30 441 days of age. However, differences in the carbon composition are significant from the 442 statistical point of view, but studies are needed to clarify if such difference can be 443 ecologically significant, and if it can be related to the differences in the trace element 444 concentrations.

445

#### 446 **Conclusions**

447 Although our study provided the first evidence of the presence of POPs in French Guiana 448 frigatebirds, PCB and DDT concentrations were generally lower compared to those found in 449 other seabird species, especially in polar seabirds, and they are not likely to be a threat for this 450 population. However, even if concentrations of these pollutants are low, they may have a 451 combined effect with trace elements and especially Hg. Our study clearly shows that this 452 frigatebird population is bearing high Hg burden, and there is an urgent need to evaluate 453 whether increased blood Hg concentrations may affect endocrine and fitness aspects in this 454 top predator bird, as has been documented in other seabird species. Other essential and non-455 essential trace elements showed different accumulation in adults and nestlings, but values 456 were in the range of previous studies on other seabirds. Since the trophic position did not 457 differ between adults and nestlings (same nitrogen isotope value), an explanation for the 458 different POPs and metal profiles between adults and nestlings might lie with the foraging 459 area of adults (carbon isotope values), which appeared to change over the breeding season. 460 Furthermore, in our study population, a previous study has reported the occurrence of herpes 461 virus outbreaks in this colony (De Thoisy et al. 2009), which is causing high mortality of 462 nestlings. These herpes virus outbreaks make this population a highly relevant biological 463 model for investigating the interactions between pollutant exposure and impact of virus 464 activity of population viability. Indeed, previous studies have underlined that there might be a 465 strong relation among exposure to trace elements and virus infections (Koller 1975; Gainer 466 1977), and, more specifically, Hg is highly suspected to aggravate herpes simplex virus-2 467 infection in mice (Christensen et al. 1996). Our data indicate that future studies may be 468 warranted to better understand if the herpes virus outbreaks in this population are favoured by 469 the high Hg contamination.

470

#### 471 Acknowledgments

472 We are grateful to DEAL Guyane, the University of la Rochelle, the University of Antwerp 473 and the FWO (Fonds Wetenschappelijk Onderzoek) and Ornis italica, who funded field 474 operations, GPS tracking, and contaminant and stable isotope analyses. Moreover, we are 475 grateful to the GEPOG (Groupe d'Etude et de Protection des Oiseaux en Guyane) and to the 476 ONCFS (Office national de la Chasse et de la Faune Sauvage), who manage the Grand 477 Connétable Nature reserve since 2008, for their logistic support and access to the Grand 478 Connétable Nature Reserve. We are especially grateful to Grand Connétable Nature staff 479 (Antoine Hauselmann, Alain Alcide and Louise Bétremieux) for their great help in the field 480 and we are also grateful to M. Brault-Favrou from the platform 'Analyses Elémentaires' 481 (University La Rochelle - LIENSs) for her assistance with laboratory work and to G. Guillou 482 and B. Lebreton from the platform 'Analyses Isotopiques' (University La Rochelle ----483 LIENSs) for their assistance in stable isotope analysis. The Contrat de Projet Etat Région

484 (CPER 13) is also acknowledged for funding the AMA and the IR-MS. We also thank three
485 anonymous reviewers that provided valuable comments that helped us to improve the
486 presentation of the work.

#### 487 **References**

- Bearhop S., Waldron S., Thompson D. and Furness R. 2000 Bioamplification of mercury in
  Great Skua *Catharacta skus* nestlings: the influence of trophic status as determined by
  stable isotopes signatures of blood and feathers. Marine Pollution Bulletin 40, 181185.
- Beckett G.J. and Arthur J.R. 2005 Selenium and endocrine systems. J. Endocrinol. 183, 455493 465.
- Beyer W.N., Heinz G.H. and Redmon-Norwood A.W. 1996 Environmental Contaminants in
  Wildlife: Interpreting Tissue Concentrations. CRC press.
- 496 Braune B.M., Outridge P.M., Fisk A.T., Muir D.C.G., Helm P.A., Hobbs K., Hoekstra P.F.,
- 497 Kuzyk Z.A., Kwan M., Letcher R.J., Lockhart W.L., Norstrom R.J., Stern G.A. and
- 498 Stirling I. 2005 Persistent organic pollutants and mercury in marine biota of the
  499 Canadian Arctic: an overview of spatial and temporal trends. Sci. Total Environ. 351,
  500 4e56.
- 501 Braune B.M. and Noble D.G. 2009 Environmental contaminants in Canadian shorebirds.
  502 Environ. Monit. Assess. 148, 185-204.
- Bugge J., Barrett R.T. and Pedersen T. 2011 Optimal foraging in chick-raising Common
  Guillemots Uria aalge. Journal of Ornithology 152, 253–259.
- Burger J. 2008 Assessment and management of risk to wildlife from cadmium. Science of
  The Total Environment 389, 37-45.
- Bustamante P., Gonzàlez A.F., Rocha F., Miramand P. and Guerra A. 2008 Metal and
  mettaloid concentrations in the giant squid *Architeuthis dux* from Iberian waters. Mar.
  Environ. Res. 66, 278-287.
- Bustnes J.O., Tveraa T., Henden J.A., Varpe Ø., Janssen K. and Skaare J.U. 2006
  Organochlorines in Antarctic and Arctic avian top predators: a comparison between

- the south polar skua and two species of Northern Hemisphere gulls. Environ. Sci.
  Technol. 40, 2826–2831.
- Bustnes J.O., Moe B., Hanssen S.A., Herzke D., Fenstad A.A., Nordstad T., Borga K. and
  Gabrielsen G.W. 2012 Temporal dynamics of circulating persistent organic pollutants
  in a fasting seabird under different environmental conditions. Environ. Sci. Technol.
  46, 10287-10294.
- Bustnes J.O., Bourgeon S., Leat E.H.K., Magnusdóttir E., Strøm H., Hanssen S.A., Petersen
  A., Olafsdóttir K., Borgå K., Gabrielsen G.W. and Furness R.W. 2015 Multiple
  Stressors in a top predator seabird: potential ecological consequences of environmental
  contaminants, population health and breeding conditions. PLoS One 10(7): e0131769.
- Carravieri A., Bustamante P., Tartu S., Meillère A., Labadie P., Budzinski H., Peluhet L.,
  Barbraud C., Weimerskirch H., Chastel O. and Cherel Y. 2014 Wandering albratrosses
  document latitudinal variations in the transfer of persistent organic pollutants and
  mercury to southern ocean predators. Environmental Science and Technology 48,
  14746-14755.
- 527 Carvalho P.C., Bugoni L., McGill R.A. and Bianchini A. 2013 Metal and selenium
  528 concentrations in blood and feathers of petrels of the genus Procellaria. Environ.
  529 Toxicol. Chem. 32, 1641-1648.
- 530 Chastel O, Barbraud C, Weimerskirch H, Lormee H, Lacroix A, Tostain O. 2005. High levels
- of LH and testosterone in a tropical seabird with an elaborate courtship display. Generaland Comparative Endocrinology 140, 33-40.
- 533 Cherel Y. and Hobson K.A. 2007 Geographical variation in carbon stable isotope signatures
  534 of marine predators: a tool to investigate their foraging areas in the Southern Ocean.
  535 Mar. Ecol. Prog. Ser. 329, 281-287.

- Christensen M.M., Ellermann-Eriksen S., Rungby J. and Morgensen S.C. 1996 Influence of
  mercury chloride on resistance to generalized infection with herpes simplex virus type
  2 in mice. Toxicology 114, 57-66.
- Costantini D. 2014 Oxidative Stress and Hormesis in Evolutionary Ecology and Physiology.
  Springer-Verlag, Berlin and Heidelberg SBN 978-3-642-54663-1,pp. 13-20.
- 541 Costantini D., Meillere A., Carravieri A., Lecomte V., Sorci G., Faivre B., Weimerskirch H.,
- 542 Bustamante P., Labadie P., Budzinski H. and Chastel O. 2014 Oxidative stress in 543 relation to reproduction, contaminants, gender and age in a long-lived seabird. 544 Oecologia 175, 1107–1116.
- 545 Dauwe T., Bervoets L., Pinxten R., Blust R. and Eens M. 2003 Variation of heavy metals
  546 within and among feathers of birds of prey: effect of molt and external contamination.
  547 Environmental pollution 124, 429-436.
- de Oliveira S.M.B., Melfi A.J., Fostier A.H., Forti M.C., Favaro D.I.T. and Boulet R. 2001
  Soils as an important sink for mercury in the Amazon, Water Air Soil Pollut. 26, 321337.
- de Thoisy B., Lavergne A., Semelin J., Pouliquen J.F., Blanchard F., Hansen E., and Lacoste
  V. 2009 Outbreaks of diseases possibly due to a natural avian herpesvirus infection in
  a colony of young Magnificent Frigatebirds (*Fregata magnificens*) in French Guiana.
  Journal of Wildlife diseases 45, 802-807.
- 555 Dujardin J.L. and Tostain O. 1990 Les oiseaux de mer nicheurs de Guyane Française. Alauda
  556 58, 107-134.
- Eisler R. 1988 Arsenic hazards to fish, wildlife, and invertebrates: a synoptic review.
  Biological Report 85 (1.12).
- Eisler R. 1994 A review of arsenic hazards to plants and animals with emphasis on fisheryand wildlife resources. John Wiley & Sons, Inc. New York.

- 561 Erikstad K.E., Sandvik T., Reinertsen T.K., Bustnes J.O. and Strom H. 2013 Persistent
  562 organic pollution in a high-artic top predator: sex dependent thresholds in adult
  563 survival. Proc. Roy. Soc. London Ser. B 280(1769)20131483.
- Eulaers I., Covaci A., Hofman J., Nygard T., Halley D.J., Pinxten R., Eens M. and Jaspers
  V.L.B. 2011 A comparison of non-destructive sampling strategies to assess the
  exposure of white-tailed eagle nestling (*Haliaeetus albicilla*) to persistent organic
  pollutants. Sci. Total Environ. 410, 258-265.
- Eulaers I., Jaspers V.L.B., Halley D.J., Lepoint G., Nygård T., Pinxten R., Covaci A. and
  Eens M. 2014 Brominated and phosphorus flame retardants in White-tailed Eagle *Haliaeetus albicilla* nestlings: Bioaccumulation and associations with dietary proxies
  (δ13C, δ15N and δ34S). Sci. Total Environ. 478, 48-57.
- Fenstad A.A., Jenssen B.M., Moe B., Hanssen S.A., Bingham C., Herzke D., Bustnes J.O. and
  Krokje A. 2014 DNA double-strand breaks in relation to persistent organic pollutants
  in a fasting seabird. Ecotoxicology and Environmental Safety 106, 68-75.
- 575 Frederick P. and Jayasena N. 2010 Altered pairing behaviour and reproductive success in
  576 white ibises exposed to environmentally relevant concentrations of methylmercury.
  577 Proceedings of the Royal Society B 278, 1851–1857.
- 578 Fréry N., Maury-Brachet R., Maillot E., Deheeger M., de Mérona B. and Boudou A. 2001
  579 Gold-mining activities and mercury contamination of native amerindian communities
  580 in French Guiana: key role of fish in dietary uptake. Environ. Health Perspect. 109,
  581 449-456.
- Fromant A., Carravieri A., Bustamante P., Labadie P., Budzinski H., Peluhet L., Churlaud C.,
  Chastel O. and Cherel Y. 2016 Wide range of metallic and organic contaminants in
  various tissues of the antartic prion, a planktonophagous seabird from the Southern
  Ocean. Sci. Total Environ. 544, 754-764.

- Frontier S. 1976 Étude de la décroissance des valeurs propres dans une analyze en
  composantes principales: comparison avec le modèle de baton brisé. J. Exp. Mar. Biol.
  Ecol. 25, 67-75.
- Fujimura M., Matsuyama A., Harvard J.P., Bourdineaud J.P. and Nakamura K. 2012 Mercury
  contamination in humans in upper Maroni, French Guiana between 2004 and 2009.
  Bull. Environ. Contam. Toxicol. 88, 135-139.
- Gabrielsen G.W. and Henriksen E. 2001 Persistent organic pollutants in Arctic animals in the
  Barents Sea area and at Svalbard: levels and effects. Mem. Natl. Inst. Polar Res. 54,
  349e364.
- Gainer J.H. 1977 Effects of heavy metals and the deficiency of zinc on mortality rates in mice
  infected with encephalomyocarditis virus. Am. J. Vet. Res. 38, 869-672.
- 597 Girlet E., Das K., Thomé J.P., Girondot M. 2010 Maternal transfer of chlorinated
  598 contaminants in the leatherback turtles, *Dermochelys coriacea*, nesting in French
  599 Guiana. Chemosphere 79, 720–726.
- Goutte A., Bustamante P., Barbraud C., Delord K., Weimerskirch H. and Chastel O. 2014a.
   Demographic responses to mercury exposure in two closely-related Antarctic top
   predators. Ecology 95, 1075-1086.
- Goutte A., Barbraud C., Meillère A., Carravieri A., Bustamante P., Labadie P., Budzinski H.,
  Delord K., Cherel Y., Weimerskirch H. and Chastel O. 2014b Demographic
  consequences of heavy metals and persistent organic pollutants in a vulnerable longlived bird, the wandering albatross. Proceedings of the Royal Society B 281(1787),
  20133313.
- Goutte A., Barbraud C., Herzke D., Bustamante P., Angelier F., Tartu S., Clément-Chastel C.,
  Moe B., Bech C., Gabrielsen G.W., Bustnes J.O. and Chastel O. 2015 Survival rate

- and breeding outputs in a high Arctic seabird exposed to legacy persistent organicpollutants and mercury. Environmental Pollution 200, 1-9.
- Hobson K.A. 1993 Trophic relationship among high Artic seabirds: insights from tissuedependent stable-isotope models. Mar. Ecol. Progr. Ser. 95, 7-18.
- Hobson K.A. and Clark R.G. 1992 Assessing avian diets using stable isotopes I: turnover of
  13C in tissues. Condor 94, 181–188.
- Hobson K.A. and Clark R.G. 1993 Turnover of 13C in cellular and plasma fractions of blood:
  Implications for nondestructive sampling in avian dietary studies. Auk 110, 638–641

618 Ikemoto T., Kunito T., Tanaka H., Baba N., Miyazaky N. and Tanabe S. 2004 Detoxification

- mechanism of heavy metals in marine mammals and seabirds: interaction of selenium
  with mercury, silver, copper, zinc, and cadmium in liver. Arch. Environ. Contam.
  Toxicol. 47, 402-413.
- Kaiser H.F. 1960 The Application of Electronic Computers to Factor Analysis. Educationaland Psychological Measurement 20, 141-151.
- Kelly J.F. 2000 Stable isotopes of carbon and nitrogen in the study of avian and mammalian
  trophic ecology. Can. J. Zool. 78, 1-27.
- Koller L.D. 1975 Methylmercury: effect on oncogenic and non-oncogenic viruses in mice.
  Am. J. Vet. Res. 36, 1501-1504.
- Kunito T., Kubota R., Fujihara J., Agusa T. and Tanabe S. 2008 Arsenic in marine mammals,
  seabirds, and sea turtles. Rev. Environ. Contam. Toxicol. 195, 31-69.Martoja R. and
  Berry J.P. 1981 Identification of tiemannite as a probable product of delmethylation of
- 631 mercury by selenium in cetaceans. Vie Milieu 30, 7-10.
- Larison J.R., Likens G.E., Fitzpatrick J.W. and Crock J.G. 2000 Cadmium toxicity among
  wildlife in the Colorado Rocky Mountains. Nature 406, 181-183.

- Lavoie R.A., Jardine T.D., Chumchal M.M., Kidd K.A. and Campbell L.M. 2013
  Biomagnification of mercury in aquatic food webs: a worldwide meta-analysis.
  Environ. Sci. Technol. 47(23): 13385-94.
- 637 Lewis S.A., Becker P.H. and Furness R.W. 1993 Mercury levels in eggs, tissues, and feathers
  638 of herring gulls *Larus argentatus* from the German Wadden Sea Coast. Environ.
  639 Pollut. 80,293-299.
- Mansillon Y., Allain Y.M., de Chalvron J.G., Hirtzman P. 2009 Proposition de schéma
  departmental d'orientation minière de la Guyane, rapport au president de la
  République. Ministère de l'environnement, de l'énergie et de la mer.
- Neff J.M. 1997 Ecotoxicology of arsenic in the marine environment. Environ. Toxicol. Chem
  16, 917-927.
- Newsome S.D., Martinez del Rio C., Bearhop S., and Phillips D.L. 2007 A niche for isotopic
  ecology. Front. Ecol. Environ. 5, 429–436
- Nigro M. and Leonzio C. 1996 Intracellular storage of mercury and selenium in different
   marine vertebrates. Marine Ecology Progress Series 135, 137-143.
- Overman N.C. and Parrish D.L. 2001 Stable isotope composition of walleye: δ<sup>15</sup>N
   accumulation with age and area specific differences in δ<sup>13</sup>C. Can. J. Fish. Aquat. Sci.
   58, 1253-1260.
- Raymond L.J. and Ralston N.V.C. 2009 Selenium's importance in regulatory issues regarding
  mercury. Fuel Process. Technol. 90, 1333–1338.
- Roulet M., Lucotte M., Farella N., Serique G., Coelho H., Sousa-Passos C.J., da Silva J.E., de
  Andrade S.P., Mergler D., Guimaraes J.R.D. and Amorim M. 1999 Effects of recent
- 656 human colonization on the presence of mercury in Amazonian ecosystems. Water Air
- 657 Soil Pollut 112, 297-313.

- Rowe C.L. 2008 The calamity of so long life: life histories, contaminants, and potential
  emerging threats to long -lived vertebrates. Bioscience 58, 623-631.
- Sánchez-Virosta P., Espín S., García-Fernanández A.J. and Eeva T. 2015 A review on
  exposure and effects of arsenic in passerine birds. Sci. Tot. Environ. 512-513, 506525.
- Scheuhammer A.M. 1987 The chronic toxicity of aluminium, cadmium, mercury, and lead in
  birds: a review. Environmental Pollution 46, 263-295.
- Schmutz J.A. and Hobson K.A. 1998 Geographic, temporal and age-specific variation in diets
  of glaucous gulls in Western Alaska. Condor 100, 119-130.
- 667 Sebastiano M., Chastel O., de Thoisy B., Eens M. and Costantini D. 2016 Oxidative stress
  668 favours herpes virus infection in vertebrates: a meta-analysis. Current Zoology (in
  669 press).
- 670 Siscar R., Koenig S., Torreblanca A. and Solè M. 2013 The role of metallothionein and
  671 selenium in metal detoxification in the liver of deep-sea fish from the NW
  672 Mediterranean Sea. Sci. Total Environ. 466-467, 898-905
- Song Y., Leonard S.W., Traber M.G. and Ho E. 2009 Zinc deficiency affects DNA damage,
  oxidative stress, antioxidant defences, and DNA repair in rats. Journal of Nutrition
  139, 1626-1631.
- 676 Sørmo E.G., Ciesielski T.M., Øverjordet I.B., Lierhagen S., Eggen G.S., Berg T. and Jenssen
  677 B.M. 2011 Selenium moderates mercury toxicity in free-ranging freshwater fish.
  678 Environ. Sci. Technol. 45, 6561–6566.
- Summers C.F., Bowerman W.W., Parsons N., Chao W.Y. and Bridges W.C.Jr. 2014 Lead and
  Cadmium in the Blood of nine species of seabirds, Marion Island, South Africa. Bull.
  Environ. Contam. Toxicol. 93, 417-422.

- Tan S.W., Meiller J.C., and Mahaffey K.R.. 2009 The endocrine effects of mercury in humans
  and wildlife. Critica Reviews in Toxicology 39, 228–269.
- Tartu S., Goutte A., Bustamante P., Angelier F., Moe B., Clément-Chastel C., Bech C.,
  Gabrielsen G.W., Bustnes J.O. and Chastel O. 2013 To breed or not to breed:
  endocrine response to mercury contamination by an arctic seabird. Biology Letters
  9:20130317.
- Tartu S., Bustamante P., Goutte A., Cherel Y., Weimerskirch H., Bustnes J.O. and Chastel O.
  2014 Age-related mercury contamination and relationship with luteinizing hormone in
  a long-lived Antartic bird. PLoS One 29, 9(7).
- Tartu S., Angelier F., Wingfield J.C., Bustamante P., Labadie P., Budzinski H., Weimerskirch
  H., Bustnes J.O. and Chastel O. 2015a Corticosterone, prolactin and egg neglect
  behavior in relation to mercury and legacy POPs in a long-lived Antarctic bird. Sci.
  Total Environ. 505, 180-188.
- Tartu S., Angelier F., Bustnes J.O., Moe B., Hanssen S.A., Herzke D., Gabrielsen G.W.,
  Verboven N., Verreault J., Labadie P., Budzinski H., Wingfield J.C., Chastel O. 2015b
  Polychlorinated biphenyl (PCB) exposure and adrenocortical function in seven polar
  seabirds. Environmental Pollution 197, 173-180.
- Tartu S., Bustamante P., Angelier F., Lendvai A.Z., Moe B., Blévin P., Bech C., Gabrielsen
  G.W., Bustnes J.O. and Chastel O. 2016 Mercury exposure, stress and prolactin
  secretion in an Arctic seabird: an experimental study. Functional Ecology
  DOI:10.1111/1365-2435.12534.
- Thompson D.R. and Furness R.W. 1989 The chemical form of mercury stored in SouthAtlantic seabirds. Environ. Pollut. 60, 305-317.
- Trefry S.A., Diamond W.A., Spencer N.C., Mallory M.L. 2013 Contaminants in Magnificent
   frigatebird eggs from Barbuda, West Indies. Marine Pollution Bulletin 75, 317-321.

- Tudesque L., Grenouillet G., Gevrey M., Khazraie K., Brosse S. 2012 Influence of small scale gold mining on French Guiana streams: Are diatom assemblages valid
   disturbance sensors? Ecological Indicators 14, 100-106.
- Verreault J, Bustnes J.O., Gabrielsen G.W. 2010 The Svalbard glaucous gull (Larus hyperboreus) as bioindicator species in the Norwegian Arctic: a review of 35 years of contaminant research. Rev. Environ. Contam. Toxicol. 205, 77–116. (doi:10. 1007/978-1-4419-5623-1\_2).
- Weimerskirch H., Le Corre M., Marsac F., Barbraud C., Tostain O. and Chastel O. 2006
  Postbreeding movements of frigatebirds tracked with satellite telemetry. The Condor
  108, 220-225.
- Wolfe M.F., Schwarzbach S. and Sulaiman R.A. 1998 Effects of mercury on wildlife: a
  comprehensive review. Environmental Toxicology and Chemistry 17, 146–160.
- 719
- 720

7	2	1
1	4	T

722	Figure captions
723	Figure 1. Plot of the PC1 correlation coefficients on trace elements. **Correlation is
724	significant at the 0.01 level (2-tailed). *Correlation is significant at the 0.05 level (2-tailed).
725	
726	Figure 2. Scatter plot of the principal component analysis for trace elements for adult
727	individuals (black squares) and nestlings (red circles).
728	
729	Figure 3. Plot of the PC1 correlation coefficients on POPs. **Correlation is significant at the
730	0.01 level (2-tailed).
731	
732	Figure 4. Scatter plot of the principal component analysis for POPs for adult individuals
733	(black squares) and nestlings (red circles).
734	
735	Figure 5. Stable carbon and nitrogen isotope values (mean $\pm$ SD) of red blood cells of adults
736	and nestlings of the Magnificent frigatebird from French Guiana.
737	
738	Figure 6. GPS tracks of seven Magnificent frigatebirds adults recorded during the breeding
739	season of 2011. The white dotted line represents the political border between French Guiana
740	and Brazil.
741	

# 742 Figures







748 Figure 2







754 Figure 4











- 762 Tables
- **Table 1**. Concentrations ( $\mu g g^{-1} dw$ .) of trace elements in red blood cells of adult and nestling
- 764 Magnificent frigatebirds. df = detection frequency.

		Adults			Nestlings		P
	$mean \pm SD$	median (range)	df (%)	$mean \pm SD$	median (range)	df (%)	
Non-esse	ential trace elemen	ts					
Ag	-	-	0	-	-	0	-
Cd	-	-	0	-	-	0	-
Hg	$5.81 \pm 1.27$	5.62 (3.78 - 7.83)	100	$0.99 \pm 0.23$	0.96(0.68 - 1.68)	100	< 0.01
Pb	$0.02\pm0.01$	0.02(0.02 - 0.04)	100	$0.02\pm0.005$	0.02 (0.01 - 0.03)	100	< 0.01
Essential	l trace elements						
As	$2.35 \pm 1.44$	2.15 (0.58 - 7.33)	100	$1.55\pm0.67$	1.51 (0.67 – 3.61)	100	0.04
Co	-	-	0	-	_	0	-
Cr	-	-	0	-	-	0	-
Cu	$0.78\pm0.07$	0.80(0.65 - 0.90)	100	$0.74 \pm 0.07$	0.73(0.60 - 0.86)	100	0.06
Fe	$2413\pm68$	2411 (2235 - 2503)	100	$2330\pm80$	2337 (2146-2477)	100	< 0.01
Mn	$0.12 \pm 0.03$	0.11(0.09 - 0.19)	100	$0.21 \pm 0.05$	0.19(0.13 - 0.19)	100	< 0.01
Ni	-	-	0	-	-	0	-
Se	$9.09 \pm 1.91$	8.74 (6.67 – 13.09)	100	$5.75 \pm 0.63$	5.82 (4.57 - 6.57)	100	< 0.01
Zn	$19.44 \pm 0.90$	19.36 (18.29 - 22.08)	100	$26.93 \pm 2.95$	26.80 (22.49 - 32.62)	100	< 0.01
V	-	-	0	-	-	0	-

770	Table 2. POP concentrations in adults and nestlings of Magnificent frigatebirds for all
771	congeners analysed. Concentrations are expressed as $pg g^{-1}$ of wet weight. ND = not detected.

	Adults		Nestlin	gs	P
Congener	median (range)	mean $\pm$ SD	median (range)	mean ± SD	
CB 28	ND	ND	ND	ND	ND
CB 52	ND	ND	ND	ND	ND
<b>CB 49</b>	ND	ND	ND	ND	ND
<b>CB 74</b>	ND	ND	ND	ND	ND
CB 101	ND	ND	ND	ND	ND
CB 99	<4 (<4 - 68)	$7\pm16$	ND	ND	ND
CB 105	<2 (<2 - 29)	$45\pm 6$	ND	ND	ND
CB 118	22 (6 - 122)	$29\pm25$	<1 (<1 - 8)	$2.15\pm1.88$	< 0.01
CB 128	<1 (<1 - 4)	$1 \pm 1$	ND	ND	ND
CB 138	56 (26 - 277)	$78\pm60$	6 (2 - 38)	$7\pm7$	< 0.01
<b>CB 146</b>	28 (<1 - 134)	$35 \pm 38$	ND	ND	ND
CB 153	268 (114 - 869)	$333\pm211$	17 (<1 - 84)	$19\pm16$	< 0.01
CB 156	7 (<1 - 21)	$9\pm5$	ND	ND	ND
CB 170	47 (23 - 217)	$68\pm50$	<3 (<1 - 13)	3 ± 3	< 0.01
CB 171	<1 (<1 - 2)	$1 \pm < 1$	ND	ND	ND
CB 174	ND	ND	ND	ND	ND
CB 177	<1 (<1 - 4)	$1 \pm < 1$	ND	ND	ND
CB 180	165 (78 - 879)	$240\pm189$	8 (3 - 40)	$10\pm 8$	< 0.01
CB 183	32 (14 - 122)	$42 \pm 31$	<1 (<1 - 11)	$2\pm 2$	< 0.01
CB 187	34 (15 - 141)	$43 \pm 33$	3 (<2 - 25)	$4\pm5$	< 0.01
CB 194	21 (7 - 154)	$32 \pm 32$	<1 (<1 - 3)	$1 \pm < 1$	< 0.01
CB 196/203	23 (7 - 108)	$30 \pm 25$	<1 (<1 - 6)	$1 \pm 1$	< 0.01
CB 199	9 (<4 - 30)	$11 \pm 8$	<1 (<1 - 3)	$1 \pm < 1$	< 0.01
CB 206	<1 (<1 - 14)	<3 ± 3	ND	ND	ND
CB 209	ND	ND	ND	ND	ND
ΣPCBs	673 (336 - 2801)	$967\pm688$	41 (19 - 232)	$51 \pm 44$	< 0.01
OvC	<1 (<1 7)	<2 + <2	ND	ND	ND
UXC TN	<1(<1-7)	$10 \pm 3$	ND (2) (2) (2)	$\Lambda \pm 7$	<0.01
	(3 - 10)	$10 \pm 3$	<1 (<1 5)	4 ± /	<0.01 0.10
	<1(<1-3)	$\sim 2 \pm 1$	<1(<1-3)	$1 \pm < 1$	0.10 <0.01
2CHLs	14 (7 - 22)	$14 \pm 5$	4 (3 - 37)	0 ± 7	<0.01
нсв	7 (2 - 41)	$12 \pm 11$	11 (<2 - 33)	$11 \pm 6$	0.38

<i>p,p</i> <b>'-DDE</b>	220 (75 - 2342)	$426\pm561$	25 (13 - 206)	$40\pm45$	< 0.01
<i>p,p</i> <b>'-DD</b> T	ND	ND	ND	ND	ND
ΣDDTs	220 (75 - 2342)	$426\pm561$	25 (13 - 206)	$40 \pm 45$	< 0.01
β-НСН	2 (2 - 19)	$8\pm 6$	2 (2 - 11)	$3\pm 2$	0.38
γ-НСН	2 (2 - 82)	$8 \pm 18$	12 (2 - 20)	$11 \pm 7$	0.01
ΣΗCHs	14 (5 - 84)	$16\pm18$	14 (5-23)	$14 \pm 7$	0.72
					ND
<b>BDE 28</b>	ND	ND	ND	ND	ND
<b>BDE 47</b>	ND	ND	ND	ND	ND
<b>BDE 100</b>	ND	ND	ND	ND	ND
<b>BDE 99</b>	ND	ND	ND	ND	ND
BDE 154	ND	ND	ND	ND	ND
BDE 153	ND	ND	ND	ND	ND
BDE 183	ND	ND	ND	ND	ND
ΣPBDEs	ND	ND	ND	ND	ND

## SUPPORTING INFORMATION

# High levels of mercury and low levels of persistent organic pollutants in a tropical seabird in French Guiana, the Magnificent frigatebird, *Fregata magnificens*.

Manrico Sebastiano<sup>1,\*</sup>, Paco Bustamante<sup>2</sup>, David Costantini<sup>1</sup>, Igor Eulaers<sup>1,3</sup>, Govindan Malarvannan<sup>4</sup>, Paula Mendez-Fernandez<sup>2</sup>, Carine Churlaud<sup>2</sup>, Pierre Blévin<sup>5</sup>, Giacomo Dell'Omo<sup>6</sup>, Adrian Covaci<sup>4</sup>, Marcel Eens<sup>1</sup>, Olivier Chastel<sup>5</sup>

- Behavioural Ecology & Ecophysiology group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
- Littoral Environnement et Sociétés, UMR 7266 CNRS-Université La Rochelle, La Rochelle, France
- Department of Bioscience, Aarhus University, Frederiksborgsvej 399, PO Box 358, 4000, Roskilde, Denmark
- Toxicological Centre, Department of Pharmaceutical Sciences, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk, Belgium
- Centre d'Etudes Biologiques de Chizé (CEBC), UMR7372- CNRS/Univ. La Rochelle, F-79360, France
- 6. Ornis italica, Piazza Crati 15, I-00199 Roma, Italy
- \* Corresponding author:

Manrico.Sebastiano@uantwerpen.be Tel. 0032/32652285

**Table S1**. Total Hg concentrations (mean  $\pm$  SD) in the blood of seabirds worldwide. Data are mean  $\pm$  SD of  $\mu$ g g<sup>-1</sup> of dry weight or wet weight. Please note that this table is not exhaustive, but is a comparison of selected studies on seabirds, both in contaminated and non-contaminated areas, in order to have an idea of the amount of total Hg in frigatebirds.

Species	Location	Adults	Nestlings/Juv.	Weight	Reference
Fregata magnificens	French Guiana	$5.81 \pm 1.27$	$0.99\pm0.23$	dw	This study
Pagodroma nivea	Antarctica	$2.70 \pm 1.10$		dw	Tartu et al. 2014
Diomedea exulans	(Indian Ocean)	$10.70\pm0.50$		dw	Goutte et al. 2014b
Pachyptila desolata	Georgia	$0.53\pm0.21$		dw	Anderson et al. 2009
Halobaena caerulea	Georgia	$0.56\pm0.28$		dw	Anderson et al. 2009
Pelacanoides urinatrix	Georgia	$0.31\pm0.15$		dw	Anderson et al. 2009
Pelacanoides georgicus	Georgia	$0.41\pm0.14$		dw	Anderson et al. 2009
Procellaria aequinoctialis	Georgia	$5.37 \pm 1.18$		dw	Anderson et al. 2009
Macronectes giganteus	Georgia	$2.74 \pm 1.05$		dw	Anderson et al. 2009
Thalassarche melanophrys	Georgia	$4.38~\pm~1.1$		dw	Anderson et al. 2009
Thalassarche chrysostoma	Georgia	$6.57 \hspace{0.1in} \pm 1.11$		dw	Anderson et al. 2009
Macronectes halli	Georgia	$3.93 \hspace{0.1in} \pm 1.37$		dw	Anderson et al. 2009
Diomedea exulans	Georgia	$11.15\ \pm 3.38$		dw	Anderson et al. 2009
Diomedea exulans	Georgia	$9.57 \hspace{0.2cm} \pm 4.29 \hspace{0.2cm}$	$0.84\pm0.36$	dw	Tavares et al. 2013
Sterna hirundo	USA	$\textbf{0.36} \pm \textbf{0.40}$	$0.04\pm0.05$	ww	Nisbet 2002
Catharacta maccormicki	Antarctica	$2.15\ \pm 0.17$		dw	Goutte et al. 2014a
Catharacta lonnbergi	Antarctica	$8.22\ \pm 0.24$		dw	Goutte et al. 2014a
Ptychoramphus aleuticus	Canada	$0.63\ \pm 0.07$	$0.23\pm0.07$	dw	Hipfner et al 2011
Cerorhinca monocerata	Canada	$1.75~\pm~0.11$	$0.41\pm0.12$	dw	Hipfner et al 2011
Rissa tridactyla	Norway	1.80		dw	Tartu et al. 2013
Oceanodroma leucorhoa	USA	$0.54 \pm 0.37$	$\textbf{0.03} \pm \textbf{0.04}$	ww	Huntington et al. 1996
Sterna hirundo	USA	$0.44\ \pm 0.26$		ww	Gochfeld 1980
Rissa tridactyla	Norway	$2.33\ \pm 0.44$		dw	Goutte et al. 2015
Diomedea exulans	(Indian Ocean)	$7.70\pm3.60$		dw	Carravieri et al. 2014
Fratercula arctica	(Middle Europe)	$\textbf{7.12} \pm \textbf{2.58}$		dw	Fort et al. 2015
Uria aalge	(Middle Europe)	$6.32 \pm 5.17$		dw	Fort et al. 2015
Rissa tridactyla	(Middle Europe)	$\textbf{8.58} \pm \textbf{2.82}$		dw	Fort et al. 2015
Alca torda	(Middle Europe)	$\textbf{9.38} \pm \textbf{4.12}$		dw	Fort et al. 2015
Somateria mollissima	USA	0.05		ww	Bond and Diamond 2009
Sterna paradisaea	USA	$0.16\ \pm 0.03$		ww	Bond and Diamond 2009
Oceanodroma leucorhoa	USA	$0.12\ \pm 0.02$		ww	Bond and Diamond 2009
Fratercula arctica	USA	$0.17\ \pm 0.03$		ww	Bond and Diamond 2009
Sterna hirundo	USA	$0.24\ \pm 0.04$		ww	Bond and Diamond 2009
Uria aalge	USA	$0.27\ \pm 0.05$		ww	Bond and Diamond 2009
Alca torda	USA	$0.36\ \pm 0.07$		ww	Bond and Diamond 2009
Macronectes halli	Georgia	$\textbf{2.9} \pm \textbf{0.97}$		dw	Gonzales-Solis et al. 2002
Macronectes giganteus	Georgia	$\textbf{4.9} \pm \textbf{5.1}$		dw	Gonzales-Solis et al. 2002
Sterna forsteri	USA		$0.33\pm0.01$	ww	Ackerman et al. 2008
Stercorarius skua	(Northern Europe)	$3.67 \hspace{0.1in} \pm 2.03$	$0.27\pm0.16$	dw	Bearhop et al. 2000
Stercorarius skua	Scotland	$7.37\ \pm 3.1$	$1.12\pm0.41$	dw	Bearhop et al. 2000

Procellaria aequinoctialis	Brazil	$\textbf{3.20} \pm \textbf{3.67}$		dw	Carvalho et al. 2013
Procellaria conspicillata	Brazil	$3.41 \pm 2.14$		dw	Carvalho et al. 2013
Recurvirostra americana	USA	$1.49\ \pm 0.13$	$2.02\pm0.29$	dw	Eagles-Smith et al. 2008
Himantopus mexicanus	USA	$5.05 \ \pm 0.46$	$0.99 \pm 0.09$	dw	Eagles-Smith et al. 2008
Hydroprogne caspia	USA	$6.83 \hspace{0.1cm} \pm \hspace{0.1cm} 0.89 \hspace{0.1cm}$		dw	Eagles-Smith et al. 2008
Sterna forsteri	USA	$7.06 \ \pm 0.62$	$1.71\pm0.18$	dw	Eagles-Smith et al. 2008
Acaena adscendens	(Indian Ocean)	$0.67 \pm 0.11$		dw	Fromant et al. 2016



Figure S1. Contribution of congener classes to the  $\sum$ PCBs in adults (blue line) and nestlings (red line). Data are presented as mean  $\pm$  SD.

#### References

- Ackerman J.T., Eagles-Smith C.A., Takekawa J.Y. and Iverson S.A. 2008 Survival of postfledging Forster's terns in relation to mercury exposure in San Francisco Bay. Ecotoxicology 17, 789-801.
- Anderson O., Phillips R., McDonald R., Shore R., McGill R. and Bearhop S. 2009 Influence of trophic position and foraging range on mercury levels within a seabird community. Marine Ecology Progress Series 375, 277–288 doi:10.3354/meps07784.
- Bearhop S., Waldron S., Thompson D. and Furness R. 2000 Bioamplification of mercury in Great Skua *Catharacta skus* chicks: the influence of thropic status as determined by stable isotopes signatures of blood and feathers. Marine Pollution Bulletin 40, 181-185.
- Bond A.L. and Diamond A.W. 2009 Mercury concentrations in seabird tissues from Machias Seal Island, New Brunswick, Canada. Sci. Total Environ. 407, 4330-4347.
- Carvalho P.C., Bugoni L., McGill R.A. and Bianchini A. 2013 Metal and selenium concentrations in blood and feathers of petrels of the genus Procellaria. Environ. Toxicol. Chem. 32, 1641-1648.
- Eagles-Smith C.A., Ackerman J.T., Adelsbach T.L., Takekawa J.Y., Miles A.K. and Keister R.A. 2008 Mercury correlations among six tissues for four waterbird species breeding in San Francisco Bay, California, USA. Environ. Toxicol. Chem. 27, 2136-2153.
- Fort J., Lacoue-Labarthe T., Nguyen H.L., Boué A., Spitz J. and Bustamante P. 2015 Mercury in wintering seabirds, an aggravating factor to winter wrecks? Sci. Total Environ. 527-528, 448-454.
- Fromant A., Carravieri A., Bustamante P., Labadie P., Budzinski H., Peluhet L., Churlaud C., Chastel O. And Cherel Y. 2016 Wide range of metallic and organic contaminants in

various tissues of the Antarctic prion, a planktonophagous seabird from the Southern Ocean. Sci. Total Environ. 544, 754-764.

- Gochfeld M. 1980 Tissue distribution of mercury in normal and abnormal young Common Terns. Marine Pollution Bulletin 11, 362-366.
- Gonzales-Solis J., Sanpera C. and Ruix X. 2002 Metals and selenium as bioindicators of geographic and trophic segregation in giant petrels Macronectes spp. Mar. Ecol. Prog. Ser. 244, 257-264.
- Goutte A., Bustamante P., Barbraud C., Delord K., Weimerskirch H. and Chastel O. 2014a Demographic responses to mercury exposure in two closely related Antartic top predators. Ecology 95, 1075-1086.
- Goutte A., Barbraud C., Meillère A., Carravieri A., Bustamante P., Labadie P., Budzinsky H.,Delord K., Cherel Y., Weimerskirch H. and Chastel O. 2014b Demographicconsequences of heavy metals and persistent organic pollutants in a vulnerable long-lived bird, the wandering albatross. Proceeding of the Royal Society B 281, 20133313.
- Goutte A., Barbraud C., Herzke D., Bustamante P., Angelier F., Tartu S., Clement-Chastel C., Moe B., Bech C., Gabrielsen G.W., Bustnes J.O. and Chastel O. 2015 Survival rate and breeding outputs in a high Artic seabird exposed to legacy persistent organic pollutants and mercury. Environ. Pollut. 200, 1-9.
- Hipfner J.M., Hobson K.A. and Elliott J.E. 2011 Ecological factors differentially affect mercury levels in two species of sympatric marine birds of the North Pacific. Science of the Total Environment 409, 1328-1335.
- Huntington C.E., Butler R.G. and Mauck R.A. 1996 Leach's stormpetrel. In: The Birds of North America Online, Poole A (editor), Ithaca, NY: Cornell Laboratory of Ornithology; retrieved from The Birds of North America Online database.

- Nisbet I.C.T. 2002 Common tern. In: The Birds of North America Online, Poole A (editor), Ithaca, NY: Cornell Laboratory of Ornithology; retrieved from The Birds of North America Online database.
- Tartu S., Goutte A., Bustamante P., Angelier F., Moe B., Clément-Chastel C., Bech C., Gabrielsen G.W., Bustnes J.O. and Chastel O. 2013 To breed or not to breed: endocrine response to mercury contamination by an Arctic seabird. Biology Letters 9. doi:10.1098/rsbl.2013.0317.
- Tartu S., Bustamante P., Goutte A., Cherel Y., Weimerskirch H., Bustnes J.O. and Chastel O.2014 Age-related mercury contamination and relationship with luteinizing hormone in a long-lived Antartic bird. PLoS One 29, 9(7).
- Tavares S., Xavier J.C., Pjillips R.A., Pereira M.E. and Pardal M.A. 2013 Influence of age, sex and breeding status on mercury accumulation patterns in the wandering albatross *Diomeda exulans*. Environmental Pollution 181, 315-320.