



Gadre, V. and Maher, J. (2018) The stratum of random mapping classes.  
*Ergodic Theory and Dynamical Systems*, 38(7), pp. 2666-2682.  
(doi: [10.1017/etds.2016.132](https://doi.org/10.1017/etds.2016.132))

This is the author's final accepted version.

There may be differences between this version and the published version.  
You are advised to consult the publisher's version if you wish to cite from  
it.

<http://eprints.gla.ac.uk/131384/>

Deposited on: 18 November 2016

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

# THE STRATUM OF RANDOM MAPPING CLASSES.

VAIBHAV GADRE AND JOSEPH MAHER

ABSTRACT. We consider random walks on the mapping class group that have finite first moment with respect to the word metric, whose support generates a non-elementary subgroup and contains a pseudo-Anosov map whose invariant Teichmüller geodesic is in the principal stratum. For such random walks, we show that mapping classes along almost every infinite sample path are eventually pseudo-Anosov, with invariant Teichmüller geodesics in the principal stratum. This provides an answer to a question of Kapovich-Pfaff [KP15].

## 1. INTRODUCTION

Let  $S$  be an orientable surface of finite type, and let  $\text{Mod}(S)$  denote the mapping class group of  $S$ , consisting of orientation preserving diffeomorphisms on  $S$  modulo isotopy. Let  $\mathcal{T}(S)$  be the Teichmüller space of marked conformal structures on  $S$ , and let  $\mathcal{Q}(S)$  be the space of quadratic differentials, which may be identified with the cotangent bundle of  $\mathcal{T}(S)$ . For punctured surfaces, the quadratic differentials in  $\mathcal{Q}(S)$  are assumed to be meromorphic with poles only at punctures and with every puncture a simple pole. The space  $\mathcal{Q}(S)$  is stratified by the order of the zeros of the quadratic differential; the principal stratum consists of those quadratic differentials whose zeros are all simple. By the Thurston classification, mapping classes are periodic, reducible or pseudo-Anosov. A pseudo-Anosov map has a unique invariant Teichmüller geodesic. If the invariant Teichmüller geodesic is given by a quadratic differential that lies in the principal stratum, then we say that the pseudo-Anosov map is in the principal stratum.

We consider random walks on the mapping class group  $\text{Mod}(S)$  that have finite first moment with respect to word metric and whose support generates a non-elementary subgroup of  $\text{Mod}(S)$ , i.e. the subgroup generated by the support of the initial distribution contains a pair of pseudo-Anosov maps with distinct stable and unstable measured foliations. In independent work, Maher [Mah11] and Rivin [Riv08] showed that the probability that a random walk gives a pseudo-Anosov map tends to 1 in the length of the sample path. As a refinement, Kapovich-Pfaff raise the following question: what is the stratum of quadratic differentials for the invariant Teichmüller geodesic of a random pseudo-Anosov element? See [KP15, Question 1.5], [DHM15, Question 6.1].

As a step towards answering the question, we prove the following result. We shall write  $d_{\text{Mod}}$  for the word metric on  $\text{Mod}(S)$  with respect to a choice of finite generating set.

**Theorem 1.1.** *Let  $S$  be a connected orientable surface of finite type, whose Teichmüller space  $\mathcal{T}(S)$  has complex dimension at least two. Let  $\mu$  be a probability distribution on  $\text{Mod}(S)$  such that*

- (1)  $\mu$  has finite first moment with respect to  $d_{\text{Mod}}$ ,
- (2)  $\text{Supp}(\mu)$  generates a non-elementary subgroup  $H$  of  $\text{Mod}(S)$ , and
- (3) The semigroup generated by  $\text{Supp}(\mu)$  contains a pseudo-Anosov  $g$  such that the invariant Teichmüller geodesic  $\gamma_g$  for  $g$  lies in the principal stratum of quadratic differentials.

---

2010 Mathematics Subject Classification. 30F60, 32G15.

Key words and phrases. Teichmüller theory, Moduli of Riemann surfaces.

The first author acknowledges support from the GEAR Network (U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties”).

The second author acknowledges support from the Simons Foundation and PSC-CUNY.

*Then, for almost every bi-infinite sample path  $\omega = (w_n)_{n \in \mathbb{Z}}$ , there is positive integer  $N$  such that for all  $n \geq N$  the mapping class  $w_n$  is a pseudo-Anosov map in the principal stratum, that is its invariant Teichmüller geodesic is given by a quadratic differential with simple zeros and poles. Furthermore, almost every bi-infinite sample path determines a unique Teichmüller geodesic  $\gamma_\omega$  with the same limit points as the bi-infinite sample path, and this geodesic also lies in the principal stratum.*

In the statement of Theorem 1.1, hypothesis (3) for the distribution  $\mu$  is necessary. There are random walks for which  $\mu$  satisfies hypothesis (1) and (2) but the conclusion of Theorem 1.1 is false. One such example is a random walk supported on a non-principal Teichmüller curve, i.e. on a Veech lattice whose Teichmüller disc is not contained in the principal stratum. See [Wri15] for the precise definition of a Teichmüller curve.

We only consider surfaces whose Teichmüller space has complex dimension at least two. This excludes the sphere with four or fewer punctures, and the torus with one or fewer punctures. In case of the four-punctured sphere, and the torus with at most one puncture,  $\mathcal{T}(S)$  has complex dimension one, and the principal stratum consists of all of  $\mathcal{T}(S)$ . So the result holds trivially; the remaining surfaces have finite  $\text{Mod}(S)$  and the result does not apply.

We have described the principal stratum in terms of the zeros of quadratic differentials, but it may also be described in terms of flat structures, foliations, or laminations, as we now discuss. A quadratic differential equips the surface with a flat structure with a pair of measured foliations (or measured laminations), corresponding to the vertical and horizontal directions. The condition of having simple poles at punctures is equivalent to the flat metric having cone angle  $\pi$  around every puncture. The foliations thus have one-pronged singularities at the punctures, and the laminations have a complementary region which is a punctured monogon for each singularity. In the principal stratum all zeros are simple. This corresponds to the flat metric having all other cone points of angle  $3\pi$ ; the corresponding measured foliations will have all other singularities trivalent, and the measured laminations will have all of the other complementary regions consisting of triangles.

If we consider  $\text{PMF}(S)$  as Thurston's boundary for  $\mathcal{T}(S)$ , then there is a natural Lebesgue measure class on  $\text{PMF}(S)$ , defined in terms of train track coordinate charts. Also, Hubbard-Masur [HM79] showed that at each point  $X$  in  $\mathcal{T}(S)$ , the unit cotangent space may also be identified with  $\text{PMF}(S)$ , and then the Masur-Veech measure [Mas82, Vee82] on the space of unit area quadratic differentials gives a conditional measure  $s_X$  on  $\text{PMF}(S)$ , which Athreya-Bufetov-Eskin-Mirzakhani [ABEM12] showed is absolutely continuous with respect to the Lebesgue measure class. As we now explain, the final statement of Theorem 1.1 is known for bi-infinite geodesics chosen according to this Lebesgue measure class. Kerckhoff-Masur-Smillie [KMS86] showed that the uniquely ergodic foliations have full measure with respect to the Lebesgue measure class. So any two foliations chosen according to the Lebesgue measures almost surely determine a unique bi-infinite Teichmüller geodesic. Finally, the principal stratum foliations have full measure in  $\text{PMF}(S)$  with respect to these measures. This is most easily seen using train track coordinates: the collection of non-principal strata foliations is contained in a countable union of positive co-dimension sets, and so has Lebesgue measure zero.

Kaimanovich-Masur [KM96] showed that for any basepoint  $X$  in  $\mathcal{T}(S)$  and for almost every infinite sample path  $\omega = (w_n)$  the sequence  $w_n X$  converges to the Thurston boundary  $\text{PMF}(S)$ . This defines a harmonic measure on  $\text{PMF}(S)$  giving the distribution of the limit points of infinite sample paths. Furthermore, Kaimanovich-Masur showed that the measure is non-atomic, and the set of uniquely ergodic foliations have full measure. However, such harmonic measures are singular with respect to the Lebesgue measure class on  $\text{PMF}(S)$ . See [Gad14] for singularity for finitely supported random walks and [GMT] for singularity for random walks with finite first moment with respect to the word metric. Therefore the final statement of Theorem 1.1 does not follow immediately from the Lebesgue measure case.

There are countably many pseudo-Anosov elements, so the union of their endpoints has measure zero with respect to any of the measures discussed above. Hence, the results for pseudo-Anosov elements do not follow immediately from the results for bi-infinite geodesics. Apart from random walks, there are other ways to study generic foliations of mapping class group elements. For example, every conjugacy class of a pseudo-Anosov element corresponds to a closed geodesic in moduli space. Eskin-Mirzakhani [EM11] showed that the number of closed geodesics in moduli space of Teichmüller length at most  $R$  grows asymptotically as  $e^{hR}/hR$ , and Eskin-Mirzakhani-Rafi [EMR12] showed that the proportion of the closed geodesics of length at most  $R$  which do not lie in the principal stratum tends to zero exponentially quickly in  $R$ . They also show that the number of lattice points in the ball of radius  $R$  in Teichmüller space grows asymptotically as  $e^{hR}$ , and the proportion of these lattice points which lie close to a geodesic contained in non-principal stratum also tends to zero exponentially fast.

The proof of Theorem 1.1 follows the following strategy. Let  $g$  be a pseudo-Anosov element whose invariant Teichmüller geodesic  $\gamma_g$  lies in the principal stratum. We show that any Teichmüller geodesic that fellow travels  $\gamma_g$  for a sufficiently large distance  $D$ , depending on  $g$ , also lies in the principal stratum. Next, we show that if  $g$  lies in the semigroup generated by the support of  $\mu$ , there is a positive probability that the geodesic  $\gamma_\omega$  tracked by a sample path  $\omega$ , fellow travels the invariant geodesic  $\gamma_g$  for distance at least  $D$ . Ergodicity of the shift map on  $\text{Mod}(S)^\mathbb{Z}$  then implies that a positive proportion of subsegments of  $\gamma_\omega$  of length  $D$  fellow travel some translate of  $\gamma_g$ . We then use work of Dahmani-Horbez [DH15] which shows that for almost all sample paths  $\omega$ , for sufficiently large  $n$ , all elements  $w_n$  are pseudo-Anosov, with invariant geodesics  $\gamma_{w_n}$  which fellow travel  $\gamma_\omega$  for a distance which grows linearly in  $n$ . In particular, this implies that  $\gamma_{w_n}$  fellow travels a sufficiently long subsegment of a translate of  $\gamma_g$ , and so lies in the principal stratum.

## 2. TEICHMÜLLER PRELIMINARIES

Let  $S$  be an orientable surface of finite type, whose Teichmüller space has complex dimension at least two, i.e. not a sphere with at most four punctures, or a torus with at most two punctures. The Teichmüller metric is given by

$$d_{\mathcal{T}}(X, Y) = \frac{1}{2} \inf_f \log K(f),$$

where the infimum is taken over all quasiconformal maps  $f: X \rightarrow Y$  in the given homotopy class, and  $K(f)$  is the quasiconformal constant of  $f$ . As there is a unique Teichmüller geodesic connecting any pair of points in Teichmüller space, we may sometimes write  $[X, Y]$  to denote the Teichmüller geodesic segment from  $X$  to  $Y$ . For detailed background about the Teichmüller metric and the geometry of quadratic differentials, see for example [Wri15].

The complex of curves  $\mathcal{C}(S)$  is an infinite graph with vertices isotopy classes of simple closed curves on  $S$ . Two vertices  $[\alpha], [\beta]$  are separated by an edge if the curves  $\alpha$  and  $\beta$  can be isotoped to be disjoint. The graph  $\mathcal{C}(S)$  is locally infinite and has infinite diameter, and Masur-Minsky showed that  $\mathcal{C}(S)$  is  $\delta$ -hyperbolic in the sense of Gromov [MM99].

By the uniformization theorem, a conformal class  $X$  determines a unique hyperbolic metric on  $S$ , which we shall also denote by  $X$ . For a hyperbolic surface  $X$ , a systole of  $X$  is a simple closed curve that has the shortest length in the hyperbolic metric. The set of systoles of  $X$  is a finite set whose diameter in  $\mathcal{C}(S)$  is bounded above by a constant that depends only on the topology of  $S$ . Thus, the systole defines a coarse projection map  $\pi: \mathcal{T}(S) \rightarrow \mathcal{C}(S)$ . For notational simplicity, we will use upper case letters for points  $X$  in  $\mathcal{T}(S)$ , and the corresponding lower case letters  $x = \pi(X)$  for their projections to the curve complex. Masur-Minsky [MM99, 6.1] showed that  $\pi$  is coarsely Lipschitz, i.e. there are constants  $M_1 > 0, A_1 > 0$  that depend only on  $S$ , such that for any pair of

points  $X, Y \in \mathcal{T}(S)$

$$(2.1) \quad d_{\mathcal{C}}(x, y) < M_1 d_{\mathcal{T}}(X, Y) + A_1.$$

Moreover, Masur-Minsky also show that Teichmüller geodesics  $\gamma$  project to uniformly unparameterised quasigeodesics in  $\mathcal{C}(S)$ . Let  $(M_2, A_2)$ -be the quasigeodesicity constants for the projection of a Teichmüller geodesic, and these constants depend only on  $S$ .

The set of hyperbolic surfaces  $X \in \mathcal{T}(S)$  for which the length of the systole is less than  $\epsilon$  form the  $\epsilon$ -thin part  $\mathcal{T}(S)_\epsilon$  of Teichmüller space. The complement  $K_\epsilon = \mathcal{T}(S) \setminus \mathcal{T}(S)_\epsilon$  is called the thick part. By Mumford compactness,  $\text{Mod}(S) \setminus K_\epsilon$  is compact, and furthermore a metric regular neighbourhood of the thick part is contained in a larger thick part. More precisely, for any  $\epsilon > 0$ , and any  $D \geq 0$ , there is a constant  $\epsilon'$ , depending on  $\epsilon, D$  and the surface  $S$ , such that a metric  $D$ -neighbourhood of  $K_\epsilon$ , in the Teichmüller metric, is contained in  $K_{\epsilon'}$ .

Let  $\gamma$  and  $\gamma'$  be two geodesics in a metric space  $(M, d)$ . If there are choices of (not necessarily unit speed) parameterizations  $\gamma(t)$  and  $\gamma'(t)$  such that there is a constant  $E$  with  $d(\gamma(t), \gamma'(t)) \leq E$  for all  $t$ , then we say that  $\gamma$  and  $\gamma'$  are fellow travellers with fellow travelling constant  $E$ , or  $E$ -fellow travel. If  $d(\gamma(t), \gamma'(t)) \leq E$ , for all  $t$ , for the unit speed parameterizations of  $\gamma$  and  $\gamma'$ , then we say that  $\gamma$  and  $\gamma'$  are parameterized  $E$ -fellow travellers.

Let  $\gamma$  and  $\gamma'$  be two Teichmüller geodesics whose projections to the curve complex  $\pi(\gamma)$  and  $\pi(\gamma')$  fellow travel. In general, this does not imply that the original Teichmüller geodesics fellow travel in Teichmüller space. However, we now show in the following lemma that if  $\gamma$  is contained in a thick part  $K_\epsilon$ , and  $\pi(\gamma')$  fellow travels  $\pi(\gamma)$  for a sufficiently long distance in  $\mathcal{C}(S)$ , then  $\gamma'$  contains a point that is close to  $\gamma$  in Teichmüller space.

**Lemma 2.2.** *For any constants  $\epsilon > 0$  and  $E \geq 0$ , there are constants  $L > 0$  and  $F > 0$ , depending on  $\epsilon, E$  and the surface  $S$ , such that if  $\gamma = [X, Y]$  is a Teichmüller geodesic segment contained in the thick part  $K_\epsilon$ , of length at least  $L$ , and  $\gamma' = [X', Y']$  is a Teichmüller geodesic segment, whose endpoints  $x', y'$  in  $\mathcal{C}(S)$  are distance at most  $E$  from the endpoints  $x, y$  of  $\pi(\gamma)$ , i.e.  $d_{\mathcal{C}(S)}(x, x') \leq E$  and  $d_{\mathcal{C}(S)}(y, y') \leq E$ , then there is a point  $Z$  on  $\gamma'$  such that  $d_{\mathcal{T}}(Z, \gamma) \leq F$ .*

This result may also be deduced from work of Horbez [Hor15, Proposition 3.10] and Dowdall-Duchin-Masur [DDM14, Theorem A], extending Rafi [Raf14], but for the convenience of the reader, we provide a direct proof of this result in Section 4, relying only on Rafi [Raf14]. In particular, we will make extensive use of the following fellow travelling result for Teichmüller geodesics whose endpoints are close together in the thick part.

**Theorem 2.3.** [Raf14, Theorem 7.1] *For any constants  $\epsilon > 0$  and  $A \geq 0$ , there is a constant  $B$ , depending only on  $\epsilon, A$  and the surface  $S$ , such that if  $[X, Y]$  and  $[X', Y']$  are two Teichmüller geodesics, with  $X$  and  $Y$  in the  $\epsilon$ -thick part, and*

$$d_{\mathcal{T}}(X, X') \leq A \text{ and } d_{\mathcal{T}}(Y, Y') \leq A,$$

*then  $[X, Y]$  and  $[X', Y']$  are parameterized  $B(\epsilon, A)$ -fellow travellers.*

We now continue with the proof of Theorem 1.1 assuming Lemma 2.2. Recall that the Gromov product based at a point  $u \in \mathcal{C}(S)$  is defined to be

$$(x, y)_u = \frac{1}{2} (d_{\mathcal{C}}(u, x) + d_{\mathcal{C}}(u, y) - d_{\mathcal{C}}(x, y)).$$

Given points  $x, y \in \mathcal{C}(S)$  and a constant  $R > 0$ , the  $R$ -shadow of  $y$  is defined to be

$$S_x(y, R) = \{z \in \mathcal{C}(S) \mid (y, z)_x \geq d_{\mathcal{C}}(x, y) - R\}.$$

The definition we use here for shadows follows [MT14], and may differ slightly from other sources. The following lemma follows from the thin triangles property of Gromov hyperbolic spaces, and we give a proof for the convenience of the reader.

**Lemma 2.4.** *There is a constant  $D$ , which only depends on  $\delta$ , and a constant  $E$ , which only depends on  $R$  and  $\delta$ , such that if  $d_C(x, y) \geq 2R + D$ , then for any  $x' \in S_y(x, R)$  and any  $y' \in S_x(y, R)$ , any geodesic segment  $[x', y']$  contains a subsegment which  $E$ -fellow travels  $[x, y]$ .*

*Proof.* We shall write  $O(\delta)$  to denote a constant which only depends (not necessarily linearly) on  $\delta$ .

Let  $p$  be the nearest point projection of  $x'$  to  $[x, y]$ , and let  $q$  be the nearest point projection of  $y'$  to  $[x, y]$ . The nearest point projection of the shadow  $S_x(y, R)$  is contained in an  $(R + O(\delta))$ -neighbourhood of  $y$ , see for example [MT14, Proposition 2.4], so  $d_C(x, p) \leq R + O(\delta)$  and  $d_C(y, q) \leq R + O(\delta)$ . Recall that if  $d_C(p, q) \geq O(\delta)$  then any geodesic from  $x'$  to  $y'$  passes within an  $O(\delta)$ -neighborhood of both  $p$  and  $q$ , see for example [MT14, Proposition 2.3]. Therefore, if  $d(x, y) \geq 2R + O(\delta)$ , then this implies that if  $p'$  is the closest point on  $[x', y']$  to  $p$ , and  $q'$  is the closest point on  $[x', y']$  to  $q$ , then  $[p', q']$   $E$ -fellow travels  $[x, y]$ , where  $E$  is a constant which only depends on  $R$  and  $\delta$ , as required.  $\square$

**Remark 2.5.** *One can replace the geodesic segments  $[x, y]$  and  $[x', y']$  by  $(M_2, A_2)$ -quasigeodesic segments. The constants  $D$  and  $E$  now change, and in addition to  $R$  and  $\delta$ , they now depend on the quasigeodesicity constants.*

We shall write  $\text{PMF}(S)$  for the set of projective measured foliations on the surface  $S$ , which is Thurston's boundary for Teichmüller space. A projective measured foliation is uniquely ergodic if the foliation supports a unique projective measure class. Let  $\text{UE}(S)$  be the subset of  $\text{PMF}(S)$  consisting of uniquely ergodic foliations. We shall give  $\text{UE}(S)$  the corresponding subspace topology. A uniquely ergodic foliation determines a class of mutually asymptotic geodesic rays in  $\mathcal{T}(S)$ , as shown by Masur [Mas80]. These rays project to a class of mutually asymptotic quasigeodesic rays in  $\mathcal{C}(S)$ , and so determines a point in the Gromov boundary of the curve complex. This boundary map is injective on uniquely ergodic foliations, see for example Hubbard-Masur [HM79]. Thus,  $\text{UE}(S)$  is also a subset of  $\partial\mathcal{C}(S)$ . Klarreich [Kla] showed that  $\partial\mathcal{C}(S)$  is homeomorphic to the quotient of the set of minimal foliations in  $\text{PMF}(S)$  by the equivalence relation which forgets the measure. In particular, this implies that the two subspace topologies on  $\text{UE}(S)$ , induced from inclusions in  $\text{PMF}(S)$  and  $\partial\mathcal{C}(S)$ , are the same.

Let  $\gamma$  be a Teichmüller geodesic in a thick part  $K_\epsilon$ . Let  $\lambda^+$  and  $\lambda^-$  be the projective classes of vertical and horizontal measured foliations of  $\gamma$ . By the work of Kerckhoff-Masur-Smillie [KMS86, Theorem 3], vertical foliations of Teichmüller rays that are recurrent to a thick part are uniquely ergodic, so the foliations  $\lambda^+$  and  $\lambda^-$  are uniquely ergodic, and by Hubbard-Masur [HM79] such a pair  $(\lambda^-, \lambda^+)$  determines a unique bi-infinite Teichmüller geodesic. Given two points  $X$  and  $Y$  in Teichmüller space, and a constant  $r \geq 0$ , define  $\Gamma_r(X, Y)$  to be the set of all oriented geodesics with uniquely ergodic vertical and horizontal foliations, which intersect both  $B_r(X)$  and  $B_r(Y)$ , and furthermore, whose first point of intersection with either  $B_r(X)$  or  $B_r(Y)$  lies in  $B_r(X)$ . A Teichmüller geodesic with uniquely ergodic vertical foliation  $\lambda^+$  and uniquely ergodic horizontal foliation  $\lambda^-$  determines a point  $(\lambda^-, \lambda^+)$  in  $\text{UE}(S) \times \text{UE}(S)$ . Therefore  $\Gamma_r(X, Y)$  determines a subset of  $\text{UE}(S) \times \text{UE}(S)$ , which, by abuse of notation, we shall also denote by  $\Gamma_r(X, Y)$ .

**Proposition 2.6.** *For any Teichmüller geodesic  $\gamma$  contained in a thick part  $K_\epsilon$ , with vertical foliation  $\lambda^+$  and horizontal foliation  $\lambda^-$ , there is a constant  $r > 0$ , depending on  $\epsilon$ , such that for any pair of points  $X$  and  $Y$  on  $\gamma$ , the set  $\Gamma_r(X, Y)$  contains an open neighbourhood of  $(\lambda^-, \lambda^+)$  in  $\text{UE}(S) \times \text{UE}(S)$ .*

*Proof.* As  $\text{Mod}(S)$  acts coarsely transitively on the curve complex  $\mathcal{C}(S)$ , there is a constant  $R > 0$ , depending only on  $S$ , such that for all  $x$  and  $y$  in  $\mathcal{C}(S)$ , the limit set of the shadow  $\overline{S_x(y, R)}$  contains a non-empty open set in  $\partial\mathcal{C}(S)$ , see for example [MT14, Propositions 3.18–19]. Given such an  $R$ , let  $D$  and  $E$  be the constants in Lemma 2.4, such that if  $d(x, y) \geq D$  then for any  $x' \in S_y(x, R)$  and  $y' \in S_x(y, R)$ , a geodesic  $[x', y']$  has a subsegment which  $E$ -fellow travels with  $[x, y]$ . Given  $\epsilon$  and




FIGURE 2.7. Shadows in  $\mathcal{C}(S)$ .

$E$ , let  $L$  and  $F$  be the constants in Lemma 2.2, i.e. if  $\gamma$  and  $\gamma''$  are two Teichmüller geodesics of length at least  $L$ , whose endpoints in  $\mathcal{C}(S)$  are distance at most  $E$  apart, then the distance from  $\gamma$  to  $\gamma'$  is at most  $F$ .

As  $\gamma$  lies in the thick part  $K_\epsilon$ , there is a constant  $D'$ , depending only on  $\epsilon$ , such that if  $d_{\mathcal{T}}(X, Y) \geq D'$ , then  $d_{\mathcal{C}}(x, y) \geq D$ . Let  $Z_1$  and  $Z_2$  be points along  $\gamma$  such that  $[X, Y] \subset [Z_1, Z_2]$ , the orientations of the segments agree,  $d_{\mathcal{T}}(X, Y) \geq D'$ ,  $d_{\mathcal{T}}(Z_1, X) > L$  and  $d_{\mathcal{T}}(Y, Z_2) > L$ . Consider the limit sets  $\overline{S_{z_1}(z_2, R)}$  and  $\overline{S_{z_2}(z_1, R)}$  in  $\partial\mathcal{C}(S)$ , and let  $\xi^+$  and  $\xi^-$  be uniquely ergodic foliations in  $\overline{S_{z_1}(z_2, R)}$  and  $\overline{S_{z_2}(z_1, R)}$ , respectively. Let  $\gamma'$  be the Teichmüller geodesic with vertical foliation  $\xi^+$  and the horizontal foliation  $\xi^-$ . By Lemma 2.4, the projection  $\pi(\gamma')$  fellow travels  $\pi(\gamma)$  with constant  $E$  between  $z_1$  and  $z_2$ . For clarity, denote by  $Z'_1, X', Y'$  and  $Z'_2$  the points of  $\gamma'$  whose projections  $z'_1, x', y'$  and  $z'_2$  are coarsely the closest points to  $z_1, x, y$  and  $z_2$  respectively, i.e. the distances  $d_{\mathcal{C}}(z'_1, z_1), d_{\mathcal{C}}(x', x), d_{\mathcal{C}}(y', y)$  and  $d_{\mathcal{C}}(z'_2, z_2)$  are all at most  $E$ . By Lemma 2.2 applied to the segments  $[Z'_1, X']$  and  $[Z_1, X]$  there is a point  $W_1 \in [Z'_1, X']$  such that  $d_{\mathcal{T}}(W_1, [Z_1, X]) \leq F$ . Similarly, there is a point  $W_2 \in [Y', Z'_2]$  such that  $d_{\mathcal{T}}(W_2, [Y, Z_2]) \leq F$ .

By the fellow travelling result, Theorem 2.3, the Teichmüller geodesic segment  $[W_1, W_2] \subset \gamma'$  fellow travels  $\gamma$  with the constant  $r = B(\epsilon, F)$ . In particular,  $\gamma'$  passes through  $B_r(X)$  and  $B_r(Y)$ , and hence lies in  $\Gamma_r(X, Y)$ , and so this set contains an open neighbourhood of  $(\lambda^-, \lambda^+)$ . We have shown this as long as  $d_{\mathcal{T}}(X, Y) \geq D'$ , but for  $r' = 2r + D'$ , every pair of balls  $B_{r'}(X')$  and  $B_{r'}(Y')$  contain smaller balls  $B_r(X)$  and  $B_r(Y)$  with  $d_{\mathcal{T}}(X, Y) \geq D'$ , so the stated result follows.  $\square$

### 3. FELLOW TRAVELLING OF INVARIANT AND TRACKED GEODESICS

In this section, we establish that along almost every sample path  $\omega$ , for sufficiently large  $n$ , the invariant Teichmüller geodesic for the pseudo-Anosov element  $w_n$ , has a subsegment, whose length grows linearly in  $n$ , which fellow travels the Teichmüller geodesic sublinearly tracked by  $\omega$ . This uses a result of Dahmani-Horbez [DH15] and the fellow travelling result, Theorem 2.3. We fix a basepoint  $X \in \mathcal{T}(S)$ .

We require a slight rephrasing of a result of Dahmani-Horbez. Let  $\ell$  be the drift of the random walk in the Teichmüller metric. Kaimanovich-Masur [KM96] showed that almost every bi-infinite

sample path  $\omega$  converges to distinct uniquely ergodic measured foliations  $\lambda_\omega^+$  and  $\lambda_\omega^-$ , with  $w_n X$  converging to  $\lambda_\omega^+$ , and  $w_{-n} X$  converging to  $\lambda_\omega^-$  as  $n \rightarrow \infty$ . Let  $\gamma_\omega$  be the unique bi-infinite Teichmüller geodesic determined by these foliations, and we shall give  $\gamma_\omega$  a unit speed parameterization, such that  $\gamma_\omega(0)$  is a closest point on  $\gamma_\omega$  to  $X$ , and as  $t \rightarrow \infty$  the geodesic  $\gamma_\omega(t)$  converges to  $\lambda^+$ . If  $w_n$  is pseudo-Anosov, then we shall write  $\gamma_{w_n}$  for its invariant Teichmüller geodesic.

To compare notation with [DH15, Theorem 2.6], we shall choose their group  $G$  to be  $\text{Mod}(S)$ . The hyperbolic  $G$ -electrification  $(Y, X)$  they use is in our case the electrification with  $Y = \mathcal{T}(S)$  and  $X = \mathcal{C}(S)$ . We do not have to worry about metric symmetrisation as the Teichmüller metric is symmetric. The space  $\text{PMF}(S)$  of projective measured foliations on  $S$  gives the bordification of  $\mathcal{T}(S)$ . Then steps 1 and 3 in the proof of [DH15, Theorem 2.6], stated in the context of Teichmüller space, can be rephrased as follows:

**Proposition 3.1.** *Given  $\epsilon > 0$ , there are constants  $F > 0$  and  $0 < e < \frac{1}{2}$ , such that for almost every  $\omega$ , there exists  $N$ , such that for all  $n \geq N$ , there are points  $Y_0$  and  $Y_1$  of  $\gamma_{w_n}$  and points  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$  of  $\gamma_\omega$ , such that*

- (1)  $d_{\mathcal{T}}(\gamma_\omega(T_0), Y_0) \leq F$ ,
- (2)  $d_{\mathcal{T}}(\gamma_\omega(T_1), Y_1) \leq F$ ,
- (3)  $0 \leq T_0 \leq e\ell n \leq (1 - e)\ell n \leq T_1 \leq \ell n$ , and
- (4)  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$  are in the thick part  $K_\epsilon$ .

For completeness, we summarize specifically in the Teichmüller context the key ideas behind [DH15, Theorem 2.6]. The summary differs slightly from the proof given in [DH15], as we describe the special case which applies to Proposition 3.1. First, the Teichmüller translation length for a pseudo-Anosov map  $f$  is the infimum (which is realised in  $\mathcal{T}(S)$ ) of  $d_{\mathcal{T}}(Y, fY)$  as  $Y$  varies over  $\mathcal{T}(S)$ . The definition directly implies that for almost every sample path  $\omega$  the quantity  $\ell n$  is an upper bound for the Teichmüller translation length of  $w_n$  as  $n \rightarrow \infty$ . Next, we summarize the argument for the analogous lower bound.

Almost every sample path  $\omega$  makes linear progress in  $\mathcal{T}(S)$ . By the work of Tiozzo [Tio15], almost every  $\omega$  sub-linearly tracks a Teichmüller geodesic  $\gamma_\omega$ . Thus, given  $0 < e < 1/2$ , for all  $n$  large enough depending on  $\omega$  the Teichmüller distance of  $w_n$  from  $\gamma_\omega(\ell n)$  is at most  $e\ell n$ . By the ergodicity of the shift map on  $\text{Mod}(S)^\mathbb{Z}$ , almost every  $\omega$  recurs infinitely often to a fixed neighbourhood of  $\gamma_\omega$ . Furthermore, the recurrence has an asymptotic frequency. The point of  $\gamma_\omega$  closest to such a recurrent point is also in the thick part because of its proximity to a point of the sample path. In particular, similar arguments imply that given  $0 < e < 1/2$ , there is  $n$  large enough depending on  $\omega$  such that inside the segment  $[\gamma_\omega(0), \gamma_\omega(e\ell n)]$  one can find a subsegment of a definite size contained entirely in the  $\epsilon$ -thick part. A thick Teichmüller segment of a definite size makes definite progress under projection to the curve complex. The proximity of  $w_n$  to  $\gamma_\omega(\ell n)$  and the hyperbolicity of  $\mathcal{C}(S)$  implies that the vertical foliation for  $\gamma_{w_n}$  as a point in  $\partial\mathcal{C}(S)$  is contained in the limit set of some shadow set of the point  $\pi(\gamma_\omega(\ell n))$ . In particular, this implies that the projection  $\pi(\gamma_{w_n})$  of the invariant geodesic for  $w_n$  fellow travels  $\pi(\gamma_\omega)$  over a segment that contains this segment of definite progress. Hence, by [DH15, Proposition 3.7] attributed to Dowdall-Duchin-Masur [DDM14], the fellow traveling in the curve complex lifts to a fellow traveling of Teichmüller geodesics  $\gamma_{w_n}$  and  $\gamma_\omega$  over the thick subsegment. In particular, there is a point  $Y_0$  of  $\gamma_{w_n}$  that is a bounded Teichmüller distance away from some point  $\gamma_\omega(T_0)$  with  $0 \leq T_0 \leq e\ell n$ . A symmetric argument shows that there is a point  $Y_1$  of  $\gamma_{w_n}$  that is a bounded Teichmüller distance away from some point  $\gamma_\omega(T_1)$  with  $(1 - e)\ell n \leq T_1 \leq \ell n$ . Proposition 3.1 gives a precise version of these statements. By a simple triangle inequality, it then follows that  $\ell n$  is asymptotically a lower bound for the Teichmüller translation length of  $w_n$ .

Dahmani-Horbez state condition (4) in terms of a ‘‘contraction’’ property that they define:  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$  are ‘‘contraction’’ points on  $\gamma_\omega$  for the projection map to the curve complex. In effect,



the property being used by them is that under the projection to the curve complex  $\gamma_\omega$  makes definite progress at  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$ . See the discussion related to [DH15, Propositions 3.6 and 3.7]. We recall their precise definition [DH15, Definition 3.5] for definite progress here:

**Definition 3.2.** *Given constants  $B, C > 0$ , a Teichmüller geodesic  $\gamma$  makes  $(B, C)$ -progress at a point  $Y = \gamma(T)$  if the image under  $\pi$  of the subsegment of  $\gamma$  of length  $B$  starting at  $Y$  has diameter at least  $C$  in the curve complex.*

For completeness, we prove that definite progress implies thickness.

**Lemma 3.3.** *If  $\gamma$  makes  $(B, C)$ -progress at  $Y$ , then there is a constant  $\epsilon > 0$ , which depends on  $B$  and  $C$ , such that  $Y$  lies in the thick part  $K_\epsilon$ .*

*Proof.* Let  $\alpha$  be the systole for the hyperbolic surface  $Y$ . For any point  $Y'$  on the subsegment, Wolpert's lemma implies

$$\ell_{Y'}(\alpha) \leq e^B \ell_Y(\alpha).$$

We will use the following version of the Collar Lemma, due to Matelski [Mat76], which states that a simple closed geodesic of length  $\ell$  is contained in an embedded annular collar neighbourhood of width at least  $w_\ell$ , where a lower bound for  $w_\ell$  is given by

$$\sinh^{-1} \left( \frac{1}{\sinh(\ell/2)} \right),$$

and furthermore, this lower bound holds for all  $\ell > 0$ . Thus the width of the collar neighbourhood for  $\alpha$  in the hyperbolic metric corresponding to  $Y'$  is bounded below by

$$\sinh^{-1} \left( \frac{1}{\sinh(e^B \ell_Y(\alpha)/2)} \right),$$

and the bound tends to infinity monotonically as  $\ell_Y(\alpha)$  tends to zero. Suppose  $\beta$  is the systole at  $Y'$ , and  $d_C(\alpha, \beta) \geq C$ . This implies that the intersection number satisfies

$$i(\alpha, \beta) \geq \frac{C-1}{2}.$$

From the lower bound on the width of the collar, the length of  $\beta$  has to satisfy

$$\ell_{Y'}(\beta) \geq \frac{C-1}{2} \sinh^{-1} \left( \frac{1}{\sinh(e^B \ell_Y(\alpha)/2)} \right).$$

Since  $\beta$  is the systole at  $Y'$ , the length of  $\beta$  at  $Y'$  is at most the length of  $\alpha$  at  $Y'$ , so one obtains

$$e^B \ell_Y(\alpha) \geq \frac{C-1}{2} \sinh^{-1} \left( \frac{1}{\sinh(e^B \ell_Y(\alpha)/2)} \right).$$

Note that  $\sinh$  is monotonically increasing, zero at zero, and unbounded, so this gives a lower bound  $\epsilon$  on how small  $\ell_Y(\alpha)$  can be, which depends on  $B$  and  $C$ .  $\square$

**Remark 3.4.** *Lemma 3.3 implies that the points  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$  in Proposition 3.1 are in a thick part  $K_\epsilon$ . By the fellow travelling result, Theorem 2.3 the geodesics  $\gamma_\omega$  and  $\gamma_{w_n}$  fellow travel between  $\gamma_\omega(T_0)$  and  $\gamma_\omega(T_1)$ . Let  $s = B(\epsilon, F)$  be the constant for fellow traveling of  $\gamma_\omega$  and  $\gamma_{w_n}$ .*

We now show that for a pseudo-Anosov element  $g$  in the support of  $\mu$ , there is a positive probability that the geodesic  $\gamma_\omega$  fellow travels the invariant geodesic  $\gamma_g$ . We shall write  $\nu$  for the harmonic measure on  $\text{UE}(S)$ , and  $\check{\nu}$  for the reflected harmonic measure, i.e. the harmonic measure arising from the random walk generated by the probability distribution  $\check{\mu}(g) = \mu(g^{-1})$ .

**Lemma 3.5.** *Let  $g$  be a pseudo-Anosov element contained in the support of  $\mu$  with invariant Teichmüller geodesic  $\gamma_g$ . Then there is a constant  $r > 0$  such that  $\check{\nu} \times \nu(\Gamma_r(X, Y)) > 0$  for all  $X$  and  $Y$  on  $\gamma_g$ .*

*Furthermore, there is a constant  $\rho > 0$ , depending on  $g$ , such that for all constants  $D \geq 0$ , there is a positive probability (that depends on  $D$ ) for the subsegment of  $\gamma_\omega$  of length  $D$ , centered at a closest point on  $\gamma_\omega$  to the basepoint, to  $\rho$ -fellow travel with  $\gamma_g$ .*

*Proof.* Let  $\lambda^+$  and  $\lambda^- \in \text{PMF}(S)$  be the vertical and horizontal foliations of  $\gamma_g$ . Fix an  $\epsilon > 0$  such that the thick part  $K_\epsilon$  contains the geodesic  $\gamma_g$ . Let  $r$  be the constant in Proposition 2.6, i.e. for any points  $X$  and  $Y$  on  $\gamma_g$ , the set  $\Gamma_r(X, Y)$  contains an open neighbourhood of  $(\lambda^-, \lambda^+)$ . We recall:

**Proposition 3.6.** [MT14, Proposition 5.4] *Let  $G$  be a non-elementary, countable group acting by isometries on a separable Gromov hyperbolic space  $X$ , and let  $\mu$  be a non-elementary probability distribution on  $G$ . Then there is a number  $R_0$  such that for any group element  $g$  in the semigroup generated by the support of  $\mu$ , the closure of the shadow  $S_{x_0}(gx_0, R_0)$  has positive hitting measure for the random walk determined by  $\mu$ .*

Let  $x_0 = \pi(X_0)$  be the projection of the basepoint  $X_0$  into the curve complex. We may assume that  $\Gamma_r(X, Y)$  contains an open neighbourhood of  $(\lambda^-, \lambda^+)$  of the form  $U^- \times U^+$ , where  $U^-$  is an open neighbourhood of  $\lambda^-$  in  $\text{UE}(S)$ , and  $U^+$  is an open neighbourhood of  $\lambda^+$  in  $\text{UE}(S)$ . As

$$\bigcap_{i \in \mathbb{N}} \overline{S_{x_0}(g^{-i}x_0, R_0)} = \lambda^- \quad \text{and} \quad \bigcap_{i \in \mathbb{N}} \overline{S_{x_0}(g^i x_0, R_0)} = \lambda^+,$$

there is an integer  $i$ , such that the limit sets of the shadows are contained in the open neighbourhoods of  $\lambda^+$  and  $\lambda^-$ , i.e.

$$\overline{S_{x_0}(g^{-i}x_0, R_0)} \cap \text{UE}(S) \subset U^- \quad \text{and} \quad \overline{S_{x_0}(g^i x_0, R_0)} \cap \text{UE}(S) \subset U^+.$$

The element  $g^{-1}$  is in the semigroup generated by the inverses of  $\text{Supp}(\mu)$ , i.e.  $g^{-1} \in \text{Supp}(\check{\mu})$ . Hence, by Proposition 3.6,

$$\check{\nu} \times \nu \left( \overline{S_{x_0}(g^{-i}x_0, R_0)} \times \overline{S_{x_0}(g^i x_0, R_0)} \right) > 0,$$

and so  $\check{\nu} \times \nu(\Gamma_r(X, Y)) > 0$ , as required.

The final statement then follows from Theorem 2.3, which implies that there is a  $\rho > 0$  such that any geodesic in  $\Gamma_r(X, Y)$  must  $\rho$ -fellow travel  $[X, Y]$ , as required. Here we may choose  $X$  and  $Y$  on  $\gamma_g$  such that the geodesic  $[X, Y]$  contains a subsegment of length  $D$  centered at any closest point on  $\gamma_g$  to the basepoint  $X_0$ ; as  $\gamma_g$  is contained in a thick part  $K_\epsilon$ , the set of closest points on  $\gamma_g$  to  $X_0$  has bounded diameter, depending only on  $\epsilon$  and the surface  $S$ .  $\square$

We now make use of the principal stratum assumption, i.e. that the semigroup generated by  $\text{Supp}(\mu)$  contains a pseudo-Anosov  $g$  whose invariant Teichmüller geodesic  $\gamma_g$  lies in the principal stratum. We first prove the following proposition:

**Proposition 3.7.** *Let  $g$  be a pseudo-Anosov element of  $\text{Mod}(S)$ , whose invariant Teichmüller geodesic is contained in the principal stratum. For any  $\rho > 0$ , there is a constant  $D > 0$ , depending on  $\rho$  and  $g$ , such that for any pair of points  $X, Y$  on  $\gamma_g$  with  $d_{\mathcal{T}}(X, Y) \geq D$ , any Teichmüller geodesic in  $\Gamma_\rho(X, Y)$  lies in the principal stratum.*

*Proof.* The invariant geodesic  $\gamma_g$  projects to a closed geodesic in moduli space, and so lies in the thick part  $K_\epsilon$ , for some  $\epsilon$  depending on  $g$ . If a geodesic  $\gamma$  passes through  $B_\rho(X)$  and  $B_\rho(Y)$  for  $X, Y \in \gamma_g$  then by the fellow travelling result, Theorem 2.3 it  $B(\epsilon, \rho)$ -fellow travels  $[X, Y]$ .

To derive a contradiction, suppose that there is a sequence  $\phi_n$  of geodesic segments in non-principal strata such that the  $\phi_n$  fellow travel  $\gamma_g$  for distances  $d_n$  with  $d_n \rightarrow \infty$  as  $n \rightarrow \infty$ . As the cyclic group generated by  $g$  acts coarsely transitively on  $\gamma_g$ , we may assume that the midpoints of the  $\phi_n$  are all a bounded distance from the basepoint  $X$  in Teichmüller space. By convergence on compact sets we can pass to a limiting geodesic  $\phi$  which lies in a non-principal strata, as the principal stratum is open. The geodesics  $\phi$  and  $\gamma_g$  fellow travel in the forward direction for all times. By [Mas80, Theorem 2], this implies that  $\phi$  and  $\gamma_g$  have the same vertical foliation. This is a contradiction since  $\phi$  is in a non-principal stratum.  $\square$

We now complete the proof of Theorem 1.1.

*Proof of Theorem 1.1.* We fix a pseudo-Anosov element  $g$  in the support of  $\mu$  for which the invariant Teichmüller geodesic  $\gamma_g$  is contained in the principal stratum. Without loss of generality, we fix the basepoint  $X$  to be on  $\gamma_g$ .

Let  $\epsilon > 0$  be sufficiently small such that  $\gamma_g$  is contained in the thick part  $K_\epsilon$ . Given this  $\epsilon$ , let  $F_0 > 0$  and  $0 < e_0 < \frac{1}{2}$  be the constants from Proposition 3.1. Let  $\rho > 0$  be the constant in Lemma 3.5 that ensures  $\rho$ -fellow travelling for any length  $D > 0$  between  $\gamma_\omega$  and  $\gamma_g$  with a positive probability, depending on  $D$ . By Proposition 3.7, there is a  $D_0$  such that any Teichmüller geodesic which  $(\rho + F_0)$ -fellow travels with  $\gamma_g$  distance at least  $D_0$  is contained in the principal stratum. We shall set  $D = D_0 + 2F_0$ .

Let  $k > 0$  be the smallest positive integer such that  $d_{\mathcal{T}}(g^{-k}X, g^kX) \geq D$ . By Theorem 2.3, any geodesic in  $\Gamma_r(g^{-k}X, g^kX)$   $\rho$ -fellow travels the subsegment  $[g^{-k}X, g^kX]$  of  $\gamma_g$ . Let  $\Omega \subset \text{Mod}(S)^{\mathbb{Z}}$  consist of those sample paths  $\omega$  such that the sequences  $w_{-n}X$  and  $w_nX$  converge to distinct uniquely ergodic foliations  $(\lambda^-, \lambda^+) \in \Gamma_r(g^{-k}X, g^kX)$ . Lemma 3.5 implies that the subset  $\Omega$  has positive probability  $p > 0$ .

Let  $\sigma : \text{Mod}(S)^{\mathbb{Z}} \rightarrow \text{Mod}(S)^{\mathbb{Z}}$  be the shift map. Ergodicity of  $\sigma$  implies that for almost every  $\omega$ , there is some  $n \geq 0$  such that  $\sigma^n(\omega) \in \Omega$ . For such  $n$ , the subsegment of  $\gamma_\omega$  of length  $D$ , centered at the closest point on  $\gamma_\omega$  to the point  $w_nX$ ,  $\rho$ -fellow travels with a translate of  $w_n\gamma_g$ . In particular, this implies that  $\gamma_\omega$  lies in the principal stratum, giving the final claim in Theorem 1.1.

For almost every  $\omega$ , the proportion of times  $1 \leq n \leq N$  such that  $\sigma^n(\omega) \in \Omega$  tends to  $p$  as  $N \rightarrow \infty$ . Choose numbers  $e_1$  and  $e_2$  such that  $e_0 < e_1 < e_2 < \frac{1}{2}$ , then this also holds for  $N$  replaced with either  $e_1N$  or  $(1 - e_1)N$ . So this implies that the proportion of times  $e_1N \leq n \leq (1 - e_1)N$  with this property also tends to  $p$  as  $N \rightarrow \infty$ . This implies that given  $\omega$ , there is an  $N_0$  such that for all  $N \geq N_0$ , there is an  $n$  with  $e_1N \leq n \leq (1 - e_1)N$  and  $\sigma^n(\omega) \in \Omega$ .

Recall that by sublinear tracking in Teichmüller space, due to Tiozzo [Tio15], there is a constant  $\ell > 0$  such that for almost all  $\omega$ ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} d_{\mathcal{T}}(w_nX, \gamma_\omega(\ell n)) = 0,$$

where  $\gamma_\omega$  is parameterized such that  $\gamma_\omega(0)$  is a closest point on  $\gamma_\omega$  to the basepoint. Therefore, possibly replacing  $N_0$  with a larger number, we may also assume that  $d_{\mathcal{T}}(x_NX, \gamma_\omega(\ell N)) \leq (e_2 - e_1)N$  for all  $N \geq N_0$ .

Choose numbers  $\ell_1$  and  $\ell_2$ , with  $\ell_1 < \ell < \ell_2$ , and choose them sufficiently close to  $\ell$  so that  $e_0\ell < e_1\ell_1$  and  $(1 - e_1)\ell_2 < (1 - e_0)\ell$ . Therefore the geodesic  $[\gamma_\omega(e_2\ell_1N - \rho), \gamma_\omega((1 - e_2)\ell_2N + \rho)]$  contains a subsegment of length at least  $D$  which  $\rho$ -fellow travels with a translate of  $\gamma_g$ . By our choice of  $\ell_1$  and  $\ell_2$ , the geodesic  $[\gamma_\omega(e_0\ell_1N - \rho), \gamma_\omega((1 - e_0)\ell_2N + \rho)]$  is contained in  $[\gamma_\omega(e_2\ell N), \gamma_\omega((1 - e_2)\ell N)]$  for  $N$  sufficiently large. Now using Proposition 3.1, this implies that

the invariant geodesic  $\gamma_{w_n}$  ( $\rho + F_0$ )-fellow travels with  $\gamma_g$  for a distance at least  $D - 2F_0 \geq D_0$ . Then by Proposition 3.7,  $\gamma_{w_n}$  is contained in the principal stratum, as required.  $\square$

#### 4. FELLOW TRAVELLING IN TEICHMÜLLER SPACE

We now provide a direct proof of Lemma 2.2, relying only on results from Rafi [Raf14]. The first result we shall use is the fellow travelling result for Teichmüller geodesics with endpoints in the thick part, Theorem 2.3. The second result is a thin triangles theorem for triangles in Teichmüller space, where one side has a large segment contained in the thick part.

**Theorem 4.1.** [Raf14, Theorem 8.1] *For every  $\epsilon > 0$ , there are constants  $C$  and  $L$ , depending only on  $\epsilon$  and  $S$ , such that the following holds. Let  $X, Y$  and  $Z$  be three points in  $\mathcal{T}(S)$ , and let  $[X', Y']$  be a segment of  $[X, Y]$  with  $d_{\mathcal{T}}(X', Y') > L$ , such that  $[X', Y']$  is contained in the  $\epsilon$ -thick part of  $\mathcal{T}(S)$ . Then, there is a point  $W \in [X', Y']$ , such that*

$$\min\{d_{\mathcal{T}}(W, [X, Z]), d_{\mathcal{T}}(W, [Y, Z])\} \leq C.$$

We now prove Lemma 2.2.

*Proof.* The projection of an  $\epsilon_i$ -thick Teichmüller geodesic makes definite progress in the curve complex, i.e. there exist constants  $P_i$  and  $Q_i$ , depending on  $\epsilon_i$  and the surface  $S$ , such that for any points  $X, Y$  on  $\gamma$  we have the estimate

$$(4.2) \quad d_{\mathcal{C}}(x, y) \geq P_i d_{\mathcal{T}}(X, Y) - Q_i.$$

Set  $\epsilon_1 = \epsilon$ . Let  $L_1$  and  $C_1$  be the corresponding constants from the thin triangle result, Theorem 4.1. Let  $B_1 = B(\epsilon_1, C_1 + L_1/2)$  be the constant in the fellow travelling theorem, Theorem 2.3. Set  $\epsilon_2 = \epsilon(\epsilon_1, B_1)$ , i.e. the  $B_1$ -neighbourhood of  $K_{\epsilon_1}$  is contained in  $K_{\epsilon_2}$ . Given this  $\epsilon_2$ , let  $L_2$  and  $C_2$  be the corresponding constants from the thin triangle result, Theorem 4.1. Now that all the constants we need are defined, we shall choose  $L$  to be the maximum of the following three terms

$$(4.3) \quad \begin{aligned} & \frac{3}{P_1} (M_1 C_1 + Q_1 + M_2 E + A_2 + A_1) + \frac{3}{2} L_1, \\ & 3L_2 + 3L_1 + 6C_1, \\ & \frac{3}{P_2} (M_1 C_2 + Q_2 + M_2 E + A_2 + A_1) + \frac{3}{2} L_1 + 3B_1. \end{aligned}$$

Let  $Z_1$  be the point that is  $1/3$  of the way along  $[X, Y]$ . Let  $\gamma_1$  be the geodesic segment of  $\gamma$  centered at  $Z_1$  with length  $L_1$ . Similarly, let  $Z_2$  be the point that is  $2/3$  of the way along  $[X, Y]$ . Let  $\gamma_2$  be the geodesic segment of  $\gamma$  centered at  $Z_2$  with length  $L_1$ . The second term of (4.3) implies that  $L > 3L_1$ . Figure 4.4 illustrates this setup.

Applying the thin triangles result, Theorem 4.1, to  $X, Y$  and  $Y'$ , there is a point  $W_1$  on  $[X, Y'] \cup [Y, Y']$  within distance  $C_1$  of  $\gamma_1$ . Similarly, there is a point  $W_2$  on  $[X, Y'] \cup [Y, Y']$  within distance  $C_1$  of  $\gamma_2$ .

We now show that there is a lower bound on the distance of  $\gamma_2$  from  $[Y, Y']$ . In particular, the same is true for the distance of  $\gamma_1$ , from  $[Y, Y']$ .

**Claim 4.5.** *The Teichmüller distance from  $\gamma_2$  to  $[Y, Y']$  is at least  $C_1$ .*

*Proof.* The Teichmüller distance of  $Y$  from  $\gamma_2$  is at least  $\frac{1}{3}L - \frac{1}{2}L_1$ , i.e.

$$d_{\mathcal{T}}(\gamma_2, Y) \geq \frac{1}{3}L - \frac{1}{2}L_1.$$

As  $\epsilon_1$ -thick geodesics make definite progress in  $\mathcal{C}(S)$ , (4.2), this implies

$$d_{\mathcal{C}}(\pi(\gamma_2), y) \geq P_1(\frac{1}{3}L - \frac{1}{2}L_1) - Q_1.$$




FIGURE 4.4. Fellow travelling geodesics in  $\mathcal{T}(S)$ .

Teichmüller geodesic segments project to  $(M_2, A_2)$ -quasigeodesics in  $\mathcal{C}(S)$ . Since the endpoints of  $\gamma$  and  $\gamma'$  are distance at most  $E$  apart in  $\mathcal{C}(S)$ , this implies,

$$d_{\mathcal{C}}(\pi(\gamma_2), \pi([Y, Y'])) \geq P_1(\frac{1}{3}L - \frac{1}{2}L_1) - Q_1 - M_2E - A_2.$$

As the curve complex distance is a coarse lower bound on the Teichmüller distance, (2.1), this implies

$$d_{\mathcal{T}}(\gamma_2, [Y, Y']) \geq \frac{1}{M_1}(P_1(\frac{1}{3}L - \frac{1}{2}L_1) - Q_1 - M_2E - A_2 - A_1).$$

Finally, a comparison with the first term of (4.3) shows that

$$d_{\mathcal{T}}(\gamma_2, [Y, Y']) > C_1,$$

as required.  $\square$

This implies that  $W_2$  lies on  $[X, Y']$  and not on  $[Y, Y']$ . As  $\gamma_1$  is further away from  $[Y, Y']$  along  $\gamma$  than  $\gamma_2$ , the same argument implies that  $W_1$  lies on  $[X, Y']$ . Furthermore,  $d_{\mathcal{T}}(W_1, Z_1) \leq C_1 + L_1/2$ . Similarly  $d_{\mathcal{T}}(W_2, Z_2) \leq C_1 + L_1/2$ .

The segment  $[X, Z_2]$  is in the  $\epsilon_1$ -thick part. The endpoints of  $[X, W_2]$  are within distance  $C_1 + L_1/2$  of the endpoints of  $[X, Z_2]$ . So by the fellow travelling result, i.e. Theorem 2.3,  $[X, W_2]$  and  $[X, Z_2]$  are  $B_1$ -fellow travellers, where  $B_1 = B(\epsilon_1, C_1 + L_1/2)$ . Recall that  $B_1$  depends on  $\epsilon_1, C_1 + L_1/2$ , and the surface  $S$ .

Recall that  $\epsilon_2 = \epsilon'(\epsilon_1, B_1)$ , i.e. the  $B_1$ -neighbourhood of  $K_{\epsilon_1}$  is contained in  $K_{\epsilon_2}$ . Note that  $\epsilon_2$  depends only on the constants  $\epsilon = \epsilon_1, B_1$  and the surface  $S$ . In particular, the geodesic  $[X, W_2]$  is contained in the  $\epsilon_2$ -thick part. Given  $\epsilon_2$ , recall that  $L_2$  and  $C_2$  are the corresponding constants from the thin triangle result, Theorem 4.1.

By the triangle inequality,

$$d_{\mathcal{T}}(Z_1, W_1) + d_{\mathcal{T}}(W_1, W_2) + d_{\mathcal{T}}(W_2, Z_2) \geq d_{\mathcal{T}}(Z_1, Z_2).$$

Thus, the Teichmüller distance between  $W_1$  and  $W_2$  is at least

$$d_{\mathcal{T}}(W_1, W_2) \geq \frac{1}{3}L - 2C_1 - L_1,$$

The second term of (4.3) implies that the right hand side above is at least  $L_2$ . So we may apply the thin triangles result, Theorem 4.1, to  $X, X'$  and  $Y'$  to conclude that there is a point  $Z$  on  $[X, X'] \cup [X', Y']$  within distance  $C_2$  of  $[W_1, W_2]$ .

We now show a lower bound for the distance between  $[W_1, W_2]$  and  $[X, X']$ .

**Claim 4.6.** *The distance between  $[W_1, W_2]$  and  $[X, X']$  is at least  $C_2$ .*

*Proof.* Let  $W$  be a point of  $[W_1, W_2]$  that is closest to  $X$ . Let  $V$  be the point of  $\gamma$  that is closest to  $W$ . Then

$$B_1 \geq d_{\mathcal{T}}(W, V) \quad \text{and} \quad d_{\mathcal{T}}(X, V) \geq \frac{1}{3}L - \frac{1}{2}L_1$$

Thus, by the triangle inequality

$$d_{\mathcal{T}}(X, W) \geq d_{\mathcal{T}}(X, V) - d_{\mathcal{T}}(W, V) > \frac{1}{3}L - \frac{1}{2}L_1 - B_1,$$

or equivalently

$$d_{\mathcal{T}}([W_1, W_2], X) \geq \frac{1}{3}L - \frac{1}{2}L_1 - B_1.$$

As  $\epsilon_2$ -thick geodesics make definite progress in  $\mathcal{C}(S)$ , (4.2) implies

$$d_{\mathcal{C}}(\pi([W_1, W_2]), x) \geq P_2(\frac{1}{3}L - \frac{1}{2}L_1 - B_1) - Q_2.$$

As the distance between  $x$  and  $x'$  in  $\mathcal{C}(S)$  is at most  $E$ , this implies,

$$d_{\mathcal{C}}(\pi([W_1, W_2]), \pi([X, X'])) \geq P_2(\frac{1}{3}L - \frac{3}{2}L_1 - C_1) - Q_2 - M_2E - A_2.$$

As the curve complex distance is a coarse lower bound on the Teichmüller metric (2.1), this implies

$$d_{\mathcal{T}}([W_1, W_2], [X, X']) \geq \frac{1}{M_1}(P_2(\frac{1}{3}L - \frac{1}{2}L_1 - B_1) - Q_2 - M_2E - A_2 - A_1).$$

A comparison with the third term in (4.3) then shows that

$$d_{\mathcal{T}}([W_1, W_2], [Y, Y']) > C_2,$$

as required. □

Claim 4.6 implies that  $Z$  lies on  $[X', Y']$  and not on  $[X, X']$ . The segments  $[W_1, W_2]$  and  $[Z_1, Z_2]$  are  $B_1$ -fellow travellers. As  $Z$  lies within distance  $C_2$  of  $[W_1, W_2]$ , the distance of  $Z$  from  $\gamma$  is at most  $C_2 + B_1$ . To conclude the proof of Lemma 2.2, we may set  $F = C_2 + B_1$ , which depends only on  $\epsilon, A$  and the surface  $S$ , as required. □

## REFERENCES

- [ABEM12] Jayadev Athreya, Alexander Bufetov, Alex Eskin, and Maryam Mirzakhani, *Lattice point asymptotics and volume growth on Teichmüller space*, *Duke Math. J.* **161** (2012), no. 6, 1055–1111.
- [DH15] F. Dahmani and C. Horbez, *Spectral theorems for random walks on mapping class groups and  $\text{Out}(F_N)$*  (2015), available at [arXiv:1506.06790](https://arxiv.org/abs/1506.06790).
- [DHM15] K. Delp, D. Hoffoss, and J. Manning, *Problems in groups, geometry and 3-manifolds* (2015), available at [arXiv:1512.04620](https://arxiv.org/abs/1512.04620).
- [DDM14] Spencer Dowdall, Moon Duchin, and Howard Masur, *Statistical hyperbolicity in Teichmüller space*, *Geom. Funct. Anal.* **24** (2014), no. 3, 748–795.
- [EM11] Alex Eskin and Maryam Mirzakhani, *Counting closed geodesics in moduli space*, *J. Mod. Dyn.* **5** (2011), no. 1, 71–105.
- [EMR12] Alex Eskin, Maryam Mirzakhani, and Kasra Rafi, *Counting geodesics in a stratum*, *Invent. Math.*, to appear (2012).
- [Gad14] Vaibhav Gadre, *Harmonic measures for distributions with finite support on the mapping class group are singular*, *Duke Math. J.* **163** (2014), no. 2, 309–368, DOI 10.1215/00127094-2430368. MR3161316
- [GMT] Vaibhav Gadre, Joseph Maher, and Giulio Tiozzo, *Word length statistics for Teichmüller geodesics and singularity of harmonic measure*, to appear in *Comment. Math. Helv.*
- [Hor15] Camille Horbez, *Central limit theorems for mapping class groups and  $\text{Out}(F_N)$*  (2015), available at [arXiv:1506.07244](https://arxiv.org/abs/1506.07244).
- [HM79] John Hubbard and Howard Masur, *Quadratic differentials and foliations*, *Acta Math.* **142** (1979), no. 3-4, 221–274.
- [KM96] Vadim A. Kaimanovich and Howard Masur, *The Poisson boundary of the mapping class group*, *Invent. Math.* **125** (1996), no. 2, 221–264.
- [KP15] Ilya Kapovich and Catherine Pfaff, *A train track directed random walk on  $\text{Out}(F_r)$* , *Internat. J. Algebra Comput.* **25** (2015), no. 5, 745–798.
- [KMS86] Steven Kerckhoff, Howard Masur, and John Smillie, *Ergodicity of billiard flows and quadratic differentials*, *Ann. of Math. (2)* **124** (1986), no. 2, 293–311.
- [Kla] Erica Klarreich, *The boundary at infinity of the curve complex and the relative Teichmüller space*.
- [Mah11] Joseph Maher, *Random walks on the mapping class group*, *Duke Math. J.* **156** (2011), no. 3, 429–468.
- [MT14] Joseph Maher and Giulio Tiozzo, *Random walks on weakly hyperbolic groups*, *Journal für die reine und angewandte Mathematik (Crelle’s Journal)*, to appear (2014), available at [arXiv:1410.4173](https://arxiv.org/abs/1410.4173).
- [Mas80] Howard Masur, *Uniquely ergodic quadratic differentials*, *Comment. Math. Helv.* **55** (1980), no. 2, 255–266, DOI 10.1007/BF02566685. MR576605
- [Mas82] Howard Masur, *Interval exchange transformations and measured foliations*, *Ann. of Math. (2)* **115** (1982), no. 1, 169–200.
- [MM99] Howard A. Masur and Yair N. Minsky, *Geometry of the complex of curves. I. Hyperbolicity*, *Invent. Math.* **138** (1999), no. 1, 103–149.
- [Mat76] J. Peter Matelski, *A compactness theorem for Fuchsian groups of the second kind*, *Duke Math. J.* **43** (1976), no. 4, 829–840.
- [Raf14] Kasra Rafi, *Hyperbolicity in Teichmüller space*, *Geom. Topol.* **18** (2014), no. 5, 3025–3053.
- [Riv08] Igor Rivin, *Walks on groups, counting reducible matrices, polynomials, and surface and free group automorphisms*, *Duke Math. J.* **142** (2008), no. 2, 353–379.
- [Tio15] Giulio Tiozzo, *Sublinear deviation between geodesics and sample paths*, *Duke Math. J.* **164** (2015), no. 3, 511–539.
- [Vee82] William A. Veech, *Gauss measures for transformations on the space of interval exchange maps*, *Ann. of Math. (2)* **115** (1982), no. 1, 201–242.
- [Wri15] Alex Wright, *Translation surfaces and their orbit closures: an introduction for a broad audience*, *EMS Surv. Math. Sci.* **2** (2015), no. 1, 63–108.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF GLASGOW, 15 UNIVERSITY GARDENS, GLASGOW G12 8QW UK

*E-mail address:* [Vaibhav.Gadre@glasgow.ac.uk](mailto:Vaibhav.Gadre@glasgow.ac.uk)

DEPARTMENT OF MATHEMATICS, COLLEGE OF STATEN ISLAND, CUNY, 2800 VICTORY BOULEVARD, STATEN ISLAND, NY 10314, USA, AND DEPARTMENT OF MATHEMATICS, 4307 GRADUATE CENTER, CUNY, 365 5TH AVENUE, NEW YORK, NY 10016, USA

*E-mail address:* [joseph.maher@csi.cuny.edu](mailto:joseph.maher@csi.cuny.edu)