



Busetto, F., Codognato, G. and Ghosal, S. (2017) Asymptotic equivalence between Cournot–Nash and Walras equilibria in exchange economies with atoms and an atomless part. *International Journal of Game Theory*, 46(4), pp. 975-990. (doi:[10.1007/s00182-017-0566-x](https://doi.org/10.1007/s00182-017-0566-x))

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/133462/>

Deposited on: 04 January 2017

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

# Asymptotic Equivalence between Cournot-Nash and Walras Equilibria in Exchange Economies with Atoms and an Atomless Part\*

Francesca Busetto<sup>†</sup>, Giulio Codognato<sup>‡</sup>, Sayantan Ghosal<sup>§</sup>

## Abstract

In this paper, we consider an exchange economy à la Shitovitz (1973), with atoms and an atomless set. We associate with it a strategic market game of the kind first proposed by Lloyd S. Shapley, known as the *Shapley window model*. We analyze the relationship between the set of the Cournot-Nash allocations of the strategic market game and the Walras allocations of the exchange economy with which it is associated. We show, with an example, that even when atoms are countably infinite, any Cournot-Nash allocation of the game is not a Walras allocation of the underlying exchange economy. Accordingly, in the original spirit of Cournot (1838), we partially replicate the mixed exchange economy by increasing the number of atoms, without affecting the atomless part, and ensuring that the measure space of agents remains finite. Our main theorem shows that any sequence of Cournot-Nash allocations of the strategic market games associated with the partial replications of the exchange economy has a limit point for each trader and that the assignment determined by these limit points is a Walrasian allocation of the original economy.

*Journal of Economic Literature* Classification Numbers: C72, D51.

---

\*We would like to thank Walter Bossert, Simone Tonin, and an anonymous referee for their comments and suggestions. Francesca Busetto and Giulio Codognato gratefully acknowledge financial support from MIUR (PRIN 20103S5RN3).

<sup>†</sup>Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, Via Tomadini 30, 33100 Udine, Italy.

<sup>‡</sup>Dipartimento di Scienze Economiche e Statistiche, Università degli Studi di Udine, Via Tomadini 30, 33100 Udine, Italy, and EconomiX, Université Paris Ouest-Nanterre La Défense, 200 Avenue de la République, 92001 Nanterre Cedex, France.

<sup>§</sup>Adam Smith Business School, University of Glasgow, Main Building, Glasgow, G128QQ, United Kingdom.

## 1 Introduction

A large literature has been developed in the past decades, aiming at extending to a general equilibrium framework the classical Cournot's theory of oligopoly (see Cournot (1838)), where oligopolistic agents that interact noncooperatively among themselves face a sector of consumers taking prices as given. Most of the contributions on this issue belongs to two main lines of research: the Cournot-Walras equilibrium approach, initiated by Gabszewicz and Vial (1972), and the strategic market game approach, initiated by Shapley and Shubik (1977).

In their 1972's paper, Gabszewicz and Vial transposed to a general equilibrium setting Cournot's original idea of an asymmetric economy with production, in which firms with oligopolistic power that interact strategically on quantities face a sector of consumers behaving à la Walras. Nevertheless, in the same paper, they pointed out two major difficulties inherent in the standard Cournotian approach with strategic firms: first, profit maximization may not be a rational objective for firms that have influence on prices; second, the equilibrium is not independent from the rule chosen to normalize prices. These difficulties have been overcome within the Cournot-Walras approach by moving to the analysis of pure exchange economies (see Codognato and Gabszewicz (1993), d'Aspremont, Dos Santos Ferreira, and Gérard-Varet (1997), Gabszewicz and Michel (1997), Shitovitz (1997), among others). However, this latter class of models do not avoid a well-known problem inherent in the Cournot-Walras approach: there, an equilibrium may not exist, even in mixed strategies, since the Walras price correspondence may fail to admit a continuous selection (see Dierker and Grodal (1986)). A further fundamental problem common to the whole Cournot-Walras approach is that it leaves unexplained why some agents behave strategically while other agents behave competitively.

A different approach has been developed (still in pure exchange), that uses strategic market games à la Shapley and Shubik with the aim at providing a formal explanation of perfectly and imperfectly competitive behavior. A fundamental contribution in this line is the paper of Okuno, Postlewaite and Roberts (1980). In particular, since the work of these authors, results in this direction have been obtained by incorporating within the framework of strategic market games a mixed measure space of traders à la Shitovitz (1973), composed by large traders, represented as atoms, and small traders, represented by an atomless part. In this setting, while all agents have a priori the same strategic position, some of them turn out to have influence

on prices and some other turn out to be Walrasian, depending on their characteristics and their weight in the economy. Then, the asymmetric structure typical of the Cournotian theory is endogenously generated.

Busetto, Codognato, and Ghosal (2011) obtained a generalization of Okuno et al.'s work by using a mixed version of the Cournot-Nash equilibrium model first proposed by Lloyd S. Shapley, and known as the *Shapley window model* (this model was analyzed, in the case of finite economies, by Sahi and Yao (1989)). Busetto et al. (2011) provided an endogenous explanation of oligopolistic and competitive behavior. Moreover, working within this strategic market game framework permitted them to show the existence of an equilibrium in pure strategies, and then to overcome the non-continuity problem which characterizes the Cournot-Walras approach.

This paper studies the link between Cournot and Walras equilibrium, with the aim of providing a noncooperative foundation to the theory of perfect competition. A mixed exchange economy à la Shitovitz is associated with the same strategic market game à la Shapley proposed by Busetto et al. (2011). Within this framework, the relationship between the set of the Cournot-Nash allocations of the strategic market game and the Walras allocations of the underlying exchange economy is examined. We show, with an example, that even when the set of atoms is countably infinite, there is the robust possibility that no Cournot-Nash allocation of the strategic market game is a Walras allocation of the underlying exchange economy, because some atoms remain non-negligible in size. This non equivalence result then motivates us to analyze the asymptotic relationship between appropriately defined sequences of Cournot-Nash allocations of the strategic market game and the Walras allocations of the exchange economy. We do this by introducing a concept of replication which we call *à la Cournot*, since it extends to a general equilibrium context the original Cournotian idea of replication: that is, we partially replicate the economy by increasing only the number of atoms, this way making them asymptotically negligible, without affecting the atomless part. At the same time, the mechanism of replication of atoms is constructed in such a way that the measure space of traders remains finite. If this requirement was not satisfied, the general equilibrium model of oligopoly studied in the paper would not be well-defined. Our main theorem establishes that any sequence of Cournot-Nash allocations of the strategic market games associated with the partial replications of the exchange economy has a limit point for each trader and that the assignment determined by these limit points is a Walrasian allocation of the original economy.

To prove the limit theorem, we use analytical tools introduced by Sahi and Yao (1989) to show their convergence result for finite economies, and by Codognato and Ghosal (2000) to show their equivalence theorem à la Aumann (1964). However, we have to tackle new technical issues due to the fact that the space of traders has a mixed nature. To this end, we exploit some tools previously applied to the Shapley window model for mixed economies by Busetto et al. (2011). In particular, in order to determine the limit points of the sequences of Cournot-Nash allocations, we use a version of the Fatou's lemma in several dimensions proved by Artstein (1979). Moreover, a key point in our paper is that since the Walrasian price taking behavior of small traders is not assumed a priori, as in the Cournot-Walras equilibrium models, but endogenously generated, we do not need to impose any continuity assumption on the selections from the Walras price correspondence.

The general equilibrium approach adopted here distinguishes our limit result from the well-known results, obtained within the Cournotian tradition in partial equilibrium, establishing that the Cournot equilibrium approaches the competitive equilibrium as the number of oligopolists goes to infinity (see Frank (1965), Ruffin (1971), Novshek (1980), among others).

Our limit result is also different from existing results in the strategic market game literature: within their Shapley window model for finite economies, Sahi and Yao (1989) showed the convergence of sequences of Cournot-Nash allocations to a Walras allocation by using a replication concept *à la Edgeworth*, in which all types of agents are replicated (on this kind of replication see Debreu and Scarf (1963)).

Other limit results were obtained within the line opened by Gabszewicz and Vial: see Roberts (1980), Mas-Colell (1983), and Novshek and Sonnenschein ((1983), (1987)), among others. Also these results are obtained by using replication concepts in which all types of agents are replicated. Moreover, as stressed by Mas-Colell (1982) (see, in particular, pp. 203-204), they depend in an essential way on the assumption that there exists a continuous selection from the Walras price correspondence.

The paper is organized as follows. In Section 2, we build the mathematical model. In Section 3, we introduce the notion of a  $\delta$ -positive Cournot-Nash equilibrium used in the construction of the limit theorem, and we state a theorem on its existence (the proof is in the Appendix). In Section 4, we provide the example on the non-equivalence between the sets of Cournot-Nash and Walras allocations when the atoms are countably infinite. In Section 5, we introduce the replication à la Cournot. In Section 6,

we state the existence of a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium (the proof is in the Appendix). In Section 7, we state and prove the limit theorem.

## 2 The mathematical model

We consider a pure exchange economy,  $\mathcal{E}$ , with large traders, represented as atoms, and small traders, represented by an atomless part. The space of traders is denoted by the measure space  $(T, \mathcal{T}, \mu)$ , where  $T$  is the set of traders,  $\mathcal{T}$  is the  $\sigma$ -algebra of all  $\mu$ -measurable subsets of  $T$ , and  $\mu$  is a real valued, non-negative, countably additive measure defined on  $\mathcal{T}$ . We assume that  $(T, \mathcal{T}, \mu)$  is finite, i.e.,  $\mu(T) < \infty$ . This implies that the measure space  $(T, \mathcal{T}, \mu)$  contains at most countably many atoms. Let  $T_1$  denote the set of atoms and  $T_0 = T \setminus T_1$  the atomless part of  $T$ . A null set of traders is a set of measure 0. Null sets of traders are systematically ignored throughout the paper. Thus, a statement asserted for “each” trader in a certain set is to be understood to hold for all such traders except possibly for a null set of traders. The word “integrable” is to be understood in the sense of Lebesgue.

In the exchange economy, there are  $l$  different commodities. A commodity bundle is a point in  $R_+^l$ . An assignment (of commodity bundles to traders) is an integrable function  $\mathbf{x}: T \rightarrow R_+^l$ . There is a fixed initial assignment  $\mathbf{w}$ , satisfying the following assumption.

**Assumption 1.**  $\mathbf{w}(t) > 0$ , for each  $t \in T$ .

An allocation is an assignment  $\mathbf{x}$  for which  $\int_T \mathbf{x}(t) d\mu = \int_T \mathbf{w}(t) d\mu$ . The preferences of each trader  $t \in T$  are described by a utility function  $u_t: R_+^l \rightarrow R$ , satisfying the following assumptions.

**Assumption 2.**  $u_t: R_+^l \rightarrow R$  is continuous, strongly monotone, and quasi-concave, for each  $t \in T$ .

Let  $\mathcal{B}$  denote the Borel  $\sigma$ -algebra of  $R_+^l$ . Moreover, let  $\mathcal{T} \otimes \mathcal{B}$  denote the  $\sigma$ -algebra generated by all the sets  $E \times F$  such that  $E \in \mathcal{T}$  and  $F \in \mathcal{B}$ .

**Assumption 3.**  $u: T \times R_+^l \rightarrow R$ , given by  $u(t, x) = u_t(x)$ , for each  $t \in T$  and for each  $x \in R_+^l$ , is  $\mathcal{T} \otimes \mathcal{B}$ -measurable.

We also need the following assumption (see Sahi and Yao (1989)).

**Assumption 4.** There are at least two traders in  $T_1$  for whom  $\mathbf{w}(t) \gg 0$ ,  $u_t$  is continuously differentiable in  $R_{++}^l$ , and  $\{x \in R_+^l : u_t(x) = u_t(\mathbf{w}(t))\} \subset R_{++}^l$ .

A price vector is a nonnull vector  $p \in R_+^l$ .

A Walras equilibrium of  $\mathcal{E}$  is a pair  $(p^*, \mathbf{x}^*)$ , consisting of a price vector  $p^*$  and an allocation  $\mathbf{x}^*$ , such that, for each  $t \in T$ ,  $u_t(\mathbf{x}^*(t)) \geq u_t(y)$ , for all  $y \in \{x \in R_+^l : p^*x = p^*\mathbf{w}(t)\}$ . A Walras allocation of  $\mathcal{E}$  is an allocation  $\mathbf{x}^*$  for which there exists a price vector  $p^*$  such that the pair  $(p^*, \mathbf{x}^*)$  is a Walras equilibrium of  $\mathcal{E}$ .

We define now the strategic market game,  $\Gamma$ , associated with  $\mathcal{E}$ .

Let  $\mathcal{P}(R_+^{l^2})$  denote the power set of  $R_+^{l^2}$ . A strategy correspondence is a correspondence  $\mathbf{B} : T \rightarrow \mathcal{P}(R_+^{l^2})$  such that, for each  $t \in T$ ,  $\mathbf{B}(t) = \{(b_{ij} \in R_+^{l^2} : \sum_{j=1}^l b_{ij} \leq \mathbf{w}^i(t), i = 1, \dots, l)\}$ , where  $b_{ij}$ ,  $i, j = 1, \dots, l$ , represents the amount of commodity  $i$  that trader  $t$  offers in exchange for commodity  $j$ . Moreover, we denote, with some abuse of notation, by  $b(t) \in \mathbf{B}(t)$  a strategy of trader  $t$ . A strategy selection is an integrable function  $\mathbf{b} : T \rightarrow R_+^{l^2}$ , such that, for each  $t \in T$ ,  $\mathbf{b}(t) \in \mathbf{B}(t)$ . Given a strategy selection  $\mathbf{b}$ , we call aggregate matrix the matrix  $\bar{\mathbf{B}}$  such that  $\bar{\mathbf{b}}_{ij} = (\int_T \mathbf{b}_{ij}(t) d\mu)$ ,  $i, j = 1, \dots, l$ . Moreover, we denote by  $\mathbf{b} \setminus b(t)$  the strategy selection obtained from  $\mathbf{b}$  by replacing  $\mathbf{b}(t)$  with  $b(t) \in \mathbf{B}(t)$  and by  $\bar{\mathbf{B}} \setminus b(t)$  the corresponding aggregate matrix.

Then, we introduce two further definitions (see Sahi and Yao (1989)).

**Definition 1.** A nonnegative square matrix  $A$  is said to be irreducible if, for every pair  $(i, j)$ , with  $i \neq j$ , there is a positive integer  $k$  such that  $a_{ij}^{(k)} > 0$ , where  $a_{ij}^{(k)}$  denotes the  $ij$ -th entry of the  $k$ -th power  $A^k$  of  $A$ .

**Definition 2.** Given a strategy selection  $\mathbf{b}$ , a price vector  $p$  is said to be market clearing if

$$p \in R_{++}^l, \sum_{i=1}^l p^i \bar{\mathbf{b}}_{ij} = p^j (\sum_{i=1}^l \bar{\mathbf{b}}_{ji}), j = 1, \dots, l. \quad (1)$$

By Lemma 1 in Sahi and Yao (1989), there is a unique, up to a scalar multiple, price vector  $p$  satisfying (1) if and only if  $\bar{\mathbf{B}}$  is irreducible. Then, we denote by  $p(\mathbf{b})$  a function which associates with each strategy selection  $\mathbf{b}$  the unique, up to a scalar multiple, price vector  $p$  satisfying (1), if  $\bar{\mathbf{B}}$  is irreducible, and is equal to 0, otherwise.

Given a strategy selection  $\mathbf{b}$  and a price vector  $p$ , consider the assignment determined as follows:

$$\mathbf{x}^j(t, \mathbf{b}(t), p) = \mathbf{w}^j(t) - \sum_{i=1}^l \mathbf{b}_{ji}(t) + \sum_{i=1}^l \mathbf{b}_{ij}(t) \frac{p^i}{p^j}, \text{ if } p \in R_{++}^l,$$

$$\mathbf{x}^j(t, \mathbf{b}(t), p) = \mathbf{w}^j(t), \text{ otherwise,}$$

$j = 1, \dots, l$ , for each  $t \in T$ .

Given a strategy selection  $\mathbf{b}$  and the function  $p(\mathbf{b})$ , the traders' final holdings are determined according to this rule and consequently expressed by the assignment

$$\mathbf{x}(t) = \mathbf{x}(t, \mathbf{b}(t), p(\mathbf{b})),$$

for each  $t \in T$ .<sup>1</sup> It is straightforward to show that this assignment is an allocation satisfying  $p(\mathbf{b})\mathbf{x}(t, \mathbf{b}(t), p(\mathbf{b})) = p(\mathbf{b})\mathbf{w}(t)$ , for each  $t \in T$ .

We are now able to define a Cournot-Nash equilibrium of  $\Gamma$  (see Codognato and Ghosal (2000) and Busetto et al. (2011)).

**Definition 3.** *A strategy selection  $\hat{\mathbf{b}}$  such that  $\bar{\mathbf{B}}$  is irreducible is a Cournot-Nash equilibrium of  $\Gamma$  if*

$$u_t(\mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))) \geq u_t(\mathbf{x}(t, b(t), p(\hat{\mathbf{b}} \setminus b(t)))),$$

for each  $b(t) \in \mathbf{B}(t)$  and for each  $t \in T$ .<sup>2</sup>

A Cournot-Nash allocation of  $\Gamma$  is an allocation  $\hat{\mathbf{x}}$  such that  $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$ , for each  $t \in T$ , where  $\hat{\mathbf{b}}$  is a Cournot-Nash equilibrium of  $\Gamma$ .

### 3 The existence of a $\delta$ -positive Cournot-Nash equilibrium of $\Gamma$

In this section, we define the notion of a  $\delta$ -positive Cournot-Nash equilibrium, which was first used by Sahi and Yao (1989) to prove their existence theorem for the finite version of the Shapley model and their limit result. We will use it, in this paper, for analogous purposes.

Let  $\bar{T}_1 \subset T_1$  be a set consisting of two traders in  $T_1$  for whom Assumption 4 holds. Moreover, let  $\delta = \min_{t \in \bar{T}_1} \{\frac{1}{l} \min\{\mathbf{w}^1(t), \dots, \mathbf{w}^l(t)\}\}$ . We say that the correspondence  $\mathbf{B}^\delta : T \rightarrow \mathcal{P}(R_+^{l^2})$  is a  $\delta$ -positive strategy correspondence if  $\mathbf{B}^\delta(t) = \mathbf{B}(t) \cap \{(b_{ij}) \in R_+^{l^2} : \sum_{i \notin J} \sum_{j \in J} (b_{ij} + b_{ji}) \geq \delta, \text{ for each } J \subseteq \{1, \dots, l\}\}$ , for each  $t \in \bar{T}_1$  and if  $\mathbf{B}^\delta(t) = \mathbf{B}(t)$ , for the remaining traders

<sup>1</sup>In order to save in notation, with some abuse, we denote by  $\mathbf{x}$  both the function  $\mathbf{x}(t)$  and the function  $\mathbf{x}(t, \mathbf{b}(t), p(\mathbf{b}))$ .

<sup>2</sup>Let us notice that, as this definition of a Cournot-Nash equilibrium explicitly refers to irreducible matrices, it applies only to active equilibria of  $\Gamma$  (on this point, see Sahi and Yao (1989)).

$t \in T$ . Moreover, we say that a strategy selection  $\mathbf{b}$  is  $\delta$ -positive if  $\mathbf{b}(t) \in \mathbf{B}^\delta(t)$ , for each  $t \in T$ . Finally, we say that a Cournot-Nash equilibrium  $\hat{\mathbf{b}}$  of  $\Gamma$  is  $\delta$ -positive if  $\hat{\mathbf{b}}$  is a  $\delta$ -positive strategy selection.

The following theorem shows the existence of a  $\delta$ -positive Cournot-Nash equilibrium of  $\Gamma$ . It is a straightforward consequence of the existence theorem in Busetto et al. (2011).

**Theorem 1.** *Under Assumptions 1, 2, 3, and 4, there exists a  $\delta$ -positive Cournot-Nash equilibrium  $\hat{\mathbf{b}}$  of  $\Gamma$ .*

**Proof.** See the Appendix. ■

## 4 An example

Sahi and Yao (1989) formalized the Shapley window model in the context of an exchange economy with a finite set of traders. Codognato and Ghosal (2000) reconsidered the Sahi and Yao's model within an exchange economy with an atomless continuum of traders. In this framework, they showed an equivalence result à la Aumann (1964) between the set of the Cournot-Nash allocations of the window model and the set of the Walras allocations of the underlying exchange economy. The mixed measure space we are considering here may contain countably infinite atoms. This raises the question whether an equivalence result can be obtained also in this case. The following example provides a negative answer to this question.

Consider an exchange economy  $\mathcal{E}$  where  $l = 2$ , which satisfies Assumptions 1, 2, 3, and 4. Moreover, assume that  $T_1$  contains countably infinite atoms and that there is an atom  $\tau \in T_1$  such that  $\mathbf{w}^1(\tau) = 0$ ,  $\mathbf{w}^2(\tau) > 0$ ,  $u_\tau(x) = \sum_{i=1}^2 v_\tau^i(x^i)$ , for each  $x \in \mathbb{R}_+^2$ ,  $v_\tau^i(x^i)$  is differentiable, and  $\frac{dv_\tau^i(0)}{dx^i} = +\infty$ ,  $i = 1, 2$ .<sup>3</sup> Suppose that  $\hat{\mathbf{b}}$  is a Cournot-Nash equilibrium of  $\Gamma$  and that the allocation  $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))$ , for each  $t \in T$ , is a Walras allocation of  $\mathcal{E}$ . Then, there is a price vector  $\hat{p}$  such that the pair  $(\hat{p}, \hat{\mathbf{x}})$  is a Walras equilibrium of  $\mathcal{E}$ . We have that  $\hat{p} \gg 0$  as  $u_t(\cdot)$  is strongly monotone, for each  $t \in T$ , and  $\hat{\mathbf{x}}(\tau) \gg 0$  as  $\frac{dv_\tau^i(0)}{dx^i} = +\infty$ ,  $i = 1, 2$ . Then, we have that

$$\frac{\hat{p}^1}{\hat{p}^2} = \frac{\mathbf{w}^2(\tau) - \mathbf{x}^2(\tau)}{\mathbf{x}^1(\tau)} = \frac{p^1(\hat{\mathbf{b}})}{p^2(\hat{\mathbf{b}})}, \quad (2)$$

---

<sup>3</sup>For instance, the utility function  $u(x) = \sqrt{x^1} + \sqrt{x^2}$  satisfies these conditions.

as  $\hat{p}\hat{\mathbf{x}}(\tau) = \hat{p}\mathbf{w}(\tau)$  and  $p(\hat{\mathbf{b}})\hat{\mathbf{x}}(\tau) = p(\hat{\mathbf{b}})\mathbf{w}(\tau)$ . Moreover, we have that  $0 < \hat{\mathbf{b}}_{21}(\tau) < \mathbf{w}^2(\tau)$  as  $\hat{\mathbf{x}}(\tau) \gg 0$ , and  $\hat{\mathbf{b}}_{12}(\tau) = 0$  as  $\mathbf{w}^1(\tau) = 0$ . Since

$$p^1(\hat{\mathbf{b}} \setminus b(\tau)) = \alpha \int_{T \setminus \tau} \hat{\mathbf{b}}_{21}(t) d\mu + b_{21}(\tau)\mu(\tau) = \alpha(\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)),$$

and

$$p^2(\hat{\mathbf{b}} \setminus b(\tau)) = \alpha \int_{T \setminus \tau} \hat{\mathbf{b}}_{12}(t) d\mu + b_{12}(\tau)\mu(\tau) = \alpha \int_{T \setminus \tau} \hat{\mathbf{b}}_{12}(t) d\mu = \alpha\bar{\hat{\mathbf{b}}}_{12},$$

with  $\alpha > 0$ , by (1) and Lemma 2 in Sahi and Yao (1989), trader  $\tau$ 's final holding, given the Cournot-Nash equilibrium strategies of the other traders, is

$$\mathbf{x}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))) = (b_{21}(\tau) \frac{\bar{\hat{\mathbf{b}}}_{12}}{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)}, \mathbf{w}^2(\tau) - b_{21}(\tau)).$$

Then, trader  $\tau$ 's payoff as a function of his own strategy, given the Cournot-Nash equilibrium strategies of the other traders, is

$$u_\tau(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau)))) = v_\tau^1(b_{21}(\tau) \frac{\bar{\hat{\mathbf{b}}}_{12}}{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)}) + v_\tau^2(\mathbf{w}^2(\tau) - b_{21}(\tau)).$$

The derivative of this function with respect to trader  $\tau$ 's strategy is

$$\begin{aligned} \frac{\partial u_\tau(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))))}{\partial b_{21}(\tau)} &= \frac{\partial v_\tau^1(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))))}{\partial x^1} \\ &\times \frac{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau) - b_{21}(\tau)\mu(\tau)}{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)} \frac{\bar{\hat{\mathbf{b}}}_{12}}{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)} - \frac{\partial v_\tau^2(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))))}{\partial x^2} \\ &= \frac{\partial v_\tau^1(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))))}{\partial x^1} \times \frac{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau) - b_{21}(\tau)\mu(\tau)}{\bar{\hat{\mathbf{b}}}_{21} \setminus b_{21}(\tau)} \frac{p^2(\hat{\mathbf{b}} \setminus b(\tau))}{p^1(\hat{\mathbf{b}} \setminus b(\tau))} \\ &\quad - \frac{\partial v_\tau^2(\hat{\mathbf{x}}(\tau, b(\tau), p(\hat{\mathbf{b}} \setminus b(\tau))))}{\partial x^2}. \end{aligned}$$

Computing this derivative at  $b(\tau) = \hat{\mathbf{b}}(\tau)$ , we obtain

$$\frac{\partial v_\tau^1(\hat{\mathbf{x}}(\tau))}{\partial x^1} \frac{\bar{\hat{\mathbf{b}}}_{21} - \hat{\mathbf{b}}_{21}(\tau)\mu(\tau)}{\bar{\hat{\mathbf{b}}}_{21}} \frac{p^2(\hat{\mathbf{b}})}{p^1(\hat{\mathbf{b}})} - \frac{\partial v_\tau^2(\hat{\mathbf{x}}(\tau))}{\partial x^2} = 0,$$

as  $\hat{\mathbf{b}}(\tau) = \operatorname{argmax}\{u_\tau(\mathbf{x}(\tau, b(\tau), p(\mathbf{b} \setminus b(\tau)))) : b(\tau) \in \mathbf{B}(\tau)\}$  and  $0 < \hat{\mathbf{b}}_{21}(\tau) < \mathbf{w}^2(\tau)$ . Consequently, we have

$$\frac{\frac{\partial v_\tau^1(\hat{\mathbf{x}}(\tau))}{\partial x^1}}{\frac{\partial v_\tau^2(\hat{\mathbf{x}}(\tau))}{\partial x^2}} = \frac{\tilde{\mathbf{b}}_{21}}{\hat{\mathbf{b}}_{21} - \hat{\mathbf{b}}_{21}(\tau)\mu(\tau)} \frac{\hat{p}^1}{\hat{p}^2} = \frac{\hat{p}^1}{\hat{p}^2}$$

as  $\frac{\hat{p}^1}{\hat{p}^2} = \frac{p^1(\hat{\mathbf{b}})}{p^2(\hat{\mathbf{b}})}$  by (2) and the pair  $(\hat{p}, \hat{\mathbf{x}})$  is a Walras equilibrium of  $\mathcal{E}$ . But then, we must have that  $\hat{\mathbf{b}}_{21}(\tau) = 0$ , a contradiction. Hence,  $\hat{\mathbf{x}}$  is not a Walras allocation of  $\mathcal{E}$ .

This shows that the condition that  $\mathcal{E}$  contains a countably infinite number of atoms is not sufficient to guarantee that any Cournot-Nash allocation of  $\Gamma$  is a Walras allocation of  $\mathcal{E}$ .

## 5 The replication à la Cournot of $\mathcal{E}$

The negative result obtained in the previous section leads us to deal with the question whether a limit result can be instead obtained by replicating the exchange economy  $\mathcal{E}$ .

We address this question by introducing a concept of replication in the original spirit of Cournot (1838). By analogy with the replication proposed by Cournot in a partial equilibrium framework, the concept we propose is obtained in fact by replicating only the atoms of  $\mathcal{E}$ , while making them asymptotically negligible, and without affecting the atomless part.

This partial replication à la Cournot of  $\mathcal{E}$  can be formalized as follows. Let  $\mathcal{E}^n$  be an exchange economy characterized as in Section 2, where each atom is replicated  $n$  times. For each  $t \in T_1$ , let  $tr$  denote the  $r$ -th element of the  $n$ -fold replication of  $t$ . We assume that, for each  $t \in T_1$ ,  $\mathbf{w}(tr) = \mathbf{w}(ts) = \mathbf{w}(t)$ ,  $u_{tr}(\cdot) = u_{ts}(\cdot) = u_t(\cdot)$ ,  $r, s = 1, \dots, n$ , and  $\mu(tr) = \frac{\mu(t)}{n}$ ,  $r = 1, \dots, n$ . Clearly,  $\mathcal{E}^1$  coincides with  $\mathcal{E}$ .

Then, the strategic market game  $\Gamma^n$  associated with  $\mathcal{E}^n$  can be characterized, *mutatis mutandis*, as in Section 2. Clearly,  $\Gamma^1$  coincides with  $\Gamma$ . A strategy selection  $\mathbf{b}$  of  $\Gamma^n$  is said to be atom-type-symmetric if  $\mathbf{b}^n(tr) = \mathbf{b}^n(ts)$ ,  $r, s = 1, \dots, n$ , for each  $t \in T_1$ .

We provide now the definition of an atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$ .

**Definition 4.** A strategy selection  $\hat{\mathbf{b}}$  such that  $\tilde{\mathbf{B}}$  is irreducible is an atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$  if  $\hat{\mathbf{b}}$  is atom-type-symmetric

and if

$$u_{tr}(\mathbf{x}(tr, \hat{\mathbf{b}}(tr), p(\hat{\mathbf{b}}))) \geq u_{tr}(\mathbf{x}(tr, b(tr), p(\hat{\mathbf{b}} \setminus b(tr)))),$$

for each  $b(tr) \in \mathbf{B}(tr)$ ,  $r = 1, \dots, n$ , and for each  $t \in T_1$ ;

$$u_t(\mathbf{x}(t, \hat{\mathbf{b}}(t), p(\hat{\mathbf{b}}))) \geq u_t(\mathbf{x}(t, b(t), p(\hat{\mathbf{b}} \setminus b(t)))),$$

for each  $b(t) \in \mathbf{B}(t)$  and for each  $t \in T_0$ .

## 6 The existence of a $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of $\Gamma^n$

In this section, we introduce the notion of a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$ . Moreover, we state and prove that an equilibrium of this kind exists under Assumptions 1, 2, 3, and 4. This result is needed to show our limit theorem by using the replication à la Cournot introduced in Section 5.

Let  $\delta$  be determined as in Section 3. Also a  $\delta$ -positive strategy correspondence,  $\mathbf{B}^\delta$ , is defined, *mutatis mutandis*, as in Section 3. Notice that  $\mathbf{B}^\delta(tr) = \mathbf{B}^\delta(ts)$ ,  $r, s = 1, \dots, n$ , for each  $t \in T_1$ . We say that a strategy selection  $\mathbf{b}$  is  $\delta$ -positive if  $\mathbf{b}(tr) \in \mathbf{B}^\delta(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and  $\mathbf{b}(t) \in \mathbf{B}^\delta(t)$ , for each  $t \in T_0$ . Then, we say that an atom-type-symmetric Cournot-Nash equilibrium  $\hat{\mathbf{b}}$  of  $\Gamma^n$  is  $\delta$ -positive if  $\hat{\mathbf{b}}$  is a  $\delta$ -positive strategy selection.

The following theorem establishes the existence of a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$ . The proof of the theorem adapts to our context tools and arguments developed by Sahi and Yao (1989) and Busetto et al. (2011) to show the existence of a Cournot-Nash equilibrium of the Shapley window model, respectively in the case of a finite set of traders and in the case of a mixed measure space of traders à la Shitovitz.

**Theorem 2.** *Under Assumptions 1, 2, 3, and 4, there exists a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium  $\hat{\mathbf{b}}$  of  $\Gamma^n$ .*

**Proof.** See the Appendix. ■

## 7 The limit theorem

In this section, we state and prove the limit theorem. It establishes that, given a sequence of atom-type-symmetric Cournot-Nash allocations of  $\Gamma^n$ ,

for  $n = 1, 2, \dots$ , there exists a Walras allocation of  $\mathcal{E}$  with this property: for each trader  $t \in T$ , the value of this Walras allocation at  $t$  is a limit point of the sequence of final holdings of  $t$  associated with the sequence of atom-type-symmetric Cournot-Nash equilibria of  $\Gamma^n$ , for  $n = 1, 2, \dots$

**Theorem 3.** *Under Assumptions 1, 2, 3, and 4, let  $\{\hat{\mathbf{b}}^n\}$  be a sequence of strategy selections of  $\Gamma$  and let  $\{\hat{p}^n\}$  be a sequence of prices such that  $\hat{\mathbf{b}}^n(t) = \hat{\mathbf{b}}^{\Gamma^n}(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ ,  $\hat{\mathbf{b}}^n(t) = \hat{\mathbf{b}}^{\Gamma^n}(t)$ , for each  $t \in T_0$ , and  $\hat{p}^n = p(\hat{\mathbf{b}}^{\Gamma^n})$ , where  $\hat{\mathbf{b}}^{\Gamma^n}$  is a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$ , for  $n = 1, 2, \dots$ . Then,*

(i) *there exists a subsequence  $\{\hat{\mathbf{b}}^{k_n}\}$  of the sequence  $\{\hat{\mathbf{b}}^n\}$ , a subsequence  $\{\hat{p}^{k_n}\}$  of the sequence  $\{\hat{p}^n\}$ , a strategy selection  $\hat{\mathbf{b}}$  of  $\Gamma$ , and a price vector  $\hat{p}$ , with  $\hat{p} \gg 0$ , such that  $\hat{\mathbf{b}}(t)$  is the limit of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for each  $t \in T_1$ ,  $\hat{\mathbf{b}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for each  $t \in T_0$ , the sequence  $\{\bar{\mathbf{B}}^{k_n}\}$  converges to  $\bar{\mathbf{B}}$ , and the sequence  $\{\hat{p}^{k_n}\}$  converges to  $\hat{p}$ ;*

(ii)  *$\hat{\mathbf{x}}(t)$  is the limit of the sequence  $\{\hat{\mathbf{x}}^{k_n}(t)\}$ , for each  $t \in T_1$ , and  $\hat{\mathbf{x}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{x}}^{k_n}(t)\}$ , for each  $t \in T_0$ , where  $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), \hat{p})$  for each  $t \in T$ ,  $\hat{\mathbf{x}}^{k_n}(t) = \mathbf{x}(t, \hat{\mathbf{b}}^{k_n}(t), \hat{p}^{k_n})$ , for each  $t \in T$ , and for  $n = 1, 2, \dots$ ;*

(iii) *the pair  $(\hat{p}, \hat{\mathbf{x}})$  is a Walras equilibrium of  $\mathcal{E}$ .*

**Proof.** (i) Let  $\{\hat{\mathbf{b}}^n\}$  be a sequence of strategy selections of  $\Gamma$  and let  $\{\hat{p}^n\}$  be a sequence of prices such that  $\hat{\mathbf{b}}^n(t) = \hat{\mathbf{b}}^{\Gamma^n}(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ ,  $\hat{\mathbf{b}}^n(t) = \hat{\mathbf{b}}^{\Gamma^n}(t)$ , for each  $t \in T_0$ , and  $\hat{p}^n = p(\hat{\mathbf{b}}^{\Gamma^n})$ , where  $\hat{\mathbf{b}}^{\Gamma^n}$  is a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^n$ , for  $n = 1, 2, \dots$ . The facts that the sequence  $\{\bar{\mathbf{B}}^n\}$  belongs to the compact set  $\{(b_{ij}) \in R_+^{l^2} : b_{ij} \leq \int_T \mathbf{w}^i(t) d\mu, i, j = 1, \dots, l, \sum_{i \notin J} \sum_{j \in J} (b_{ij} + b_{ji}) \geq \int_{\bar{T}_1} \delta d\mu, \text{ for each } J \subseteq \{1, \dots, l\}\}$ , the sequence  $\{\hat{\mathbf{b}}^n(t)\}$  belongs to the compact set  $\mathbf{B}^\delta(t)$ , for each  $t \in T_1$ , and the sequence  $\{\hat{p}^n\}$ , belongs, by Lemma 9 in Sahi and Yao, to a compact set  $P$ , imply that there is a subsequence  $\{\bar{\mathbf{B}}^{k_n}\}$  of the sequence  $\{\bar{\mathbf{B}}^n\}$  which converges to an element of the set  $\{(b_{ij}) \in R_+^{l^2} : b_{ij} \leq \int_T \mathbf{w}^i(t) d\mu, i, j = 1, \dots, l, \sum_{i \notin J} \sum_{j \in J} (b_{ij} + b_{ji}) \geq \int_{\bar{T}_1} \delta d\mu, \text{ for each } J \subseteq \{1, \dots, l\}\}$ , a subsequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$  of the sequence  $\{\hat{\mathbf{b}}^n(t)\}$  which converges to an element of the set  $\mathbf{B}^\delta(t)$ , for each  $t \in T_1$ , and a subsequence  $\{\hat{p}^{k_n}\}$  of the sequence  $\{\hat{p}^n\}$  which converges to an element  $\hat{p}$  of the set  $P$ . Moreover, by Lemma 9 in Sahi and Yao,  $\hat{p} \gg 0$ . Since the sequence  $\{\hat{\mathbf{b}}^{k_n}\}$  satisfies the assumptions of Theorem A in Artstein (1979), there is a function  $\hat{\mathbf{b}}$  such that  $\hat{\mathbf{b}}(t)$  is the limit of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for

each  $t \in T_1$ ,  $\hat{\mathbf{b}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for each  $t \in T_0$ , and such that the sequence  $\{\tilde{\mathbf{B}}^{k_n}\}$  converges to  $\tilde{\mathbf{B}}$ .

(ii) Let  $\hat{\mathbf{x}}(t) = \mathbf{x}(t, \hat{\mathbf{b}}(t), \hat{p})$  for each  $t \in T$ ,  $\hat{\mathbf{x}}^{k_n}(t) = \mathbf{x}(t, \hat{\mathbf{b}}^{k_n}(t), \hat{p}^{k_n})$ , for each  $t \in T$ , and for  $n = 1, 2, \dots$ . Then,  $\hat{\mathbf{x}}(t)$  is the limit of the sequence  $\{\hat{\mathbf{x}}^{k_n}(t)\}$ , for each  $t \in T_1$ , as  $\hat{\mathbf{b}}(t)$  is the limit of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for each  $t \in T_1$ , and the sequence  $\{\hat{p}^{k_n}\}$  converges to  $\hat{p}$ ,  $\hat{\mathbf{x}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{x}}^{k_n}(t)\}$ , for each  $t \in T_0$ , as  $\hat{\mathbf{b}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for each  $t \in T_0$ , and the sequence  $\{\hat{p}^{k_n}\}$  converges to  $\hat{p}$ .

(iii)  $\tilde{\mathbf{B}}^{\Gamma^n} = \tilde{\mathbf{B}}^n$  as  $\tilde{\mathbf{b}}_{ij}^{\Gamma^n} = \sum_{t \in T_1} \sum_{r=1}^n \hat{\mathbf{b}}_{ij}^{\Gamma^n}(tr) \mu(tr) + \int_{t \in T_0} \hat{\mathbf{b}}_{ij}^{\Gamma^n}(t) d\mu = \sum_{t \in T_1} n \hat{\mathbf{b}}_{ij}^n(t) \frac{\mu(t)}{n} + \int_{t \in T_0} \hat{\mathbf{b}}_{ij}^n(t) d\mu = \sum_{t \in T_1} \hat{\mathbf{b}}_{ij}^n(t) \mu(t) + \int_{t \in T_0} \hat{\mathbf{b}}_{ij}^n(t) d\mu = \tilde{\mathbf{b}}_{ij}^n$ ,  $i, j = 1, \dots, l$ , for  $n = 1, 2, \dots$ . Then,  $\hat{p}^n = p(\hat{\mathbf{b}}^n)$  as  $\hat{p}^n$  and  $\hat{\mathbf{b}}^n$  satisfy (1), for  $n = 1, 2, \dots$ . But then, by continuity,  $\hat{p}$  and  $\hat{\mathbf{b}}$  must satisfy (1). Therefore, Lemma 1 in Sahi and Yao implies that  $\tilde{\mathbf{B}}$  is completely reducible. Moreover,  $\hat{\mathbf{b}}(t) \in \mathbf{B}^\delta(t)$  since  $\hat{\mathbf{b}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{b}}^{k_n}(t)\}$ , for all  $t \in T$ . Then,  $\hat{\mathbf{b}}$  is  $\delta$ -positive. But then, by Remark 3 in Sahi and Yao,  $\tilde{\mathbf{B}}$  must be irreducible. Consider the pair  $(\hat{p}, \hat{\mathbf{x}})$ . It is straightforward to show that the assignment  $\hat{\mathbf{x}}$  is an allocation as  $\hat{p}$  and  $\hat{\mathbf{b}}$  satisfy (1) and that  $\hat{\mathbf{x}}(t) \in \{x \in R_+^l : \hat{p}x = \hat{p}\mathbf{w}(t)\}$ , for all  $t \in T$ . Suppose that  $(\hat{p}, \hat{\mathbf{x}})$  is not a Walras equilibrium of  $\mathcal{E}$ . Then, there exists a trader  $\tau \in T$  and a commodity bundle  $\tilde{x} \in \{x \in R_+^l : \hat{p}x = \hat{p}\mathbf{w}(\tau)\}$  such that  $u_\tau(\tilde{x}) > u_\tau(\hat{\mathbf{x}}(\tau))$ . By Lemma 5 in Codognato and Ghosal (2000), there exist  $\tilde{\lambda}^j \geq 0$ ,  $\sum_{j=1}^l \tilde{\lambda}^j = 1$ , such that

$$\tilde{x}^j = \tilde{\lambda}^j \frac{\sum_{i=1}^l \hat{p}^i \mathbf{w}^i(\tau)}{\hat{p}^j}, \quad j = 1, \dots, l.$$

Let  $\tilde{b}_{ij} = \mathbf{w}^i(\tau) \tilde{\lambda}^j$ ,  $i, j = 1, \dots, l$ . Then, it is straightforward to verify that

$$\tilde{x}^j = \mathbf{w}^j(\tau) - \sum_{i=1}^l \tilde{b}_{ji} + \sum_{i=1}^l \tilde{b}_{ij} \frac{\hat{p}^i}{\hat{p}^j},$$

for each  $j = 1, \dots, l$ . Consider the following cases.

**Case 1.**  $\tau \in T_1$ . Let  $\{h_n\}$  denote a sequence such that  $h_n = k_n$ , if  $k_1 > 1$ ,  $h_n = k_{n+1}$ , otherwise, for  $n = 1, 2, \dots$ . Let  $\rho$  denote the  $k_1$ -th element of the  $n$ -fold replication of  $\mathcal{E}$  and let  $\tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$  be the aggregate matrix corresponding to the strategy selection  $\hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$ , where  $\tilde{b}(\tau\rho) = \tilde{b}$ , for  $n = 1, 2, \dots$ . Let  $\Delta \tilde{\mathbf{B}}^{\Gamma^{h_n}}$ ,  $\Delta \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$ , and  $\Delta \tilde{\mathbf{B}}^{\Gamma^{h_n}}$  denote the diagonal matrices of row sums of, respectively,  $\tilde{\mathbf{B}}^{\Gamma^{h_n}}$ ,  $\tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$ , and  $\tilde{\mathbf{B}}^{\Gamma^{h_n}}$ , for  $n = 1, 2, \dots$

Moreover, let  $q^{\Gamma^{h_n}}$ ,  $q_{\tau\rho}^{\Gamma^{h_n}}$ , and  $q^{h_n}$  denote the vectors of the cofactors of the first column of, respectively,  $\Delta\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}}$ ,  $\Delta\tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho) - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$ , and  $\Delta\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}}$ , for  $n = 1, 2, \dots$ . Clearly,  $q^{\Gamma^{h_n}} = q^{h_n}$  as  $\tilde{\mathbf{B}}^{\Gamma^{h_n}} = \tilde{\mathbf{B}}^{\Gamma^{h_n}}$ , for  $n = 1, 2, \dots$ . Let  $\Delta\tilde{\mathbf{B}}$  be the diagonal matrix of row sums of  $\tilde{\mathbf{B}}$  and  $q$  be the cofactors of the first column of  $\Delta\tilde{\mathbf{B}} - \tilde{\mathbf{B}}$ . The sequences  $\{q^{\Gamma^{h_n}}\}$  and  $\{q^{h_n}\}$  converge to  $q$  as the sequence  $\tilde{\mathbf{B}}^{\Gamma^{h_n}}$  converges to  $\tilde{\mathbf{B}}$  and  $q^{\Gamma^{h_n}} = q^{h_n}$ , for  $n = 1, 2, \dots$ . Let  $\bar{w} = \max\{\mathbf{w}^1(\tau), \dots, \mathbf{w}^l(\tau)\}$ . Consider the matrix  $\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)$ , for  $n = 1, 2, \dots$ . Then,  $\hat{\mathbf{b}}_{ij}^{\Gamma^{h_n}} - \tilde{\mathbf{b}}_{ij}^{\Gamma^{h_n}} \setminus \tilde{b}_{ij}(\tau\rho) = (\frac{1}{n}\hat{\mathbf{b}}_{ij}^{\Gamma^{h_n}}(\tau\rho) - \frac{1}{n}\tilde{b}_{ij}(\tau\rho))$ ,  $i, j = 1, \dots, l$ , for  $n = 1, 2, \dots$ . But then, the sequence of Euclidean distances  $\{\|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)\|\}$  converges to 0 as  $|\frac{1}{n}\hat{\mathbf{b}}_{ij}^{\Gamma^{h_n}}(\tau\rho) - \frac{1}{n}\tilde{b}_{ij}(\tau\rho)| = \frac{1}{n}|\hat{\mathbf{b}}_{ij}^{\Gamma^{h_n}}(\tau\rho) - \tilde{b}_{ij}(\tau\rho)| \leq \frac{1}{n}\bar{w}$ ,  $i, j = 1, \dots, l$ ,  $n = 1, 2, \dots$ . The sequence  $\{\tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)\}$  converges to  $\tilde{\mathbf{B}}$  as, by the triangle inequality,  $\|\tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho) - \tilde{\mathbf{B}}\| \leq \|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)\| + \|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}\| = \|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)\| + \|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}\|$ , for  $n = 1, 2, \dots$ , and the sequences  $\{\|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)\|\}$  and  $\{\|\tilde{\mathbf{B}}^{\Gamma^{h_n}} - \tilde{\mathbf{B}}\|\}$  converge to 0. Then, the sequence  $\{q_{\tau\rho}^{\Gamma^{h_n}}\}$  converges to  $q$ .  $u_{\tau\rho}(\mathbf{x}(\tau\rho, \hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau\rho), p(\hat{\mathbf{b}}^{\Gamma^{h_n}}))) \geq u_{\tau\rho}(\mathbf{x}(\tau\rho, \tilde{b}(\tau\rho), p(\hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho))))$  as  $\hat{\mathbf{b}}^{\Gamma^{h_n}}$  is a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^{h_n}$ , for  $n = 1, 2, \dots$ . Let  $\hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau)$  be a strategy selection obtained by replacing  $\hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau)$  in  $\hat{\mathbf{b}}^{\Gamma^{h_n}}$  with  $\tilde{b}$ , for  $n = 1, 2, \dots$ . Then,  $u_{\tau}(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau), q^{\Gamma^{h_n}})) \geq u_{\tau}(\mathbf{x}(\tau, \tilde{b}(\tau), q_{\tau\rho}^{\Gamma^{h_n}}))$  as  $\hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau\rho) = \hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau)$ ,  $p(\hat{\mathbf{b}}^{\Gamma^{h_n}}) = \alpha_{h_n} q^{\Gamma^{h_n}}$ , with  $\alpha_{h_n} > 0$ , by Lemma 2 in Sahi and Yao,  $\hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho) = \hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau)$ , and  $p(\hat{\mathbf{b}}^{\Gamma^{h_n}} \setminus \tilde{b}(\tau\rho)) = \beta_{h_n} q_{\tau\rho}^{\Gamma^{h_n}}$ , with  $\beta_{h_n} > 0$ , by Lemma 2 in Sahi and Yao, for  $n = 1, 2, \dots$ . But then,  $u_{\tau}(\hat{\mathbf{x}}(\tau)) \geq u_{\tau}(\tilde{x})$ , by Assumption 2, as the sequence  $\{\hat{\mathbf{b}}^{\Gamma^{h_n}}(\tau)\}$  converges to  $\hat{\mathbf{b}}(\tau)$ , the sequence  $\{q^{\Gamma^{h_n}}\}$  converges to  $q$ , the sequence  $\{q_{\tau\rho}^{\Gamma^{h_n}}\}$  converges to  $q$ , and  $\hat{p} = \theta q$ , with  $\theta > 0$ , by Lemma 2 in Sahi and Yao, a contradiction.

**Case 2.**  $\tau \in T_0$ . Let  $\{\hat{\mathbf{b}}^{h_{k_n}}(\tau)\}$  be a subsequence of the sequence  $\{\hat{\mathbf{b}}^{k_n}(\tau)\}$  which converges to  $\hat{\mathbf{b}}(\tau)$ . Moreover, let  $\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}} \setminus \tilde{b}(\tau)$  be a strategy selection obtained by replacing  $\hat{\mathbf{b}}^{h_{k_n}}(\tau)$  in  $\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}}$  with  $\tilde{b}$ , for  $n = 1, 2, \dots$ .  $u_{\tau}(\mathbf{x}(\tau, \hat{\mathbf{b}}^{\Gamma^{h_{k_n}}}(\tau), p(\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}})) \geq u_{\tau}(\mathbf{x}(\tau, \tilde{b}(\tau), p(\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}} \setminus \tilde{b}(\tau))))$  as  $\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}}$  is a  $\delta$ -positive atom-type-symmetric Cournot-Nash equilibrium of  $\Gamma^{h_{k_n}}$ , for  $n = 1, 2, \dots$ . Let  $\hat{\mathbf{b}}^{h_{k_n}} \setminus \tilde{b}(\tau)$  be a strategy selection obtained by replacing

$\hat{\mathbf{b}}^{h_{k_n}}(\tau)$  in  $\hat{\mathbf{b}}^{h_{k_n}}$  with  $\tilde{b}$ , for  $n = 1, 2, \dots$ . Then,  $u_\tau(\mathbf{x}(\tau, \hat{\mathbf{b}}^{h_{k_n}}(\tau), \hat{p}^{h_{k_n}})) \geq u_\tau(\mathbf{x}(\tau, \tilde{b}(\tau), \hat{p}^{h_{k_n}}))$  as  $\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}}(\tau) = \hat{\mathbf{b}}^{h_{k_n}}(\tau)$ ,  $p(\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}}) = \hat{p}^{h_{k_n}}$ ,  $\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}} \setminus \tilde{b}(\tau) = \hat{\mathbf{b}}^{h_{k_n}} \setminus \tilde{b}(\tau)$ , and  $p(\hat{\mathbf{b}}^{\Gamma^{h_{k_n}}} \setminus \tilde{b}(\tau)) = \hat{p}^{h_{k_n}}$ . But then,  $u_\tau(\hat{\mathbf{x}}(\tau)) \geq u_\tau(\tilde{x})$ , by Assumption 2, as the sequence  $\{\hat{\mathbf{b}}^{h_{k_n}}(\tau)\}$  converges to  $\hat{\mathbf{b}}(\tau)$  and the sequence  $\{\hat{p}^{h_{k_n}}\}$  converges to  $\hat{p}$ , a contradiction.

Hence, the pair  $(\hat{p}, \hat{\mathbf{x}})$  is a Walras equilibrium of  $\mathcal{E}$ .  $\blacksquare$

## 8 Appendix

**Proof of Theorem 1.** Busetto et al. (2011) showed that, under Assumptions 1, 2, 3, and 4, there exists a Cournot-Nash equilibrium of  $\Gamma$ ,  $\hat{\mathbf{b}}$ , such that, for each  $t \in T$ ,  $\hat{\mathbf{b}}(t) \in \mathbf{B}^\delta(t)$ . This implies that  $\hat{\mathbf{b}}$  is a  $\delta$ -positive Cournot-Nash equilibrium of  $\Gamma$ .  $\blacksquare$

**Proof of Theorem 2.** Let us introduce a slightly perturbed version of the game  $\Gamma^n$ , denoted by  $\Gamma^n(\epsilon)$ . Given  $\epsilon > 0$  and a strategy selection  $\mathbf{b}$ , we define the aggregate matrix  $\bar{\mathbf{B}}_\epsilon$  to be the matrix such that  $\bar{\mathbf{b}}_{\epsilon ij} = (\mathbf{b}_{ij} + \epsilon)$ ,  $i, j = 1, \dots, l$ . Clearly, the matrix  $\bar{\mathbf{B}}_\epsilon$  is irreducible. The interpretation is that an outside agency places fixed bids of  $\epsilon$  for each pair of commodities  $(i, j)$ . Given  $\epsilon > 0$ , we denote by  $p^\epsilon(\mathbf{b})$  the function which associates, with each strategy selection  $\mathbf{b}$ , the unique, up to a scalar multiple, price vector which satisfies

$$\sum_{i=1}^l p^i (\bar{\mathbf{b}}_{ij} + \epsilon) = p^j (\sum_{i=1}^l \bar{\mathbf{b}}_{ji} + \epsilon), \quad j = 1, \dots, l.$$

Then, let us introduce the following notion of equilibrium for  $\Gamma^n(\epsilon)$ .

**Definition 5.** Given  $\epsilon > 0$ , a strategy selection  $\hat{\mathbf{b}}^\epsilon$  is an atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$  if  $\hat{\mathbf{b}}^\epsilon$  is atom-type-symmetric and

$$u_{tr}(\mathbf{x}(tr, \hat{\mathbf{b}}^\epsilon(tr), p^\epsilon(\hat{\mathbf{b}}^\epsilon))) \geq u_{tr}(tr, b(tr), p^\epsilon(\hat{\mathbf{b}}^\epsilon \setminus b(tr))),$$

for each  $b(tr) \in \mathbf{B}(tr)$ ,  $r = 1, \dots, n$ , and for each  $t \in T_1$ ;

$$u_t(\mathbf{x}(t, \hat{\mathbf{b}}^\epsilon(t), p^\epsilon(\hat{\mathbf{b}}^\epsilon))) \geq u_t(t, b(t), p^\epsilon(\hat{\mathbf{b}}^\epsilon \setminus b(t))),$$

for each  $b(t) \in \mathbf{B}(t)$  and for each  $t \in T_0$ .

Moreover, we say that an atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium  $\hat{\mathbf{b}}^\epsilon$  of  $\Gamma^n(\epsilon)$  is  $\delta$ -positive if  $\hat{\mathbf{b}}^\epsilon$  is a  $\delta$ -positive strategy selection.

To show Theorem 2, we first need to prove the existence of a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$ . To do so, we apply, as in Busetto et al. (2011), the Kakutani-Fan-Glicksberg Theorem (see Theorem 17.55 in Aliprantis and Border (2006), p. 583).

We neglect, as usual, the distinction between integrable functions and equivalence classes of such functions and denote by  $L_1(\mu, R^{l^2})$  the set of integrable functions taking values in  $R^{l^2}$ , by  $L_1(\mu, \mathbf{B}(\cdot))$  the set of strategy selections, and by  $L_1(\mu, \mathbf{B}^*(\cdot))$  the set of atom-type-symmetric strategy selections. Note that the locally convex Hausdorff space we shall be working in is  $L_1(\mu, R^{l^2})$ , endowed with its weak topology.

The proof of existence of a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$  is articulated in three lemmas.

The first lemma establishes the properties of  $L_1(\mu, \mathbf{B}^*(\cdot))$  required to apply the Kakutani-Fan-Glicksberg Theorem.

**Lemma 1.** *The set  $L_1(\mu, \mathbf{B}^*(\cdot))$  is nonempty, convex and weakly compact.*

**Proof.**  $L_1(\mu, \mathbf{B}^*(\cdot))$  is nonempty, convex and it has a weakly compact closure by the same argument used by Busetto et al. (2011) to prove their Lemma 1. Now, let  $\{\mathbf{b}^m\}$  be a convergent sequence of  $L_1(\mu, \mathbf{B}^*(\cdot))$ . Since  $L_1(\mu, R^{l^2})$  is complete,  $\{\mathbf{b}^m\}$  converges in the mean to an integrable function  $\mathbf{b}$ . But then, there exists a subsequence  $\{\mathbf{b}^{k_m}\}$  of  $\{\mathbf{b}^m\}$  such that  $\mathbf{b}^{k_m}(tr)$  converges to  $\mathbf{b}(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and  $\mathbf{b}^{k_m}(t)$  converges to  $\mathbf{b}(t)$ , for each  $t \in T_0$  (see Theorem 25.5 in Aliprantis and Burkinshaw (1998), p. 203). The compactness of  $\mathbf{B}(t)$ , for each  $t \in T$ , and the fact that  $\mathbf{b}^{k_m}(tr) = \mathbf{b}^{k_m}(ts)$ ,  $r, s = 1, \dots, n$ , for each  $t \in T_1$ , implies that  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$ . Hence  $L_1(\mu, \mathbf{B}^*(\cdot))$  is norm closed and, since it is also convex, it is weakly closed (see Corollary 4 in Diestel (1984), p. 12). ■

Now, given  $\epsilon > 0$ , let  $\alpha_{tr}^\epsilon : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow \mathbf{B}(tr)$  be a correspondence such that  $\alpha_{tr}^\epsilon(\mathbf{b}) = \operatorname{argmax}\{u_{tr}(\mathbf{x}(t, b(tr)), p^\epsilon(\mathbf{b} \setminus b(tr))) : b(tr) \in \mathbf{B}(tr)\}$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and let  $\alpha_t^\epsilon : L_1(\mu, \mathbf{B}(\cdot)) \rightarrow \mathbf{B}(t)$  be a correspondence such that  $\alpha_t^\epsilon(\mathbf{b}) = \operatorname{argmax}\{u_t(\mathbf{x}(t, b(t)), p^\epsilon(\mathbf{b} \setminus b(t))) : b(t) \in \mathbf{B}(t)\}$ , for each  $t \in T_0$ . Let  $\alpha^\epsilon : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow L_1(\mu, \mathbf{B}(\cdot))$  be a correspondence such that  $\alpha^\epsilon(\mathbf{b}) = \{\mathbf{b} \in L_1(\mu, \mathbf{B}(\cdot)) : \mathbf{b}(tr) \in \alpha_{tr}^\epsilon(\mathbf{b}), r = 1, \dots, n, \text{ for each } t \in T_1, \text{ and } \mathbf{b}(t) \in \alpha_t^\epsilon(\mathbf{b}), \text{ for each } t \in T_0\}$ .

The second lemma provides us with the properties of the correspondence  $\alpha^\epsilon$ . The proof is obtained by readapting to our context the arguments used to show Lemma 2 in Busetto et al. (2011), and we omit it here.

**Lemma 2.** *Given  $\epsilon > 0$ , the correspondence  $\alpha^\epsilon : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow L_1(\mu, \mathbf{B}(\cdot))$*

is such that the set  $\alpha^\epsilon(\mathbf{b})$  is nonempty and convex, for all  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$ , and it has a weakly closed graph.

Now, given  $\epsilon > 0$ , let  $\alpha_{tr}^{\epsilon\delta} : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow \mathbf{B}(tr)$  be a correspondence such that  $\alpha_{tr}^{\epsilon\delta}(\mathbf{b}) = \alpha_{tr}^\epsilon(\mathbf{b}) \cap \mathbf{B}^\delta(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and let  $\alpha_t^\epsilon : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow \mathbf{B}(t)$  be a correspondence such that  $\alpha_t^{\epsilon\delta}(\mathbf{b}) = \alpha_t^\epsilon(\mathbf{b}) \cap \mathbf{B}^\delta(t)$ , for each  $t \in T_0$ . Let  $\alpha^{\epsilon\delta} : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow L_1(\mu, \mathbf{B}^*(\cdot))$  be a correspondence such that  $\alpha^{\epsilon\delta}(\mathbf{b}) = \{\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot)) : \mathbf{b}(tr) \in \alpha_{tr}^{\epsilon\delta}(\mathbf{b}), r = 1, \dots, n, \text{ for each } t \in T_1, \text{ and } \mathbf{b}(t) \in \alpha_t^{\epsilon\delta}(\mathbf{b}), \text{ for each } t \in T_0\}$ . Moreover, let  $\alpha^{\epsilon\delta*} : L_1(\mu, \mathbf{B}^*(\cdot)) \rightarrow L_1(\mu, \mathbf{B}^*(\cdot))$  be a correspondence such that  $\alpha^{\epsilon\delta*}(\mathbf{b}) = \alpha^{\epsilon\delta}(\mathbf{b}) \cap L_1(\mu, \mathbf{B}^*(\cdot))$ .

We are ready to prove the existence of a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$ .

**Lemma 3.** *Given  $\epsilon > 0$ , there exists a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium  $\hat{\mathbf{b}}^\epsilon$  of  $\Gamma^n(\epsilon)$ .*

**Proof.** Let  $\epsilon > 0$  be given. By Lemma 6 in Sahi and Yao (1989), we know that, for each  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$ ,  $\alpha_{tr}^{\epsilon\delta}(\mathbf{b})$  is nonempty,  $r = 1, \dots, n$ , for each  $t \in \bar{T}_1$ . Moreover, for each  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$  and for each  $t \in T_1$ , there exists  $\tilde{b} \in \mathbf{B}(t)$  such that  $\tilde{b} \in \alpha_{tr}^{\epsilon\delta}(\mathbf{b})$ ,  $r = 1, \dots, n$  as  $\mathbf{b}$  is an atom-type-symmetric strategy profile. But then, by the same argument of Lemma 2 in Busetto et al. (2011),  $\alpha^{\epsilon\delta*}(\mathbf{b})$  is nonempty. The convexity of  $\alpha^{\epsilon\delta}(\mathbf{b})$ , for each  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$ , is a straightforward consequence of the convexity of  $\alpha_{tr}^\epsilon(\mathbf{b})$  and  $\mathbf{B}^\delta(t)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and of  $\alpha_t^\epsilon(\mathbf{b})$  and  $\mathbf{B}^\delta(t)$ , for each  $t \in T_0$ . But then,  $\alpha^{\epsilon\delta*}$  is convex valued as  $L_1(\mu, \mathbf{B}^*(\cdot))$  is convex.  $\alpha_{tr}^{\epsilon\delta}$  is upper hemicontinuous and compact valued,  $r = 1, \dots, n$ , for each  $t \in T_1$ , as it is the intersection of the correspondence  $\alpha_{tr}^\epsilon$ , which is upper hemicontinuous and compact valued by Lemma 2 in Busetto et al. (2011), and the continuous and compact valued correspondence which assigns to each strategy selection  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$  the strategy set  $\mathbf{B}^\delta(tr)$  (see Theorem 17.25 in Aliprantis and Border (2006), p. 567). Moreover,  $\alpha_t^{\epsilon\delta}$  is upper hemicontinuous and compact valued, for each  $t \in T_0$ , using the same argument. Therefore,  $\alpha^{\epsilon\delta}$  has a weakly closed graph, by the same argument used in the proof of Lemma 2. Finally,  $\alpha^{\epsilon\delta*}$  has a weakly closed graph as it is the intersection of the correspondence  $\alpha^{\epsilon\delta}$  and the continuous correspondence which assigns to each strategy selection  $\mathbf{b} \in L_1(\mu, \mathbf{B}^*(\cdot))$  the weakly closed set  $L_1(\mu, \mathbf{B}^*(\cdot))$  which, by the Closed Graph Theorem (see Theorem 17.11 in Aliprantis and Border (2006), p. 561), has a weakly closed graph (see Theorem 17.25 in Aliprantis and Border (2006), p. 567). But then, by the Kakutani-Fan-Glicksberg Theorem (see Theorem 17.55 in Aliprantis and Border (2006), p.

583), there exists a fixed point  $\hat{\mathbf{b}}^\epsilon$  of the correspondence  $\alpha^{\epsilon\delta^*}$  and hence a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$ . ■

To complete the proof of Theorem 2, we have to show that there exists the limit of a sequence of  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibria and that this limit is a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium of  $\Gamma^n$ . Following Busetto et al. (2011), in this part of the proof we essentially refer to Lemma 9 in Sahi and Yao (1989) and a generalization of the Fatou's lemma in several dimensions provided by Artstein (1979).

Then, let  $\epsilon_m = \frac{1}{m}$ ,  $m = 1, 2, \dots$ . By Lemma 3, for each  $m = 1, 2, \dots$ , there is a  $\delta$ -positive atom-type-symmetric  $\epsilon$ -Cournot-Nash equilibrium  $\hat{\mathbf{b}}^{\epsilon_m}$ . The facts that the sequence  $\{\bar{\mathbf{B}}^{\epsilon_m}\}$  belongs to the compact set  $\{(b_{ij}) \in R_+^{l^2} : b_{ij} \leq n \int_{T_1} \mathbf{w}^i(t) d\mu + \int_{T_0} \mathbf{w}^i(t) d\mu, i, j = 1, \dots, l, \sum_{i \notin J} \sum_{j \in J} (b_{ij} + b_{ji}) \geq n \int_{T_1} \delta d\mu, \text{ for each } J \subseteq \{1, \dots, l\}\}$ , the sequence  $\{\hat{\mathbf{b}}^{\epsilon_m}(tr)\}$  belongs to the compact set  $\mathbf{B}^\delta(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and the sequence  $\{\hat{p}^{\epsilon_m}\}$ , where  $\hat{p}^{\epsilon_m} = p^{\epsilon_m}(\hat{\mathbf{b}}^{\epsilon_m})$ , for each  $m = 1, 2, \dots$ , belongs, by Lemma 9 in Sahi and Yao (1989), to a compact set  $P$ , imply that there is a subsequence  $\{\bar{\mathbf{B}}^{\epsilon_{k_m}}\}$  of the sequence  $\{\bar{\mathbf{B}}^{\epsilon_m}\}$  which converges to an element of the set  $\{(b_{ij}) \in R_+^{l^2} : b_{ij} \leq n \int_{T_1} \mathbf{w}^i(t) d\mu + \int_{T_0} \mathbf{w}^i(t) d\mu, i, j = 1, \dots, l, \sum_{i \notin J} \sum_{j \in J} (b_{ij} + b_{ji}) \geq n \int_{T_1} \delta d\mu, \text{ for each } J \subseteq \{1, \dots, l\}\}$ , a subsequence  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}(tr)\}$  of the sequence  $\{\hat{\mathbf{b}}^{\epsilon_m}(tr)\}$  which converges to an element of the set  $\mathbf{B}^\delta(tr)$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ , and a subsequence  $\{\hat{p}^{\epsilon_{k_m}}\}$  of the sequence  $\{\hat{p}^{\epsilon_m}\}$  which converges to an element of the set  $P$ . Since the sequence  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}\}$  satisfies the assumptions of Theorem A in Artstein (1979), there is a function  $\hat{\mathbf{b}}$  such that  $\hat{\mathbf{b}}(tr)$  is the limit of the sequence  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}(tr)\}$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ ,  $\hat{\mathbf{b}}(t)$  is a limit point of the sequence  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}(t)\}$ , for each  $t \in T_0$ , and such that the sequence  $\{\bar{\mathbf{B}}^{\epsilon_{k_m}}\}$  converges to  $\bar{\mathbf{B}}$ . Then,  $\hat{\mathbf{b}}(tr) = \hat{\mathbf{b}}(ts)$  as  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}(tr)\} = \{\hat{\mathbf{b}}^{\epsilon_{k_m}}(ts)\}$ ,  $r, s = 1, \dots, n$ , for each  $t \in T_1$ , and  $\hat{\mathbf{b}}(tr)$  is the limit of the sequence  $\{\hat{\mathbf{b}}^{\epsilon_{k_m}}(tr)\}$ ,  $r = 1, \dots, n$ , for each  $t \in T_1$ . Hence, it can be proved, by the same argument used by Busetto et al. (2011) to show their existence theorem, that  $\hat{\mathbf{b}}$  is an atom-type-symmetric  $\delta$ -positive Cournot-Nash equilibrium of  $\Gamma^n(\epsilon)$ . ■

## References

- [1] Aliprantis C.D., Border K.C. (2006), *Infinite dimensional analysis*, Springer, New York.
- [2] Aliprantis C.D., Burkinshaw O. (1998), *Principles of real analysis*, Academic Press, Boston.
- [3] Amir R., Sahi S., Shubik M., Yao S. (1990), “A strategic market game with complete markets,” *Journal of Economic Theory* **51**, 126-143.
- [4] Artstein Z. (1979), “A note on Fatou’s lemma in several dimensions,” *Journal of Mathematical Economics* **6**, 277-282.
- [5] Aumann R.J. (1964), “Markets with a continuum of traders,” *Econometrica* **32**, 39-50.
- [6] Busetto F., Codognato G., Ghosal S. (2011), “Noncooperative oligopoly in markets with a continuum of traders,” *Games and Economic Behavior*, **72**, 38-45.
- [7] Codognato G., Gabszewicz J.J. (1993), “Cournot-Walras equilibria in markets with a continuum of traders,” *Economic Theory* **3**, 453-464.
- [8] Codognato G., Ghosal S. (2000), “Cournot-Nash equilibria in limit exchange economies with complete markets and consistent prices,” *Journal of Mathematical Economics* **34**, 39-53.
- [9] Cournot A. (1838), *Recherches sur les principes mathématiques de la théorie des richesses*, Hachette, Paris.
- [10] Debreu G., Scarf H. (1963), “A limit theorem on the core of an economy,” *International Economic Review* **4**, 235-246.
- [11] d’Aspremont C., Dos Santos Ferreira R., Gérard-Varet L.-A. (1997), “General equilibrium concepts under imperfect competition: a Cournotian approach,” *Journal of Economic Theory* **73**, 199-230.
- [12] Dierker H., Grodal B. (1986), “Nonexistence of Cournot-Walras equilibrium in a general equilibrium model with two oligopolists,” in Hildenbrand W., Mas-Colell A. (eds), *Contributions to mathematical economics in honor of Gérard Debreu*, North-Holland, Amsterdam.

- [13] Diestel J. (1984), *Sequences and series in Banach spaces*, Springer, New York.
- [14] Dubey P., Shubik M. (1978), "The noncooperative equilibria of a closed trading economy with market supply and bidding strategies," *Journal of Economic Theory* **17**, 1-20.
- [15] Frank C.R. (1965), "Entry in a Cournot market," *Review of Economic Studies* **32**, 245-250.
- [16] Gabszewicz J.J., Michel P. (1997), "Oligopoly equilibrium in exchange economies," in Eaton B.C., Harris R. G. (eds), *Trade, technology and economics. Essays in honour of Richard G. Lipsey*, Edward Elgar, Cheltenham.
- [17] Gabszewicz J.J., Vial J.-P. (1972), "Oligopoly 'à la Cournot-Walras' in a general equilibrium analysis," *Journal of Economic Theory* **4**, 381-400.
- [18] Okuno M., Postlewaite A., Roberts J. (1980), "Oligopoly and competition in large markets," *American Economic Review* **70**, 22-31.
- [19] Mas-Colell A. (1982), "The Cournotian foundations of Walrasian equilibrium theory," in Hildenbrand W. (ed), *Advances in economic theory*, Cambridge University Press, Cambridge.
- [20] Mas-Colell A. (1983), "Walrasian equilibria as limits of noncooperative equilibria. Part I: mixed strategies," *Journal of Economic Theory* **30**, 153-170.
- [21] Novshek W. (1980), "Cournot equilibrium with free entry," *Review of Economic Studies* **47**, 473-486.
- [22] Novshek W., Sonnenschein H. (1983), "Walrasian equilibria as limits of noncooperative equilibria. Part I: pure strategies," *Journal of Economic Theory* **30**, 171-187.
- [23] Novshek W., Sonnenschein H. (1987), "General equilibrium with free entry: a synthetic approach to the theory of perfect competition," *Journal of Economic Literature* **25**, 1281-1306.
- [24] Peck J., Shell K., Spear S.E. (1992), "The market game: existence and structure of equilibrium," *Journal of Mathematical Economics* **21**, 271-299.

- [25] Roberts K. (1980), "The limit points of monopolistic competition," *Journal of Economic Theory* **22**, 256-278.
- [26] Ruffin R.J. (1971), "Cournot oligopoly and competitive behavior," *Review of Economic Studies* **38**, 493-502.
- [27] Sahi S., Yao S. (1989), "The noncooperative equilibria of a trading economy with complete markets and consistent prices," *Journal of Mathematical Economics* **18**, 325-346.
- [28] Shapley L.S., Shubik M. (1977), "Trade using one commodity as a means of payment," *Journal of Political Economy* **85**, 937-968.
- [29] Shitovitz B. (1973), "Oligopoly in markets with a continuum of traders," *Econometrica* **41**, 467-501.
- [30] Shitovitz B. (1997), "A comparison between the core and the monopoly solutions in a mixed exchange economy," *Economic Theory* **10**, 559-563.