

Supplemental Table 1. Comparison between patients with and without available clinical outcome after stroke.

Variables	Total N=259	Available outcome N=143	Unavailable outcome N=116	p
Age, mean (±SD)	69.7±15.5	71.1±14.3	67.9±16.8	0.101
Sex, male	131 (49)	65 (52)	66 (46)	0.048
Baseline NIHSS, median (IQR)	10 (6-17)	11 (6-18)	9 (4-16)	0.033
OTT, mean (±SD)	176.32±62.4	186.6±75.4	161.9±32.6	0.002
Glucose, mean, mg/dl (±SD)	133.1±51.6	132.1±49.1	134.3±54.8	0.729
Systolic BP, mean (±SD)	152.2±26.2	148.1±23.5	156.2±28.2	0.039
Diastolic BP, mean (±SD)	81.9±15.2	80.4±14.2	83.5±15.7	0.171
Diabetes	57 (21)	26 (17)	31 (26)	0.082
Hypertension	185 (69)	102 (68)	87 (70)	0.759
Atrial fibrillation	79 (29)	50 (33)	29 (24)	0.109
Hyperlipidemia	123 (46)	68 (45)	55 (46)	0.885
Smoke	73 (27)	42 (28)	31 (26)	0.721

SD=Standard deviation; NIHSS=National Institutes of Health Stroke Scale; IQR=Interquartile Range; OTT=Onset To Treatment time; BP=Blood Pressure. Data are numbers (%) unless otherwise stated.