



Vidal-Petiot, E., Greenlaw, N., Ford, I. , Ferrari, R., Fox, K. M., Tardif, J.-C., Tendera, M., Parkhomenko, A., Bhatt, D. L. and Steg, P. G. (2018) Relationships between components of blood pressure and cardiovascular events in patients with stable coronary artery disease and hypertension. *Hypertension*, 71, pp. 168-176. (doi:[10.1161/HYPERTENSIONAHA.117.10204](https://doi.org/10.1161/HYPERTENSIONAHA.117.10204))

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/151238/>

Deposited on: 08 November 2018

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

**RELATIONSHIPS BETWEEN COMPONENTS OF BLOOD PRESSURE AND
CARDIOVASCULAR EVENTS IN PATIENTS WITH STABLE CORONARY
ARTERY DISEASE AND HYPERTENSION**

Emmanuelle Vidal-Petiot, MD, PhD^{1,2}, Nicola Greenlaw, MSc³, Ian Ford, PhD³, Roberto Ferrari, MD⁴, Kim M. Fox, MD⁵, Jean-Claude Tardif, MD⁶, Michal Tendera, MD⁷, Alexander Parkhomenko, MD⁸, Deepak L. Bhatt, MD, MPH⁹, Philippe Gabriel Steg, MD^{1,5,10}, for the CLARIFY Investigators§

From the ¹Cardiology and Physiology Departments, Département Hospitalo-Universitaire FIRE, Assistance Publique-Hôpitaux de Paris, Hôpital Bichat, and Paris Diderot University, Sorbonne Paris Cité, Paris, France; ²Inserm U1149, Paris, France; ³University of Glasgow, Glasgow, UK; ⁴Maria Cecilia Hospital, GVM Care & Research, ES Health Science Foundation, and Department of Cardiology and LTTA Centre, University of Ferrara, Cotignola, Italy; ⁵National Heart and Lung Institute, Imperial College, Institute of Cardiovascular Medicine and Science, Royal Brompton Hospital, London, UK; ⁶Montreal Heart Institute, Université de Montréal, Montréal, Canada; ⁷Medical University of Silesia, Katowice, Poland; ⁸Institute of Cardiology, Kiev, Ukraine; ⁹Brigham and Women's Hospital Heart and Vascular Center, Harvard Medical School, Boston, MA, USA; ¹⁰French Alliance for Cardiovascular Trials, an F-CRIN network, INSERM U1148, Paris, France.

§A list of the CLARIFY Investigators is included in the [online-only Data Supplement](#).

Short title: Blood pressure in hypertensive CAD patients

Total Word Count: 7013

Abstract word count: 249

Number of Figures: 2

Number of Tables: 3

Corresponding author:

Prof Ph. Gabriel STEG, MD

Department of Cardiology, Hôpital Bichat, 46 rue Henri Huchard, 75018 Paris, France.

Tel. +331-40-25-86-68, Fax. +331-40-25-88-65, Email: gabriel.steg@aphp.fr

ABSTRACT

Observational studies have shown a J-shaped relationship between diastolic blood pressure (BP) and cardiovascular events in hypertensive patients with coronary artery disease (CAD). We investigated whether the increased risk associated with low diastolic BP reflects elevated pulse pressure. In 22 672 hypertensive CAD patients from the CLARIFY registry, followed for a median of 5.0 years, BP was measured annually and averaged. The relationships between pulse pressure and diastolic BP, alone or combined, and the primary composite outcome (cardiovascular death or myocardial infarction) were analyzed using multivariable Cox proportional hazards models. Adjusted hazard ratios (HRs) for the primary outcome were 1.62 (95%CI 1.40–1.87), 1.00 (ref), 1.07 (0.94–1.21), 1.54 (1.32–1.79), and 2.34 (1.95–2.81) for PP<45, 45–54 (reference), 55–64, 65–74, and ≥75 mmHg, respectively, and 1.50 (1.31–1.72), 1.00 (reference), and 1.58 (1.42–1.77) for diastolic BPs of <70, 70–79 (ref), and ≥80 mmHg, respectively. In a cross-classification analysis between diastolic BP and pulse pressure, the relationship between diastolic BP and the primary outcome remained J-shaped when the analysis was restricted to patients with the lowest-risk pulse pressure (45–64 mmHg), with adjusted HRs of 1.53 (1.27–1.83), 1.00 (ref), and 1.54 (1.34–1.75) in the <70, 70–79 (reference), and ≥80 mmHg subgroups, respectively. The J-shaped relationship between diastolic BP and cardiovascular events in hypertensive CAD patients persists in patients within the lowest-risk pulse pressure range and is therefore unlikely to be solely the consequence of an increased pulse pressure reflecting advanced vascular disease.

Key Words: hypertension, J-curve, pulse pressure, coronary artery disease, CLARIFY registry

INTRODUCTION

Elevated blood pressure (BP) is a well-established risk factor for cardiovascular events, and lowering elevated BP has been demonstrated to reduce risk.^{1,2} However, post-hoc analyses of randomized controlled trials^{3,4} and observational studies^{5,6} have shown that – in hypertensive patients with coronary artery disease (CAD) – the relationship between BP and cardiovascular events is J-shaped, particularly for diastolic BP, with an increased risk of cardiovascular events (except stroke) among patients with diastolic BP <70 mmHg.

The increased cardiovascular risk observed at low diastolic BP may be a direct consequence of altered myocardial perfusion.^{6–8} However, a causal link between low diastolic BP and cardiovascular events has not been demonstrated in randomized controlled trials. Rather, such trials have shown no benefit^{9,10} or a decreased risk¹¹ in the lowest BP groups, although it should be borne in mind that the mean BP in the lowest BP subgroup was never below the inflection point of the J-curve when considering office BP levels, either measured or extrapolated.^{12,13} Alternatively, the observed association may not be causal, but rather reflect reverse causality, whereby low diastolic BP would be an epiphenomenon of underlying poor health, itself leading to increased morbidity and mortality.^{14,15} Although indirect evidence argues against reverse causality being the sole or major explanation for the J-curve,^{5,14} irrefutable evidence will require future dedicated randomized interventional trials. Lastly, the association between low diastolic BP and cardiovascular events may be an epiphenomenon of increased pulse pressure, itself a cardiovascular risk marker,^{16–18} which is associated with diminished diastolic BP in patients with stiffened large arteries. Among 2207 patients from a hypertension control program, Madhavan et al. reported a J-shaped relation of diastolic BP to myocardial infarction that occurred only among patients with a PP>63 mmHg.¹⁹ Similarly, Kannel et al. showed – in 7798 subjects from the Framingham study and offspring cohort – that the increased risk observed at low diastolic BP was confined to patients with increased systolic BP and, therefore, increased PP, whereas the relationship

between diastolic BP and outcome remained linear among patients with systolic BP<140 mmHg.²⁰ More recently, Franklin et al observed in 791 individual with a previous cardiovascular event from the Framingham study that a diastolic BP <70 mmHg was associated with a greater risk than a diastolic BP of 70-79 mmHg only in patients with PP \geq 68 mmHg.²¹

The purpose of this study was to explore whether the J-curve observed for diastolic BP is restricted to patients with concomitant high systolic BP or PP, or whether it persists for patients with systolic BP and PP values within the lowest-risk range. We assessed the relationship between PP or diastolic BP and cardiovascular outcomes in a population of CAD patients treated for hypertension from the CLARIFY registry. We then performed cross-classifications to assess the relationship between diastolic BP and outcomes, stratified by PP or systolic BP subgroup.

METHODS

Details of the prospective, observational, longitudinal registry of patients with stable CAD (CLARIFY) have been reported.^{5,22} Briefly, 32 703 outpatients with stable CAD were recruited in 45 countries between November 2009 and June 2010. Exclusion criteria were hospital admission for cardiovascular reasons in the past 3 months, planned revascularization, or any health condition compromising 5-year follow-up, including severe other cardiovascular diseases (e.g. advanced heart failure, severe valve disease, or history of valve repair or replacement). Patients received standard clinical care; enrolment did not mandate any specific treatment or procedure.

Data were collected using standardized electronic case report forms at baseline and at every yearly visit for up to 5 years. At each yearly visit, symptoms, clinical examination, results of the main clinical and biological tests, treatment, and clinical outcomes were recorded; office BP was measured in seated subjects after a rest of 5 minutes, using the same

arm throughout the study, with no pre-specified device.

This analysis was restricted to patients treated for hypertension [Figure S1 in the Supplement](#), defined by “treated hypertension” on the baseline form and the use of at least one antihypertensive drug.

The study was performed in accordance with the Declaration of Helsinki. Local ethical approval was obtained in all countries. All patients gave written informed consent. This study is registered with clarify-registry.com, number ISRCTN43070564.

Blood pressure subgroups.

Pulse pressure was calculated as the difference between systolic BP and diastolic BP. Mean arterial pressure was calculated as diastolic BP + 1/3 PP. Analyses were performed using the arithmetic mean of all BP values measured throughout follow-up, from the baseline visit to the visit before an event (the event depending on the outcome), or all available visits if no event occurred.

For the relationship between a single BP component and outcome, patients were categorized into five subgroups by 10-mmHg increments, from <45 to ≥75 mmHg for PP (with the 45–54-mmHg subgroup as the reference), and from <60 to ≥90 mmHg for diastolic BP (with the 70–79-mmHg subgroup as the reference).

For cross-classification analyses, each BP component was divided in three subgroups, defined from the relationship with the primary outcome, using the thresholds below and above which event rates increased compared to the reference group: <70, 70–79 (reference), and ≥80 mmHg for diastolic BP, <45, 45–64 (reference), and ≥65 mmHg for PP, and <120, 120–139 (reference), and ≥140 mmHg for systolic BP. For systolic BP, these thresholds were defined from the previously established relationship between systolic BP and cardiovascular events in this population.⁵ In the cross-classification analyses, patients were further categorized into one of nine groups, using the combination of the three-level diastolic BP groupings and the three-level PP or systolic BP groupings.

Study outcomes.

The primary outcome was the composite of cardiovascular death or myocardial infarction.

Secondary outcomes included cardiovascular death, myocardial infarction, stroke, and hospitalization for heart failure. For patients with multiple events, the time to the first applicable event was considered in each analysis.

Statistical analyses.

Baseline characteristics are summarized according to the average PP categories prior to a primary outcome. Continuous variables are presented as means±standard deviations (SDs) or medians (interquartile ranges [IQRs]), depending on the distribution of the data; categorical data are presented as numbers and percentages. Comparisons between the average PP categories were made using either one-way analysis of variance or the Kruskal-Wallis test for continuous data, depending on the distribution of the data, or the chi-squared test for categorical data.

Cox proportional hazards models, both adjusted and unadjusted, were used to evaluate the relationship between BP categories (either for a single BP component or for combined BP components) and outcomes.

Covariates used for multivariable-adjustment were selected a priori as potential confounders and included (model 1) age, sex, geographic region, smoking status, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, diabetes, low- and high-density lipoprotein cholesterol levels, body mass index, glomerular filtration rate (GFR), peripheral artery disease, hospitalization for or symptoms of heart failure, left ventricular ejection fraction, ethnicity, stroke, transient ischemic attack, and baseline medications (aspirin, statins, angiotensin-converting enzyme inhibitors, angiotensin-receptor blockers, beta-blockers, calcium channel blockers, diuretics, and other antihypertensive medications).

Analyses performed on PP as a single BP component were also adjusted for mean

arterial pressure, in a separate model. Analyses performed on diastolic BP as a single BP component were also adjusted for PP, in a separate model.

No imputation was performed for missing data. Covariates with a large amount of missing data were categorized, including a category for missing data to minimize the loss of data in the analysis.

The relationship between PP subgroups and all outcomes was further assessed after excluding patients with heart failure, as defined by previous hospitalization for heart failure, symptoms of heart failure (then excluded from covariates), or a left ventricular ejection fraction <45%.

The models were further adjusted by including an interaction term between diastolic BP category and PP category or systolic BP category to determine whether any observed relationship was consistent across PP and systolic BP subgroups, respectively.

In the event of significant interactions with diastolic BP, the relationship of diastolic BP and outcome was further examined in the relevant PP or systolic BP subgroups.

Statistical analyses were performed using SAS (version 9.3). A *P*-value <0.05 was used to signify statistical significance using two-sided testing, with no correction for multiple comparisons.

RESULTS

Baseline characteristics of the patients in the total population (22 672 patients with CAD and hypertension) and by subgroups of PP are reported in [Table 1](#) and [Table S1 \(baseline medications\)](#). Mean age was 65.2 ± 10.0 years, 17 019 (75.1%) were men, and 7591 (33.5%) had diabetes. Mean systolic BP was 133.7 ± 16.7 mmHg, mean diastolic BP was 78.2 ± 10.1 mmHg, and mean PP was 55.4 ± 14.0 mmHg. Compared to patients with low PP, those with higher PP tended to be older, more likely to be women, have diabetes, be non-smokers, less likely to have had a myocardial infarction or percutaneous coronary intervention, had less

symptoms of heart failure, and had a higher prevalence of stroke. Baseline characteristics of the patients by subgroups of systolic and diastolic BP have been reported.⁵

Pulse pressure.

After a median (IQR) follow-up of 5.0 (4.5–5.1) years, the primary outcome had occurred in 1746 patients (7.7%). Cardiovascular death occurred in 1209 patients (5.3%), myocardial infarction (fatal/nonfatal) in 827 (3.6%), stroke (fatal/nonfatal) in 526 (2.3%), and hospital admission for heart failure in 1306 (5.8%). Event rates and adjusted hazard ratios (HRs) for PP subgroups are indicated in [Table 2](#). The relationship between average PP and crude and adjusted risk of the primary outcome followed a J-shaped curve. A similar J-shaped relationship was found for cardiovascular death, myocardial infarction, stroke, and hospitalization for heart failure ([Table 2](#)). The increased risk for elevated PP was more pronounced for myocardial infarction than for the other outcomes, with progressively increasing HRs as average PP increased. Similar results were found after exclusion of patients with heart failure, except for lost and attenuated associations of elevated PP with stroke and hospitalization for heart failure, respectively ([Table 2](#)).

Diastolic BP.

As previously demonstrated for cardiovascular death, myocardial infarction, and hospitalization for heart failure, but not for stroke,⁵ the relationship between average diastolic BP and the primary composite outcome was J-shaped, even after multiple adjustments for potential confounders. A similar J-shaped pattern was seen after further adjusting for PP ([Figure S2](#)).

Diastolic BP cross-classified with PP.

Diastolic BP, categorized in the three subgroups used for cross-classifications, had similar relationships to cardiovascular events as when categorized in five subgroups, namely J-shaped for all endpoints but stroke ([Table S2](#)). Event rates and HRs for the nine BP subgroups defined by cross-classifications between diastolic BP and PP are shown in the

[Figure 1](#) (primary outcome) and [Table 3](#) (secondary outcomes). The J-shaped relationship observed between diastolic BP and all endpoints but stroke remained, with very similar patterns, in patients with a PP within the lowest-risk range (45–64 mmHg). Interactions between PP categories and diastolic BP categories were significant for the primary outcome and cardiovascular death, but non-significant for myocardial infarction, stroke, and hospitalization for heart failure ([Table S3](#)). Analyses by PP subgroup in case of a significant interaction are shown in [Table S4](#). The significant interaction for the primary outcome and cardiovascular death revealed a steeper J-curve, with an even greater risk associated with low diastolic BP (<70 mmHg) in patients with PP<45 mmHg. Conversely, in patients with PP≥65 mmHg, there was no increase in the risk of cardiovascular events in patients with diastolic BP<70 versus 70–79 mmHg. The non-significant interaction for other endpoints showed that the relationship between diastolic BP and cardiovascular events was consistent across PP subgroups. Mean BPs in each subgroup are shown in [Table S5](#).

Diastolic BP cross-classified with systolic BP.

Crude and adjusted HRs between BP subgroups, defined by cross-classifications between diastolic BP and systolic BP categories, are indicated in [Figure 2](#) (primary endpoint) and [Table 3](#) (secondary endpoints). The J-shaped relationship observed between diastolic BP and cardiovascular events remained when restricting the analysis to patients with systolic BP in the lowest-risk range of 120–139 mmHg. Interaction between diastolic BP and systolic BP was non-significant for most endpoints ([Table S3](#)), hence the relationship between diastolic BP and outcome was consistent across systolic BP subgroups. In the case of hospitalization for heart failure, there was a significant interaction between diastolic BP and systolic BP ($p=0.0199$), and the risk associated with a low diastolic BP increased as systolic BP increased ([Table S4](#)). Mean BP values in each subgroup are indicated in [Table S5](#).

DISCUSSION

This large international study evaluated the cardiovascular risk associated with single or combined components of BP in 22 672 CAD patients treated for hypertension to decipher the underlying mechanisms of the J-shaped relationship observed between diastolic BP and cardiovascular events (except stroke) in this population. Even though elevated PP was strongly associated with all cardiovascular endpoints, this phenomenon did not account for the increased risk observed at low diastolic BP. Indeed, the J-shaped relationship between diastolic BP and cardiovascular outcomes (primary endpoint and all secondary endpoints but stroke) persisted in patients with PP or systolic BP in the lowest-risk range (45–64 and 120–139 mmHg, respectively).

In this large population of CAD patients treated for hypertension and followed according to routine clinical practice, elevated PP was associated with an increased risk of all outcomes, even after adjustment for multiple covariates, including mean arterial pressure, confirming previous studies conducted in various populations that have shown that PP is an independent cardiovascular risk marker.^{16–18} PP is an indicator of left ventricle ejection volume and velocity, and viscoelastic properties of large arteries. Therefore, elevated PP is correlated with vascular ageing, both through a weakened Windkessel effect and an increased pulse wave velocity, with an earlier reflection wave increasing systolic BP and reducing diastolic BP. Furthermore, by definition, as PP increases for a given mean arterial pressure, systolic BP – and thus the afterload of the left ventricle – increases, and diastolic BP – and thus myocardial perfusion – decreases, both factors also potentially accounting for the increased risk associated with high PP. However, whether reducing PP reduces cardiovascular risk has not yet been established, unlike the clear beneficial effect of reducing elevated systolic BP.^{1,23,24}

In addition to the increased risk associated with elevated PP, we also found an increased risk associated with low PP (<45 mmHg). Such a J-shaped pattern for the link

between PP and the rate of cardiovascular events had not been shown in all previous studies.^{16,25} However, in a post-hoc analysis of the INVEST trial,¹⁸ which included 22 576 CAD patients with hypertension, the relationship between PP and cardiovascular death and myocardial infarction was J-shaped, with an increased risk below a nadir value of 54 mmHg (95% CI, 42–60 mmHg), in line with our results. In addition, in one of the largest studies examining the risk associated with PP, conducted in the REACH registry (45 087 high-risk subjects), we recently showed that not only was elevated PP associated with multiple adverse cardiovascular outcomes, but patients in the first quartile of PP (<50 mmHg) also displayed an increased risk of cardiovascular death.¹⁷ Reverse causality, associated with low stroke volume,²⁶ may at least in part explain this phenomenon. However, patients with severe aortic stenosis were excluded from the study, this association persisted after multiple adjustments for confounding factors and in a sensitivity analysis excluding patients with heart failure. Alternatively, the increased risk observed for low PP, which was highest in the lower end of the BP spectrum, may be driven by the possibly additive risks of low diastolic BP (compromising myocardial perfusion) combined with low mean arterial pressure (compromising perfusion of all other organs). Because the relationship between PP and the primary outcome was J-shaped in our study population, we divided PP in three subgroups for cross-classifications, the middle subgroup being that associated with the lowest risk (45–65 mmHg).

The increased risk associated with elevated PP has led many authors to hypothesize that the J-shaped relation to cardiovascular risk associated with diastolic BP largely reflects increased pulse pressure, an indicator of advanced vascular disease and stiffened large arteries.^{16,27–29} However, our results do not confirm previous studies, which had suggested that the increased risk associated with low diastolic BP was restricted to patients with increased PP^{19,21} or increased systolic BP.²⁰ Importantly, these studies were conducted in much smaller cohorts, and did not specifically include CAD patients. We showed that, even

in patients within the lowest-risk range of PP or systolic BP, the J-shaped relationship between diastolic BP and cardiovascular events remained, which strongly argues against increased PP being the sole explanation for the increased risk observed at low diastolic BP. Similarly, the J-curve relationship between diastolic BP and the primary outcome persisted after adjustment for PP, as previously shown in 10001 CAD patients enrolled in the TNT trial.⁴ In line with our results, in a cohort of 331 frail elderly patients, Protogerou et al had found that a diastolic BP ≤ 60 mmHg was associated with increased mortality during a two-year follow-up, independently of large artery stiffness as measured by pulse wave analysis.³⁰

Not only was the J-shaped relationship between diastolic BP and cardiovascular events not restricted to patients with increased PP, it was actually attenuated as PP increased, at least for the primary outcome and cardiovascular death, for which the interaction between diastolic BP and PP was significant. The increased risk observed at low values of diastolic BP was highest when accompanied by a low PP, as discussed above. Conversely, in patients with elevated PP (≥ 65 mmHg), the risk was higher than in patients with intermediate levels of PP, but there was no further increase in the risk when diastolic BP was < 70 mmHg. A likely explanation for that is that in this subgroup (PP ≥ 65 mmHg and diastolic BP < 70 mmHg), although the lower diastolic BP is expected to be deleterious, this may be compensated by the lower systolic BP accompanying these lower diastolic BP values (138 mmHg in the lowest diastolic BP subgroup versus 147 mmHg in the 70–79-mmHg subgroup), itself clearly beneficial.

The main limitation of our study is that it is an observational registry, and we cannot therefore draw conclusions on whether associations between single or combined components of BP and cardiovascular risk are causal or would be reversed by interventions to diminish PP or increase low diastolic BP. Our results suggest that low myocardial perfusion may be a more likely explanation for the J-curve than increased vascular ageing associated with high PP and low diastolic BP, but do not demonstrate that it is the sole or major mechanism, nor

rule out some reverse causality. Assumptions based on observational studies are the basis for future randomized trials to define optimal BP targets. Amongst other limitations of our study, our results were obtained in hypertensive patients with stable CAD free from other severe conditions, and cannot be extrapolated to healthier hypertensive subjects, or to elderly frail patients. Similarly, these data should not be extrapolated to non-hypertensive CAD patients, who are commonly treated with BP-lowering drugs and in whom the potential deleterious effects of low BP remain to be studied. Finally, the CLARIFY registry reflects routine clinical practice, and although our results may have wider external validity than randomized trials, measurements of BP were less standardized and outcome identification possibly less accurate than in randomized trials.

PERSPECTIVES

Even though elevated PP is associated with increased cardiovascular risk and is closely intertwined with decreased diastolic pressure, it does not appear to be the major determinant of the increased risk associated with low diastolic BP in this large cohort of CAD patients. A compromised myocardial perfusion associated with low diastolic BP in patients with CAD appears to be a more plausible explanation, although reverse causation cannot be ruled out. Our data, although observational, suggest caution when lowering diastolic BP below 70 mmHg, and even more so 60 mmHg, in patients with CAD treated for hypertension. However, only randomized trials will provide irrefutable evidence for a causal link between low diastolic BP and adverse cardiovascular events. BP target trials comparing levels of achieved BP, especially levels of diastolic BP in patients with CAD, are needed to define optimal BP targets in this population.

ACKNOWLEDGEMENT

Editorial support limited to editing, checking content and language, and formatting was provided by Jenny Lloyd and Sophie Rushton-Smith (MedLink Healthcare Communications Ltd) and was funded by Servier.

SOURCE OF FUNDING

The CLARIFY registry was supported by Servier.

DISCLOSURES

Dr Vidal-Petiot reports travel support from Servier. Ms Greenlaw reports grants from Servier. Dr Ford reports grants and personal fees from Servier and Amgen. Dr Ferrari reports receiving honorarium from Servier for steering committee membership consulting and speaking, support for travel to study meetings from Servier., personal fees from Boehringer-Ingelheim, Novartis, Merck Serono and Irbtech, and he is a stockholder in Medical Trials Analysis. Dr Fox reports personal fees and non-financial support from Servier and Broadview Ventures, personal fees from AstraZeneca, TaurX and CellAegis, and non-financial support from Armgo. He is Director of Heart Research Ltd and Vesalius Trials Ltd and a minimal stockholder of Armgo and CellAegis. Dr Tardif reports grants and personal fees from Amarin, AstraZeneca, Pfizer, Sanofi, and Servier, and grants, personal fees and other (minor equity interest) from DalCor, grants from Esperion, Ionis, and Merck. In addition, Dr. Tardif has a patent Pharmacogenomics-guided therapy with CETP inhibition pending. Dr Tendera reports personal fees from Servier, Bayer, Janssen Cilag, KOWA, and the Perfuse Study Group, and grants from EU Framework Program VII and the from Polish National Center for Research and Development. Dr Parkhomenko reports grants and personal fees from Servier. Dr Bhatt discloses the following relationships: Advisory Board: Cardax, Elsevier Practice

Update Cardiology, Medscape Cardiology, Regado Biosciences; Board of Directors: Boston VA Research Institute, Society of Cardiovascular Patient Care; Chair: American Heart Association Quality Oversight Committee; Data Monitoring Committees: Cleveland Clinic, Duke Clinical Research Institute, Harvard Clinical Research Institute, Mayo Clinic, Mount Sinai School of Medicine, Population Health Research Institute; Honoraria: American College of Cardiology (Senior Associate Editor, Clinical Trials and News, ACC.org), Belvoir Publications (Editor in Chief, Harvard Heart Letter), Duke Clinical Research Institute (clinical trial steering committees), Harvard Clinical Research Institute (clinical trial steering committee), HMP Communications (Editor in Chief, Journal of Invasive Cardiology), Journal of the American College of Cardiology (Guest Editor; Associate Editor), Population Health Research Institute (clinical trial steering committee), Slack Publications (Chief Medical Editor, Cardiology Today's Intervention), Society of Cardiovascular Patient Care (Secretary/Treasurer), WebMD (CME steering committees); Other: Clinical Cardiology (Deputy Editor), NCDR-ACTION Registry Steering Committee (Chair), VA CART Research and Publications Committee (Chair); Research Funding: Amarin, Amgen, AstraZeneca, Bristol-Myers Squibb, Chiesi, Eisai, Ethicon, Forest Laboratories, Ironwood, Ischemix, Lilly, Medtronic, Pfizer, Roche, Sanofi Aventis, The Medicines Company; Royalties: Elsevier (Editor, Cardiovascular Intervention: A Companion to Braunwald's Heart Disease); Site Co-Investigator: Biotronik, Boston Scientific, St. Jude Medical (now Abbott); Trustee: American College of Cardiology; Unfunded Research: FlowCo, Merck, PLx Pharma, Takeda. Dr Steg reports research grants from Merck, Sanofi, and Servier and speaking or consulting fees from Amarin, Amgen, AstraZeneca, Bayer, Boehringer-Ingelheim, Bristol-Myers-Squibb, Janssen, Lilly, Merck, Novartis, Pfizer, Regeneron, Sanofi, and Servier

REFERENCES

1. Ettehad D, Emdin CA, Kiran A, Anderson SG, Callender T, Emberson J, Chalmers J, Rodgers A, Rahimi K. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet* 2016;387(10022):957-967.
2. Mancia G, Fagard R, Narkiewicz K, et al. 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *J Hypertens.* 2013;31:1281-1357.
3. Messerli FH, Mancia G, Conti CR, Hewkin AC, Kupfer S, Champion A, Kolloch R, Benetos A, Pepine CJ. Dogma disputed: can aggressively lowering blood pressure in hypertensive patients with coronary artery disease be dangerous? *Ann Intern Med.* 2006;144:884-893.
4. Bangalore S, Messerli FH, Wun C-C, Zuckerman AL, DeMicco D, Kostis JB, LaRosa JC. J-curve revisited: An analysis of blood pressure and cardiovascular events in the Treating to New Targets (TNT) Trial. *Eur Heart J.* 2010;31:2897-2908.
5. Vidal-Petiot E, Ford I, Greenlaw N, Ferrari R, Fox KM, Tardif J-C, Tendera M, Tavazzi L, Bhatt DL, Steg PG, CLARIFY Investigators. Cardiovascular event rates and mortality according to achieved systolic and diastolic blood pressure in patients with stable coronary artery disease: an international cohort study. *Lancet* 2016;388(10056):2142-2152.
6. McEvoy JW, Chen Y, Rawlings A, Hoogeveen RC, Ballantyne CM, Blumenthal RS, Coresh J, Selvin E. Diastolic Blood Pressure, Subclinical Myocardial Damage, and Cardiac Events: Implications for Blood Pressure Control. *J Am Coll Cardiol.* 2016;68:1713-1722.

7. Bhatt DL. Troponin and the J-Curve of Diastolic Blood Pressure: When Lower Is Not Better. *J Am Coll Cardiol.* 2016;68:1723-1726.
8. Rabkin SW, Shiekh IA, Wood DA. The Impact of Left Ventricular Mass on Diastolic Blood Pressure Targets for Patients With Coronary Artery Disease. *Am J Hypertens.* 2016;29:1085-1093.
9. ACCORD Study Group, Cushman WC, Evans GW, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med.* 2010;362:1575-1585.
10. SPS3 Study Group, Benavente OR, Coffey CS, Conwit R, Hart RG, McClure LA, Pearce LA, Pergola PE, Szychowski JM. Blood-pressure targets in patients with recent lacunar stroke: the SPS3 randomised trial. *Lancet* 2013;382:507-515.
11. SPRINT Research Group, Wright JT, Williamson JD, et al. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med.* 2015;373:2103-2116.
12. Kjeldsen SE, Lund-Johansen P, Nilsson PM, Mancia G. Unattended Blood Pressure Measurements in the Systolic Blood Pressure Intervention Trial: Implications for Entry and Achieved Blood Pressure Values Compared With Other Trials. *Hypertens* 2016;67:808-812.
13. Kjeldsen SE, Mancia G. Unobserved automated office blood pressure measurement in the Systolic Blood Pressure Intervention Trial (SPRINT): systolic blood pressure treatment target remains below 140 mmHg. *Eur Heart J Cardiovasc Pharmacother.* 2016;2:79-80.
14. Messerli FH, Panjrath GS. The J-curve between blood pressure and coronary artery disease or essential hypertension: exactly how essential? *J Am Coll Cardiol.* 2009;54:1827-1834.

15. Ravindrarajah R, Hazra NC, Hamada S, Charlton J, Jackson SHD, Dregan A, Gulliford MC. Systolic Blood Pressure Trajectory, Frailty, and All-Cause Mortality >80 Years of Age: Cohort Study Using Electronic Health Records. *Circulation*. 2017;135:2357-2368.
16. Blacher J, Staessen JA, Girerd X, Gasowski J, Thijs L, Liu L, Wang JG, Fagard RH, Safar ME. Pulse pressure not mean pressure determines cardiovascular risk in older hypertensive patients. *Arch Intern Med*. 2000;160:1085-1089.
17. Selvaraj S, Steg PG, Elbez Y, Sorbets E, Feldman LJ, Eagle KA, Ohman EM, Blacher J, Bhatt DL, REACH Registry Investigators. Pulse Pressure and Risk for Cardiovascular Events in Patients With Atherothrombosis: From the REACH Registry. *J Am Coll Cardiol*. 2016;67:392-403.
18. Bangalore S, Messerli FH, Franklin SS, Mancia G, Champion A, Pepine CJ. Pulse pressure and risk of cardiovascular outcomes in patients with hypertension and coronary artery disease: an INternational VErapamil SR-trandolapril STudy (INVEST) analysis. *Eur Heart J*. 2009;30:1395-1401.
19. Madhavan S, Ooi WL, Cohen H, Alderman MH. Relation of pulse pressure and blood pressure reduction to the incidence of myocardial infarction. *Hypertens* 1994;23:395-401.
20. Kannel WB, Wilson PWF, Nam B-H, D'Agostino RB, Li J. A likely explanation for the J-curve of blood pressure cardiovascular risk. *Am J Cardiol*. 2004;94:380-384.
21. Franklin SS, Gokhale SS, Chow VH, Larson MG, Levy D, Vasan RS, Mitchell GF, Wong ND. Does low diastolic blood pressure contribute to the risk of recurrent hypertensive cardiovascular disease events? The Framingham Heart Study. *Hypertens* 2015;65:299-305.

22. Steg PG, Greenlaw N, Tendera M, Tardif J-C, Ferrari R, Al-Zaibag M, Dorian P, Hu D, Shalnova S, Sokn FJ, Ford I, Fox KM, Prospective Observational Longitudinal Registry of Patients With Stable Coronary Artery Disease (CLARIFY) Investigators. Prevalence of anginal symptoms and myocardial ischemia and their effect on clinical outcomes in outpatients with stable coronary artery disease: data from the International Observational CLARIFY Registry. *JAMA Intern Med.* 2014;174:1651-1659.
23. Bangalore S, Toklu B, Ganos E, Schwartzbard A, Weintraub H, Ogedegbe G, Messerli FH. Optimal Systolic Blood Pressure Target After SPRINT: Insights from a Network Meta-Analysis of Randomized Trials. *Am J Med.* 2017;130:707-719.e8.
24. Bundy JD, Li C, Stuchlik P, Bu X, Kelly TN, Mills KT, He H, Chen J, Whelton PK, He J. Systolic Blood Pressure Reduction and Risk of Cardiovascular Disease and Mortality: A Systematic Review and Network Meta-analysis. *JAMA Cardiol.* 2017;2:775-81.
25. Elliott HL, Lloyd SM, Ford I, Meredith PA. Prognostic importance of pretreatment and on-treatment blood pressure: Further analysis of the ACTION database and the effect of nifedipine gastrointestinal therapeutic system. *Blood Press.* 2016;25:67-73.
26. Jackson CE, Castagno D, Maggioni AP, et al. Differing prognostic value of pulse pressure in patients with heart failure with reduced or preserved ejection fraction: results from the MAGGIC individual patient meta-analysis. *Eur Heart J.* 2015;36:1106-1114.
27. Franklin SS, Lopez VA, Wong ND, Mitchell GF, Larson MG, Vasan RS, Levy D. Single versus combined blood pressure components and risk for cardiovascular disease: the Framingham Heart Study. *Circulation.* 2009;119:243-250..
28. Millar JA, Lever AF. Implications of pulse pressure as a predictor of cardiac risk in patients with hypertension. *Hypertens* 2000;36:907-911.

29. Kannel WB, Vasan RS. Aggressive lowering of blood pressure. *Lancet* 2006;368:627-628.
30. Protoperou AD, Safar ME, Iaria P, Safar H, Le Dugal K, Filipovsky J, Henry O, Ducimetière P, Blacher J. Diastolic blood pressure and mortality in the elderly with cardiovascular disease. *Hypertens* 2007; 50:172-80

NOVELTY AND SIGNIFICANCE

What is new?

- In this large population of patients with coronary artery disease and treated hypertension, we show for the first time that the J-shaped relationship between diastolic BP and cardiovascular events persists in patients with the lowest-risk pulse pressure.

What is relevant?

- The increased risk observed at low diastolic BP is not an epiphenomenon of increased pulse pressure.
- Although reverse causality cannot be ruled out by our observational study, the alternative hypothesis of a compromised myocardial perfusion associated with low diastolic BP appears to be a likely explanation for the J-curve of diastolic

Summary

In 22 672 hypertensive patients from the CLARIFY registry, the J-shaped relationship between diastolic BP and the primary outcome (cardiovascular death or myocardial infarction) remained in patients within the lowest-risk pulse pressure range (45-65 mmHg), with adjusted HRs of 1.53, 1.00, and 1.54 in the <70, 70–79 (reference), and ≥80 mmHg diastolic BP subgroups, respectively.

Figure legends

FIGURE 1 Forest plots of adjusted hazard ratios for the primary outcome (cardiovascular death or myocardial infarction) for diastolic blood pressure subgroups cross-classified with pulse pressure subgroups.

Analyses are adjusted as defined for model 1.

HR, hazard ratio; CI, confidence interval; PP, pulse pressure; DBP, diastolic blood pressure.

FIGURE 2 Forest Plots of Adjusted Hazard Ratios of the Primary Outcome (Cardiovascular Death or Myocardial Infarction) for Diastolic Blood Pressure Subgroups Cross-Classified with Systolic Blood Pressure Subgroups.

Analyses are adjusted as defined for model 1.

HR, hazard ratio; CI, confidence interval; DBP, diastolic blood pressure; SBP, systolic blood pressure.

TABLE 1 Demographic and Baseline Characteristics of the Patients, for the Total Population and Each Average On-Treatment PP**Subgroup**

Parameter	n	Total population (n=22 672)	Average PP categories					p Value
			<45 mmHg (n=3088)	45–54 mmHg (n=9013)	55–64 mmHg (n=6907)	65–74 mmHg (n=2611)	≥75 mmHg (n=1053)	
Age, y	22 666	65.2±10.0	61.0±10.3	63.4±9.8	66.9±9.4	69.2±8.7	71.5±8.6	<0.0001
Male	22 672	17 019 (75.1)	2505 (81.1)	6974 (77.4)	5070 (73.4)	1826 (69.9)	644 (61.2)	<0.0001
BMI, kg/m ²	22 654	27.7 (25.2–30.9)	27.2 (24.7–30.1)	27.8 (25.4–30.9)	27.8 (25.1–31.1)	27.8 (25.2–31.2)	27.6 (24.8–30.8)	<0.0001
Diabetes	22 670	7591 (33.5)	818 (26.5)	2750 (30.5)	2461 (35.6)	1089 (41.7)	473 (44.9)	<0.0001
Smoking status	22 672							<0.0001
Current		2569 (11.3)	478 (15.5)	1100 (12.2)	683 (9.9)	234 (9.0)	74 (7.0)	
Former		10 158 (44.8)	1404 (45.5)	4104 (45.5)	3044 (44.1)	1131 (43.3)	475 (45.1)	
Never		9945 (43.9)	1206 (39.1)	3809 (42.3)	3180 (46.0)	1246 (47.7)	504 (47.9)	
PP, mmHg	22 658	55.4±14.0	40.7±7.5	50.3±8.8	59.7±10.2	68.5±12.0	82.3±14.6	—
SBP, mmHg	22 659	133.7±16.7	118.6±11.8	129.3±12.8	137.8±14.2	145.9±16.0	158.1±18.4	—

Parameter	n	Total population (n=22 672)	Average PP categories					p Value
			<45 mmHg (n=3088)	45–54 mmHg (n=9013)	55–64 mmHg (n=6907)	65–74 mmHg (n=2611)	≥75 mmHg (n=1053)	
DBP, mmHg	22 659	78.2±10.1	77.9±9.2	79.0±9.6	78.1±10.2	77.4±11.4	75.8±12.1	—
Heart rate, bpm	22 660	68.5±10.6	68.7±10.4	68.6±10.4	68.6±10.7	68.0±11.5	67.0±11.1	<0.0001
Myocardial infarction	22 670	13 258 (58.5)	2081 (67.4)	5542 (61.5)	3785 (54.8)	1341 (51.4)	509 (48.3)	<0.0001
PCI	22 670	12 962 (57.2)	1863 (60.3)	5182 (57.5)	3930 (56.9)	1441 (55.2)	546 (51.9)	<0.0001
CABG	22 670	5691 (25.1)	695 (22.5)	2073 (23.0)	1804 (26.1)	770 (29.5)	349 (33.1)	<0.0001
TIA	22 670	801 (3.5)	86 (2.8)	291 (3.2)	273 (4.0)	110 (4.2)	41 (3.9)	0.0047
Stroke	22 670	1089 (4.8)	132 (4.3)	415 (4.6)	341 (4.9)	132 (5.1)	69 (6.6)	0.035
Hospitalization for HF	22 670	1211 (5.3)	202 (6.5)	445 (4.9)	343 (5.0)	158 (6.1)	63 (6.0)	0.0019
LVEF, %	15 969	56.1±11.0	54.2±12.1	55.9±10.8	56.6±10.6	57.1±10.9	57.8±10.9	<0.0001
HbA1C, %	6173	6.9±1.8	6.7±1.4	6.8±1.7	6.9±1.4	7.0±1.5	7.4±4.2	<0.0001
Creatinine, µmol/L	17 165	88 (76–104)	88 (77–103)	88 (76–102)	88 (76–103)	88 (75–106)	90 (76–110)	0.0010

Parameter	n	Total population (n=22 672)	Average PP categories					p Value
			<45 mmHg (n=3088)	45–54 mmHg (n=9013)	55–64 mmHg (n=6907)	65–74 mmHg (n=2611)	≥75 mmHg (n=1053)	
HDL-cholesterol, mmol/L	16 054	1.1 (1.0–1.4)	1.1 (0.9–1.3)	1.1 (1.0–1.3)	1.2 (1.0–1.4)	1.2 (1.0–1.4)	1.1 (1.0–1.4)	<0.0001
LDL-cholesterol, mmol/L	15 257	2.4 (1.9–3.0)	2.3 (1.8–2.9)	2.4 (1.9–3.0)	2.4 (1.9–2.9)	2.3 (1.8–2.8)	2.4 (1.9–3.0)	<0.0001
Fasting triglycerides, mmol/L	16 806	1.4 (1.0–2.0)	1.4 (1.0–2.0)	1.5 (1.1–2.0)	1.4 (1.0–1.9)	1.4 (1.0–2.0)	1.4 (1.0–1.9)	0.0018

Values are mean±SD, n (%), or median (IQR).

BMI=body mass index; CABG=coronary artery bypass graft; HbA1c=hemoglobin A1c; HF=heart failure; IQR=interquartile range; LVEF=left ventricular ejection fraction; NYHA>New York Heart Association; PCI=percutaneous coronary intervention; PP=pulse pressure; SD=standard deviation; TIA=transient ischemic attack.

TABLE 2 Event rates and Adjusted Hazard Ratios for PP Subgroups

Parameter	HR (95% CI) for average PP subgroups					p Value
	<45 mmHg	45–54 mmHg	55–64 mmHg	65–74 mmHg	≥75 mmHg	
Cardiovascular death or myocardial infarction (primary composite outcome)						
Event rate, %	9.3%	6.0%	6.8%	10.4%	16.3%	<0.0001
HR (Model 1)	1.62 (1.40–1.87)	1.00	1.07 (0.94–1.21)	1.54 (1.32–1.79)	2.34 (1.95–2.81)	<0.0001
HR (Model 2)	1.62 (1.40–1.88)	1.00	1.07 (0.94–1.21)	1.54 (1.31–1.79)	2.33 (1.92–2.83)	<0.0001
Excluding HF	1.63 (1.33–2.00)	1.00	0.99 (0.84–1.17)	1.38 (1.14–1.69)	2.15 (1.70–2.73)	<0.0001
Cardiovascular death						
Event rate, %	7.0%	4.0%	4.6%	7.4%	11.7%	<0.0001
HR (Model 1)	1.80 (1.51–2.13)	1.00	1.02 (0.87–1.19)	1.48 (1.24–1.78)	2.20 (1.77–2.74)	<0.0001
HR (Model 2)	1.79 (1.50–2.13)	1.00	1.02 (0.87–1.19)	1.49 (1.24–1.80)	2.23 (1.77–2.80)	<0.0001
Excluding HF	1.90 (1.48–2.44)	1.00	0.94 (0.77–1.16)	1.40 (1.10–1.79)	2.04 (1.52–2.73)	<0.0001
Myocardial infarction						
Event rate, %	3.9%	2.9%	3.4%	4.9%	7.8%	<0.0001
HR (Model 1)	1.39 (1.12–1.73)	1.00	1.25 (1.04–1.50)	1.81 (1.45–2.25)	2.94 (2.25–3.83)	<0.0001

HR (Model 2)	1.42 (1.14– 1.78)	1.00	1.23 (1.02– 1.47)	1.74 (1.39– 2.20)	2.79 (2.10– 3.69)	<0.0001
Excluding HF	1.41 (1.06– 1.88)	1.00	1.18 (0.95– 1.48)	1.55 (1.18– 2.04)	2.32 (1.65– 3.25)	<0.0001
Stroke						
Event rate, %	2.1%	1.9%	2.2%	3.3%	4.4%	<0.0001
HR (Model 1)	1.26 (0.94– 1.68)	1.00	1.06 (0.85– 1.32)	1.56 (1.20– 2.04)	1.98 (1.41– 2.80)	0.0001
HR (Model 2)	1.39 (1.04– 1.86)	1.00	0.98 (0.78– 1.23)	1.35 (1.03– 1.79)	1.56 (1.09– 2.25)	0.0044
Excluding HF	1.55 (1.09– 2.21)	1.00	0.78 (0.59– 1.03)	1.06 (0.76– 1.49)	1.24 (0.81– 1.91)	0.0032
Hospitalization for HF						
Event rate, %	7.7%	5.3%	5.4%	6.5%	8.6%	<0.0001
HR (Model 1)	1.47 (1.25– 1.73)	1.00	1.17 (1.02– 1.35)	1.40 (1.16– 1.68)	2.08 (1.63– 2.65)	<0.0001
HR (Model 2)	1.56 (1.32– 1.83)	1.00	1.12 (0.97– 1.29)	1.27 (1.05– 1.54)	1.82 (1.41– 2.35)	<0.0001
Excluding HF	1.51 (1.11– 2.06)	1.00	0.98 (0.78– 1.25)	0.92 (0.66– 1.29)	1.65 (1.13– 2.41)	0.0019

Model 1, see methods. Model 2: additional adjustment for mean arterial pressure.

CI=confidence interval; GFR=glomerular filtration rate; HF=heart failure; HR=hazard ratio;

PP=pulse pressure.

TABLE 3 Cross-Classifications Analysis: Event Rates and adjusted Hazard Ratios for Secondary Outcomes

	PP	Diastolic BP subgroups			Systolic BP	Diastolic BP subgroups		
		<70 mmHg	70–79 mmHg	≥80 mmHg		<70 mmHg	70–79 mmHg	≥80 mmHg
Cardiovascular death								
n/N	mmHg	50/332	84/1509	81/1230	mmHg	121/1073	93/1472	13/148
Event rate, %		15.1	5.6	6.6		11.3	6.3	8.8
HR (95% CI)		3.49 (2.56–4.77)	1.64 (1.27–2.10)	2.80 (2.17–3.61)		2.51 (2.02–3.14)	1.77 (1.39, 2.26)	4.08 (2.32, 7.15)
n/N	mmHg	126/1807	278/7805	275/6318	mmHg	105/1671	270/7826	218/5101
Event rate, %		7.0	3.6	4.4		6.3	3.5	4.3
HR (95% CI)		1.55 (1.25–1.93)		1.52 (1.28–1.81)		1.43 (1.14–1.81)		1.59 (1.32–1.91)
n/N	mmHg	81/913	113/1581	120/1144	mmHg	31/308	112/1597	245/3443
Event rate, %		8.9	7.1	10.5		10.1	7.0	7.1
HR (95% CI)		1.77 (1.37–2.29)	1.67 (1.34–2.09)	3.10 (2.49–3.84)		1.85 (1.26–2.71)	1.72 (1.37–2.14)	2.43 (2.04–2.90)

	PP	Diastolic BP subgroups			Systolic BP	Diastolic BP subgroups		
		<70 mmHg	70–79 mmHg	≥80 mmHg		<70 mmHg	70–79 mmHg	≥80 mmHg
myocardial infarction								
n/N	<45 mmHg	23/338	49/1504	49/1242	<120 mmHg	61/1075	48/1464	6/149
Event rate, %		6.8	3.3	3.9		5.7	3.3	4.0
HR (95% CI)		2.75 (1.77– 4.27)	1.37 (0.99– 1.88)	1.91 (1.38– 2.64)		2.06 (1.54– 2.77)	1.28 (0.93– 1.76)	2.13 (0.94– 4.83)
n/N	45–64 mmHg	76/1803	185/7755	234/6341	120–139 mmHg	74/1665	195/7783	162/5108
Event rate, %		4.2	2.4	3.7		4.4	2.5	3.2
HR (95% CI)		1.61 (1.23– 2.12)		1.68 (1.38– 2.05)		1.62 (1.23– 2.14)		1.38 (1.12– 1.71)
n/N	≥65 mmHg	49/905	77/1577	84/1174	≥140 mmHg	13/306	68/1589	199/3500
Event rate, %		5.4	4.9	7.2		4.2	4.3	5.7
HR (95% CI)		2.07 (1.49– 2.87)	2.03 (1.55– 2.66)	3.46 (2.66– 4.49)		1.41 (0.80– 2.50)	1.69 (1.28– 2.23)	2.60 (2.12– 3.18)
stroke								

	PP	Diastolic BP subgroups			Systolic BP	Diastolic BP subgroups		
		<70 mmHg	70–79 mmHg	≥80 mmHg		<70 mmHg	70–79 mmHg	≥80 mmHg
n/N	<45 mmHg	11/339	25/1508	29/1245	<120 mmHg	26/1076	26/1469	1/147
Event rate, %		3.2	1.7	2.3		2.4	1.8	0.7
HR (95% CI)		2.17 (1.16– 4.05)	1.22 (0.79– 1.89)	2.10 (1.39– 3.18)		1.38 (0.90– 2.11)	1.19 (0.78– 1.82)	0.67 (0.09– 4.83)
n/N	45–64 mmHg	43/1804	127/7761	157/6327	120–139 mmHg	46/1669	131/7789	108/5109
Event rate, %		2.4	1.6	2.5		2.8	1.7	2.1
HR (95% CI)		1.32 (0.93– 1.87)		1.77 (1.39– 2.26)		1.42 (1.01– 2.01)		1.53 (1.17– 1.98)
n/N	≥65 mmHg	31/912	55/1588	47/1155	≥140 mmHg	13/310	50/1599	124/3471
Event rate, %		3.4	3.5	4.1		4.2	3.1	3.6
HR (95% CI)		1.66 (1.11– 2.49)	1.93 (1.40– 2.66)	2.82 (2.01– 3.95)		1.80 (1.01– 3.22)	1.74 (1.25– 2.42)	2.47 (1.92– 3.18)
Hospitalization for heart failure								
n/N	<45	28/299	79/1444	121/1200	<120	84/1016	87/1408	16/135

	PP	Diastolic BP subgroups			Systolic BP	Diastolic BP subgroups		
		<70 mmHg	70–79 mmHg	≥80 mmHg		<70 mmHg	70–79 mmHg	≥80 mmHg
Event rate, %	mmHg	9.4	5.5	10.1	mmHg	8.3	6.2	11.9
HR (95% CI)		2.07 (1.39– 3.07)	1.40 (1.09– 1.81)	2.52 (2.02– 3.14)		2.12 (1.65– 2.72)	1.47 (1.15– 1.88)	4.67 (2.80– 7.79)
n/N	45–64	109/1754	274/7576	444/6105	120–139	86/1618	265/7590	302/4915
Event rate, %		6.2	3.6	7.3		5.3	3.5	6.1
HR (95% CI)	mmHg	1.82 (1.45– 2.29)		1.66 (1.42– 1.94)	mmHg	1.68 (1.31– 2.15)		1.43 (1.21– 1.70)
n/N		55/874	77/1539	118/1120		22/293	78/1561	365/3375
Event rate, %	≥65	6.3	5.0	10.5	≥140	7.5	5.0	10.8
HR (95% CI)		1.89 (1.40– 2.55)	1.41 (1.09– 1.83)	2.63 (2.11– 3.28)		2.38 (1.53– 3.71)	1.45 (1.12– 1.87)	2.60 (2.21– 3.07)

Covariates at those indicated for model 1.

BP=blood pressure; CI=confidence interval; HR=hazard ratio; PP=pulse pressure



