



SYSTEMATIC REVIEW

Long term cognitive outcomes of early term (37-38 weeks) and late preterm (34-36 weeks) births: A systematic review [version 1; referees: 1 approved, 2 approved with reservations]

Sarah R. Murray ¹, Susan D. Shenkin², Kirsten McIntosh ³, Jane Lim³, Benjamin Grove⁴, Jill P. Pell⁵, Jane E. Norman ¹, Sarah J. Stock¹

¹Tommy's Centre for Maternal and Fetal Health, MRC Centre for Reproductive Health, University of Edinburgh Queen's Medical Research Institute, Edinburgh, EH16 4TJ, UK

²Geriatric Medicine Unit, Royal Infirmary of Edinburgh and Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, EH16 4TJ, UK

³Royal Infirmary of Edinburgh, University of Edinburgh, Edinburgh, EH16 4TJ, UK

⁴Department of Psychology, University of Edinburgh, Edinburgh, EH16 4T, UK

⁵Section of Public Health, University of Glasgow, Glasgow, G12 8RZ, UK

v1 First published: 17 Oct 2017, 2:101 (doi: [10.12688/wellcomeopenres.12783.1](https://doi.org/10.12688/wellcomeopenres.12783.1))
Latest published: 17 Oct 2017, 2:101 (doi: [10.12688/wellcomeopenres.12783.1](https://doi.org/10.12688/wellcomeopenres.12783.1))

Abstract

Background: There is a paucity of evidence regarding long-term outcomes of late preterm (34-36 weeks) and early term (37-38 weeks) delivery. The objective of this systematic review was to assess long-term cognitive outcomes of children born at these gestations.

Methods: Four electronic databases (Medline, Embase, clinicaltrials.gov and PsycINFO) were searched. Last search was 5th August 2016. Studies were included if they reported gestational age, IQ measure and the ages assessed. The protocol was registered with the International prospective register of systematic reviews (PROSPERO Record [CRD42015015472](https://doi.org/10.1136/2016015015472)). Two independent reviewers assessed the studies. Data were abstracted and critical appraisal performed of eligible papers.

Results: Of 11,905 potential articles, seven studies reporting on 41,344 children were included. For early term births, four studies (n = 35,711) consistently showed an increase in cognitive scores for infants born at full term (39-41 weeks) compared to those born at early term (37-38 weeks) with increases for each week of term (difference between 37 and 40 weeks of around 3 IQ points), despite differences in age of testing and method of IQ/cognitive testing. Four studies (n = 5644) reporting childhood cognitive outcomes of late preterm births (34 – 36 weeks) also differed in study design (cohort and case control); age of testing; and method of IQ testing, and found no differences in outcomes between late preterm and term births, although risk of bias was high in included studies.

Conclusion: Children born at 39-41 weeks have higher cognitive outcome scores compared to those born at early term (37-38 weeks). This should be considered when discussing timing of delivery. For children born late preterm, the data is scarce and when compared to full term (37-42 weeks) did not show any difference in IQ scores.

Open Peer Review

Referee Status:

	Invited Referees		
	1	2	3
version 1 published 17 Oct 2017	 report	 report	 report

- Jennifer M. Walsh**, Murdoch Children's Research Institute, Australia
Royal Women's Hospital, Australia
Royal Children's Hospital, Australia
- Marjolein Kok** , University of Amsterdam, Netherlands
- Nana Hyldig** , University of Southern Denmark, Denmark
Gitte Zachariassen, University of Southern Denmark, Denmark

Discuss this article

Comments (0)

Corresponding author: Sarah R. Murray (sarah.murray@ed.ac.uk)

Author roles: **Murray SR:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Shenkin SD:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **McIntosh K:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation; **Lim J:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Grove B:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Writing – Review & Editing; **Pell JP:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Writing – Review & Editing; **Norman JE:** Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Writing – Review & Editing; **Stock SJ:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

How to cite this article: Murray SR, Shenkin SD, McIntosh K *et al.* **Long term cognitive outcomes of early term (37-38 weeks) and late preterm (34-36 weeks) births: A systematic review [version 1; referees: 1 approved, 2 approved with reservations]** Wellcome Open Research 2017, 2:101 (doi: [10.12688/wellcomeopenres.12783.1](https://doi.org/10.12688/wellcomeopenres.12783.1))

Copyright: © 2017 Murray SR *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Grant information: This work was supported by a Wellcome Trust PhD studentship [104490] to Sarah Murray.

First published: 17 Oct 2017, 2:101 (doi: [10.12688/wellcomeopenres.12783.1](https://doi.org/10.12688/wellcomeopenres.12783.1))

Introduction

Globally, preterm birth rates are rising with 10% of neonates born less than 37 weeks gestation¹. Late preterm births (34–36 weeks) account for three quarters of all preterm births². Early term births (37–38 weeks gestation) have also increased and contribute substantially to an overall decrease in gestational age at delivery. In the US, the average gestational age at delivery has decreased from 40 weeks in 1994 to 39 weeks of gestation in 2004³.

Early term delivery is associated with increased short term adverse physical morbidity, including respiratory distress syndrome, transient tachypnoea of the neonate and ventilator use, as well as an increased risk of infant mortality at 37 weeks compared to full-term delivery^{4–6}. It is for this reason that both the Royal College of Obstetricians and Gynaecologists UK (RCOG)⁷ and the American college of Obstetricians and Gynecologists (ACOG)⁸ endorse the policy of elective birth after 39 weeks in order to reduce the risk of adverse outcome in infants born before full term (39–40 weeks gestation). There is a paucity of evidence regarding the long-term morbidity of this group, in particular the impact on cognitive function. Advanced gestational age is associated with a lower risk of having special educational need at school⁹. Davis *et al.*¹⁰ has also shown that even amongst the weeks of term advanced gestational age is associated with better neurodevelopment as demonstrated by magnetic resonance imaging (MRI). As obstetric efforts worldwide continue to attempt to reduce stillbirth amongst term deliveries, induction of labour at an earlier gestational age is becoming more common, despite the guidance above, and therefore it is imperative to consider the long-term outcomes of deliveries before term to guide clinicians and parents on optimum timing of delivery.

The association between preterm birth and long-term neurological morbidity is better established with the risk increasing with decreasing gestational age, with extremely preterm babies (≤ 26 weeks) having the worst neurological outcomes^{11,12}. The aetiology of this is hypothesized to be due to the disruption of the pathways of dendritic arborisation, synaptogenesis and the thickening of the developing cortex¹³. However, there is less evidence regarding long-term cognitive outcomes of late preterm/early term infants, and given they account for the largest proportion of singleton preterm births more research is necessary. A systematic review of 29,375,675 late preterm infants (34–36 weeks)¹⁴, found increased risks of cerebral palsy (RR 3.1, 95% CI 2.3–4.2) and lower likelihood of finishing school in the late preterm born infants (RR 0.96, 95% CI 0.95–0.97), but we could find no prior reviews on cognitive outcomes for early term births.

The aims of this systematic review are to describe the objectively measured cognitive outcomes in childhood up to the age of 18 years i) within each gestational week of term (37–42 weeks) and ii) of late-preterm (34–36 weeks) compared to term (37–42 weeks) deliveries. The results are necessary for informed decision making regarding timing of delivery.

Methods

This systematic review of the literature was conducted according to the STROBE guidelines¹⁵ and reported according to the recommendations of the Preferred Reporting Items for Systematic

Reviews and Meta-Analyses (PRISMA) guidelines¹⁶ (see [Supplementary File 1](#)). The study protocol was registered with the University of York Centre for Reviews and Dissemination International prospective register of systemic reviews (PROSPERO Record [CRD42015015472](#)). MEDLINE (1946–2016), EMBASE (1947–2016) and PsycINFO (1945–2016) were searched using a search strategy developed and tested in collaboration with a librarian experienced in literature searching ([Supplementary File 2](#)). The searches were supplemented with a manual search through the reference lists of selected primary articles. A forward citation search was performed on all included studies. The first search date was 12th January 2015 and the last search was 5th of August 2016. A subsequent search on Clinicaltrials.gov was performed on 2nd June 2017.

Study selection

One reviewer (JL) screened all titles and abstracts and a second reviewer (BG) independently screened through a 10% sample of the 10,882 articles, by reading the title and abstracts of the first 100 articles of every 1000. The search was updated in August 2016, which yielded an additional 1023 titles and abstracts screened independently by two reviewers (SM, KM). After a consensus was reached, the full texts were retrieved and critically appraised by both reviewers independently (SM, KM). We contacted the individual authors of the included studies to obtain the data necessary to complete the results table. Reasons for exclusion were recorded.

Late-preterm birth was defined as a live birth from 34 to 36+6 completed weeks of gestation. The primary outcome was the results of standardised general intelligence quotient (IQ) tests before age 18 rather than specific domains of cognition. General cognitive ability of a physically and neurologically normal, healthy population of individuals was the key outcome measure recorded. Term birth was defined as a live birth from 37 to 42 completed weeks of gestation.

Studies were included if they reported the range of the participants gestational age, assessment of IQ using a validated score; and the age when IQs was assessed. There were no restrictions by study design, language or method of gestational age assessment. Preterm participants were included as long as there was a clear subgroup of gestational age of 34–36 weeks. Excluded studies included those with: unclear method of cognitive testing; if only selected domains of cognition (e.g. verbal intelligence) were tested; if educational outcomes rather than IQ reported; studies involving high-risk or atypical groups as controls (e.g. multiple births, intra-uterine growth restriction, those with bronchopulmonary dysplasia or brain haemorrhage). Full details of inclusion and exclusion criteria are presented in [Supplementary File 3](#).

The quality of studies was assessed based on the representativeness of the general population, the method of measurement of gestational age and the recording of intelligence testing using the Risk of Bias Assessment Tool for nonrandomized studies (RoBANS) tool¹⁷ ([Supplementary File 4](#)). Two independent reviewers extracted data from each paper on study location, design, population, IQ score used and the main results.

The studies differed widely in outcome measures of cognition used, and due to the large heterogeneity between the study designs and methods a meta-analysis was not possible.

Results

The database and additional record search identified 11,905 articles after removal of duplicates. After exclusions (see flow diagram, [Figure 1](#)), six studies and one conference abstract (which reported on both late preterm and outcomes within term and is therefore included in both groups), reporting on 41,344 children/adolescents, were included in the review; four studies comparing the outcomes within term (37–42 weeks) and three studies comparing the outcomes of late preterm delivery (34–36 weeks) with term delivery.

The studies comparing the outcomes within term differed in a number of ways, including: age of testing (1 year, 4 years and 6 years); method of IQ testing (Bayley Scales of Infant Development (BSID), the Stanford-Binet general IQ test and the Wechsler Abbreviated Scale of Intelligence (WASI); details in [Supplementary File 5](#)); and the categories of gestational age

investigated (37–41 weeks in one study, 37–42 weeks in two studies and 37–43 weeks in one study).

The studies comparing the outcomes of late preterm delivery (34–36 weeks) also differed in a number of ways, including: study design (three prospective cohort studies and one case control study); age of testing (2 years and 13–14 years); and the method of IQ testing (Bayley Scales of Infant Development (BSID) and the Wechsler Intelligence Scale for Children (WISC); details in [Supplementary File 5](#)).

[Table 1](#) provides details on the characteristics of the included studies. Three studies^{18–20} and one study, which was only available as a conference abstract (on contacting the author no further information was available as it is not yet published)²¹, reporting on term deliveries (35,711 children/adolescents) and three studies^{22–24} (as well as the conference abstract) on late preterm deliveries (5,644 children/adolescents).

[Table 2](#) shows the cognitive outcomes of each week of gestation among term births (range 37–43 weeks). In general, although the




Figure 1. Flowchart showing the study selection process (adapted from the PRISMA flow diagram).

Table 1. Characteristics of the included studies, ordered by gestational age categories and age of participant testing.

Author, Year (country)	Study design, total n	GA (n)	Control (n)	Age assessed (yrs)	Exclusions	Exposure measurement	Cognitive outcomes*	Adjustment
Comparison within term (37-42 weeks)								
Espel <i>et al.</i> 2014 ¹⁸ (USA)	Longitudinal prospective study N = 232	37-41 (232)	-	1	Preterm birth (<37 weeks), Post term birth (>41 weeks)	GA measured from LMP And Ultrasound	BSID II assessed by trained clinicians, no blinding	Maternal ethnicity Marital status Fetal Sex Obstetric complications Maternal education Parity Socio-economic status Birth weight percentile
Rose <i>et al.</i> 2013 ¹⁹ (Chile)	Prospective cohort from an RCT N = 1,562	37-42 (1562)	-	1	Infants born at >42 weeks gestation, birth weight <3kg	GA based on LMP	BSID II performed by trained psychologists, blinding not reported	Fetal sex Socio-economic status Birthweight
Gyamfi Bannerman <i>et al.</i> 2014 ²¹ (USA)	Prospective cohort study N = 20,093	37-42 (18,242)		4	Parents with mental retardation or alcohol or drug use	Not reported	Full-scale IQ scores from the Stanford-Binet IQ	Maternal age Maternal education Socio-economic status Smoking
Yang <i>et al.</i> 2009 ²⁰ (Belarus)	Prospective cohort from an RCT N = 13,824	37 (469) 38 (2100) 42 (171) 43 (10)	39-41 (11, 074)	6.5	Birth weight <2500g	GA from hospital records, 94% confirmed by ultrasound	Full-scale IQ from WASI scores performed by trained pediatricians	Maternal age Marital status Maternal education Parity Smoking Maternal height Parental occupation

Author, Year (country)	Study design, total n	GA (n)	Control (n)	Age assessed (yrs)	Exclusions	Exposure measurement	Cognitive outcomes*	Adjustment
Late preterm vs term (34 – 36+6)								
Nepomnyaschy <i>et al.</i> 2011 ²² (USA)	Prospective cohort study N = 5450	34-36 (400)	37-41 (5050)	2	Mothers < 15 yrs, congenital anomalies, multiple births	GA measured by maternal estimated LMP	General, BSID II (converted to a percentile) mental ability, Language by trained interviewer blinded to GA	Maternal ethnicity Maternal age Marital status Fetal sex Obstetric complications Maternal Education Parity Socio-economic status Smoking
Romeo <i>et al.</i> 2015 ²³ (Italy)	Prospective cohort study N = 119	34-36 (71)	37-41 (48)	2	Abnormal cranial Ultrasound, congenital malformations, SGA	Not reported	MDI of BSID II at 2 yrs Blinding and interviewer training not reported	Fetal sex
Narberhaus <i>et al.</i> 2007 ²⁴ (Spain)	Case Control study N = 64	34-36 (11)	37-41 (53)	13-14	Mental or physical disability, perinatal complications/hypoxic risk	Not reported	General WISC full scale assessed by the same psychologist	Matched by: Maternal ethnicity Maternal age Fetal sex Maternal education Socio-economic status

*Cognitive Outcomes (details in [Supplementary File 4](#)): BSID-II = Bayley Scales of Infant Development (2nd Edition); MDI = Mental Developmental Index, PDI = Psychomotor Developmental Index, WISC = Wechsler Intelligence Scale for Children, WASI = Wechsler Abbreviated Scale of Intelligence

Table 2. Results of individual studies comparing cognitive outcomes of children born within term (37–42 weeks gestation), ordered by participant age at cognitive testing.

Study (n)	Cognitive test and overall scores	Age at testing (years)	Main findings			Effect size (95%CI)/significance level	Risk of bias ^a	
				Scores				
Espel <i>et al.</i> 2014 ¹⁸ (n = 232)	BSID-II (Bayley scores of infant development) Range 50-150 MDI and PDI ^e	1	GA ^b	MDI mean ± SD ^c	PDI mean±SD ^d	Gestational age as a continuous variable Adj MDI ^f b = 2.0 (0.45-3.51), p <0.05 Adj PDI ^g b = 3.9 (1.52 - 6.05), p <0.01	Moderate	
			37-38	92±NA ^e	92.5±NA			
			39-40	95±NA	97±NA			
			41	96±NA	106±NA			
Rose <i>et al.</i> 2013 ¹⁹ (n = 1562)	Original BSID (Bayley scores of infant development) MDI and PDI	1	GA (n)	MDI mean ± SD	PDI mean (SD)	Gestational age as a continuous variable Adj MDI ^f b = 0.8 (0.2-1.4), p = 0.025 Adj PDI ^g b = 1.4 (0.6-2.1), p = 0.036	Moderate	
			37 (45)	102.6±11.4	94.4±14.9			
			38 (260)	103.4±12.3	94.4±14.6			
			39 (469)	103.3±12.9	96.6±15.4			
			40 (604)	105.1±11.5	98.5±15.1			
41 (184)	105.4±12.2	97.6±14.8						
Gyamfi-Bannerman <i>et al.</i> 2014 ²¹ (n = 20,093)	Stanford-Binet IQ Full scale intelligence Quotient (IQ) Range 40-160	4	GA (n)	Mean IQ Scores (95%CI)		Test of linear trend P <0.001	High	
			37 (1290)	95.9 (93.6 - 95.3)				
			38 (2361)	97.6 (95.2 - 96.5)				
			39 (4040)	98.6 (97.1 - 98.1)				
			40 (4816)	99.8 (98.1 - 99.0)				
			41 (3782)	99.8 (99.3 - 100.4)				
42 (1953)	98.1 (99.0 - 100.5)							
Yang <i>et al.</i> 2009 ²⁰ (n = 13,824)	WASI (Wechsler Abbreviated Scale of Intelligence) Full scale intelligence quotient (IQ) Range 40–160	6.5	GA (n)		IQ scores	Unadj mean diff (95%CI)	Adj ^h mean diff (95%CI)	Moderate
			37 (469)		NA	-2.6 (-3.7 - -1.4)	-1.7 (-2.7 - -0.7)	
			38 (2100)		NA	0.6 (-1.1 - -0.01)	-0.4 (-1.1 - 0.2)	
			39 (11074)		NA	REF ^k	REF	
			42 (171)		NA	-1.4 (-3.5 - 0.6)	-0.4 (-2.5 - 1.7)	
			43 (10)		NA	-5.8 (-14 - -2.5)	-5.9 (-15 - 3.3)	

^bGA = Gestational age^cMDI = Mental Developmental Index, SD = standard deviation^dPDI = Psychomotor Developmental Index^eNA = not available^{f,g}β coefficients represent the actual change in score with each additional week of term, Adj = adjusted for factors ethnicity, parity, maternal age, obstetric risk, birth weight percentile^hMeans difference coefficients represent the actual change in score with each additional week of term, Adj = adjusted for factors fetal sex, birth weight centile, socio-economic status^kREF = Reference category^lAdjusted for cluster, birth weight, sex, maternal age, maternal height, parental education, and parental occupation

studies differed in age at assessment and the IQ test used, all four studies (18–21) showed an increase in cognitive outcome scores for Infants born at full term (39–41 weeks) compared to those children born at early term (37–38 weeks), with statistically significant increases for each week of term. Three of the studies were classed at moderate risk of bias^{18–20} and one was at high risk because it did not have any clear information on how gestational age was measured²¹. Yang *et al.*²⁰ was the only study to measure outcomes up to post-term (43 weeks gestation) and found an inverse U-shaped relationship of IQ score and gestational age. In this large study (n = 13,824), with a moderate risk of bias, full-term (39–41 weeks) was used as a reference group with mean differences in

IQ scores reported at early term (37–38 weeks) weeks that were lower than full term and also post term (42–43 weeks), which had a higher mean difference from full term than early term (for full risk of bias see [Supplementary File 4](#)). The effect size cannot be summarised due to differences between the studies, but the IQ difference between 37 and 40 weeks was approximately 3 IQ points. This may not be clinically significant at an individual level, but would have an impact at a population level.

Table 3 shows the results of the three studies included reporting childhood cognitive outcomes of late preterm birth (34–36 weeks) compared to term birth (37–42 weeks). The abstract

Table 3. Results of individual studies comparing cognitive outcomes of children born late preterm (34-36 weeks gestation) to term-born infants (37-41 weeks gestation), ordered by participant age at cognitive testing.

Study (n)	Cognitive test and overall scores	Age at testing (years)	Main findings Late preterm Mean±SD	Term Mean±SD	Difference in means (95%CI)/ significance level	Risk of bias
Nepomnyaschy <i>et al.</i> 2011 ²² n = 5,450 (400 late preterm, 5050 term)	BSID-II (Bayley scores of infant development) Bayley short form – mental ability (MDI) Bayley short form –psychomotor ability (PDI) Scores standardised as short form only used (unadjusted range 50-150)	2	MDI 48.9±10 (Range 92.3 – 174.14) PDI 50.2±9.9 (Range 56.43 – 108.53)	50.3±10 49.9±10	Unadj MDI -1.43 (-2.70 - -0.16), p <0.05 Adj MDI -0.35 (-1.52 - 0.83), p >0.10 Unadj PDI -0.38 (-1.61 - 0.85), p >0.10 Adj PDI -0.33 (-1.58 - 0.91), p >0.10	Moderate
Romeo <i>et al.</i> 2015 ²³ n = 119 (71 late preterm, 48 term)	BSID-II (Bayley scores of infant development) MDI only (range 50-150)	2	96.7±9.3	97.1±6.5	p > 0.05	High
Gyamfi-Bannerman <i>et al.</i> 2014 ²¹ n = 20,093 (1,951 late preterm, 18,242 term)	Stanford-Binet IQ+ Full scale intelligence Quotient (IQ) Range 40-160	4	92.5±NA	98.3±NA	NA	High
Naberhaus <i>et al.</i> 2007 ²⁴ n = 64 (11 late preterm, 53 term)	WISC (Wechsler Intelligence Scale for Children) Full scale intelligence quotient Range 40-160	13-14	112.7±13.8	113.6±11.5	p > 0.05	High

¹Risk of bias is a summary using the RoBANS tool, full details in [Supplementary File 4](#)

²MDI = Mental Developmental Index, PDI = Psychomotor Developmental Index

³Coefficient represent the actual change in score associated with being late-preterm versus full term and Adj. = adjusted for race/ethnicity, age, education, marital birth, father co-residence, household residence, household below poverty

⁴Coefficient represents the actual change in score associated with being late-preterm versus full term and Adj. = adjusted for race/ethnicity, age, education, marital birth, father co-residence, household residence, household below poverty

⁵NA = not available

by Gyamfi-Bannerman *et al.*²¹ did not specifically compare late preterm and term deliveries statistically; however, the results were available for each week of gestation and have therefore been recorded in the table. This was the only study that showed a difference in IQ scores between late preterm (mean IQ 92.5) compared to term born children (mean IQ 98.3); however this was not statistically analysed in the published abstract and standard deviations were not available on contacting the authors. This was a large study (n = 20,093), but was assessed as having a high risk of bias as there was no information on the method of gestational age measurement. The two studies using the Bayley scores of infant development did not show a difference in scores between late preterm and term born infants; however this was only done at age 2 and there was no further follow up of the infants. The study by Romeo *et al.*²³ was assessed as having a high risk of bias as there was no mention of how gestational age was measured. The study by Narberhaus *et al.*²⁴ provided the longest follow up of the late preterm-born children, testing IQ using the WISC score (Wechsler Intelligence Scale for Children²⁵) at ages 13–14. No statistically significant difference was found in the IQ scores between late

preterm (mean IQ 112.7, SD [standard deviation] 13.8) and term born children (mean IQ 113.6, SD 11.5). However, these results should be interpreted with caution as the risk of bias was high (no way to determine selective outcome reporting, only some mentioning blinding of outcome assessments and no clear indication of how gestational age was calculated).

Discussion

Main findings

In this systematic review of seven studies (reporting on 41,433 children), the four studies investigating IQ scores within term deliveries found that children born at early term (37–38 weeks) had lower IQ scores at ages one, four and six compared to those born at full term (39–41 weeks). One study (n = 13,824)²⁰ found a decrease in IQ score at >42 weeks. In the four studies comparing late-preterm deliveries (34–36 weeks gestation) to their term counterparts there were no differences in cognitive outcomes at ages two, four and 14. Studies were heterogeneous and several were at high risk of bias, and therefore summary effect sizes cannot be reported. No studies were identified comparing

outcomes between the ages of four and 14. However, it is useful to consider individual study effect sizes and the potential effect on clinical practice. For example, a three point difference in the Stanford-Binet IQ test between children born at 37 weeks and those born at 41 weeks²¹ may not be important at an individual level, but this can have important implications at a population level and should be considered along with other factors (estimated birthweight, obstetric risk factors) when clinicians are discussing timing of delivery with parents.

Strengths and limitations

The strengths of this review include the comprehensive and extensive search strategy, with no language restriction, combined with a detailed pre-defined eligibility criteria for study selection. At the screening stage, to reduce reporter bias, two reviewers independently screened a selected sample to check for accuracy and agreement regarding inclusion of studies. Two reviewers critically appraised all included studies independently. A wide range of cognitive assessments was used in the included studies providing a good overview of the various tests available, but this does make comparison between studies more difficult.

Despite the comprehensive nature of the search, the possibility of missing relevant papers cannot be excluded. We did not have the resources to translate the papers in foreign languages; however we did non-expertly translate to see if any papers fitted the inclusion criteria and none were thought to be relevant. Another limitation was the problems encountered with categorisations of gestational age. A number of studies (22 studies reporting cognitive outcomes of 3,357 infants) only listed <37 weeks of gestational age (all preterm births), which inevitably included those <34 weeks and therefore the whole study was excluded. This may have potentially exacerbated the risk of publication bias as we excluded these studies. Due to limited resources, attempts to contact the individual authors of these studies to see if data was available for 34–36 weeks of gestation was not performed. At delivery, birthweight and gestational age are highly correlated. There is a small but statistically significant correlation between birthweight and cognition in childhood and adulthood (each 1kg increase is associated with 0.13 standard deviation test score increase)²⁶. Some studies did account for birthweight in analyses, and some did not. This may not be appropriate due to their high correlation, and birth weight may be a mediator of the relationship between gestational age and cognition, rather than a confounder. Future studies should report both birth weight and gestational age as a continuous measure, to allow their relative contributions to be measured. Structural equation modelling or similar techniques could be used to model the potential competing causal pathways. We excluded studies with intra-uterine growth restriction (IUGR) because we wanted to study normal healthy singletons, appropriate for gestational age and IUGR may be associated with adverse cognitive outcomes. This review is based on observational data with high levels of between-study heterogeneity, and therefore statistical analysis of the studies was not possible given that the studies were not directly comparable. Limited conclusions can be made regarding the mechanism of action of gestational age on long-term cognitive outcomes because of the nature of the

observational data. There were a number of potential sources of bias across the included studies. Although most studies stated how the participants in the studies were chosen in an attempt to reduce selection bias, it is difficult to determine generalizability of the results outwith the populations that were studied. There was a large variation in the number of confounders (Table 1) adjusted for the various studies, and many did not account for indication for delivery and some did not account for socio-economic factors (strongly associated with cognitive outcomes); therefore there is a risk of residual confounding among the studies.

Interpretation

Comparing the outcomes within the weeks of term (37–42 weeks), this review has shown that cognitive scores in childhood differ throughout the weeks of term delivery and are lowest in those individuals born in early term gestation (37–38 weeks) when measured at ages one, four and six. Although this review specifically set out to look at cognitive outcomes, the findings are in line with those studies of school performance of individuals born within the different weeks of term. Two large population based studies^{27,28} have recently published school outcomes of individuals born within the weeks of term. Smithers *et al.*²⁸ (n = 12,601) showed that children born at 40–41 weeks gestation had the lowest risk of vulnerability at school aged five compared to those born early term (37–39 weeks) or post term (42–45 weeks). Figlio *et al.*²⁷ (n = 1,536,482) showed that children born late term (41 weeks) performed better in school at the age of five through to 18 compared to those born at full term (39 or 40 weeks). Only one of the studies in the review looked at the effect of post-term delivery (>42 weeks) on cognitive outcomes and found those individuals to have a lower score compared to full term (39–40 weeks). This U-shaped relationship has previously been described in the study by MacKay *et al.* (n = 407,503), which found the lowest risk of special educational need at school in those born at 41 weeks gestation compared to those born <41 weeks and >41 weeks⁹. We identified a previous systematic review published in 2015 that also found a reduction in long term cognitive outcomes of children born early term compared to those born full term, but we were unable to reconcile data included in this previous review with source data²⁹.

The mechanism of early term (37–38 week) delivery leading to lower cognitive outcome scores compared to full term deliveries (39–41 weeks) is likely to be multifactorial. Vohr *et al.* described how brain weight increases rapidly in the last trimester of pregnancy with brain weight at 38 weeks 90% of the weight at full term, which may account for the increased vulnerability of early term infants at school⁵. For those born post term (>42 weeks), it is thought the increased vulnerability at school age is due to poorer placental perfusion³⁰.

Only three studies were included in the review comparing the cognitive outcomes of children born late preterm (34–36 weeks) with those born at term (37–41 weeks). There were no (statistically or clinically) significant differences in cognition found between these groups at ages two, four and 13. However, the quality of evidence from these observational studies is poor due to high risk of bias

(high chance of residual confounding, no outcome assessor blinding and no way to ascertain if selective outcome reporting took place). We therefore do not make any clinical recommendations relating to the timing of delivery, as these observational data cannot be used for this purpose. The included studies all used term deliveries as the control group, but as this includes early term (which, as described above, have lower cognitive outcomes than 39+ weeks) there is a possibility that differences between <37 weeks and later were masked. The conference abstract, although it did not specifically compare the results for late preterm versus term delivery, if we compared late preterm deliveries (mean IQ 92.5) with only full term deliveries (39–41 weeks, mean IQ 98.3) the difference is large and provides further evidence of a partial dilution of the results in treating term deliveries as a continuum. This was a large study; however the data was taken from an old cohort study performed between 1959 and 1966, and many of the variables, including method of gestational age measurement, were not available. Three out of four of the studies had a high risk of bias, and they all assumed homogeneity between the term cases, which, as shown above, is not the case. Although a previous systematic review has shown a clear increase in physical morbidity associated with late preterm delivery¹⁴ (34–36 weeks) compared to 37+ weeks, there remains a paucity of evidence regarding long term cognitive outcomes in this group. Future studies should use a full-term delivery group (39–41 weeks) as the control group and adopt uniform gestational age categorizations, ideally with similar outcome measures, to allow for easy comparison between studies. Individual level data should be made available as soon as possible to allow large scale individual participant meta-analysis.

Supplementary material

Supplementary File 1: PRISMA checklist.

[Click here to access the data.](#)

Supplementary File 2: Search strategies.

[Click here to access the data.](#)

Supplementary File 3: Eligibility criteria for study selection.

[Click here to access the data.](#)

Supplementary File 4: Study quality assessed using the RoBANS tool.

[Click here to access the data.](#)

Supplementary File 5: Description of the cognitive tests included in the review.

[Click here to access the data.](#)

Conclusion

Overall, this systematic review has found that children born at full term (39–41 weeks) have the highest cognitive outcome scores compared to those born at early term (37–38 weeks). Given the high prevalence of early term deliveries (the fastest growing proportion of singleton births in the US), small differences at an individual level in cognitive outcomes are likely to have a large impact at a population level. Further research is required to look at the potential reasons for this, and to consider outcomes of late-preterm delivery using a suitable control group of full term (39–41 weeks). The findings from this review have important implications for clinicians and the long-term cognitive outcomes based on gestation at delivery should be discussed with parents regarding optimum timing of delivery.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported by a Wellcome Trust PhD studentship [104490] to Sarah Murray.

Acknowledgements

The authors would like to thank Sheila Fisker for her contribution to the database search strategies and the corresponding authors who provided more information about their studies (Lisa Smithers, David Figlio, Cynthia Gyamfi-Bannerman and Lenna Nepomnyaschy).

References

1. Blencowe H, Cousens S, Oestergaard MZ, *et al.*: **National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications.** *Lancet.* 2012; **379**(9832): 2162–72.
[PubMed Abstract](#) | [Publisher Full Text](#)
2. Blencowe H, Cousens S, Chou D, *et al.*: **Born too soon: the global epidemiology of 15 million preterm births.** *Reprod Health.* BioMed Central; 2013; **10** Suppl 1: S2.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
3. Gyamfi-Bannerman C: **The scope of the problem: the epidemiology of late preterm and early-term birth.** *Semin Perinatol.* 2011; **35**(5): 246–8.
[PubMed Abstract](#) | [Publisher Full Text](#)
4. Reddy UM, Ko CW, Raju TN, *et al.*: **Delivery indications at late-preterm gestations and infant mortality rates in the United States.** *Obstet Gynecol Surv.* 2009; **64**(12): 780–1.
[Publisher Full Text](#)
5. Vohr B: **Long-term outcomes of moderately preterm, late preterm, and early term infants.** *Clin Perinatol.* 2013; **40**(4): 739–51.
[PubMed Abstract](#) | [Publisher Full Text](#)
6. Martínez-Nadal S, Demestre X, Raspall F, *et al.*: **[Neonatal morbidity in early-term newborns].** *An Pediatr (Barc).* 2014; **81**(1): 39–44.
[PubMed Abstract](#) | [Publisher Full Text](#)
7. Royal College of Obstetricians and Gynaecologists: **Greentop Guideline No. 7: Antenatal Corticosteroids to Reduce Neonatal Morbidity and Mortality.** *London: RCOG.* 2010.
[Reference Source](#)
8. American College of Obstetricians and Gynecologists: **ACOG committee opinion no. 559: Cesarean delivery on maternal request.** *Obstet Gynecol.* 2013; **121**(4): 904–907.
[PubMed Abstract](#) | [Publisher Full Text](#)
9. MacKay DF, Smith GC, Dobbie R, *et al.*: **Gestational age at delivery and special educational need: retrospective cohort study of 407,503 schoolchildren.** Lau TK, editor. *PLoS Med.* Public Library of Science; 2010; **7**(6): e1000289.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
10. Davis EP, Buss C, Muftuler LT, *et al.*: **Children's Brain Development Benefits from Longer Gestation.** *Front Psychol.* 2011; **2**: 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. Moore T, Hennessy EM, Myles J, *et al.*: **Neurological and developmental outcome in extremely preterm children born in England in 1995 and 2006: the EPICure studies.** *BMJ.* British Medical Journal Publishing Group; 2012; **345**: e7961.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
12. Costeloe KL, Hennessy EM, Haider S, *et al.*: **Short term outcomes after extreme preterm birth in England: comparison of two birth cohorts in 1995 and 2006 (the EPICure studies).** *BMJ.* British Medical Journal Publishing Group; 2012; **345**: e7976.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
13. Huttenlocher PR, Dabholkar AS: **Regional differences in synaptogenesis in human cerebral cortex.** *J Comp Neurol.* 1997; **387**(2): 167–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
14. Teune MJ, Bakhuizen S, Gyamfi Bannerman C, *et al.*: **A systematic review of severe morbidity in infants born late preterm.** *Am J Obstet Gynecol.* 2011; **205**(4): 374.e1–374.e9.
[PubMed Abstract](#) | [Publisher Full Text](#)
15. von Elm E, Altman DG, Egger M, *et al.*: **The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies.** *J Clin Epidemiol.* 2008; **61**(4): 344–9.
[PubMed Abstract](#) | [Publisher Full Text](#)
16. Moher D, Liberati A, Tetzlaff J, *et al.*: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement.** *Ann Intern Med.* 2009; **151**(4): 264–9, W64.
[PubMed Abstract](#) | [Publisher Full Text](#)
17. Kim SY, Park JE, Lee YJ, *et al.*: **Testing a tool for assessing the risk of bias for nonrandomized studies showed moderate reliability and promising validity.** *J Clin Epidemiol.* 2013; **66**(4): 408–14.
[PubMed Abstract](#) | [Publisher Full Text](#)
18. Espel EV, Glynn LM, Sandman CA, *et al.*: **Longer gestation among children born full term influences cognitive and motor development.** *PLoS One.* 2014; **9**(11): e113758.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
19. Rose O, Bianco E, Martinez SM, *et al.*: **Developmental scores at 1 year with increasing gestational age, 37–41 weeks.** *Pediatrics.* 2013; **131**(5): e1475–81.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
20. Yang S, Platt RW, Kramer MS: **Variation in child cognitive ability by week of gestation among healthy term births.** *Am J Epidemiol.* 2010; **171**(4): 399–406.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
21. Gyamfi-Bannerman C, Son M, Ananth C: **696: Long-term neurodevelopmental outcomes of late preterm and early term neonates.** *Am J Obstet Gynecol.* 2014; **210**(1): S342–3.
[Publisher Full Text](#)
22. Nepomnyaschy L, Hegyi T, Ostfeld BM, *et al.*: **Developmental outcomes of late-preterm infants at 2 and 4 years.** *Matern Child Health J.* 2012; **16**(8): 1612–24.
[PubMed Abstract](#) | [Publisher Full Text](#)
23. Romeo DM, Brogna C, Sini F, *et al.*: **Early psychomotor development of low-risk preterm infants: influence of gestational age and gender.** *Eur J Paediatr Neurol.* 2016; **20**(4): 518–23.
[PubMed Abstract](#) | [Publisher Full Text](#)
24. Narberhaus A, Segarra D, Caldú X, *et al.*: **Gestational age at preterm birth in relation to corpus callosum and general cognitive outcome in adolescents.** *J Child Neurol.* 2007; **22**(6): 761–5.
[PubMed Abstract](#) | [Publisher Full Text](#)
25. Benson DF: **Neuropsychological Assessment By Muriel Lezak.** Oxford University Press: London. 1976. *Psychol Med.* 1978; **8**(02): 347–349.
[Publisher Full Text](#)
26. Grove BJ, Lim SJ, Gale CR, *et al.*: **Birth weight and cognitive ability in adulthood: A systematic review and meta-analysis.** *Intelligence.* 2017; **61**: 146–58.
[Publisher Full Text](#)
27. Figlio DN, Guryan J, Karbownik K, *et al.*: **Long-term Cognitive and Health Outcomes of School-Aged Children Who Were Born Late-Term vs Full-Term.** *JAMA Pediatr.* American Medical Association; 2016; **170**(8): 758–64.
[PubMed Abstract](#) | [Publisher Full Text](#)
28. Smithers LG, Searle AK, Chittleborough CR, *et al.*: **A whole-of-population study of term and post-term gestational age at birth and children's development.** *BJOG.* 2015; **122**(10): 1303–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
29. Chan E, Leong P, Malouf R, *et al.*: **Long-term cognitive and school outcomes of late-preterm and early-term births: a systematic review.** *Child Care Health Dev.* 2016; **42**(3): 297–312.
[PubMed Abstract](#) | [Publisher Full Text](#)
30. Link G, Clark KE, Lang U: **Umbilical blood flow during pregnancy: evidence for decreasing placental perfusion.** *Am J Obstet Gynecol.* 2007; **196**(5): 489.e1–7.
[PubMed Abstract](#) | [Publisher Full Text](#)

Open Peer Review

Current Referee Status:



Version 1

Referee Report 07 December 2017

doi:10.21956/wellcomeopenres.13850.r27478



Nana Hyldig , **Gitte Zachariassen**

Hans Christian Andersens Hospital, Odense University Hospital, University of Southern Denmark, Odense, Denmark

Thank you for the opportunity to review this very well written manuscript on cognitive outcomes of early term and late preterm births. A few thoughts to consider are listed below:

Abstract

1. In the result section seven studies are mentioned – as in table 1 in the manuscript. For early term births – four studies are mentioned – as in the manuscript. For late preterm births – four studies are mentioned – but only three studies are mentioned in the manuscript including table 1?
2. Definitions also in abstract: late preterm (34-36 weeks), early term (37-38 weeks), full term (39-41 weeks) and term (37-40 weeks)? In conclusion “full term” are also defined as “37-42 weeks” – confusing! In the manuscript in Methods and study design “37-42 completed weeks” are defined as “term birth”.

Method section

1. The last search was more than one year ago. We recommend that an updated search before publication.
2. In the attached PRISMA statement document the authors report that the summary measure is reported on page 10. This is, however, not identifiable in the manuscript. Please report the summary measure / statistical method used to calculate the effect estimates in the method section.
3. At PROSPERO the authors state that they will assess risk of bias by using GRADE, but in the manuscript they have used RoBANS.
4. It is very impressive that one author have screened 10,882 titles and abstracts. The last part of the first paragraph in “Study selection” it says that a second author screened 10% of the articles. How was consensus reached for the remaining 90% of the articles?

Discussion

1. Exclusion of studies including infants <34 does not induce risk of publication bias as the studies are published.
2. Only one study have adjusted for obstetric complications (caesarean sections?) and no information on the fetus/infants.
Often late preterm and early term born infants are delivered early due to a sick fetus or a sick mother – both with implication on the fetal brain and post natal development.
3. Why were the infants in the studies delivered late preterm or at early term? How many caesarean sections in these groups compared to full term infants? Maybe these infants have a lower IQ due to

other circumstances than being born at early term?

It is mentioned shortly in discussion section under “the mechanism of early term.....” but need further discussion.

Conclusion in both abstract and manuscript:

...”for discussion with parents regarding optimum timing of delivery.” Do you mean counselling parents who wish or request a caesarean section? Or all parents including those with sick fetuses? Should be specified in the conclusion.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Partly

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

We have read this submission. We believe that we have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however we have significant reservations, as outlined above.

Referee Report 20 November 2017

doi:[10.21956/wellcomeopenres.13850.r27305](https://doi.org/10.21956/wellcomeopenres.13850.r27305)



Marjolein Kok 

Academic Medical Center, Department of Obstetrics and Gynecology, University of Amsterdam, Amsterdam, Netherlands

This is a well written systematic review on long term cognitive outcome of late preterm and early term infants. The strength of this systematic review exposes a weakness at the same time. The detailed pre-defined eligibility criteria of IQ scores limit the number of eligible studies and thus gives an insight in the limitations of current long term follow-up studies. There is no uniformity in report on gestational age, outcome measures and confounding factors, making it difficult to draw conclusions from the aggregated data. The authors are aware of this short-fall and suggest themselves that future studies should use a full-term delivery group (39–41 weeks) as the control group and adopt uniform gestational age categorizations, with similar outcome measures, to allow for easy comparison between studies. Although the results of this study contributes to the knowledge on this subject, the impact is limited due to above mentioned limitations and the fact that the difference in IQ is only 3 points. I agree however with the reviewers that this may not be clinically significant at an individual level, but would have an impact at a population level. I have no suggestions for changes.

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Yes

Competing Interests: No competing interests were disclosed.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Referee Report 20 November 2017

doi:[10.21956/wellcomeopenres.13850.r27265](https://doi.org/10.21956/wellcomeopenres.13850.r27265)



Jennifer M. Walsh ^{1,2,3}

¹ Murdoch Children's Research Institute, Melbourne, Vic, Australia

² Royal Women's Hospital, Melbourne, Vic, Australia

³ Royal Children's Hospital, Melbourne, Vic, Australia

This systematic review attempts to identify differences in IQ scores in 2 comparison groups. The first group are term infants, by comparing early term 37-38 weeks with full term 39-41 weeks' gestation and by identifying differences with each gestational week. The second group compares late preterm infants' IQ scores, gestation 34-36 weeks, with term infants' IQ scores, gestation 37-42 weeks' gestation.

There is clearly a paucity of data in this regard and this study highlights this.

The search criteria were well identified by the researchers. The methods well explained and the results clearly outlined. The aims of the study are clear and very important to publish in order to highlight the importance for clear decisions regarding timing of delivery if that is possible.

The difficulty I have as a clinician, is in the interpretation of the results and I think needs some further qualification in the discussion by the authors.

The study gives us a clear message that term infants are better off being delivered at 39-41 weeks' gestation in order to maximise their IQ potential. However, the biggest difficulty in being able to interpret and extrapolate the results shown, is that we have no idea for the reason for early term birth versus full term birth and if it's different or similar between the groups in these studies. The researchers can only deal with what has been published but have made no comment regarding the indications for delivery and the potential impact that might have on the results if the studies were adjusted for this. Potentially, the infants born 37-38 weeks were electively delivered/induced but more likely is that there were a mix of both clear indications for delivery and elective deliveries and this may have been very different to the 39-41 week

indications for delivery and contribute to the differences found.

Of the 4 articles looking at term infants IQ scores only Espel et al. 2014 (the smallest study) adjusted for obstetric complications. Although clearly there are differences in the IQ scores of the 2 groups and this is important, can this be helped? The only way to truly disentangle this is to look prospectively at term infants delivered electively, which is an almost impossible study to do with the numbers needed to evaluate the outcome measures. Alternatively adjustment for the reason for delivery particularly if there's an indication that can impact on the infant long term should be done.

The preterm data has smaller numbers, the largest study was adjusted for obstetric complications on analysis but similarly the delivery indications are paramount to teasing out the answer to this question. The researchers have clearly highlighted in the discussion that potentially including infants born 37-38 weeks in the comparison group have diluted the results to show no significant difference. Which in itself is interesting. Is being born at 37 weeks as bad as being born at 34 weeks? The studies are generally small making differences difficult to establish but adding the late preterm data from the Gyamfi-Bannerman et al. 2014 study has shown there are likely differences in IQ to be found.

A recent study by Cheong et al.¹ showed that moderate and late preterm (MLPT) children do worse on all domains of assessment using the BSID II compared with term born controls, though moderate preterm children were included (32-33 weeks'), within the MLPT group, there was little evidence of an association between gestational age at birth and neurodevelopment or social-emotional development at age 2. The group acknowledged that there were potential differences between the MLPT infants' in the study compared with other MLPT infants as they were all born in a tertiary setting and hence may have been sicker. The rates of induction and caesarean section are well delineated in the paper.

The question of term IQ differences is a very interesting and worthwhile subject to pursue researching.

References

1. Cheong JL, Doyle LW, Burnett AC, Lee KJ, Walsh JM, Potter CR, Treyvaud K, Thompson DK, Olsen JE, Anderson PJ, Spittle AJ: Association Between Moderate and Late Preterm Birth and Neurodevelopment and Social-Emotional Development at Age 2 Years. *JAMA Pediatr.* 2017; **171** (4): e164805 [PubMed Abstract](#) | [Publisher Full Text](#)

Are the rationale for, and objectives of, the Systematic Review clearly stated?

Yes

Are sufficient details of the methods and analysis provided to allow replication by others?

Yes

Is the statistical analysis and its interpretation appropriate?

Yes

Are the conclusions drawn adequately supported by the results presented in the review?

Partly

Competing Interests: No competing interests were disclosed.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.
