



Knudsen, M. D., Hjartåker, A., Robb, K. A., de Lange, T., Hoff, G. and Berstad, P. (2018) Improving cancer preventive behaviors: a randomized trial of tailored lifestyle feedback in colorectal cancer screening. *Cancer Epidemiology, Biomarkers and Prevention*, 27(12), pp. 1442-1449. (doi:[10.1158/1055-9965.EPI-18-0268](https://doi.org/10.1158/1055-9965.EPI-18-0268))

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/173715/>

Deposited on: 10 January 2019

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

1 **Title page**

2 **Title: Improving cancer preventive behaviors: a randomized trial of tailored lifestyle**

3 **feedback in colorectal cancer screening**

4 Running title: Improving preventive behavior in colorectal cancer screening

5 Markus Dines Knudsen*^{1,2,3}, Anette Hjartåker³, Kathryn A. Robb⁴, Thomas de Lange^{1,5}, Geir
6 Hoff^{1,2,6}, and Paula Berstad¹

7 ¹ Department of Bowel cancer screening, Cancer Registry of Norway, Norway

8 ² Department of Research and Development, Telemark Hospital, Norway

9 ³ Department of Nutrition, Institute of Basic Medical Sciences, University of Oslo, Norway

10 ⁴ Institute of Health and Wellbeing, University of Glasgow, UK.

11 ⁵ Institute of Clinical medicine, University of Oslo, Norway

12 ⁶ Department of Health Management and Health Economics, Institute of Health and Society,

13 University of Oslo, Norway

14 ***Corresponding Author:**

15 Markus Dines Knudsen

16 Ullernchausseen 64, 0379 Oslo, Norway

17 Tlf. +47 22 92 88 88

18 email: markus.knudsen@krefregisteret.no

19 Conflict of interest:

20 The authors declare no potential conflicts of interest

21 Financial support: The Norwegian Ministry of Health and Care Services funded this study
22 (project number: 2014105)

23 Word count: 3053

24 Tables: 4

25 Figures: 1

26 **Abstract:**

27 **Background:** Cancer screening provides an opportunity to increase awareness of cancer
28 preventive lifestyle behaviors such as non-smoking, physical activity, low alcohol consumption
29 and a healthy diet. We tested the effect of standardized, individually-tailored written feedback
30 and a standard leaflet on one-year lifestyle behaviors in a colorectal cancer (CRC) screening
31 setting.

32 **Methods:** Three-thousand-six-hundred-and-forty-two men and women aged 50-74 years invited
33 to sigmoidoscopy screening were randomly assigned to; i) standardized, individually-tailored
34 written feedback (TF); ii) standard leaflet (SL) for cancer preventive lifestyle behaviors; or iii)
35 control. Participants were mailed two self-reported lifestyle questionnaire (LSQ) one year apart.
36 The TF intervention was based on the prescreening LSQ answers. We analyzed differences (with
37 95% confidence intervals (CI)) by comparing prescreening to one-year follow-up of single cancer
38 preventive factors and the number of cancer preventive lifestyle behaviors (range 0-4) between
39 the groups by multivariable logistic regression and ANCOVA analyses.

40 **Results:** One-thousand-and-fifty-four screening participants without neoplastic findings (29% of
41 those invited to screening) were included in the present study. Participants in the TF group
42 increased their number of cancer preventive lifestyle behaviors significantly compared to those in
43 the control group by 0.11 (95% CI 0.02 to 0.19). Overweight/obese individuals in the TF group
44 had a -0.84 kg (95% CI -1.47 to -0.22) larger reduction in body weight compared to the control
45 group.

46 **Conclusions:** Individually-tailored written feedback at sigmoidoscopy screening led to small
47 improvements in cancer preventive behaviors.

48 **Impact:** CRC screening is a suitable setting for increasing awareness of cancer preventive
49 behavior.

50 **Keyword:** lifestyle, behavior, intervention, score, change, prevention, colorectal cancer
51 screening.

52

53

54 **Introduction:**

55 The context of cancer screening provides an opportunity for a teachable moment to increase
56 participants' awareness of cancer prevention with a healthy lifestyle (1). Cancer screening
57 programs have not yet fully utilized this opportunity (2,3). It is particularly important to increase
58 lifestyle awareness at screening for cancers that are closely related to lifestyle such as colorectal
59 neoplasia (4-7), as well as recurrent adenomas (8,9). Raising awareness of the importance of a
60 healthy lifestyle at CRC screening is also particularly relevant in light of evidence that CRC
61 screening participation may reduce participants' motivation to make healthy lifestyle choices
62 (10,11).

63 An automatized written feedback letter delivered in a screening context would be a feasible low-
64 cost strategy for increasing screening participants' awareness of their own lifestyle. Two separate
65 British intervention studies within CRC screening programs have shown beneficial effects of
66 individually-tailored written advice on consumption of fruit and vegetables in screening
67 participants in the short (six weeks) (12) and longer term (six months) (13). Because only long-
68 lasting beneficial lifestyle behaviors may impact chronic disease risk, such intervention effect
69 should be investigated by an extended follow-up.

70 The present study aimed to investigate the effect of i) standardized, individually-tailored written
71 feedback and ii) a standard leaflet for cancer preventive lifestyle on one-year follow-up of
72 lifestyle behaviors in the context of CRC sigmoidoscopy screening.

73 **Materials and Methods**

74 Study design and participants

75 The present study is a sub-study within the Bowel Cancer Screening in Norway (BCSN) trial, a
76 randomized trial piloting a national CRC screening program. The BCSN is carried out in two
77 geographically defined areas in south-eastern Norway, Moss representing a more rural area and
78 Bærum representing a more urban area. Men and women aged 50-74 years are included (14).
79 From November 2014 to September 2015, 3642 individuals invited to sigmoidoscopy were
80 additionally invited to complete a two-page lifestyle questionnaire (LSQ). We sent the
81 questionnaire along with the screening invitation to be completed prior to the screening
82 examination (prescreening LSQ). The individuals were randomized (1:1:1) at invitation based on
83 the unique Norwegian social security number to one of the three groups: i) standardized,
84 individually-tailored written feedback (TF); ii) standard leaflet (SL) for cancer preventive
85 lifestyle; or iii) control. A computer program carried out the randomization automatically. This
86 randomization was blinded to the researchers and designed by the IT developer, following the
87 consort guidelines (supplementary material 4). We mailed a second LSQ to the prescreening
88 responders 12 months after the mailing of the prescreening LSQ. The outcome change in
89 lifestyle was assessed by the follow-up LSQ. A paper version of the LSQ was included in the
90 screening invitation letter. It was also possible to complete the LSQs in an online version
91 available by personal login via a link provided in the invitation. No reminder was sent to non-
92 responders of the questionnaire.

93 Lifestyle questionnaire (LSQ)

94 The LSQ consisted of questions used in previous national surveys (15,16) and the Norwegian
95 Colorectal Cancer Prevention study (11,17). The participants were asked about demographic
96 factors as well as lifestyle behaviors.

97 Demographic factors included ethnicity - dichotomized as native (Norway) or non-native (any
98 other country), marital status - dichotomized as married/cohabiting or non-married/non-

99 cohabiting (or single), education length (primary school, high school, or a minimum of two years
100 at university/college) and working status - dichotomized as working or not working (including
101 retired, unemployed, homemakers and disabled/on rehabilitation).

102 Height was assessed by whole centimeters and weight as whole kilograms.

103 The lifestyle behaviors included smoking status, dichotomized into current smokers (daily and
104 occasional) and non-smoker (former or never smokers). Physical activity (times/week of 30 min
105 of activity) was calculated by adding the responses on frequency to the two questions on
106 physical activity “without sweating or shortness of breath” and “with sweating or getting short of
107 breath”. Frequency ranged from ‘never’ to ‘more than seven times/week’. Consumption of
108 alcoholic beverages (glasses/week) was calculated by frequency of intake multiplied by the
109 number of glasses usually consumed. Consumption of fruit, berries and vegetables was
110 calculated as a sum of reported consumption of 1) fruits and berries, 2) raw vegetables, and 3)
111 boiled vegetables (portions/day). Consumption of red and processed meat for dinner was
112 calculated as a sum of reported frequency consumption of 1) steak, pork chops or similar, 2)
113 hamburgers or other dishes with minced meat, and 3) sausages (portions/week). Six frequency
114 alternatives ranging from ‘seldom/never’ to ‘more than three portions/day’ were provided as
115 response options for the dietary questions.

116 Based on the following factors: smoking habits, physical activity, and consumption of alcoholic
117 beverages, fruit, berries and vegetables we created a scale for the number of cancer preventive
118 lifestyle behaviors (Table 1). The number of cancer preventive lifestyle behaviors ranged from
119 zero to four. Each of the single lifestyle factors was dichotomized to reflect adherence to health
120 recommendations (18-20). Change in weight was used as a separate outcome and not included in
121 the scale for number of cancer preventive lifestyle behaviors. Body mass Index (BMI, kg/m^2)

122 was calculated to identify individuals who were not following the health recommendations on
123 weight (≥ 25 kg/m²).

124

125 Intervention

126 The *control* group did not receive any intervention or information on CRC prevention.

127 One to four weeks after completion of the prescreening LSQ, responders in the *SL* group
128 received the Norwegian Cancer Society's one-page leaflet, "Good habits for a healthier life"
129 with lifestyle advice for low cancer risk (supplementary material 1) by mail. The leaflet was
130 mailed either before or after the screening examination.

131 Similarly, one to four weeks after completion of the prescreening LSQ, responders in the *TF*
132 group received a two-to-three-page letter by mail from the research team with a standardized,
133 individually-tailored written feedback letter based on their answers to the prescreening LSQ. The
134 letter addressed five lifestyle factors; smoking, consumption of alcoholic beverages,
135 consumption of fruit, berries and vegetables, physical activity and body weight. The behaviors
136 reported by the participant were compared to health recommendations. The participant was
137 praised if meeting the recommendations. If the reported behaviors did not meet the
138 recommendations, the individual was encouraged to change their behavior to meet the
139 recommended levels. This could be; "You answered that you rarely or never eat fruit, berries and
140 vegetables. This is less than recommended. The recommendation is to eat at least five
141 servings/day. One serving is approximately 100g. This equals e.g. a small bowl of salad, a carrot
142 or a medium sized fruit". All participants in the *TF* group also received the Norwegian Cancer
143 Society's one-page leaflet (Supplementary material 1 and 2). Subjects in both the *TF* and *SL*
144 groups who reported current smoking additionally received the Norwegian Cancer Society's
145 leaflet "Stop smoking without gaining weight" (supplementary material 3).

146

147 Screening

148 The sigmoidoscopy screening result was defined as positive if one of the following was detected
149 or suspected: 1) any polyp ≥ 10 mm in diameter, 2) any adenoma with villous histology or high-
150 grade dysplasia, 3) ≥ 3 adenomas or 4) cancer. Participants with a positive screening were
151 referred to a follow-up colonoscopy. The final screening result in the present study was one of
152 the following: 1) negative screening, 2) other findings or 3) neoplasia based on the
153 sigmoidoscopy and colonoscopy.

154 Exclusion criteria

155 Participants were excluded from screening due to medical reasons (e.g., severe heart, lung or
156 liver diseases, cancer with life expectancy less than one year), previous CRC, relocating out of
157 the screening municipalities or previous colonoscopy in the last 12 months. Furthermore,
158 participants were excluded from the present study if not completing the prescreening LSQ or if
159 the completion date was not possible to determine. Participants who completed the prescreening
160 LSQ after the screening examination, or who completed the one-year follow-up LSQ <10 or >14
161 months after prescreening LSQ were also excluded. Individuals with any adenomas or cancer
162 findings at screening were excluded from the present study (Figure 1) to minimize potential bias
163 of lifestyle change caused by being diagnosed with adenomas or CRC. This adds comparability
164 between the present and earlier studies, e.g. Robb *et al.* 2010

165

166 Statistical analyses

167 We used t-tests to evaluate the changes in lifestyle between prescreening and one-year follow-up
168 within each group. McNemar's test was used similarly for changes in smoking status. When

169 examining differences in changes in lifestyle variables between the intervention groups and the
170 control group at follow-up, we used an analysis of covariance for the continuous variables, and a
171 multivariable logistic regression model for smoking. The 95% confidence interval (CI) was also
172 calculated. The statistical models were adjusted for sex, age at invitation, education length,
173 working status, ethnicity, marital status, screening center (Moss or Bærum Hospital), and time
174 between completion of prescreening and follow-up questionnaires. The models were additionally
175 adjusted for the prescreening value of the examined variable and prescreening values for weight,
176 and the lifestyle variables; smoking status, level of physical activity, consumption of alcoholic
177 beverages, fruit and vegetables, and red and processed meat and fish. Self-reported chronic
178 disease was not included in the final model, because the preliminary models adjusting for this
179 variable did not differ from the presented results. We conducted analyses stratified by gender. We
180 also compared change in lifestyle between the TF and SL -groups. Furthermore, we completed
181 statistical analyses including only individuals who did not adhere to single health
182 recommendations or who had a number of cancer preventive lifestyle behaviors ≤ 2 at
183 prescreening.

184 *Sample size estimates:* We based the power calculation on predicted difference in self-reported
185 one year change in intake of fruit, berries and vegetables between the standardized, individually-
186 tailored written feedback and the control group, which we expected to be 0.26 (SD 1.53)
187 portions/day (80% power, $P=0.05$) (21). We estimated 540 subjects in each of the three groups to
188 be an adequate number for analyses. We invited 3642 subjects to participate.

189 The primary analyses were based on intention-to-treat, where if missing the one-year follow-up
190 LSQ the values in the prescreening LSQ was carried forward. Similarly, if answered the one-year
191 follow-up LSQ but values were missing, the baseline values were carried forward. Secondary

192 analyses were based on complete case analyses, meaning that individuals with missing one-year
193 follow-up LSQ or missing values in one or several variables were excluded from the analyses.

194 The analyses were carried out using STATA software, version 14.1 (Stata Corp., College
195 Station, Texas, USA).

196 **Results**

197 Overall, 3642 individuals were invited, and 1433 (39%) participated in sigmoidoscopy screening
198 and completed the prescreening LSQ. Of these, 1054 (75%) completed the prescreening LSQ
199 before the screening and had no neoplastic findings: 308 in the TF, 392 in the SL and 354 in the
200 control group (Figure 1). These were used for the primary analyses based on intention to treat.

201 The demographic characteristics of the three groups at prescreening are shown in Table 2. More
202 individuals in the SL group had a high-level education compared to the TF and control group.

203 Table 3 shows lifestyle characteristics at prescreening and one-year changes in the TF, SL and
204 control groups and adjusted differences in lifestyle changes during follow-up. There were no
205 differences in lifestyle factors at prescreening between the groups. Individuals in the SL group
206 reduced their alcohol consumption significantly by -0.54 glasses/week (95% CI -0.94 to -0.14)
207 compared to the control group. Individuals in the TF group increased their number of cancer
208 preventive lifestyle behaviors significantly by 0.11 (95% CI 0.02 to 0.19) compared to the control
209 group (Table 3). Analyses stratified by gender showed that only men in the SL group
210 significantly decreased their consumption of alcoholic beverages compared to controls -0.91
211 glasses/week, (95% CI -1.56 to -0.26). Women in the TF group significantly increased their
212 physical activity by 0.51 times/week (95% CI (0.05 to 1.98) compared to controls. Only men in

213 the SL group significantly increased their number of cancer preventive lifestyle behaviors by 0.13
214 (95% CI 0.01 to 0.24) compared to controls (See supplementary material 5, table 1).

215 Lifestyle characteristics and one-year changes for individuals who did not adhere to single health
216 recommendations and those with a number of cancer preventive lifestyle behavior ≤ 2 at
217 prescreening are shown in Table 4. Among participants with a BMI ≥ 25 kg/m², individuals in the
218 TF group reduced their weight significantly by -0.84 kg (95% CI -1.47 to -0.22) compared to the
219 controls at one-year follow-up. In the SL group, participants with an alcohol consumption higher
220 than recommended had a significant decrease in consumption by
221 -4.98 glasses/week (95% CI -7.83 to -2.13) compared to the controls at one-year follow-up (Table
222 4).

223 When comparing the TF with the SL group, a significantly higher increase in fruit and vegetable
224 intake was observed in the TF group; 0.18 portions/day (95% CI 0.01 to 0.34) (See
225 supplementary material 5, table 2).

226 The proportion of non-responders to the one-year follow-up LSQ differed between the three
227 groups, being 100/308 (32%) in the TF, 90/392 (22%) in the SL and 85/354 (24%) in the control
228 group (Figure 1). Non-responders to the follow-up LSQ were younger (mean age 62.8, 63.9 and
229 62.8 years) compared to the responders (mean age 65.9, 65.2 and 64.7 years) in the TF, SL and
230 control groups, respectively. Prescreening lifestyle variables and screening result did not differ
231 between the follow-up LSQ responders and non-responders. The secondary results by complete
232 case analyses, based on the 779 participants who completed both the prescreening LSQ and the
233 one-year follow-up LSQ showed similar trends as the primary intention-to-treat analyses. The
234 improvement in the number of cancer preventive lifestyle behaviors was some larger in the
235 complete case analyses than in the intention-to-treat analyses (see supplementary material, 5,

236 table 3 and 4). The improvement in fruit and vegetables intake in the TF compared to the control
237 group was significant only in the complete case analyses.

238 **Discussion**

239 In this randomized trial in a CRC screening setting, we found that standardized, individually-
240 tailored written feedback (TF) led to small improvements at one-year follow-up for cancer
241 preventive behaviors among participants with no neoplastic findings. There was a low overall
242 questionnaire response rate at prescreening (39%). The non-response rate to the one-year follow-
243 up LSQ was higher in the TF group (32%) compared to the SL (22%) and control (24%) –groups.
244 However, similar trends were observed for the intention-to-treat analyses (including non-
245 responders to one-year follow-up LSQ) and complete-case analyses (excluding non-responders to
246 one-year follow-up LSQ).

247 To the best of our knowledge, this is the first letter-based lifestyle intervention study with follow-
248 up time longer than six months in a population-based CRC screening context. The few studies on
249 lifestyle intervention in a CRC screening setting have suggested that an individually tailored
250 approach is more effective than delivery of standard leaflets (12,13,22,23). This has been
251 observed as increased consumption of fruit and vegetables in both short (six weeks) (12) and
252 longer term (six months) (13). The short-term trial (six weeks) intervened on fruit and vegetable
253 intake only. That study differed from the present study by including individuals who voluntarily
254 signed up to receive more information about healthy diet (12), while the present study included a
255 random sample of sigmoidoscopy invitees. The six-month-follow-up trial (13) showed that the
256 individually tailored intervention had an effect on fruit and vegetable consumption similar to our
257 study. An intervention trial including only individuals diagnosed with colorectal adenomas at
258 CRC screening found personalized advice letters and face-to-face contact to increase fiber intake

259 after three months (23). Also in a non-screening setting, eight months of telephone counseling
260 and a tailored letter intervention after removal of adenomatous polyps increased physical activity
261 level, reduced intake of red meat and increased an overall score on lifestyle (22). These studies
262 may indicate a higher success of lifestyle interventions in high-risk individuals compared to our
263 population, as earlier suggested (24), although the interventions used were also more intense
264 (face-to-face contact, telephone counseling) than the present study.

265 Some limitations have to be acknowledged in interpreting the present results. We did not have
266 information about the participants' pre-study awareness of lifestyle recommendations or their
267 knowledge of the association between lifestyle and risk of CRC at prescreening. However, this
268 might be a minor problem, as previous studies have not observed any effect of awareness of
269 lifestyle recommendations on change in lifestyle (13). Attitudes to lifestyle change were not
270 assessed in the present study and could be a confounding variable. The findings are only
271 generalizable to those attending CRC screening and completing a LSQ. People attending cancer
272 screening willing to complete questionnaires might be more motivated towards cancer preventive
273 behavior or lifestyle changes than the general population and non-participants. The sample size
274 was smaller than estimated by the power calculations, which resulted in the study being
275 underpowered. We were unable to analyze the independent effect of the standardized,
276 individually-tailored written feedback without the standard leaflet for cancer preventive lifestyle,
277 because these were both sent to the TF and SL group. Furthermore, chance findings cannot be
278 ruled out as a large number of statistical tests were carried out. The results should be interpreted
279 with caution and as indicative findings that should be tested in a fully powered trial.

280 A strength of the present study was the relatively long-term follow-up period. The intervention
281 with a multiple risk factor approach may be more effective on overall lifestyle change compared

282 to an approach with one or two individual lifestyle factors (25,26). Furthermore, being a
283 population-based randomized trial increases the generalizability of the results to population based
284 screening programs.

285 A minimal intervention such as the TF or SL in this study might not be adequate to enhance
286 lifestyle behaviors or reinforce motivation to change lifestyle behaviors over time. However, it
287 may increase awareness of the importance of lifestyle behavior to lower CRC risk when delivered
288 at CRC screening (27). To have an impact on population health, the lifestyle improvement has to
289 be long lasting. To date, there are no intervention trials with longer than one-year follow-up
290 within CRC-screening. Future trials should therefore test the effect of repeated reminders of
291 lifestyle recommendations on lifestyle behavior e.g. by standard leaflets or a smartphone app.
292 Furthermore, it should be investigated if intervention materials should be tailored to gender and
293 educational level. A previous Norwegian study showed that CRC screening increased the
294 occurrences of lifestyle related diseases among individuals with low educational levels but not for
295 people with higher levels of education (28). The present study indicates that the effect of giving a
296 leaflet on healthy behaviors at CRC screening may be almost as effective as standardized,
297 individually-tailored written feedback in promoting favorable lifestyle changes. Future studies are
298 necessary to separate the impact of individual and general feedback.

299 **Conclusion**

300 A low-cost, minimal intervention using standardized, individually-tailored written feedback and a
301 standard leaflet for cancer preventive behaviors given in a CRC screening context led to small
302 improvement in cancer preventive behaviors. The intervention appeared to be most effective in
303 over weight individuals.

304

305 **Abbreviations:**

306 Bowel Cancer Screening in Norway (BCSN), Confidence Intervals (CI), Colorectal Cancer
307 (CRC), standardized, written individually-tailored feedback (TF), lifestyle questionnaire (LSQ),
308 standard leaflet (SL)
309

310 **Declarations**

311 Ethics:

312 The Regional Ethics Committee of South-East Norway approved the study protocol (approval no.
313 2011/1272). Trial registration: <http://www.clinicaltrials.gov>, identifier: NCT 03396029,
314 retrospectively registered 10 of January 2018). The invited, randomized subjects gave their
315 consent for participation by returning the filled prescreening questionnaire.

316

317 Acknowledgments:

318 We would like to thank all the participants and the teams at the screening centers.

319 Authors' contributions:

320 MDK generated the standardized, individually-tailored written feedback letter, collected and
321 analyzed the data and drafted the manuscript. PB participated in collecting data. PB, GH, TdL,
322 KR and AH provided supervision in generating the standardized, individually-tailored written
323 feedback letter and helped to draft the manuscript. MDK, PB, GH, TdL, KR and AH read and
324 approved the final manuscript.

325 .

326

327 **References**

- 328 1. Anderson AS, Craigie AM, Caswell S, Treweek S, Stead M, Macleod M, *et al.* The impact of a
329 bodyweight and physical activity intervention (BeWEL) initiated through a national colorectal
330 cancer screening programme: randomised controlled trial. *Bmj* 2014;348:g1823 doi
331 10.1136/bmj.g1823.
- 332 2. Anderson AS, Mackison D, Boath C, Steele R. Promoting changes in diet and physical activity in
333 breast and colorectal cancer screening settings: an unexplored opportunity for endorsing
334 healthy behaviors. *Cancer prevention research* 2013;6(3):165-72 doi 10.1158/1940-6207.CAPR-
335 12-0385.
- 336 3. Stead M, Caswell S, Craigie AM, Eadie D, Anderson AS, Be WELt. Understanding the potential
337 and challenges of adenoma treatment as a prevention opportunity: insights from the BeWEL
338 formative study. *Preventive medicine* 2012;54(1):97-103 doi 10.1016/j.ypmed.2011.10.017.
- 339 4. World Cancer Research Fund International American Institute for Cancer Research.
340 Continuous Update Project Report. Food, Nutrition, Physical Actevity, and the Prevention of
341 Colorectal Cancer. London, UK: Imperial College London; 2011.
- 342 5. World Cancer Research Fund International/American Institute for Cancer Research. Cancer
343 preventability estimates for diet, nutrition, body fatness, and physical activity.

- 344 [http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-](http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-estimates-diet-nutrition)
345 [estimates-diet-nutrition](http://www.wcrf.org/int/cancer-facts-figures/preventability-estimates/cancer-preventability-estimates-diet-nutrition). Accessed 7 March 2018.
- 346 6. Aleksandrova K, Pischon T, Jenab M, Bueno-de-Mesquita H, Fedirko V, Norat T, *et al*.
347 Combined impact of healthy lifestyle factors on colorectal cancer: a large European cohort
348 study. *BMC medicine* 2014;12(1):168 doi 10.1186/s12916-014-0168-4.
- 349 7. Knudsen MD, de Lange T, Botteri E, Nguyen DH, Evensen H, Steen CB, *et al*. Favorable lifestyle
350 before diagnosis associated with lower risk of screen-detected advanced colorectal neoplasia.
351 *World journal of gastroenterology* 2016;22(27):6276-86 doi 10.3748/wjg.v22.i27.6276.
- 352 8. Botteri E, Crosta C, Bagnardi V, Tamayo D, Sonzogni AM, De Roberto G, *et al*. Predictors of
353 advanced colorectal neoplasia at initial and surveillance colonoscopy after positive screening
354 immunochemical faecal occult blood test. *Dig Liver Dis* 2016;48(3):321-6 doi
355 10.1016/j.dld.2015.11.020.
- 356 9. Sansbury LB, Wanke K, Albert PS, Kahle L, Schatzkin A, Lanza E, *et al*. The effect of strict
357 adherence to a high-fiber, high-fruit and -vegetable, and low-fat eating pattern on adenoma
358 recurrence. *American journal of epidemiology* 2009;170(5):576-84 doi 10.1093/aje/kwp169.
- 359 10. van der Aalst CM, van Klaveren RJ, de Koning HJ. Does participation to screening
360 unintentionally influence lifestyle behaviour and thus lifestyle-related morbidity? *Best Pract*
361 *Res Clin Gastroenterol* 2010;24(4):465-78 doi 10.1016/j.bpg.2010.06.001.
- 362 11. Berstad P, Loberg M, Larsen IK, Kalager M, Holme O, Botteri E, *et al*. Long-term lifestyle
363 changes after colorectal cancer screening: randomised controlled trial. *Gut* 2015;64(8):1268-76
364 doi 10.1136/gutjnl-2014-307376.
- 365 12. Baker AH, Wardle J. Increasing fruit and vegetable intake among adults attending colorectal
366 cancer screening: the efficacy of a brief tailored intervention. *Cancer epidemiology,*
367 *biomarkers & prevention : a publication of the American Association for Cancer Research,*
368 *cosponsored by the American Society of Preventive Oncology* 2002;11(2):203-6.
- 369 13. Robb KA, Power E, Kralj-Hans I, Atkin WS, Wardle J. The impact of individually-tailored lifestyle
370 advice in the colorectal cancer screening context: a randomised pilot study in North-West
371 London. *Preventive medicine* 2010;51(6):505-8 doi 10.1016/j.ypmed.2010.10.002.
- 372 14. Lange T, Randel KR, Schult AL, Knudsen MD, Kirkoen B, Botteri E, *et al*. Sigmoidoscopy and
373 faecal occult blood test - a comparative screening trial. *Tidsskrift for den Norske laegeforening*
374 *: tidsskrift for praktisk medicin, ny raekke* 2017;137(10):727-30 doi 10.4045/tidsskr.16.1031.
- 375 15. Johansson L, Solvoll K, Bjorneboe GE, Drevon CA. Under- and overreporting of energy intake
376 related to weight status and lifestyle in a nationwide sample. *The American journal of clinical*
377 *nutrition* 1998;68(2):266-74.
- 378 16. Norwegian Institute of Public Health. The Oslo Study I and II. 2016
379 <https://www.fhi.no/en/studies/regional-health-studies/the-oslo-study-i-and-ii/>. Accessed 7
380 March 2018.
- 381 17. Larsen IK, Grotmol T, Almendingen K, Hoff G. Lifestyle characteristics among participants in a
382 Norwegian colorectal cancer screening trial. *European journal of cancer prevention : the*
383 *official journal of the European Cancer Prevention Organisation* 2006;15(1):10-9.
- 384 18. Nordic Council of Ministers. *Nordic Nutrition Recommendations, 5th ed*. DK. Nordic Council of
385 Ministers; 2012.
- 386 19. The Norwegian Directory of Health. *Kostråd for å fremme folkehelsen og forebygge kroniske*
387 *sykdommer*. Andvord Grafik AS : 2011..
- 388 20. World Health Organization: *DIET, NUTRITION AND THE PREVENTION OF CHRONIC DISEASES.*
389 *916*. Geneva: WHO; 2003: 149.
- 390 21. Larsen IK, Grotmol T, Almendingen K, Hoff G. Impact of colorectal cancer screening on future
391 lifestyle choices: a three-year randomized controlled trial. *Clinical gastroenterology and*

- 392 hepatology : the official clinical practice journal of the American Gastroenterological
393 Association 2007;5(4):477-83 doi 10.1016/j.cgh.2006.12.011.
- 394 22. Emmons KM, McBride CM, Puleo E, Pollak KI, Clipp E, Kuntz K, *et al.* Project PREVENT: a
395 randomized trial to reduce multiple behavioral risk factors for colon cancer. *Cancer*
396 *epidemiology, biomarkers & prevention* : a publication of the American Association for Cancer
397 Research, cosponsored by the American Society of Preventive Oncology 2005;14(6):1453-9 doi
398 10.1158/1055-9965.EPI-04-0620.
- 399 23. Caswell S, Anderson AS, Steele RJ. Bowel health to better health: a minimal contact lifestyle
400 intervention for people at increased risk of colorectal cancer. *The British journal of nutrition*
401 2009;102(11):1541-6 doi 10.1017/S0007114509990808.
- 402 24. Ebrahim S, Taylor F, Ward K, Beswick A, Burke M, Davey Smith G. Multiple risk factor
403 interventions for primary prevention of coronary heart disease. *The Cochrane database of*
404 *systematic reviews* 2011(1):CD001561 doi 10.1002/14651858.CD001561.pub3.
- 405 25. Emmons KM, McBride CM, Puleo E, Pollak KI, Marcus BH, Napolitano M, *et al.* Prevalence and
406 predictors of multiple behavioral risk factors for colon cancer. *Preventive medicine*
407 2005;40(5):527-34 doi 10.1016/j.ypmed.2004.10.001.
- 408 26. McBride CM, Puleo E, Pollak KI, Clipp EC, Woolford S, Emmons KM. Understanding the role of
409 cancer worry in creating a "teachable moment" for multiple risk factor reduction. *Social*
410 *science & medicine* 2008;66(3):790-800 doi 10.1016/j.socscimed.2007.10.014.
- 411 27. Senore C, Giordano L, Bellisario C, Di Stefano F, Segnan N. Population based cancer screening
412 programmes as a teachable moment for primary prevention interventions. A review of the
413 literature. *Frontiers in oncology* 2012;2:45 doi 10.3389/fonc.2012.00045.
- 414 28. Aas E, Iversen T, Hoff G. The Effect of Education on Health Behavior after Screening for
415 Colorectal Cancer. In: Kristian Bolin BL, Michael Grossman , Dorte Gyrd-Hansen , Tor Iversen ,
416 Robert Kaestner , Jody L. Sindelar, editor. *Human Capital and Health Behavior*. Volume 252017.
417 p 207-42.

418

419

420

Table 1. Number of cancer preventive lifestyle behaviors and cutoff for each cancer preventive behavior		
	Prescreening	Follow-up
Smoking		
Non-smoking	1	1
Smoking	0	0
Physical activity,		
≥ 30 min times/week	1	1
< 30 min times/week	0	0
Alcoholic beverages, mean glasses/week		
≤ 14 for ♂, ≤ 7 for ♀)	1	1
> 14 for ♂, > 7 for ♀)	0	0
Fruits & vegetables, mean portions/day		
≥ 5 a day	1	1
< 5 a day	0	0
Number of cancer preventive lifestyle behaviors	1-4	1-4

421

Table 2. Demographic characteristic.			
A randomized trial of tailored lifestyle feedback in a sub study of the sigmoidoscopy arm of the bowel cancer screening in Norway: a pilot study. TF = standardized, individually-tailored written feedback, SL= standard leaflet and controls N=1054.(Intention-to-treat analyses),			
	TF (n=308)	SL (n=392)	Controls (n=354)
Age			
Mean (SD), years	64.1 (6.9)	64.9 (7.0)	63.8 (6.8)
Time between sigmoidoscopy and one-year follow-up LSQ			
Mean (SD), days	345.6 (29.4)	341.4 (25.6)	346.6 (33.9)
Sex (%)			
Female	53	50	52
Center (%)			
Moss	72	39	74
Bærum	28	61	26
Working status (%)*			
Working	48	47	49
Not working	48	49	49
Marital status (%)*			
Non-married/ non-cohabiting (or single)	19	20	18
Married/cohabiting	79	77	80
Education length (%)*			
Primary school	18	12	16
High school	39	39	43
University/≥2 years at college	41	46	37
Ethnicity (%)*			
Norwegian	92	92	95
Not Norwegian	5	7	3

N=1054, who answered the lifestyle questionnaire at prescreening before sigmoidoscopy and without neoplasia. *The percent might not add up to 100% due to missing data.

422

Table 3. Changes in cancer preventive factors.			
A randomized trial of tailored lifestyle feedback in a sub study of the sigmoidoscopy arm of the bowel cancer screening in Norway: a pilot study. TF standardized, individually-tailored written feedback, SL= standard leaflet and controls. N=1,054 (Intention-to-treat analyses)			
	TF (n=308)	SL (n=392)	Control (n=354)
Non-smoker (%)			
Prescreening	83.4	87.5	83.3
one-year follow up	86.6	88.5	86.8
Change [‡]	ns	ns	ns
Adjusted one-year outcome compared to the controls, odds ratio (95% confidence interval (CI))	2.38 (0.56 to 10.2)	1.85 (0.41 to 8.28)	1.00 (ref)
Weight, mean (kg)			
Prescreening, (S.D)	79.6 (14.7)	78.7 (14.9)	80.8 (15.1)
one-year follow-up, (S.D)	79.6 (14.8)	78.6 (15.0)	80.8 (15.2)
Change, (95% CI)	-0.08 (-0.37 to 0.22)	-0.03 (-0.28 to 0.22)	0.17(-0.07 to 0.41)
Adjusted one-year outcome compared to the controls, (95%CI)	-0.27 (-0.73 to 0.19)	-0.39 (-0.83 to 0.06)	(ref)
Physical activity, mean 30 min times/week			
Prescreening, (S.D)	4.2 (2.8)	4.7 (3.0)	4.1 (2.9)
one-year follow-up, (S.D)	4.2 (2.7)	4.7 (3.0)	4.0 (2.8)
Change, (95% CI)	-0.01 (-0.22 to 0.21)	-0.06 (-0.27 to 0.14)	-0.05 (-0.26 to 0.16)
Adjusted one-year outcome compared to the controls, (95%CI)	0.14 (-0.19 to 0.48)	0.04 (-0.29 to 0.37)	(ref)
Alcoholic beverages, mean glasses/week			
Prescreening, (S.D)	4.2 (15.4)	4.4 (9.1)	3.8 (5.0)
one-year follow-up, (S.D)	4.4 (15.6)	4.0 (5.7)	4.0 (5.2)
Change, (95% CI)	0.23 (-0.12 to 0.57)	-0.41 (-1.18 to 0.35)	0.18 (-0.09 to 0.44)
Adjusted one-year outcome compared to the controls, (95%CI)	-0.27 (-0.68 to 0.14)	-0.54 (-0.94 to -0.14)	(ref)
Fruits & vegetables, mean portions/day			
Prescreening, (S.D)	2.3 (1.3)	2.3 (1.3)	2.2 (1.4)
one-year follow-up, (S.D)	2.4 (1.5)	2.3 (1.3)	2.2 (1.4)
Change, (95% CI)	0.11 (0.00 to 0.23)	-0.04 (-0.12 to 0.05)	0.02 (-0.11 to -0.14)
Adjusted one-year outcome compared to the controls, (95%CI)	0.12 (-0.05 to 0.28)	-0.01 (-0.16 to 0.15)	(ref)
‡ Number of cancer preventive lifestyle behaviors, mean number			
Prescreening (S.D)	2.0 (0.7)	2.1 (0.7)	2.0 (0.7)
one-year follow-up (S.D)	2.1 (0.7)	2.1 (0.6)	2.0 (0.7)
Change (95% CI)	0.02 (-0.04 to 0.09)	-0.03 (-0.08 to 0.03)	-0.04 (-0.10 to 0.02)
Adjusted one-year outcome compared to the controls, (95%CI)	0.11 (0.02 to 0.19)	0.06 (-0.02 to 0.14)	(ref)
Paired t-test was used to test mean changes and 95% confidence intervals (95% CI), [‡] McNemar test was used to test for changes in smoking status, within the groups (TP, SL, control), ns =nonsignificant. Intention-to-treat analyses used.			
* In the adjusted models differences in change of lifestyle between TP vs. control and SL vs. control were tested.			

A logistic regression model was used for smoking and ANCOVA for the other lifestyle variables. The adjusted models were controlled for: age, sex, screening center, ethnicity, marital status, working status, education length, prescreening weight and prescreening value of the dependent variable along with prescreening value of the other lifestyle variables. †† the number of cancer preventive lifestyle behaviors were adjusted for age, sex, screening center, ethnicity, working status, education length, prescreening weight and the prescreening number of cancer preventive lifestyle behaviors.

	TF	SL	Controls
Smokers prescreening N=156	n=51	n=49	n=56
Non-smokers, prescreening, n	0	0	0
Non-smokers, one-year follow-up, n	7	4	6
Change	p=0.02	p=0.13	p=0.03
Adjusted one-year outcome compared to the controls, by logistic regression, odds ratio (95% confides interval (CI))	2.33 (0.31-17.5)	0.50 (0.03-7.55)	1.00 (ref)
Weight, mean (kg), N=620	n=178	n=217	n=225
prescreening, (S.D)	87.5 (12.6)	87.0 (12.1)	88.0 (12.5)
one-year follow-up, (S.D)	87.1 (12.8)	86.9 (12.5)	88.1 (12.6)
Change	-0.40 (-0.85 to 0.04)	-0.14 (-0.51 to 0.23)	0.12 (-0.17 to 0.40)
Adjusted one-year outcome compared to the controls (95% CI)	-0.84 (-1.47 to -0.22)	-0.61 (-1.22 to 0.00)	(ref)
Physical activity, mean 30 min times/week N=743	n=229	n=257	n=257
prescreening, (S.D)	3.1 (1.7)	3.2 (1.6)	2.8 (1.7)
one-year follow-up, (S.D)	3.5 (2.3)	3.5 (2.2)	3.2 (2.2)
Change	0.38 (0.17 to 0.59)	0.32 (0.10 to 0.54)	0.38 (0.19 to 0.58)
Adjusted one-year outcome compared to the controls (95% CI)	0.03 (-0.33 to 0.39)	-0.11 (-0.47 to 0.25)	(ref)
Alcoholic beverages, mean glasses/week N=77	n=22	n=29	n=26
prescreening, (S.D)	11.8 (3.9)	12.7 (5.2)	15.9 (7.8)
one-year follow-up, (S.D)	10.3 (4.3)	9.5 (5.2)	15.6 (8.5)
Change	-1.52 (-3.44 to 0.40)	-3.20 (-5.40 to -1.00)	-0.27 (-1.29 to 0.75)
Adjusted one-year outcome compared to the controls (95% CI)	-0.83 (-3.82 to 2.16)	-4.98 (-7.83 to -2.13)	(ref)
Fruits & vegetables, mean portions/day N=941	n=275	n=350	n=316
prescreening, (S.D)	2.2 (1.1)	2.2 (1.1)	2.1 (1.1)
one-year follow-up, (S.D)	2.3 (1.4)	2.2 (1.1)	2.2 (1.3)
Change	0.13 (0.02 to 0.25)	0.00 (-0.08 to 0.08)	0.08 (-0.03 to 0.19)
Adjusted one-year outcome compared to the controls (95% CI)	0.12 (-0.04 to 0.28)	-0.01 (-0.16 to 0.15)	(ref)
Number of cancer preventive lifestyle behaviors ≤ 2, mean †† N=641	n=192	n=235	n=214
prescreening (S.D)	1.8 (0.4)	1.8 (0.4)	1.7 (0.5)
one-year follow-up, (S.D)	1.9 (0.5)	1.9 (0.5)	1.8 (0.6)
Change	0.11 (0.05 to 0.17)	0.09 (0.03 to 0.15)	0.08 (0.02 to 0.13)
Adjusted one-year outcome compared to the controls (95% CI)	0.08 (-0.00 to 0.17)	0.03 (-0.06 to 0.11)	(ref)

Paired t-test was used to test mean changes and 95% confidence intervals (95% CI), ^bMcNemar test was used to test for changes in smoking status, within the groups (TP, SL, control), ns =nonsignificant. Intention-to-treat analyses used.

* In the adjusted models differences in change of lifestyle between TP vs. control and SL vs. control were tested. A logistic regression model was used for smoking and ANCOVA for the other lifestyle variables. The adjusted models were controlled for: age, sex, screening center, ethnicity, marital status, working status, education length, prescreening value of the dependent variable, prescreening weight along with prescreening value of the other lifestyle variables.

^{††} the number of cancer lifestyle preventive behaviors were adjusted for age, sex, screening center, ethnicity, working status, education length, prescreening weight and the prescreening number of cancer preventive lifestyle behaviors.

424

425 **Figure legends**

426 Figure 1. Flow-chart of participant recruitment and randomization.

427




Figure 1.