



Gadre, V. and Maher, J. (2020) Recurrence of quadratic differentials for harmonic measure. *Mathematical Proceedings of the Cambridge Philosophical Society*, 169(2), pp. 299-305.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/186648/>

Deposited on: 15 May 2019

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

## Recurrence of quadratic differentials for harmonic measure.

BY VAIBHAV GADRE

*School of Mathematics and Statistics, University of Glasgow  
University Place, Glasgow G12 8SQ, UK  
email: Vaibhav.Gadre@glasgow.ac.uk*

AND JOSEPH MAHER

*Department of Mathematics, College of Staten Island, CUNY  
2800 Victory Boulevard, Staten Island, NY 10314, USA  
and Department of Mathematics, 4307 Graduate Center, CUNY  
365 5th Avenue, New York, NY 10016, USA  
email: joseph.maher@csi.cuny.edu*

(Received )

### *Abstract*

We consider random walks on the mapping class group that have finite first moment with respect to the word metric, whose support generates a non-elementary subgroup and contains a pseudo-Anosov map whose invariant Teichmüller geodesic is in the principal stratum of quadratic differentials. We show that a Teichmüller geodesic typical with respect to the harmonic measure for such random walks, is recurrent to the thick part of the principal stratum. As a consequence, the vertical foliation of such a random Teichmüller geodesic has no saddle connections.

---

### 1. Introduction

Let  $S$  be an orientable surface of finite type. Let  $\text{Mod}(S)$  be the mapping class group of orientation preserving diffeomorphisms of  $S$  modulo isotopy. Let  $\mathcal{T}(S)$  be the Teichmüller space of marked conformal structures on  $S$ . The moduli space  $\mathcal{M}(S)$  of Riemann surfaces is the quotient  $\mathcal{T}(S)/\text{Mod}(S)$ .

#### 1.1. Quadratic differentials:

A quadratic differential on a Riemann surface  $X$  homeomorphic to  $S$  is a meromorphic section of the square of the canonical bundle with simple poles at and only at the punctures in  $S$ . By contour integration, a quadratic differential  $q$  defines charts to  $\mathbb{C}$  with transition functions of the form  $z \rightarrow \pm z + c$ . In particular, this defines a singular flat metric on  $S$ . The flat metric is in the same conformal class as the underlying Riemann surface. A quadratic differential is said to be unit area if the flat metric that it defines has area 1. This description also shows how the space of quadratic differentials admits  $SL(2, \mathbb{R})$ -action: the transition functions  $z \rightarrow \pm z + c$  are preserved under the  $SL(2, \mathbb{R})$  action on  $\mathbb{C} = \mathbb{R}^2$ .

Let  $\mathcal{Q}(S)$  be the space of unit area quadratic differentials. The space  $\mathcal{Q}$  can be iden-

tified with the unit cotangent bundle of  $\mathcal{T}(S)$ . We shall write  $\pi$  for the projection map  $\pi: \mathcal{Q}(S) \rightarrow \mathcal{T}(S)$  which sends a quadratic differential to the underlying Riemann surface.

1.2. *Strata of quadratic differentials:*

The space  $\mathcal{Q}(S)$  is stratified by the order of the zeros of the quadratic differential; the *principal stratum*  $\mathcal{Q}_{pr}(S)$  consists of those quadratic differentials whose zeros are all simple. Our results apply to all but finitely many finite type surfaces, and we now describe the exceptions. For a torus with at most one puncture or a sphere with four punctures, the Teichmüller space  $\mathcal{T}(S)$  has complex dimension one, and all quadratic differentials have simple zeros. In these cases the stratification of the space  $\mathcal{Q}(S)$  consists entirely of the single principal stratum; and so all Teichmüller geodesics are principal. In the case of a sphere with at most three punctures,  $\mathcal{T}(S)$  is either empty or a single point, and the mapping class group is finite, and our results do not apply.

For the remainder of this paper we shall assume that we have fixed the surface  $S$ , and so we shall omit it from our notation, and just write  $\mathcal{T}$  for  $\mathcal{T}(S)$ , and so on.

1.3. *Thick parts of strata:*

The  $\epsilon$ -*thick part* of Teichmüller space  $\mathcal{T}$ , which we shall denote  $\mathcal{T}(\epsilon)$ , is the collection of all conformal structures corresponding to hyperbolic metrics in which no simple closed curve has length less than  $\epsilon$ . The complement  $\mathcal{T} \setminus \mathcal{T}_\epsilon$  is called the  $\epsilon$ -thin part of Teichmüller space. The thick part  $\mathcal{T}(\epsilon)$  is mapping class group invariant, and we shall write  $\mathcal{M}(\epsilon)$  for the quotient, which is a subset of moduli space  $\mathcal{M}$  and is called the  $\epsilon$ -thick part of moduli space.

Given a component  $C$  of a stratum of quadratic differentials, a *saddle connection* is an arc embedded in  $S$  that joins a pair of (not necessarily distinct) singularities. Given a quadratic differential  $q \in C$ , the  $q$ -length of a saddle connection is the flat length of a straight (in the flat metric) representative of the arc.

**DEFINITION 1.4.** *The  $\epsilon$ -thick part of  $C$  is the set of all quadratic differentials in  $C$  for which the  $q$ -length of all saddle connections is at least  $\epsilon$ . The principal stratum is connected, and we shall write  $\mathcal{Q}_{pr}(\epsilon)$  for the  $\epsilon$ -thick part of the principal stratum.*

Maskit [Mas85] showed that the existence of a short curve in a hyperbolic metric implies that the curve is short in any compatible unit area flat metric. More precisely, given  $\epsilon > 0$ , there is an  $\epsilon' > 0$ , such that for any hyperbolic metric  $X$  in the thin part  $\mathcal{T} \setminus \mathcal{T}(\epsilon)$ , and for any flat metric in the same conformal class as  $X$ , the length of any curve in the flat metric is at most  $\epsilon'$ , i.e.  $\pi^{-1}(\mathcal{T} \setminus \mathcal{T}(\epsilon)) \subset \mathcal{Q} \setminus \mathcal{Q}(\epsilon')$ . In a flat metric, a simple closed curve  $\alpha$  either has a unique geodesic representative which is a concatenation of saddle connections, or else there is a maximal flat cylinder on  $S$  foliated by parallel closed geodesics. In the latter case, the boundary curves of the cylinder will contain singularities, and hence saddle connections. In either case, the existence of a short curve in the flat metric implies the existence of a short saddle connection. However, the converse need not be true: there may be arbitrarily short saddle connections even though there are no short simple closed curves in the flat metric. In summary, thick part of a stratum of quadratic differentials has a projection into moduli space that is contained in a thick part of moduli space.

By the discussion above, for any  $\epsilon > 0$  there is an  $\epsilon' > 0$  such that  $\pi(\mathcal{Q}_{pr}(\epsilon')) \subset \mathcal{T}(\epsilon)$ .

We remark however, that any point in  $\mathcal{T}$  has a pre-image in  $\mathcal{Q}$  which contains points which do not lie in the thick part of the principal stratum.

1.5. *Recurrent geodesics in a stratum:*

The action on  $C$  of the subgroup of  $SL(2, \mathbb{R})$  consisting of the diagonal matrices defines the Teichmüller flow: given a quadratic differential  $q$  the bi-infinite Teichmüller geodesic  $\gamma$  determined by it is the image of  $q$  under the diagonal subgroup of  $SL(2, \mathbb{R})$ . This also makes it clear that if for any Teichmüller geodesic segment  $\gamma$  if  $q(\gamma_t)$  is in  $C$  for some  $t$  then the entire segment  $\gamma$  is in  $C$ .

Let  $\gamma$  be a bi-infinite Teichmüller geodesic in  $C$ . Given  $\epsilon > 0$ , we say that  $\gamma$  is *forward recurrent* to the  $\epsilon$ -thick part of  $C$  if there exists  $\epsilon > 0$  such that in any unit speed parameterisation of  $\gamma$  there is a sequence  $t_m$  of times with  $t_m \rightarrow \infty$  as  $m \rightarrow \infty$  and  $q(\gamma_{t_m})$  is contained in the  $\epsilon$ -thick part of  $C$ . Similarly, we may define *backward recurrence* for  $\gamma$ . We say  $\gamma$  is *recurrent* in  $C$  if it is both forward and backward recurrent.

1.6. *Principal pseudo-Anosov maps:*

By the Nielsen-Thurston classification, mapping classes are periodic, reducible or pseudo-Anosov. A pseudo-Anosov map  $g$  has a unique invariant Teichmüller geodesic  $\gamma_g$ . Given a point  $X \in \gamma_g$  there is a unique quadratic differential  $q$  at  $X$  in the direction of  $\gamma_g$ . If the invariant Teichmüller geodesic is given by a quadratic differential that lies in the principal stratum, then we say that the pseudo-Anosov map is in the principal stratum.

1.7. *Random walks:*

We consider random walks on the mapping class group  $\text{Mod}(S)$  that have finite first moment with respect to word metric and whose support generates a non-elementary subgroup of  $\text{Mod}(S)$ , i.e. the subgroup generated by the support of the initial distribution contains a pair of pseudo-Anosov maps with distinct stable and unstable measured foliations. In independent work, Maher [Mah11] and Rivin [Riv08] showed that the probability that a random walk gives a pseudo-Anosov map tends to 1 in the length of the sample path, and in particular, the invariant foliations of pseudo-Anosov elements do not contain saddle connections. As a refinement of these results, we showed the following in [GM17], answering a question of Kapovich and Pfaff [KP15]:

**THEOREM 1.8.** [GM17] *Let  $S$  be a connected orientable surface of finite type, whose Teichmüller space  $\mathcal{T}(S)$  has complex dimension at least two. Let  $\mu$  be a probability distribution on  $\text{Mod}(S)$  such that*

- (i)  $\mu$  has finite first moment with respect to  $d_{\text{Mod}}$ ,
- (ii)  $\text{Supp}(\mu)$  generates a non-elementary subgroup  $H$  of  $\text{Mod}(S)$ , and
- (iii) The semigroup generated by  $\text{Supp}(\mu)$  contains a pseudo-Anosov  $g$  such that the invariant Teichmüller geodesic  $\gamma_g$  for  $g$  lies in the principal stratum of quadratic differentials.

*Then, for almost every bi-infinite sample path  $\omega = (w_n)_{n \in \mathbb{Z}}$ , there is positive integer  $N$  such that for all  $n \geq N$  the mapping class  $w_n$  is a pseudo-Anosov map in the principal stratum, that is its invariant Teichmüller geodesic is given by a quadratic differential with simple zeros and poles. Furthermore, almost every bi-infinite sample path determines a unique Teichmüller geodesic  $\gamma_\omega$  with the same limit points as the bi-infinite sample path, and this geodesic also lies in the principal stratum.*

For clarification, the backward random walk is defined with respect to the reflected distribution  $\widehat{\mu}$  defined by  $\widehat{\mu}(a) = \mu(a^{-1})$  for all group elements  $a$ .

In this note, we prove the following recurrence result, answering a further question of Algom-Kfir, Kapovich and Pfaff [AKKP19]:

**THEOREM 1.9.** *Let  $S$  and  $\mu$  satisfy the hypothesis of Theorem 1.8. Then there exists  $\epsilon(S, \mu) > 0$  such that almost every bi-infinite sample path  $\omega = (w_n)_{n \in \mathbb{Z}}$  determines a unique Teichmüller geodesic  $\gamma_\omega$  in the principal stratum of quadratic differentials with the same limit points in  $\text{PMF}(S)$  as  $\omega$ , and moreover  $\gamma_\omega$  is recurrent to the  $\epsilon$ -thick part of the principal stratum.*

Recurrence to the thick part of the moduli space  $\mathcal{M}$  is shown in Kaimanovich-Masur [KM96] and does not require the extra hypothesis that the subgroup generated by  $\text{Supp}(\mu)$  contains a pseudo-Anosov in the principal stratum. With this extra hypothesis, Theorem 1.9 is a finer recurrence statement and implies their result. A consequence of Theorem 1.9 and [Mas92, Theorem 1] is the following refinement of Theorem 1.8.

**COROLLARY 1.10.** *Let  $S$  and  $\mu$  satisfy the hypothesis of Theorem 1.8. Then almost every bi-infinite sample path  $\omega$  determines a unique Teichmüller geodesic  $\gamma_\omega$  in the principal stratum of quadratic differentials with the same limit points as  $\omega$ , and the vertical and horizontal projective measured foliations corresponding to  $\gamma_\omega$  are uniquely ergodic with no vertical and horizontal saddle connections.*

This corollary follows from the fact that if a quadratic differential has a saddle connection which is contained in a leaf of the horizontal or vertical foliations, then the length of this saddle connection tends to zero in one direction along the geodesic, and so the geodesic cannot be recurrent to the thick part of a strata.

As the vertical and horizontal foliations of the quadratic differential determined by  $\gamma_\omega$  are uniquely ergodic, they are equal to the forward and backward limits of  $\gamma_\omega$  in the Thurston boundary  $\text{PMF}$ , and the dual  $\mathbb{R}$ -trees to these foliations are trivalent. Corollary 1.10 also implies that if one passes from measured foliations to measured laminations then the lamination given by  $\gamma_\omega$  are principal i.e., all of their complementary regions are ideal triangles or once-punctured monogons.

The proof of the recurrence result, Theorem 1.9, follows from the fellow traveling discussion in Section 2 below and the ergodicity of the shift map on  $\text{Mod}(S)$ .

Finally, we remark that recent work of Kapovich, Maher, Pfaff and Taylor shows analogous results for the action of  $\text{Out}(F_n)$  on outer space. In [KMPT18] they consider elements of  $\text{Out}(F_n)$  arising from random walks, and shows that they have attracting and repelling trees which are trivalent, while [KMPT19] considers the limiting trees in the boundary of outer space arising from bi-infinite sample paths, and shows that they are also trivalent. In both of these cases, the generic trees are non-geometric, i.e. they are not realized as dual trees to a measured foliation on a 2-complex.

#### 1.11. Acknowledgements

We would like to thank Saul Schleimer for helpful conversations. We thank the referee for comments that helped to improve the paper.

### 2. Fellow traveling and thickness

Let  $\mathcal{Q}_{\text{pr}}$  be the principal stratum of quadratic differentials. Let  $\mathcal{Q}_{\text{pr}}(\epsilon)$  be the set of principal quadratic differentials  $q$  for which every saddle connection  $\beta$  on  $q$  satisfies

$\ell_q(\beta) \geq \epsilon$  in the induced unit area flat metric on  $S$ . We shall write  $\overline{\mathcal{Q}}_{pr}$  for the quotient of  $\mathcal{Q}_{pr}$  by the mapping class group.

A quadratic differential  $q$  determines a Teichmüller geodesic  $\gamma$  in  $\mathcal{T}$ , and we shall write  $\tilde{\gamma}$  for the corresponding image of  $q$  in  $\mathcal{Q}$  under the geodesic flow, which projects down to  $\gamma$ . Given a quadratic differential  $q$ , we shall parameterize the corresponding geodesic by setting  $q(0) = q$  and  $\gamma(0) = \pi(q(0))$ . We shall write  $\gamma_t$  for the point in  $\mathcal{T}$  distance  $t$  along the geodesic in  $\mathcal{T}$ , and  $q(t)$  for the corresponding point in  $\tilde{\gamma}$ , so  $\gamma(t) = \pi(q(t))$ .

We say a Teichmüller geodesic  $\gamma$  is *recurrent in  $\mathcal{M}$  in the forward direction* if there is a compact set  $K$  in  $\mathcal{M}$ , and a sequence of points  $t_n \rightarrow \infty$ , such that  $\gamma(t_n) \in K$ . For any compact set in  $\mathcal{M}$  there is an  $\epsilon > 0$  such that  $K$  is contained in the  $\epsilon$ -thick part of  $\mathcal{M}$ , so recurrent in  $\mathcal{M}$  implies recurrence to  $\mathcal{M}(\epsilon)$  for some  $\epsilon > 0$ . Masur [Mas92] showed that if  $\gamma$  is recurrent in  $\mathcal{M}$ , then  $\gamma$  has a uniquely ergodic vertical foliation. We say a Teichmüller geodesic  $\gamma$  is *recurrent in  $\overline{\mathcal{Q}}_{pr}$  in the forward direction* if there is a compact set  $K$  in  $\overline{\mathcal{Q}}_{pr}$ , and a sequence of points  $t_n \rightarrow \infty$ , such that  $q(t_n) \in K$ . Any compact set in  $\overline{\mathcal{Q}}_{pr}$  is contained in  $\overline{\mathcal{Q}}_{pr}(\epsilon)$  for some  $\epsilon > 0$ , so recurrence in  $\overline{\mathcal{Q}}_{pr}$  implies recurrence to the thick part  $\overline{\mathcal{Q}}_{pr}(\epsilon)$ , for some sufficiently small  $\epsilon$ . Recurrence in  $\overline{\mathcal{Q}}_{pr}$  implies recurrence in  $\mathcal{M}$ , and furthermore recurrence in  $\overline{\mathcal{Q}}_{pr}$  implies that the vertical foliation of  $\gamma$  contains no saddle connections, as the length of a vertical saddle connection tends to zero as  $t \rightarrow \infty$ .

**PROPOSITION 2.1.** *Suppose that a Teichmüller geodesic  $\gamma$ , determined by a quadratic differential  $q_0 \in \mathcal{Q}_{pr}(\epsilon)$ , is recurrent to  $\overline{\mathcal{Q}}_{pr}(\epsilon)$  in both the forwards and backwards directions. Suppose  $\tau_n$  is sequence of Teichmüller geodesic segments that  $R$ -follow travel  $\gamma$  for distance  $d_n$  such that the midpoints  $X_n$  of  $\tau_n$  are within Teichmüller distance  $R$  of  $X_0$  and  $d_n \rightarrow \infty$ . Let  $q_n$  be the quadratic differential at  $X_n$  corresponding to  $\gamma_n$ . Then there exists  $\epsilon' > 0$ , depending on  $q_0$  and  $R$ , and a subsequence  $n_k$  with  $k \in \mathbb{Z}$  such that  $q_{n_k} \in \mathcal{Q}_{pr}(\epsilon')$  as  $k \rightarrow \pm\infty$ .*

*Proof.* As the Teichmüller geodesic  $\gamma$  is recurrent to the thick part  $\overline{\mathcal{Q}}_{pr}(\epsilon)$ , it is also recurrent to a thick part of  $\mathcal{M}(\epsilon_1)$  for some  $\epsilon_1 > 0$ . By work of Masur [Mas92], as the Teichmüller geodesic is recurrent in both directions, this implies that both the vertical and horizontal foliations are uniquely ergodic. As  $\overline{\mathcal{Q}}_{pr}$  is open, we may choose an open neighbourhood  $U$  of  $\{q_t \mid t \in (-R, R)\}$  in  $\overline{\mathcal{Q}}_{pr}$  which is contained in  $\overline{\mathcal{Q}}_{pr}$ , and whose closure  $K = \overline{U}$  is also contained in  $\overline{\mathcal{Q}}_{pr}$ , and is compact. In particular, there is an  $\epsilon_2 > 0$  such that  $K \subset \overline{\mathcal{Q}}_{pr}(\epsilon_2)$ .

By convergence on compact sets, one can pass to a subsequence of  $\tau_n$ 's that converges to bi-infinite Teichmüller geodesic  $\gamma'$  whose vertical and horizontal foliations have intersection number zero with the vertical and horizontal foliations  $(F_s, F_u)$  of  $\gamma$ . Hence, the vertical and horizontal foliations of  $\gamma'$  are also  $F_s$  and  $F_u$ . Since a Teichmüller geodesic with this foliation data has to be unique,  $\gamma' = \gamma$ . In particular, by passing to a subsequence we get that  $q_{n_k} \rightarrow q_0(s)$  for some  $s \in (-R, R)$ . So the tail of the sequence  $q_{n_k}$  must consist of quadratic differentials in  $K \subset \overline{\mathcal{Q}}_{pr}$  and moreover in  $\overline{\mathcal{Q}}_{pr}(\epsilon')$  as  $k \rightarrow \infty$  proving the proposition.  $\square$

Let  $g$  be a pseudo-Anosov map whose invariant Teichmüller geodesic  $\gamma_g$  is in the principal stratum. Also suppose that  $\epsilon$  has been chosen small enough such that  $\gamma_g$  is contained in  $\mathcal{Q}_{pr}(\epsilon)$ .

**PROPOSITION 2.2.** *Given a pseudo-Anosov element  $g$  and a constant  $R$ , there is an  $\epsilon > 0$ , such that if  $\gamma$  is a Teichmüller geodesic which has sequences  $T_n, d_n$  for  $n \in \mathbb{N}$  such that*

- (i)  $T_n, d_n \rightarrow \infty$  as  $n \rightarrow \infty$ , and
- (ii) there are mapping classes  $h_n$  such that the geodesic  $\gamma_n = h_n(\gamma_g)$  has a segment that  $R$ -fellow travels  $\gamma_t$  over the time interval  $(T_n - d_n, T_n + d_n)$ .

Then there is a subsequence  $n_k$  such that  $q_{T_{n_k}} \in \mathcal{Q}_{pr}(\epsilon)$ .

*Proof.* Pulling back by  $h_n^{-1}$ , the sequence of geodesic segments  $g_n = h_n^{-1}(\gamma(T_n - d_n, T_n + d_n))$  satisfy the hypothesis of Proposition 2.1 with respect to the geodesic  $\gamma_g$  which is recurrent by the virtue of being thick. The proposition then follows from Proposition 2.1.  $\square$

### 3. Random walks and recurrence

We recall some terminology and results from [GM17]. For a point  $X \in \mathcal{T}(S)$  and  $r > 0$  let  $B_r(X)$  be the ball of radius  $r$  centred at  $X$ . Let  $\gamma$  be a Teichmüller geodesic. For points  $X$  and  $Y$  on  $\gamma$  let  $\Gamma_r(X, Y)$  be the set of Teichmüller geodesics that pass through  $B_r(X)$  and  $B_r(Y)$ . By work of Rafi [Raf14], if  $X$  and  $Y$  lie in the thick part  $\mathcal{M}(\epsilon)$ , then there is an  $R$ , that depends on  $r$  and  $\epsilon$ , such that every geodesic in  $\Gamma_r(X, Y)$  fellow travels with constant  $R$  the geodesic segment  $[X, Y]$  of  $\gamma$ .

Now let  $g$  be a pseudo-Anosov element in  $\text{Supp}(\mu)$  such that  $\mu^{(j)}(g) > 0$  for some  $j \in \mathbb{N}$  and the invariant Teichmüller geodesic  $\gamma_g$  is in the principal stratum of quadratic differentials. Without loss of generality, we choose a base-point  $X$  on  $\gamma_g$ . Following the proof of [GM17, Theorem 1.1], for all  $k \in \mathbb{N}$  large enough let  $\Omega_k$  be the set of bi-infinite sample paths  $\omega = (w_n)_{n \in \mathbb{Z}}$  such that the sequence  $w_n X$  converges to uniquely ergodic foliations  $F_+$  and  $F_-$  as  $n \rightarrow \infty$  and  $n \rightarrow -\infty$  respectively and the Teichmüller geodesic  $\gamma(F_-, F_+)$  is contained in  $\Gamma_r(g^{-k}X, g^kX)$ .

Let  $\nu$  be the harmonic measure and  $\hat{\nu}$  be the reflected harmonic measure. Let  $\sigma : \text{Mod}^{\mathbb{Z}} \rightarrow \text{Mod}^{\mathbb{Z}}$  be the shift map. Following the proof of [GM17, Theorem 1.1], we get the following result

**PROPOSITION 3.1.** *Let  $S$  and  $\mu$  satisfy the hypothesis of Theorem 1.8. For any large  $k$  and for almost every bi-infinite sample path  $\omega$ , there is a sequence of times  $n_j \rightarrow \infty$  as  $j \rightarrow \infty$  such that  $\sigma^{n_j}(\omega) \in \Omega_k$ .*

Since a countable intersection of full measure sets has full measure we get that

**PROPOSITION 3.2.** *Let  $S$  and  $\mu$  satisfy the hypothesis of Theorem 1.8. For almost every bi-infinite sample path  $\omega$  there is a sequence  $m_k \rightarrow \infty$  as  $k \rightarrow \infty$  such that  $\sigma^{m_k}(\omega) \in \Omega_k$  for all  $k$  large enough.*

Now we get to the proof of the main recurrence result, Theorem 1.9:

*Proof of Theorem 1.9* By Proposition 3.2, for almost every sample path  $\omega = (w_n)$  there exists a sequence  $m_k$  such that  $\gamma_\omega$  fellow travels  $w_{m_k}(\gamma_g)$  between  $w_{m_k}g^{-k}X$  and  $w_{m_k}g^kX$ . Equivalently, the geodesics  $w_{m_k}^{-1}(\gamma_\omega)$  fellow travels  $\gamma_g$  between  $[g^{-k}X, g^kX]$ . The distances  $d_{\mathcal{T}}(g^{-k}X, g^kX)$  form a sequence that tends to infinity as  $k \rightarrow \infty$ . So by Proposition 2.1, a further subsequence of quadratic differentials given by the midpoints of the fellow travelling segments of  $w_{m_k}^{-1}(\gamma_\omega)$  are in  $\overline{\mathcal{Q}}_{pr}(\epsilon)$ . Thus  $\gamma_\omega$  is recurrent to  $\overline{\mathcal{Q}}_{pr}(\epsilon)$ .  $\square$

*Proof of Corollary 1.10* By Theorem 1.9, for almost every sample path  $\omega$  the tracked Teichmüller geodesic  $\gamma_\omega$  is recurrent to the thick part  $\overline{\mathcal{Q}}_{pr}(\epsilon)$ . The projection to moduli space  $\mathcal{M}$  of  $\gamma_\omega$  is then recurrent to the thick part  $\mathcal{M}(\epsilon')$  for some  $\epsilon' > 0$ . By Masur's

theorem [Mas92], the vertical foliation  $F_s$  of  $\gamma_t$  is uniquely ergodic. Moreover, recurrence to  $\overline{Q}_{pr}(\epsilon)$  implies that  $F_s$  has no vertical saddle connections  $\square$

*Acknowledgements.* The first author acknowledges support from the GEAR Network (U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: Geometric structures And Representation varieties”). The second author acknowledges support from the Simons Foundation and PSC-CUNY.

#### References

- [AKKP19] Yael Algom-Kfir, Ilya Kapovich, and Catherine Pfaff, *Stable Strata of Geodesics in Outer Space*, International Mathematics Research Notices, to appear (2019).
- [GM17] Vaibhav Gadre and Joseph Maher, *The stratum of random mapping classes*, Ergodic Theory and Dynamical Systems, to appear (2017).
- [KM96] Vadim A. Kaimanovich and Howard Masur, *The Poisson boundary of the mapping class group*, Invent. Math. **125** (1996), no. 2, 221–264.
- [KP15] Ilya Kapovich and Catherine Pfaff, *A train track directed random walk on  $\text{Out}(F_r)$* , Internat. J. Algebra Comput. **25** (2015), no. 5, 745–798.
- [KMPT18] Ilya Kapovich, Joseph Maher, Catherine Pfaff, and Samuel J. Taylor, *Random outer automorphisms of free groups: Attracting trees and their singularity structures* (2018), available at [arXiv:1805.12382](https://arxiv.org/abs/1805.12382).
- [KMPT19] Ilya Kapovich, Joseph Maher, Catherine Pfaff, and Samuel J. Taylor, *Random trees in the boundary of Outer space* (2019), available at [arXiv:1904.10026](https://arxiv.org/abs/1904.10026).
- [Mah11] Joseph Maher, *Random walks on the mapping class group*, Duke Math. J. **156** (2011), no. 3, 429–468.
- [Mas85] Bernard Maskit, *Comparison of hyperbolic and extremal lengths*, Ann. Acad. Sci. Fenn. Ser. A I Math. **10** (1985), 381–386.
- [Mas92] Howard Masur, *Hausdorff dimension of the set of nonergodic foliations of a quadratic differential*, Duke Math. J. **66** (1992), no. 3, 387–442.
- [Raf14] Kasra Rafi, *Hyperbolicity in Teichmüller space*, Geom. Topol. **18** (2014), no. 5, 3025–3053.
- [Riv08] Igor Rivin, *Walks on groups, counting reducible matrices, polynomials, and surface and free group automorphisms*, Duke Math. J. **142** (2008), no. 2, 353–379.