



Henson, J., Edwardson, C. L., Davies, M. J., Gill, J. M.R., Heaney, L. M., Khunti, K., Ng, L., Sattar, N., Zaccardi, F. and Yates, T. (2020) Physical activity and lipidomics in a population at high risk of type 2 diabetes mellitus. *Journal of Sports Sciences*, 38(10), pp. 1150-1160. (doi: [10.1080/02640414.2020.1744836](https://doi.org/10.1080/02640414.2020.1744836)).

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/205909/>

Deposited on: 16 December 2019

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

1 **Title Page: Physical activity and lipidomics in a population at high risk of type 2**
2 **diabetes mellitus**

3 **Running title – Physical activity intensity and lipidomics**

4

5 Joseph Henson¹, Charlotte L. Edwardson¹, Melanie J. Davies¹, Jason M.R. Gill², Liam M.
6 Heaney³, Kamlesh Khunti⁴, Leong Ng⁵, Naveed Sattar², Francesco Zaccardi¹, Thomas Yates¹

7

8 ¹ NIHR Leicester Biomedical Research Centre, UK and Diabetes Research Centre, College of
9 Life Sciences, University of Leicester, UK.

10 ² Institute of Cardiovascular and Medical Sciences, University of Glasgow, UK.

11

12 ³ School of Sport, Exercise and Health Sciences, Loughborough University, UK.

13

14 ⁴ NIHR Applied Research Collaboration (ARC) East Midlands, UK and Diabetes Research
15 Centre, College of Life Sciences, University of Leicester, UK..

16

17 ⁵ NIHR Leicester Biomedical Research Centre, UK and Department of Cardiovascular
18 Sciences, University of Leicester, UK.

19

20 Main text word count =3538; Number of Tables = 1; Number of Figures = 3; Number of
21 Supplementary Tables =3; Number of references =49

22

23 **Corresponding Author:**

24 Joseph Henson

25 Leicester Diabetes Centre

26 Leicester General Hospital

27 Leicester

28 LE5 4PW

29 UK

30

31 **ORCID iD:** 0000-0002-3898-7053

32 **Email address:** jjh18@le.ac.uk

33 **Tel:** +44 116 258 8624.

34 **Fax:** +44116 258 4053.

35

36

37

38 **Abstract**

39

40 The aim was to investigate how measurements of the lipidome differ according to the level and
41 intensity of physical activity in a population at high risk of type 2 diabetes (T2DM). A targeted
42 metabolomics platform provided quantitative molecular data on lipid species. Linear regression
43 examined the associations between plasma lipid concentrations, particle size and time spent in
44 objectively measured physical activity intensity domains, in increments of 500 counts per
45 minute (cpm) (up to >4500cpm (~>5.6METs)). Results are presented as % difference in the
46 concentration (lower/higher) or particle size (smaller/larger) per 10 minutes of activity within
47 each intensity. 509 participants were included. Time spent in the lowest physical activity
48 intensity domain (<500cpm) was unfavourably associated with VLDL (2%), HDL (-2%) and
49 Apolipoprotein A-1 particle concentrations (-2%) and HDL diameter (-2%). Conversely, time
50 spent in intensities ≥ 1000 cpm were favourably associated with HDL subclass concentrations;
51 with stronger associations seen at moderate intensities (2000-3999cpm (~4.5METs)). For
52 Apolipoprotein-B concentration and VLDL particle concentration and size, a negative
53 association was consistently observed at the highest physical activity intensity only. If these
54 associations are causal, HDL subclasses appear sensitive to light-intensities whereas only the
55 high category of physical activity intensity was consistently associated with VLDL subclasses.

56

57

58 **Keywords:** Metabolomics, lipidomics, physical activity, high risk, accelerometer

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76 **Introduction**

77

78 Previous research has consistently demonstrated that individuals who engage in physical
79 activity, particularly moderate-to-vigorous intensity (MVPA), on a regular basis manifest a
80 myriad of physiological benefits related to lipid metabolism (1). For example, HDL-C
81 (including very large HDL particle concentrations) is generally responsive to physical activity
82 and increases in a dose-dependent manner with increased energy expenditure (2,3).
83 Conversely, physical inactivity (the failure to achieve the minimum activity recommendations
84 for health (4)), and sedentary behaviour (any sitting or reclining activity with low energy
85 expenditure (5)) are each independently associated with an increased risk of cardiovascular and
86 all-cause mortality (6,7), primarily driven by a worsening of atherogenic dyslipidemia, which
87 includes reduced HDL-C and so potentially greater non-HDL-C levels (7). In contrast, the
88 impact of exercise and inactivity on LDL-C, triglycerides and triglyceride rich lipoproteins are
89 less consistent (8).

90

91 Lipidomics is a sub-class of metabolomics focussing on the structure and function of lipids and
92 lipid derivatives (e.g. phospholipids). These molecules may aid in pinpointing the molecular
93 pathways linking health and disease and how they are influenced by lifestyle behaviours, such
94 as physical activity (9). Historically, many studies have focused exclusively upon the
95 metabolite response to exercise training (10,11). More recently, studies have also reported
96 associations in relation to habitual physical activity and sedentary behaviour across multiple
97 metabolite networks (3,12,13).

98

99 However, there has been limited research on lipidomics and physical activity in populations at
100 high risk of chronic disease. This is an important limitation as international recommendations
101 and policies specify that chronic disease prevention strategies should include targeted
102 interventions aimed at the identification and management of high-risk individuals (14,15).
103 Therefore, the importance of sedentary behaviour and physical activity in this group needs
104 to be better understood in order to inform the content and structure of prevention
105 programmes. Moreover, previous investigations have typically categorised sedentary
106 behaviour and physical activity (light, moderate, vigorous) using population-dependent
107 thresholds. Using a broader continuum of intensity categories allows for greater insight into
108 the dose–response relationship between physical activity intensity and health outcomes (16).
109 This is important as previous research has typically focused on MVPA, which occupies a very

110 small fraction of the day, if at all. Conversely, substantial cardiometabolic benefits may be
111 gained from light-intensity activity, particularly in those at high risk of chronic disease (17),
112 which may also represent a more feasible means to increasing overall activity volume.
113 Applying this approach to measurements of the lipidome may allow for greater understanding
114 of how lipid metabolism differs across the precise physical activity intensity spectrum.

115

116 Therefore, the aim of this study was to explore the associations between circulating lipid
117 species and various physical activity intensities in a population at high risk of type 2 diabetes
118 mellitus (T2DM).

119

120 **Materials and Methods**

121

122 **Study population**

123

124 This study reports cross-sectional baseline data from the Walking Away from Diabetes study.
125 Participants were recruited through 10 primary care practices in Leicestershire, UK (18).
126 Individuals at increased risk of impaired glucose regulation (IGR; any combination of impaired
127 glucose tolerance (IGT) and/or impaired fasting glycaemia (IFG) or undiagnosed T2DM) were
128 identified for recruitment using a modified version of the Leicester Risk Score (19). This risk
129 score applies a validated algorithm to routinely collected data within primary care; based on
130 age, sex, BMI, ethnicity, prescribed antihypertensives, and family history of diabetes. Those
131 individuals scoring within the 90th percentile in each practice were invited to take part in the
132 study. This approach has reasonable sensitivity and specificity for identifying participants with
133 IGR (19). Individuals were unaware of their diabetes risk status before entering the study.
134 Those who had previously been diagnosed with T2DM, were currently taking steroids or were
135 unable to take part in any walking were excluded. Written informed consent was obtained from
136 all eligible participants and the study had full ethical and governance approval.

137

138 **Accelerometer derived measures of physical activity**

139

140 All eligible participants were asked to wear an accelerometer, (Actigraph GT3X, Florida,
141 USA), around their waist, for seven consecutive days during waking hours. These
142 accelerometers translate raw accelerations into activity counts. Data were recorded in 15-s
143 epochs and reintegrated into 60-s epoch files for this analysis. Non-wear time was defined as a

144 minimum of 60 minutes of continuous zero counts; days with at least 600 minutes of wear time
145 were considered valid. In order to be included in the analysis, participants required a minimum
146 of any four valid days (20).

147
148 A commercially available data analysis tool (KineSoft version 3.3.76, Kinesoft,
149 Loughborough, UK; www.kinesoft.org) was used to process the accelerometer data. Activity
150 intensity was generated in increments of 500 counts per minute (cpm) from 0 to 4499cpm for
151 each participant who met the inclusion criteria for accelerometer wear time; with the
152 corresponding categories (0-499, 500-999, 1000-1499, 1500-1999, 2000-2499, 2500-2999,
153 3000-3499, 3500-3999, 4000-4499) representing a summation of all included individuals. Any
154 counts above 4500 were amalgamated, due to a lack of power at higher intensities. For
155 descriptive purposes and to aid interpretation, we used the following thresholds to group 500
156 cpm increments into: very low intensities of physical activity or sedentary behaviour
157 (<500cpm); light-intensity physical activity (≥ 500 -<2000cpm) and MVPA (≥ 2000 counts per
158 minute); these thresholds were are broadly comparable to those that have been commonly used
159 in the literature (21-23).

160

161 **Blood sample collection and lipidomics analysis**

162

163 Lipids were measured from Ethylenediaminetetraacetic acid (EDTA) plasma samples, obtained
164 following an overnight fast and avoidance of alcohol and MVPA for 48 hours previously. The
165 level of systemic lipids in the fasting state arise from a broad combination of genetic and
166 lifestyle related factors. As such, the nuclear magnetic resonance (NMR) spectroscopy
167 metabolomics platform provides a comprehensive snapshot of the individual's physiological
168 status as reflected in their systemic metabolism (9).

169

170 Analysis was performed by Nightingale Health (Helsinki, Finland), whose platform and
171 procedures have been described elsewhere (9). Given the fact that the chosen NMR spectra
172 allows significant modelling of lipoprotein subclasses (24), coupled with the previous
173 epidemiological work showing associations between sedentary time, physical activity and
174 cardiovascular outcomes (6,20,25), the targeted focus of our analysis was on lipid species.

175

176 Briefly, plasma samples were analysed using an automated high-throughput NMR workflow,
177 acquiring NMR spectra on either a Bruker AVANCE III 500 MHz or Bruker AVANCE III HD

178 600 MHz spectrometer. Following organic solvent lipid extraction, further NMR spectra were
179 acquired from the lipid extracts on the 600 MHz spectrometer. The initial data processing,
180 including the Fourier transformations to NMR spectra and automated phasing were performed
181 using computers that control the spectrometers. The spectra were then automatically transferred
182 to a centralized server, which performs various automated spectral processing steps, including
183 overall signal check for missing/extra peaks, background control, baseline removal and spectral
184 area-specific signal alignments (9). The spectral information also underwent various
185 comparisons with the spectra of 2 quality control samples; the data for which is also followed
186 and compared in a consecutive manner. For those spectral areas that passed all the quality
187 control steps, regression modelling was performed to produce the quantified molecular data.
188 Individual metabolic measures also underwent various statistical quality control steps and were
189 checked against an extensive database of quantitative molecular data (9). All analyses were
190 conducted by individuals blinded to the participants' identity and physical activity levels. As
191 traditional clinical lipid profile may not fully capture meaningful information with regards to
192 cardiometabolic risk (26), we report the concentration of particles (“number”) within
193 subclasses of VLDL, HDL, IDL and LDL, apolipoprotein concentration (Apolipoprotein-A1
194 (Apo-A1) and Apo B) and the ratio of Apo B to Apo-A1. We also report the mean diameter
195 particle size of VLDL, HDL and LDL.

196

197 **Covariates**

198

199 Information on smoking status and ethnicity was obtained following an interview administered
200 protocol conducted by a healthcare professional. We were also able to adjust for available
201 dietary biomarkers (omega-3 and omega 6 fatty acids) which are reflective of the composition
202 of ingested fatty acids (27) and act as lipid mediators in the inflammatory response (28). In
203 addition, an increasing dietary ratio of omega-3/omega 6 fatty acids has been associated with
204 a higher incidence of obesity, cardiovascular disease (CVD), metabolic syndrome and insulin
205 resistance (28-30). Conversely, diets including high amounts of seafood and fish increase the
206 dietary amount of omega-3 and have been linked to a reduced risk of CVD, T2DM and
207 metabolic syndrome (31,32). .

208

209

210

211

212 **Statistical analysis**

213

214 Linear regression was used to examine the associations between lipid type concentrations,
215 particle size and physical activity intensities. All lipid outcomes were log-transformed,
216 standardised (Z score) and centered (mean =0, standard deviation =1). Any value that was
217 below detection was set to the minimum observed value of the corresponding lipid. Time spent
218 in each of the physical activity intensity increments was entered into models individually
219 because of the high correlation between intensities (Table S1). Models were adjusted for age
220 (continuous), sex (categorical), smoking status (categorical), ethnicity (categorical),, time
221 accelerometer worn (continuous; average minutes per day) and omega-3 and 6 fatty acids
222 (continuous).

223

224 Results are presented as % difference in the lipid variable associated with 10 minutes of each
225 activity within each intensity. Two-tailed *p* values of <0.05 were considered statistically
226 significant. No further adjustment was made for multiple comparisons, therefore data were
227 viewed with caution and in relation to the overall pattern of results. All statistical analyses were
228 conducted using IBM SPSS Statistics v24.0.

229

230 **Results**

231

232 A total of 509 participants had complete lipidomic and accelerometer data (63% of total
233 sample). The main reasons for participants not having complete data was insufficient
234 accelerometer wear time over too few days and insufficient volumes of blood for additional
235 analyses. There was no difference in the proportions of males/females, ethnicity, smoking
236 status or age in those included vs. those excluded. Table 1 displays the characteristics of
237 included study participants.

238

239 **Physical activity**

240

241 The average time spent in each 500cpm intensity banding is shown in Table 1. 82.4% of total
242 accelerometer wear time was spent in the lowest physical activity category (<500cpm),
243 compared with 0.3% in the highest activity category (>4500cpm). Table S1 also displays the
244 correlations between each intensity band.

245 **Lipoprotein concentrations and particle size**

246

247 The associations between lipid sub-type particle concentrations and physical activity intensities
248 (in 500cpm increments, per 10 minutes of activity) are displayed in Figure 1 with associations
249 for apolipoprotein concentration displayed in Figure 2 (corresponding values presented in
250 Tables S2 and S3). Figure 3 displays the associations between particle size and physical activity
251 intensities.

252

253 **Lipoprotein subclass HDL**

254

255 Concentrations of both large and medium HDL particles showed negative associations with
256 time spent in the lowest intensity of physical activity, which is likely to include a significant
257 amount of sedentary behaviour (<500cpm (both -2%; 95% CI= -3% to -1%, per 10 minutes of
258 activity)) (Figure 1A, Table S3). Time spent in physical activity intensities >1000cpm were
259 favourably associated with small and medium HDL subclasses (range = 3%-24%) with results
260 displaying a dose response relationship for medium subclasses up to moderate intensities.
261 Concentration of very large HDL particles were only associated with time spent in the highest
262 intensity of physical activity (>4500cpm).

263

264 **Lipoprotein subclass VLDL**

265

266 Time spent in the lowest physical activity intensity band (<500cpm) was negatively associated
267 with the concentration of very large, large, medium and small VLDL particles (2%; 95% CI=
268 1% to 3%, per 10 minutes of activity) (Figure 2A, Figure 2C; Table S3). For higher intensities
269 of physical activity, the majority of VLDL subclasses were only found to be favourably
270 associated with time spent in the highest intensity category (>4500cpm).

271 **Concentration of IDL and LDL particles**

272

273 There was no association between physical activity intensities and IDL particle concentrations,
274 whereas LDL particle concentrations (small, medium and large) were only associated at 3500-
275 3999cpm (range =10%-11%) (Figure 1D; Table S3).

276

277

278

279 Apolipoproteins

280

281 Apo-A1 was negatively associated with physical activity <500cpm (-2%; 95% CI= -3% to -
282 1%, per 10 minutes of activity; Figure 2). Even low levels of activity (\geq 500cpm) yielded
283 positive associations, with significant results seen up to >4500cpm (range = 3-20%). For Apo
284 B, significant negative associations were seen from moderate (2500-2999cpm) through to the
285 highest physical activity intensity (>4500cpm) (range =-1% to -15%). The ratio of Apo B to
286 Apo-A1 also displayed negative associations as the physical activity intensity increased, with
287 significant results seen at >2500cpm (range = -11% to -23%).

288

289 Lipoprotein particle size

290

291 Time spent in the lowest intensity of physical activity (<500cpm) was significantly associated
292 with a higher mean diameter of VLDL (2% (95% CI = 1% to 3%, per 10 minutes of activity))
293 and lower mean diameter of HDL (-2% (95% CI= -3% to -1%, per 10 minutes of activity))
294 particles (Figure 3; Table S2). As the physical activity intensity increased, there was a dose-
295 response relationship for HDL, with greater intensity associated with a larger particle size,
296 whereas differences in VLDL particle size were observed at the lowest (<500cpm, 500-
297 999cpm, 1000-1499cpm)and highest physical activity intensities (>4500cpm). No associations
298 were seen for the mean diameter of LDL.

299

300 Discussion

301

302 This study highlights the dose-response associations between physical activity intensity and
303 lipid species involved in the underlying pathophysiology of insulin resistance, CVD and
304 physical activity in a population at high risk of T2DM. The most consistent associations were
305 seen in the HDL and VLDL subclass concentrations. Associations between VLDL subclass
306 concentrations and physical activity were consistently only evident at the lowest (<500cpm,
307 approximately <2.6 METs (33)) and highest intensity of physical activity (>4500cpm,
308 approximately >5.6 METs (33)). Conversely, although results for concentrations of very large
309 HDL particles mirrored those for VLDL, those for smaller HDL particles and Apo-A1 showed
310 significant adverse associations with time spent in the lowest category of physical activity
311 (<500cpm) and positive associations across the spectrum of light- and moderate-intensity
312 physical activity. These results suggest that engaging in different intensities of physical activity

313 may result in a differential impact on lipid metabolism, with high intensities of physical activity
314 needed to disrupt the hepatic release of VLDL.

315

316 To our knowledge, this is the first study to specifically investigate the association of a spectrum
317 of intensities of objectively assessed physical activity on the lipidome, with the findings
318 extending previous research using broad categories of physical activity. For example, a
319 previous study in twins reported that compared to inactive individuals, active individuals had
320 a shift towards lower levels of VLDL and higher levels of large and very large HDL (3). Our
321 findings give insight into how physical activity intensity contributes to these observations, with
322 most HDL subclass concentrations and Apo-A1 being sensitive to time spent in lower
323 intensities of physical activity, whereas lower levels of VLDL subclasses and Apo B were
324 consistently only associated with a moderate to high intensity of physical activity. The results
325 for VLDL and Apo B are consistent with previous research suggesting that the intensity of
326 aerobic exercise must surpass that of moderate intensity in order to have a favourable effect on
327 non HDL-lipids, with adaptations largely modulated through glucagon stimulation (34,35).

328

329 For HDL-cholesterol concentrations and Apo A-1 , associations were consistently seen across
330 light and moderate intensities of physical activity up to a threshold of between 2500–3500cpm
331 (approximately 3.7-4.4 METs (33)), after which little additional benefit was observed. This
332 intensity of physical activity is equivalent to walking at ~5km/h and is considered at the lower
333 end of the moderate intensity spectrum (36). Our finding for HDL subclasses is somewhat in
334 agreement with a recently published study of 66 metabolome measures, which found that
335 higher cardiorespiratory fitness, for which moderate intensities of physical activity are an
336 important determinant, was associated with greater concentrations of larger HDL-particles
337 (37). Our findings are also broadly consistent with a meta-analysis of exercise training studies
338 which concluded that duration, and not intensity, is a predictor of the HDL-C response (38). It
339 has also been shown that low-intensity exercise may improve reverse cholesterol transport via
340 the activation of gene transcription variables proliferator-activated receptor gamma
341 (PPARgamma) and liver X receptor alpha (LXRalpha) (39). HDL-cholesterol may also be
342 affected by other physiological processes, such as inflammation (40), which may be influenced
343 by overall volumes of physical activity (41). Therefore, these data suggest that both light-
344 intensity and moderate-intensity physical activity interventions are effective at improving
345 HDL-C concentrations, whereas engaging in higher-intensities of physical activity may not
346 provide additional benefit on HDL-C.

347

348 A further novel finding was that the time spent in moderate intensities of physical activity were
349 associated with larger average HDL-C particle size. Larger HDL particles are hypothesised to
350 be more important in promoting health benefits and thus reducing the risk of CVD (42).
351 Therefore, one of the many mechanisms linking moderate physical activity to cardiometabolic
352 health could be through altering HDL particle size. Previous studies have also reported stronger
353 associations between self-reported physical activity status and greater effects of exercise
354 intervention studies on large HDL compared to smaller particles (43). That noted, we accept
355 that recent trials and genetics have placed a question on the causal link between HDL-C and
356 CVD (44).

357

358 Our findings also extend previous findings by showing that low levels of physical activity
359 (<500cpm), indicative of sedentary behaviour, are detrimentally associated with HDL
360 concentrations (20,45). Interestingly, the time spent below 500cpm was also detrimentally
361 associated with Apo-A1 and the concentration of very large, large, medium and small VLDL
362 particles. VLDLs are substrates for lipoprotein lipase (LPL)-mediated triglyceride removal,
363 with larger VLDL particles carrying more triglycerides than smaller particles and correlating
364 with insulin resistance (46). Although the precise mechanism of sedentary behaviour and
365 (in)activity-induced lipid changes are unclear, muscle LPL regulation is thought to be one of
366 the most sensitive metabolic responses to sedentary behaviour and low-intensity contractile
367 activity and may explain why even small amounts of physical activity appear to confer
368 cardiovascular benefits (47). The mechanistic relevance of LPL to sedentary behaviour has
369 been demonstrated in animal models (48) whereas in humans moderate intensity activity was
370 shown to increase the affinity of large VLDL particles for LPL clearance (49). However,
371 further insight is needed into the precise impact of increased sedentary time and reduced
372 physical activity on LPL activity.

373

374 Strengths and limitations

375

376 Strengths of our study include the objective measurement of physical activity and examination
377 of lipids in relation to different characteristics across a range of physical activity intensities.
378 By enabling identification of the minimum intensity at which benefits may occur as well as a
379 quantifiable dose–response relationship, this information may aid in generating hypotheses to
380 be tested in future physical activity interventions. Furthermore, our targeted metabolomic

381 platform covered a wide variety of lipids with known identity and quantitative measurements.
382 Our results are strengthened by the fact that associations with all lipoprotein subclasses were
383 present after adjusting for dietary biomarkers (omega-3 and omega 6), supporting an
384 independent association of physical (in) activity *per se* on the lipoprotein subclass profile.
385 These results are in agreement with previous studies which have shown that significant changes
386 in HDL and VLDL concentrations and particle size after exercise training are independent of
387 diet (43). Our study is also accompanied by important limitations. For example, despite
388 individuals spending a reasonable amount of time in moderate activity, the time spent in higher,
389 more vigorous intensity activities is limited. However, this is likely reflective of the habitual
390 behaviour of the majority of individuals at high risk of T2DM. This coupled with the fact that
391 our analysis is observational, means that we cannot prove biological mechanisms or
392 demonstrate causality; reverse causality is also a possibility whereby those with a greater
393 burden of risk factors may be less likely to engage in greater volumes or intensities of physical
394 activity. The high risk nature of the cohort, where higher relative exercise intensities can be
395 anticipated for a given exercise compared to a healthy population, may also affect the
396 interpretation of the intensity thresholds used for this study. However, this is unlikely to affect
397 the interpretation for HDL-cholesterol, where associations were seen across sedentary time and
398 the lower intensity spectrums. Furthermore, despite adjusting for a range of potential
399 cofounders, residual confounding or confounding from unmeasured factors remains a
400 possibility (e.g. alcohol intake). Finally, although accelerometers allow for more robust
401 assessments of physical activity compared to self-report, they are not without limitations. They
402 rely on categorising movement (acceleration) strength, rather than directly distinguishing
403 between postures or modes of physical activity.

404

405 In conclusion, our data suggests potential differences in the associations between different
406 physical activity intensities and the lipidome in subjects with a high risk of T2DM, with most
407 HDL subclass and Apo A-1 concentrations appearing sensitive to light-intensities of physical
408 activity. Although structured physical activity should remain a strong focus and end point of
409 behavioural interventions, lipid related benefits may be gained through light-intensity activity
410 (whilst also reducing sedentary time). Given the limited time spent in higher intensity activities
411 in this population, this may also be the option that is best tolerated in those at high risk of
412 chronic disease. This is particularly pertinent as they are also representative of those likely to
413 be identified as being at high risk of T2DM within routine care and referred onto available
414 prevention programmes. Therefore, future interventions that encourage increases in physical

415 activity, may need to be tailored to individual characteristics and tolerability. In particular,
416 consideration should be given to the relative intensity of physical activity prescribed, as the
417 absolute values will differ considerably between individuals. The results of this analysis also
418 highlight the fact that more work is needed to elucidate the mechanisms by which different
419 physical activity intensities, particularly at the lower end of the spectrum, impact health.

420

421 **Acknowledgements**

422

423 The Walking Away trial was funded by The National Institute for Health Research (NIHR)
424 Collaboration for Leadership in Applied Health Research and Care for Leicestershire,
425 Northamptonshire and Rutland (NIHR CLAHRC – LNR) and East Midlands (NIHR CLAHRC
426 EM). The research was further supported by and the NIHR Leicester Biomedical Research
427 Centre, which is a partnership between University Hospitals of Leicester NHS Trust,
428 Loughborough University and the University of Leicester and the University of Leicester
429 Clinical Trials Unit. The views expressed are those of the author(s) and not necessarily those
430 of the NHS, the NIHR or the Department of Health. Clinical Trials Registration:
431 ClinicalTrials.gov NCT: NCT00941954. The authors would also like to the participants for
432 taking the time to take part. No financial disclosures were reported by the authors of this paper.

433

434 **Conflict of Interest**

435

436 The authors declare no conflict of interest

437

438 **Data availability**

439

440 The datasets generated during and/or analysed during the current study are not publicly
441 available but are available from the corresponding author upon reasonable request.

442

443

444

445

446

447

448 **References**

449

450 (1) Myers J. Cardiology patient pages. Exercise and cardiovascular health. *Circulation* 2003
451 Jan 7;107(1):e2-5.

452 (2) Kelley GA, Kelley KS. Aerobic exercise and HDL2-C: a meta-analysis of randomized
453 controlled trials. *Atherosclerosis* 2006 Jan;184(1):207-215.

454 (3) Kujala UM, Makinen VP, Heinonen I, Soininen P, Kangas AJ, Leskinen TH, et al. Long-
455 term leisure-time physical activity and serum metabolome. *Circulation* 2013 Jan
456 22;127(3):340-348.

457 (4) Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT, et al. Effect of
458 physical inactivity on major non-communicable diseases worldwide: an analysis of burden of
459 disease and life expectancy. *Lancet* 2012 Jul 17.

460 (5) Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, et al.
461 Sedentary Behavior Research Network (SBRN) - Terminology Consensus Project process
462 and outcome. *Int J Behav Nutr Phys Act* 2017 Jun 10;14(1):75-017-0525-8.

463 (6) Wilmot EG, Edwardson CL, Achana FA, Davies MJ, Gorely T, Gray LJ, et al. Sedentary
464 time in adults and the association with diabetes, cardiovascular disease and death: systematic
465 review and meta-analysis. *Diabetologia* 2012 Nov;55(11):2895-2905.

466 (7) Lakka TA, Laaksonen DE. Physical activity in prevention and treatment of the metabolic
467 syndrome. *Appl Physiol Nutr Metab* 2007 Feb;32(1):76-88.

468 (8) Wang Y, Xu D. Effects of aerobic exercise on lipids and lipoproteins. *Lipids Health Dis*
469 2017 Jul 5;16(1):132-017-0515-5.

470 (9) Soininen P, Kangas AJ, Wurtz P, Suna T, Ala-Korpela M. Quantitative serum nuclear
471 magnetic resonance metabolomics in cardiovascular epidemiology and genetics. *Circ*
472 *Cardiovasc Genet* 2015 Feb;8(1):192-206.

473 (10) Daskalaki E, Easton C, G Watson D. The application of metabolomic profiling to the
474 effects of physical activity. *Current Metabolomics* 2014;2(4):233-263.

475 (11) Heaney LM, Deighton K, Suzuki T. Non-targeted metabolomics in sport and exercise
476 science. *J Sports Sci* 2017 Mar 27:1-9.

477 (12) Xiao Q, Moore SC, Keadle SK, Xiang YB, Zheng W, Peters TM, et al. Objectively
478 measured physical activity and plasma metabolomics in the Shanghai Physical Activity
479 Study. *Int J Epidemiol* 2016 Apr 12.

480 (13) Floegel A, Wientzek A, Bachlechner U, Jacobs S, Drogan D, Prehn C, et al. Linking
481 diet, physical activity, cardiorespiratory fitness and obesity to serum metabolite networks:
482 findings from a population-based study. *Int J Obes (Lond)* 2014 Nov;38(11):1388-1396.

- 483 (14) Chatterton H, Younger T, Fischer A, Khunti K, Programme Development Group. Risk
484 identification and interventions to prevent type 2 diabetes in adults at high risk: summary of
485 NICE guidance. *BMJ* 2012 Jul 12;345:e4624.
- 486 (15) Paulweber B, Valensi P, Lindstrom J, Lalic NM, Greaves CJ, McKee M, et al. A
487 European evidence-based guideline for the prevention of type 2 diabetes. *Horm Metab Res*
488 2010 Apr;42 Suppl 1:S3-36.
- 489 (16) Jelleyman C, Edwardson CL, Henson J, Gray LJ, Rowlands AV, Khunti K, et al.
490 Associations of Physical Activity Intensities with Markers of Insulin Sensitivity. *Med Sci*
491 *Sports Exerc* 2017 Jul 19.
- 492 (17) Chastin SFM, De Craemer M, De Cocker K, Powell L, Van Cauwenberg J, Dall P, et al.
493 How does light-intensity physical activity associate with adult cardiometabolic health and
494 mortality? Systematic review with meta-analysis of experimental and observational studies.
495 *Br J Sports Med* 2019 Mar;53(6):370-376.
- 496 (18) Yates T, Edwardson CL, Henson J, Gray LJ, Ashra NB, Troughton J, et al. Walking
497 Away from Type 2 diabetes: a cluster randomized controlled trial. *Diabet Med* 2017
498 May;34(5):698-707.
- 499 (19) Gray LJ, Davies MJ, Hiles S, Taub NA, Webb DR, Srinivasan BT, et al. Detection of
500 impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic
501 data, in a multiethnic UK community setting. *Diabetologia* 2012 Apr;55(4):959-966.
- 502 (20) Henson J, Yates T, Biddle SJ, Edwardson CL, Khunti K, Wilmot EG, et al. Associations
503 of objectively measured sedentary behaviour and physical activity with markers of
504 cardiometabolic health. *Diabetologia* 2013 May;56(5):1012-1020.
- 505 (21) Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and
506 Applications, Inc. accelerometer. *Med Sci Sports Exerc* 1998 May;30(5):777-781.
- 507 (22) Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity
508 in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008 Jan;40(1):181-
509 188.
- 510 (23) Gorman E, Hanson HM, Yang PH, Khan KM, Liu-Ambrose T, Ashe MC.
511 Accelerometry analysis of physical activity and sedentary behavior in older adults: a
512 systematic review and data analysis. *Eur Rev Aging Phys Act* 2014;11:35-49.
- 513 (24) Mihaleva VV, van Schalkwijk DB, de Graaf AA, van Duynhoven J, van Dorsten FA,
514 Vervoort J, et al. A systematic approach to obtain validated partial least square models for
515 predicting lipoprotein subclasses from serum NMR spectra. *Anal Chem* 2014 Jan
516 7;86(1):543-550.
- 517 (25) Brocklebank LA, Falconer CL, Page AS, Perry R, Cooper AR. Accelerometer-measured
518 sedentary time and cardiometabolic biomarkers: A systematic review. *Prev Med* 2015
519 Jul;76:92-102.

- 520 (26) Wurtz P, Havulinna AS, Soininen P, Tynkkynen T, Prieto-Merino D, Tillin T, et al.
521 Metabolite profiling and cardiovascular event risk: a prospective study of 3 population-based
522 cohorts. *Circulation* 2015 Mar 3;131(9):774-785.
- 523 (27) DiNicolantonio JJ, O'Keefe JH. Importance of maintaining a low omega-6/omega-3 ratio
524 for reducing inflammation. *Open Heart* 2018 Nov 26;5(2):e000946-2018-000946. eCollection
525 2018.
- 526 (28) Dasilva G, Medina I. Lipidomic methodologies for biomarkers of chronic inflammation
527 in nutritional research: omega-3 and omega-6 lipid mediators. *Free Radic Biol Med* 2019
528 Nov 20;144:90-109.
- 529 (29) Serra-Majem L, Nissensohn M, Overby NC, Fekete K. Dietary methods and biomarkers
530 of omega 3 fatty acids: a systematic review. *Br J Nutr* 2012 Jun;107 Suppl 2:S64-76.
- 531 (30) Nuernberg K, Breier BH, Jayasinghe SN, Bergmann H, Thompson N, Nuernberg G, et
532 al. Metabolic responses to high-fat diets rich in n-3 or n-6 long-chain polyunsaturated fatty
533 acids in mice selected for either high body weight or leanness explain different health
534 outcomes. *Nutr Metab (Lond)* 2011 Aug 11;8(1):56-7075-8-56.
- 535 (31) Kris-Etherton PM, Harris WS, Appel LJ, American Heart Association. Nutrition
536 Committee. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease.
537 *Circulation* 2002 Nov 19;106(21):2747-2757.
- 538 (32) Massaro M, Scoditti E, Carluccio MA, De Caterina R. Basic mechanisms behind the
539 effects of n-3 fatty acids on cardiovascular disease. *Prostaglandins Leukot Essent Fatty Acids*
540 2008 Sep-Nov;79(3-5):109-115.
- 541 (33) Hall KS, Howe CA, Rana SR, Martin CL, Morey MC. METs and accelerometry of
542 walking in older adults: standard versus measured energy cost. *Med Sci Sports Exerc* 2013
543 Mar;45(3):574-582.
- 544 (34) Kraus WE, Houmard JA, Duscha BD, Knetzger KJ, Wharton MB, McCartney JS, et al.
545 Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 2002
546 Nov 7;347(19):1483-1492.
- 547 (35) Mann S, Beedie C, Jimenez A. Differential effects of aerobic exercise, resistance
548 training and combined exercise modalities on cholesterol and the lipid profile: review,
549 synthesis and recommendations. *Sports Med* 2014 Feb;44(2):211-221.
- 550 (36) Kozey SL, Lyden K, Howe CA, Staudenmayer JW, Freedson PS. Accelerometer output
551 and MET values of common physical activities. *Med Sci Sports Exerc* 2010 Sep;42(9):1776-
552 1784.
- 553 (37) Kujala UM, Vaara JP, Kainulainen H, Vasankari T, Vaara E, Kyrolainen H.
554 Associations of Aerobic Fitness and Maximal Muscular Strength With Metabolites in Young
555 Men. *JAMA Netw Open* 2019 Aug 2;2(8):e198265.
- 556 (38) King AC, Haskell WL, Young DR, Oka RK, Stefanick ML. Long-term effects of
557 varying intensities and formats of physical activity on participation rates, fitness, and

- 558 lipoproteins in men and women aged 50 to 65 years. *Circulation* 1995 May 15;91(10):2596-
559 2604.
- 560 (39) Butcher LR, Thomas A, Backx K, Roberts A, Webb R, Morris K. Low-intensity exercise
561 exerts beneficial effects on plasma lipids via PPARgamma. *Med Sci Sports Exerc* 2008
562 Jul;40(7):1263-1270.
- 563 (40) Feingold KR, Grunfeld C. Effect of inflammation on HDL structure and function. *Curr*
564 *Opin Lipidol* 2016 Oct;27(5):521-530.
- 565 (41) King DE, Carek P, Mainous AG,3rd, Pearson WS. Inflammatory markers and exercise:
566 differences related to exercise type. *Med Sci Sports Exerc* 2003 Apr;35(4):575-581.
- 567 (42) Gebhard C, Rhainds D, Tardif JC. HDL and cardiovascular risk: is cholesterol in particle
568 subclasses relevant? *Eur Heart J* 2015 Jan 1;36(1):10-12.
- 569 (43) Sarzynski MA, Burton J, Rankinen T, Blair SN, Church TS, Despres JP, et al. The
570 effects of exercise on the lipoprotein subclass profile: A meta-analysis of 10 interventions.
571 *Atherosclerosis* 2015 Dec;243(2):364-372.
- 572 (44) van Capelleveen JC, Bochem AE, Motazacker MM, Hovingh GK, Kastelein JJ. Genetics
573 of HDL-C: a causal link to atherosclerosis? *Curr Atheroscler Rep* 2013 Jun;15(6):326-013-
574 0326-8.
- 575 (45) Healy GN, Matthews CE, Dunstan DW, Winkler EA, Owen N. Sedentary time and
576 cardio-metabolic biomarkers in US adults: NHANES 2003-06. *Eur Heart J* 2011
577 Mar;32(5):590-597.
- 578 (46) Garvey WT, Kwon S, Zheng D, Shaughnessy S, Wallace P, Hutto A, et al. Effects of
579 insulin resistance and type 2 diabetes on lipoprotein subclass particle size and concentration
580 determined by nuclear magnetic resonance. *Diabetes* 2003 Feb;52(2):453-462.
- 581 (47) Bey L, Hamilton MT. Suppression of skeletal muscle lipoprotein lipase activity during
582 physical inactivity: a molecular reason to maintain daily low-intensity activity. *J Physiol*
583 2003 Sep 1;551(Pt 2):673-682.
- 584 (48) Hamilton MT, Hamilton DG, Zderic TW. Role of low energy expenditure and sitting in
585 obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 2007
586 Nov;56(11):2655-2667.
- 587 (49) Ghafouri K, Cooney J, Bedford DK, Wilson J, Caslake MJ, Gill JM. Moderate Exercise
588 Increases Affinity of Large Very Low-Density Lipoproteins for Hydrolysis by Lipoprotein
589 Lipase. *J Clin Endocrinol Metab* 2015 Jun;100(6):2205-2213.

590

591

592

593

594 **Table 1. Participant characteristics**

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

Variable	All
Participants (N)	509
Age (years)	64 ± 8
Female	176 (34.6)
Current smokers	35 (6.9)
Glycosylated haemoglobin (HbA1c) (%)	5.9 ± 0.4
HbA1c (mmol/mol)	41.0 ± 2.0
Total cholesterol (mmol/L)	5.1 ± 1.0
LDL (mmol/L)	3.1 ± 0.9
HDL (mmol/L)	1.4 (0.4)
Triglycerides (mmol/L)	1.3 (0.7)
Ethnicity	
White European	473 (92.9)
South Asian	33 (6.5)
Other	3 (0.6)
Cardiovascular disease*	176 (34.6)
Accelerometer variables (time in minutes per day)	
Wear-time	853.4 ± 84
<500cpm	704.3 (127.4)
500-999cpm	73.3 (40.8)
1000-1499cpm	29.8 (26.5)
1500-1999cpm	13.2 (15)
2000-2499cpm	6.8 (8.66)
2500-2999cpm	4.0 (5.5)
3000-3499cpm	2.5 (4.9)
3500-3999cpm	1.2 (3.8)
4000-4499cpm	0.3 (2.3)
>4500cpm	0 (1.5)
Average steps per day	6581 ± 3143

618 Data presented as mean ± standard deviation, median (interquartile range) or number (column
619 percent). cpm=counts per minute. *Cardiovascular Disease is defined a medical history of one
620 or more of the following: Myocardial Infarction (MI), Heart Valve Disease, Heart Failure,
621 Atrial Fibrillation, Angina, Stroke, Angioplasty/Coronary Artery Bypass Graft, Leg
622 Angioplasty/bypass, Peripheral Vascular Disease.


623

624 **Figures**

625

626 **Figure 1a. Forest plot displaying the percentage difference in HDL subclass**
627 **concentrations with a 10 minute increase in time spent in bands of 500 counts per minute**
628 **of physical activity intensities.**

629




630

631 **Figure 1b. Forest plot displaying the percentage difference in VLDL subclass**
 632 **concentrations with a 10 minute increase in time spent in bands of 500 counts per minute**
 633 **of physical activity intensities.**

634


635



636

637 **Figure 1c. Forest plot displaying the percentage difference in VLDL subclass**
 638 **concentrations with a 10 minute increase in time spent in bands of 500 counts per minute**
 639 **of physical activity intensities.**

640




641

642

643

644 **Figure 1d. Forest plot displaying the percentage difference in IDL and LDL subclass**
 645 **concentrations with a 10 minute increase in time spent in bands of 500 counts per minute**
 646 **of physical activity intensities.**

647




648

649


650

651 **Figure 2. Forest plot displaying the percentage difference in apolipoproteins with a 10**
 652 **minute increase in time spent in bands of 500 counts per minute of physical activity**
 653 **intensities.**



654
 655
 656
 657

658 **Figure 3. Forest plot displaying the percentage difference in lipoprotein particle size**
 659 **with a 10 minute increase in time spent in bands of 500 counts per minute of physical**
 660 **activity intensities.**



661

662 Colours broadly represent commonly used accelerometer cut points for low levels of physical
 663 activity, which includes sedentary behaviour (red) (<500cpm), light (blue) (≥500-<2000cpm)
 664 and MVPA (green) (≥2000 counts per minute).

665