

**Special Issue**

147-154	155-162	163-170	171-178
179-186	187-194	195-202	203-210
211-218	219-226	227-234	235-242
243-250	251-258	259-266	267-274
275-282	283-290	291-298	299-306
307-314	315-322	323-330	331-338
339-346	347-354	355-362	363-370
371-378	379-386	387-394	395-402
403-410	411-418	419-426	427-434
435-442	443-450	451-458	459-466
467-474	475-482	483-490	491-498
499-506	507-514	515-522	523-530
531-538	539-546	547-554	555-562
563-570	571-578	579-586	587-594
595-602	603-610	611-618	619-626
627-634	635-642	643-650	651-658
659-666	667-674	675-682	683-690
691-698	699-706	707-714	715-722
723-730	731-738	739-746	747-754
755-762	763-770	771-778	779-786
787-794	795-802	803-810	811-818
819-826	827-834	835-842	843-850
851-858	859-866	867-874	875-882
883-890	891-898	899-906	907-914
915-922	923-930	931-938	939-946
947-954	955-962	963-970	971-978
979-986	987-994	995-1002	1003-1010
1011-1018	1019-1026	1027-1034	1035-1042
1043-1050	1051-1058	1059-1066	1067-1074
1075-1082	1083-1090	1091-1098	1099-1106
1107-1114	1115-1122	1123-1130	1131-1138
1139-1146	1147-1154	1155-1162	1163-1170
1171-1178	1179-1186	1187-1194	1195-1202
1203-1210	1211-1218	1219-1226	1227-1234
1235-1242	1243-1250	1251-1258	1259-1266
1267-1274	1275-1282	1283-1290	1291-1298
1299-1306	1307-1314	1315-1322	1323-1330
1331-1338	1339-1346	1347-1354	1355-1362
1363-1370	1371-1378	1379-1386	1387-1394
1395-1402	1403-1410	1411-1418	1419-1426
1427-1434	1435-1442	1443-1450	1451-1458
1459-1466	1467-1474	1475-1482	1483-1490
1491-1498	1499-1506	1507-1514	1515-1522
1523-1530	1531-1538	1539-1546	1547-1554
1555-1562	1563-1570	1571-1578	1579-1586
1587-1594	1595-1602	1603-1610	1611-1618
1619-1626	1627-1634	1635-1642	1643-1650
1651-1658	1659-1666	1667-1674	1675-1682
1683-1690	1691-1698	1699-1706	1707-1714
1715-1722	1723-1730	1731-1738	1739-1746
1747-1754	1755-1762	1763-1770	1771-1778
1779-1786	1787-1794	1795-1802	1803-1810
1811-1818	1819-1826	1827-1834	1835-1842
1843-1850	1851-1858	1859-1866	1867-1874
1875-1882	1883-1890	1891-1898	1899-1906
1907-1914	1915-1922	1923-1930	1931-1938
1939-1946	1947-1954	1955-1962	1963-1970
1971-1978	1979-1986	1987-1994	1995-2002
2003-2010	2011-2018	2019-2026	2027-2034
2035-2042	2043-2050	2051-2058	2059-2066
2067-2074	2075-2082	2083-2090	2091-2098
2099-2106	2107-2114	2115-2122	2123-2130
2131-2138	2139-2146	2147-2154	2155-2162
2163-2170	2171-2178	2179-2186	2187-2194
2195-2202	2203-2210	2211-2218	2219-2226
2227-2234	2235-2242	2243-2250	2251-2258
2259-2266	2267-2274	2275-2282	2283-2290
2291-2298	2299-2306	2307-2314	2315-2322
2323-2330	2331-2338	2339-2346	2347-2354
2355-2362	2363-2370	2371-2378	2379-2386
2387-2394	2395-2402	2403-2410	2411-2418
2419-2426	2427-2434	2435-2442	2443-2450
2451-2458	2459-2466	2467-2474	2475-2482
2483-2490	2491-2498	2499-2506	2507-2514
2515-2522	2523-2530	2531-2538	2539-2546
2547-2554	2555-2562	2563-2570	2571-2578
2579-2586	2587-2594	2595-2602	2603-2610
2611-2618	2619-2626	2627-2634	2635-2642
2643-2650	2651-2658	2659-2666	2667-2674
2675-2682	2683-2690	2691-2698	2699-2706
2707-2714	2715-2722	2723-2730	2731-2738
2739-2746	2747-2754	2755-2762	2763-2770
2771-2778	2779-2786	2787-2794	2795-2802
2803-2810	2811-2818	2819-2826	2827-2834
2835-2842	2843-2850	2851-2858	2859-2866
2867-2874	2875-2882	2883-2890	2891-2898
2899-2906	2907-2914	2915-2922	2923-2930
2931-2938	2939-2946	2947-2954	2955-2962
2963-2970	2971-2978	2979-2986	2987-2994
2995-3002	3003-3010	3011-3018	3019-3026
3027-3034	3035-3042	3043-3050	3051-3058
3059-3066	3067-3074	3075-3082	3083-3090
3091-3098	3099-3106	3107-3114	3115-3122
3123-3130	3131-3138	3139-3146	3147-3154
3155-3162	3163-3170	3171-3178	3179-3186
3187-3194	3195-3202	3203-3210	3211-3218
3219-3226	3227-3234	3235-3242	3243-3250
3251-3258	3259-3266	3267-3274	3275-3282
3283-3290	3291-3298	3299-3306	3307-3314
3315-3322	3323-3330	3331-3338	3339-3346
3347-3354	3355-3362	3363-3370	3371-3378
3379-3386	3387-3394	3395-3402	3403-3410
3411-3418	3419-3426	3427-3434	3435-3442
3443-3450	3451-3458	3459-3466	3467-3474
3475-3482	3483-3490	3491-3498	3499-3506
3507-3514	3515-3522	3523-3530	3531-3538
3539-3546	3547-3554	3555-3562	3563-3570
3571-3578	3579-3586	3587-3594	3595-3602
3603-3610	3611-3618	3619-3626	3627-3634
3635-3642	3643-3650	3651-3658	3659-3666
3667-3674	3675-3682	3683-3690	3691-3698
3699-3706	3707-3714	3715-3722	3723-3730
3731-3738	3739-3746	3747-3754	3755-3762
3763-3770	3771-3778	3779-3786	3787-3794
3795-3802	3803-3810	3811-3818	3819-3826
3827-3834	3835-3842	3843-3850	3851-3858
3859-3866	3867-3874	3875-3882	3883-3890
3891-3898	3899-3906	3907-3914	3915-3922
3923-3930	3931-3938	3939-3946	3947-3954
3955-3962	3963-3970	3971-3978	3979-3986
3987-3994	3995-4002	4003-4010	4011-4018
4019-4026	4027-4034	4035-4042	4043-4050
4051-4058	4059-4066	4067-4074	4075-4082
4083-4090	4091-4098	4099-4106	4107-4114
4115-4122	4123-4130	4131-4138	4139-4146
4147-4154	4155-4162	4163-4170	4171-4178
4179-4186	4187-4194	4195-4202	4203-4210
4211-4218	4219-4226	4227-4234	4235-4242
4243-4250	4251-4258	4259-4266	4267-4274
4275-4282	4283-4290	4291-4298	4299-4306
4307-4314	4315-4322	4323-4330	4331-4338
4339-4346	4347-4354	4355-4362	4363-4370
4371-4378	4379-4386	4387-4394	4395-4402
4403-4410	4411-4418	4419-4426	4427-4434
4435-4442	4443-4450	4451-4458	4459-4466
4467-4474	4475-4482	4483-4490	4491-4498
4499-4506	4507-4514	4515-4522	4523-4530
4531-4538	4539-4546	4547-4554	4555-4562
4563-4570	4571-4578	4579-4586	4587-4594
4595-4602	4603-4610	4611-4618	4619-4626
4627-4634	4635-4642	4643-4650	4651-4658
4659-4666	4667-4674	4675-4682	4683-4690
4691-4698	4699-4706	4707-4714	4715-4722
4723-4730	4731-4738	4739-4746	4747-4754
4755-4762	4763-4770	4771-4778	4779-4786
4787-4794	4795-4802	4803-4810	4811-4818
4819-4826	4827-4834	4835-4842	4843-4850
4851-4858	4859-4866	4867-4874	4875-4882
4883-4890	4891-4898	4899-4906	4907-4914
4915-4922	4923-4930	4931-4938	4939-4946
4947-4954	4955-4962	4963-4970	4971-4978
4979-4986	4987-4994	4995-5002	5003-5010
5011-5018	5019-5026	5027-5034	5035-5042
5043-5050	5051-5058	5059-5066	5067-5074
5075-5082	5083-5090	5091-5098	5099-5106
5107-5114	5115-5122	5123-5130	5131-5138
5139-5146	5147-5154	5155-5162	5163-5170
5171-5178	5179-5186	5187-5194	5195-5202
5203-5210	5211-5218	5219-5226	5227-5234
5235-5242	5243-5250	5251-5258	5259-5266
5267-5274	5275-5282	5283-5290	5291-5298
5299-5306	5307-5314	5315-5322	5323-5330
5331-5338	5339-5346	5347-5354	5355-5362
5363-5370	5371-5378	5379-5386	5387-5394
5395-5402	5403-5410	5411-5418	5419-5426
5427-5434	5435-5442	5443-5450	5451-5458
5459-5466	5467-5474	5475-5482	5483-5490
5491-5498	5499-5506	5507-5514	5515-5522
5523-5530	5531-5538	5539-5546	5547-5554
5555-5562	5563-5570	5571-5578	5579-5586
5587-5594	5595-5602	5603-5610	5611-5618
5619-5626	5627-5634	5635-5642	5643-5650
5651-5658	5659-5666	5667-5674	5675-5682
5683-5690	5691-5698	5699-5706	5707-5714
5715-5722	5723-5730	5731-5738	5739-5746
5747-5754	5755-5762	5763-5770	5771-5778
5779-5786	5787-5794	5795-5802	5803-5810
5811-5818	5819-5826	5827-5834	5835-5842
5843-5850	5851-5858	5859-5866	5867-5874
5875-5882	5883-5890	5891-5898	5899-5906
5907-5914	5915-5922	5923-5930	5931-5938
5939-5946	5947-5954	5955-5962	5963-5970
5971-5978	5979-5986	5987-5994	5995-6002
6003-6010	6011-6		

## Actin on trafficking

### Could actin guide directed receptor transport?

Tobias Zech,\* Simon D.J. Calaminus and Laura M. Machesky\*

Beatson Institute for Cancer Research; Bearsden, UK

**H**ere, we present emerging ideas surrounding the interplay between the actin cytoskeleton and receptor transport and activation. The bulk of actin dynamics in cells is thought to contribute to architecture and mobility. Actin also contributes to trafficking, acting as a molecular scaffold, providing force to deform membranes, facilitating vesicle abscission or propelling a vesicle through the cytoplasm<sup>1,2</sup> and recent studies highlight important connections between the directed trafficking of receptors and the impact on cell migration and actin dynamics. Additionally, a number of newly described actin nucleation promoting factors, such as the vesicle associated protein WASH, reveal unexpected roles of actin in membrane traffic and suggest that the cell dedicates a significant proportion of its regulation of actin dynamics to controlling trafficking.

#### Multiple Endocytic Compartments Display Arp2/3-Mediated Actin Assembly Dynamics

Cells often communicate with their environment via transmembrane receptors that receive signals from the extracellular milieu and transmit them internally. Regulation occurs not only at the level of receptors binding to ligands, but can critically depend on the numbers of receptors displayed on the cell surface and the endocytic trafficking pathways shuttling receptors and receptor-ligand complexes away from and back to the plasma membrane (for recent reviews see refs.3–5).

As a general rule, new actin assembly in cells occurs near a membrane interface. One of the main drivers of new actin filament formation adjacent to membranes is the Arp2/3 complex (actin-related protein 2/3 complex), which is activated by the WASP-family (Wiskott-Aldrich syndrome family) proteins (for recent reviews see refs.1, 6 and 7). During cell migration, new filament assembly occurs mostly underneath the plasma membrane, but also on internal membrane systems. Actin on intracellular vesicles is often in very small patches ( $\leq 1 \mu\text{m}$  diameter) and is usually transient and dynamic, requiring very sensitive high-speed imaging. Only recently has technology developed to a point where we are able to begin to capture some of the intricacies of intracellular actin/membrane interactions.

Mammalian genomes encode at least five types of WASP family proteins and surprisingly, at least three (N-WASP, WASH and WHAMM), likely have important and evolutionarily conserved roles in membrane trafficking.<sup>3</sup> WASP proteins generate branched actin networks by bringing the Arp2/3 complex and actin monomers together to nucleate a new branch from a mother filament.<sup>1</sup> In addition to WASP proteins, formins (such as FMNL1/2 and Diaphanous), which nucleate and processively promote elongation of unbranched filaments also can polymerize actin on vesicular structures.<sup>8-12</sup> Likewise, Spire1, which contains multiple WASP-homology domains and can nucleate actin independently of Arp2/3 complex, may cooperate with Arp2/3 complex, as well as formin-2 (Cappuccino) to somehow control progression of early to late endosomes.<sup>9</sup>

**Keywords:** trafficking, endosomes, receptor trafficking, exocytosis, WASP family proteins, Arp2/3 complex

Submitted: 04/30/12

Revised: 06/15/12

Accepted: 07/05/12

<http://dx.doi.org/10.4161/cam.21373>

\*Correspondence to: Tobias Zech and Laura M. Machesky;  
Email: [t.zech@beatson.gla.ac.uk](mailto:t.zech@beatson.gla.ac.uk)  
and [l.machesky@beatson.gla.ac.uk](mailto:l.machesky@beatson.gla.ac.uk)

Commentary to: Zech T, Calaminus SD, Caswell P, Spence HJ, Carnell M, Insall RH, et al. The Arp2/3 activator WASH regulates  $\alpha 5\beta 1$ -integrin-mediated invasive migration. *J Cell Sci* 2011; 124:3753–9; <http://dx.doi.org/10.1242/jcs.080986>

Here we highlight recent research surrounding the vesicle associated WASP-family protein WASH and we propose that actin assembly has an important role in trafficking of various receptor cargoes through multiple endocytic compartments. WASH was originally identified as a gene upregulated in breast cancer cells that is located in the subtelomeric regions of multiple human chromosomes.<sup>13,14</sup> In humans, there are several WASH copies in the genome, with most of them currently curated as pseudogenes. WASH is found in a complex of five proteins, containing SWIP, FAM21, Strumpellin and CCDC53 (Fig. 1) and termed the WASH regulatory complex (SHRC).<sup>15-17</sup>

### Actin Has an Important Role in Multiple Endocytic Events/Stages and WASH Facilitates Receptor Retrieval Back to the Plasma Membrane


The most well understood aspect of actin in trafficking is probably the role of actin assembly in clathrin-mediated endocytosis. This has been extensively reviewed elsewhere,<sup>18-20</sup> so we will only briefly summarize the main ideas. Clathrin coated pits associate close to the plasma

membrane with both dendritic actin networks and actin comet tails.<sup>21</sup> Notably, the comet tails observed in a recent correlative EM study were not always visible in the matching immunofluorescence pictures suggesting that small/subtle membrane associated actin structures still need high magnification microscopy to distinguish them from the high “background” of other cellular actin structures.<sup>21</sup> N-WASP contributes to the generation of branched actin networks on clathrin-coated vesicles. Actin is thought to form a cage around the newly forming neck of a nascent vesicle together with BAR domain proteins and eventually dynamin. Actin polymerization helps to physically stabilize the tubular neck of a nascent vesicle as the clathrin coat assembles and the force from actin assembly pushes against the inherent membrane tension.<sup>22,23</sup> Once dynamin assembles onto the neck, scission can begin via dynamin’s GTPase powered squeezing mechanism, although the actual mechanical forces involved here are still debated.<sup>24</sup> Finally, when the new vesicle breaks free, the clathrin coat is disassembled and actin may then polymerize at one pole to push the new vesicle away from the plasma membrane as it journeys to become an early endosome.

Once the vesicle becomes an early endosome, it is likely to be associated with WASH rather than N-WASP. It isn’t clear how N-WASP dissociates and WASH associates or even when, but EEA1 positive early endosomal vesicles contain WASH puncta that co-localize with actin and Arp2/3 complex.<sup>15,16,25,26</sup> WASH mediated actin polymerization is maintained on endosomes until the multi-vesicular body (MVB) stage, or presumably until late endosomes (LE) lose the ability to recycle receptors and merge/mature to lysosomes.<sup>27</sup>

Receptor transport back to the plasma membrane from an endocytic vesicle is much less defined than the endocytic internalization process. The common view is that tubulin is the cytoskeletal element contributing to outward vesicular traffic via kinesins, and WASH contains a putative tubulin binding motif, which may link actin and microtubule based transport (reviewed in ref.7). But secretion, in a variety of cell types, shows little change upon tubulin depolymerization.<sup>28,29</sup> Rocketing of vesicles with endo/lysosomal content might be facilitated by N-WASP and actin comet tail formation<sup>30,31</sup> and a recent publication on vesicular transport in oocytes showed evidence for a novel mechanism of outward vesicle movement that is facilitated by actin cables assembled by Formin2 and Spire 1/2.<sup>32</sup> Thus, there are still many questions about the relative importance of actin and microtubules in delivery of vesicles back to the plasma membrane from internal endocytic compartments and likely different cell types and compartments have different mechanisms.

WASH localizes on many different endosomal compartments, including early endosomes, late endosomes and, in *Dictyostelium*, post lysosomes.<sup>15,16,25,26,33</sup> It co-localizes with the retromer complex,<sup>16</sup> in support of the idea that WASH-mediated actin networks work together with retromer in salvaging cargo such as receptors for retrograde transport. Furthermore, depletion of WASH led to a redistribution of CI-MPR (cation-independent mannose 6-phosphate receptor), a cargo of retromer that is retrieved from endosomes back to the trans Golgi.<sup>16</sup>



**Figure 1.** The WASH containing regulatory complex (SHRC) activates actin polymerization on endosomes. Schematic of the pentameric WASH containing SHRC complex in an inactivated state with sequestered VCA domain (depicted in analogy to the WAVE/Scar complex; see ref. 17 for description) and when polymerizing actin on an endosome. The complex is likely tethered to the membrane by multiple lipid-protein interactions.

WASH strongly localizes to the Rab7 positive late endosome/multivesicular body (LE/MVB),<sup>26</sup> a compartment that is also marked by the retromer complex, which is thought to interact with Rab7.<sup>34</sup> Seaman and colleagues identified the SHRC subunit strumpellin, as a retromer interacting protein and showed that the other subunits of the WASH regulatory complex also co-interact with retromer complex.<sup>35,36</sup> How WASH functions together with retromer is not fully understood, but a recent study links the FAM21 subunit of SHRC to retromer directly<sup>37</sup> and the SHRC is likely to be important in the formation and/or organization of structures known as retromer tubules.<sup>38</sup> Retromer tubules form when receptors are to be recycled back to the plasma membrane from endosomes. Temkin and colleagues found WASH in association with the retromer tubule associated sorting nexin SNX27 as an important component for recycling of the  $\beta$ -2 adrenergic receptor.<sup>38</sup> However, this is a Rab4-dependent process,<sup>38</sup> so may be distinct from the Rab7-dependent sorting of other receptors. The mechanism for WASH uncoupling from retrograde vesicles when they reach their destination is not known and it is unclear whether retromer and WASH always co-cycle together.

### Actin Has Emerging Roles in Membrane Tubulation, Sorting of Receptor/Cargo Complexes and Determining the Fate of Particular Membrane Proteins

Increased imaging capability has allowed detection of multiple actin networks on various vesicle compartments, including tubules that emerge from endosomes, which often are associated with retrieval of cargo back to the plasma membrane or in the case of the CI-MPR, to the Golgi. The nature and function of actin on these emerging tubules and the small vesicles that bud from them is just beginning to be understood.

Deforming membranes to bud vesicles requires energy to overcome membrane tension and to induce curvature, movement and scission, which could come from actin polymerization.<sup>24</sup> The Arp2/3 activator WHAMM (WASP homolog associated with actin, membranes and

microtubules) localizes to the endoplasmic reticulum and Golgi complex<sup>39</sup> and is proposed to form a link between microtubules and nascent membrane tubules forming in ER to Golgi transport. WHAMM is proposed to nucleate branched actin networks on nascent tubules and drive their active formation. In direct contrast, WASH is proposed to oppose membrane tubulation, since knockdown of WASH can result in increased appearance of tubules in cells.<sup>7,15,16</sup> WASH was proposed to establish an actin network on a nascent tubule, allowing dynamin to be recruited and thus promote vesicle scission from the tubule. This idea fits with previous studies of BAR domain proteins showing that when actin assembly was prevented, the tubules failed to be cleaved.<sup>40</sup> The arrival of branched actin allowed dynamin to be recruited and efficiently cleave nascent tubules. It is unclear if WHAMM and WASH actually work by such contrasting mechanisms, with WHAMM promoting tubulation and WASH promoting scission, but clearly further studies are needed.

Whether WASH and WHAMM promote tubulation or scission of vesicles, it is clear that at least WASH is important for efficient receptor recycling. Several endosomal receptor cargos have now been analyzed for their dependence on WASH function, actin polymerization and other members of the SHRC. All studies agreed that WASH depletion led to a redistribution of recycling receptors from the plasma membrane, Golgi or lysosomes to endosomal membranes indicating a retention of receptors at that level.<sup>15,16,26,41</sup> WASH depleted cells display slower recycling kinetics of multiple cargoes.<sup>15,26</sup> This was less pronounced for the constitutive recycling transferrin receptor, which was only modestly affected<sup>15,26</sup> or even unaffected<sup>25</sup> by WASH depletion, than for receptors that are regulated by signal transduction and degradation like the integrin  $\alpha 5 \beta 1$ .<sup>26,42-44</sup> Perhaps surprisingly, it is increasingly becoming apparent that  $\alpha 5 \beta 1$  integrins primarily recycle from a late endosomal compartment when cells are migrating in a 3D environment<sup>19,26,45</sup> in agreement with WASH's partial colocalization with Rab7 positive endosomes.<sup>15</sup> Recycling from a mature MVB/

LE compartment had been long suggested for trafficking of the MHCII in immune cells, which relies on the late endosomal environment for loading with cognate peptides.<sup>46</sup>

The mechanism by which WASH regulates sorting of receptors such as  $\alpha 5 \beta 1$  integrin and targeting for recycling rather than degradation is not yet clear, but two lines of evidence point toward a possible role in actin-based sorting of receptors. First, Dictyostelium WASH is important for the recycling of the vacuolar ATPase that allows lysosomes to be neutralized and then exocytosed from cells.<sup>33</sup> Insall and colleagues demonstrated that V-ATPase is an actin binding protein and hypothesized that actin networks created by WASH and Arp2/3 complex capture V-ATPase in patches on the surface of lysosomes and thus sort the V-ATPase for removal and recycling.<sup>33</sup> Second, in mammalian cells, a recent study by Puthenveedu and colleagues identified determinants of receptor recycling and found that  $\beta$ -2-adrenergic receptors need to be sorted into actin rich domains on endosomes to allow efficient exit from the endosome.<sup>47</sup> Actin binding, either directly by the receptor or via receptor-associated proteins, as in the case of the  $\beta$ -2-adrenergic receptor, was the only prerequisite for recycling of a subset of signaling receptors.

A systematic survey of endosomal transport discriminated between at least two separate types of receptor cargo.<sup>48</sup> The first group, represented by the transferrin receptor, had characteristics of passive cargo where receptor accumulation was positively correlated with the size of the endosome the recycling of receptors presumably commenced at a constant rate. The second group, represented in the study by the EGFR (epidermal growth factor receptor), showed signs of guidance and adaptable regulation in the endosomal system. The mean EGFR cargo content remained constant independent of endosome size indicating tight regulation of EGFR localization.<sup>48</sup> After activation, regulated receptors such as EGF receptor are ubiquitinated and subsequently sorted toward degradation.<sup>49</sup> Ubiquitinated EGFR is sorted into intraluminal vesicles (ILVs) by the ESCRT (endosomal sorting

complex required for transport) family of complexes and degraded after endosomal fusion with lysosomes.<sup>50</sup> Thus degradation works by active sorting of receptors away from the limiting membrane of the endosome, suggesting that receptors are otherwise sorted for recycling to the plasma membrane or any other cellular destination.


The ESCRT complex and WASH can localize to the same endosome and our preliminary data do not indicate direct colocalization (Fig. 2). Actin binding of some receptors is necessary for efficient recycling from a distal endosomal compartment<sup>33,47</sup> and we found that WASH overexpression correlated with increased cell surface expression of  $\alpha 5 \beta 1$  integrin.<sup>26</sup> Recycling endosomes, a term used here for the aforementioned endosomal population still capable of retrograde receptor traffic, have been shown to consist of distinct domains and protrusions enriched in specialized sets of molecules required for transport away from endosomes.<sup>51</sup>

We propose that actin and the WASH complex might act as a positive recycling guide opposing ESCRT mediated sorting into ILVs of MVBs. Reversible actin binding—extrinsic or intrinsic—would form an ideal way to sequester receptors away from each other. Whenever receptors are destined for degradation the actin binding would be inhibited and sorting into ESCRT rich region could proceed. The continuous occurrence of actin-rich domains from the plasma membrane to late endosomes would be an ideal method to hand over receptors that are destined for reuse.

In summary, it is emerging that mammalian cells devote a considerable component of their actin nucleation promoting proteins to vesicle trafficking functions. Actin is likely to be important in more than its currently established role in pushing against membranes to drive protrusion and movement and these likely include receptor sorting and directed

## References

1. Insall RH, Machesky LM. Actin dynamics at the leading edge: from simple machinery to complex networks. *Dev Cell* 2009; 17:310-22; PMID: 19758556; <http://dx.doi.org/10.1016/j.devcel.2009.08.012>
2. Anitei M, Hoflack B. Bridging membrane and cytoskeleton dynamics in the secretory and endocytic pathways. *Nat Cell Biol* 2012; 14:11-9; PMID: 22193159; <http://dx.doi.org/10.1038/ncb2409>
3. Caswell P, Norman J. Endocytic transport of integrins during cell migration and invasion. *Trends Cell Biol* 2008; 18:257-63; PMID:18456497; <http://dx.doi.org/10.1016/j.tcb.2008.03.004>



**Figure 2.** Actin mediated receptor sorting. (A) Immunofluorescence staining of the ESCRT-0 component Hrs and WASH on an endosomal membrane of a senescent IMR90 fibroblast. Scale bar 5  $\mu$ M. Blue line indicates endosomal outline. E, endosome. (B) Model of multivesicular body with WASH and actin rich domains leading to recycling of receptors and ESCRT rich regions that lead to the sorting of ubiquitinated receptor into intraluminal vesicles.

trafficking. The actin nucleation promoting protein WASH has emerged as an important regulator of receptor trafficking. The roles of proteins such as WASH and the actin networks that they generate are still only beginning to be understood and likely are complex.

A recent study demonstrated the presence of short cortical actin filaments on the plasma membrane and predicted that they would spontaneously form aster-shaped clusters on the membrane surface, due to myosins or to attractive forces between the filaments.<sup>52</sup> If receptors with engineered actin binding motifs are added

to the system, nano-clusters spontaneously formed.<sup>52</sup> This showed that actin networks on membrane surfaces are in general able to organize receptors into clusters and suggests a novel mechanism by which they could alter sorting and/or signaling.

## Acknowledgments

L.M.M., S.D.J.C. and T.Z. are funded by a core grant from Cancer Research UK to L.M.M. and a pilot grant from Breast Cancer Campaign to L.M.M. and T.Z. Thanks to Peter van den Berghe and Jim Norman (Beatson Institute) for providing the IMR90 fibroblasts.

4. Jones MC, Caswell PT, Norman JC. Endocytic recycling pathways: emerging regulators of cell migration. *Curr Opin Cell Biol* 2006; 18:549-57; PMID: 16904305; <http://dx.doi.org/10.1016/j.ceb.2006.08.003>
5. Sorkin A, von Zastrow M. Endocytosis and signalling: intertwining molecular networks. *Nat Rev Mol Cell Biol* 2009; 10:609-22; PMID:19696798; <http://dx.doi.org/10.1038/nrm2748>
6. Campellone KG, Welch MD. A nucleator arms race: cellular control of actin assembly. *Nat Rev Mol Cell Biol* 2010; 11:237-51; PMID:20237478; <http://dx.doi.org/10.1038/nrm2867>
7. Rottner K, Hänisch J, Campellone KG, WASH, WHAMM and JMY: regulation of Arp2/3 complex and beyond. *Trends Cell Biol* 2010; 20:650-61; PMID:20888769; <http://dx.doi.org/10.1016/j.tcb.2010.08.014>
8. Kitzing TM, Wang Y, Pertz O, Copeland JW, Grosse R. Formin-like 2 drives amoeboid invasive cell motility downstream of RhoC. *Oncogene* 2010; 29:2441-8; PMID:20101212; <http://dx.doi.org/10.1038/onc.2009.515>
9. Morel E, Parton RG, Gruenberg J. Annexin A2-dependent polymerization of actin mediates endosome biogenesis. *Dev Cell* 2009; 16:445-57; PMID: 19289089; <http://dx.doi.org/10.1016/j.devcel.2009.01.007>
10. Gardberg M, Talvinen K, Kaipio K, Iljin K, Kampf C, Uhlen M, et al. Characterization of Diaphanous-related formin FMNL2 in human tissues. *BMC Cell Biol* 2010; 11:55; PMID:20633255; <http://dx.doi.org/10.1186/1471-2121-11-55>
11. Colón-Franco JM, Gomez TS, Billadeau DD. Dynamic remodeling of the actin cytoskeleton by FMNL1 $\gamma$  is required for structural maintenance of the Golgi complex. *J Cell Sci* 2011; 124:3118-26; PMID: 21868368; <http://dx.doi.org/10.1242/jcs.083725>
12. Fernandez-Borja M, Janssen L, Verwoerd D, Hordijk P, Neeffes J. RhoB regulates endosome transport by promoting actin assembly on endosomal membranes through Dia1. *J Cell Sci* 2005; 118:2661-70; PMID: 15944396; <http://dx.doi.org/10.1242/jcs.02384>
13. Linardopoulou EV, Parghi SS, Friedman C, Osborn GE, Parkhurst SM, Trask BJ. Human subtelomeric WASH genes encode a new subclass of the WASP family. *PLoS Genet* 2007; 3:e237; PMID:18159949; <http://dx.doi.org/10.1371/journal.pgen.0030237>
14. Leirdal M, Shadidy M, Rosok O, Sioud M. Identification of genes differentially expressed in breast cancer cell line SKBR3: potential identification of new prognostic biomarkers. *Int J Mol Med* 2004; 14:217-22; PMID:15254768
15. Derivery E, Sousa C, Gautier JJ, Lombard B, Loew D, Gautreau A. The Arp2/3 activator WASH controls the fission of endosomes through a large multiprotein complex. *Dev Cell* 2009; 17:712-23; PMID: 19922875; <http://dx.doi.org/10.1016/j.devcel.2009.09.010>
16. Gomez TS, Billadeau DDA. A FAM21-containing WASH complex regulates retromer-dependent sorting. *Dev Cell* 2009; 17:699-711; PMID:19922874; <http://dx.doi.org/10.1016/j.devcel.2009.09.009>
17. Jia D, Gomez TS, Metlagel Z, Umetani J, Orwinowski Z, Rosen MK, et al. WASH and WAVE actin regulators of the Wiskott-Aldrich syndrome protein (WASP) family are controlled by analogous structurally related complexes. *Proc Natl Acad Sci U S A* 2010; 107:10442-7; PMID:20498093; <http://dx.doi.org/10.1073/pnas.0913293107>
18. McMahon HT, Boucrot E. Molecular mechanism and physiological functions of clathrin-mediated endocytosis. *Nat Rev Mol Cell Biol* 2011; 12:517-33; PMID: 21779028; <http://dx.doi.org/10.1038/nrm3151>
19. Tu C, Ortega-Cava CF, Winograd P, Stanton MJ, Reddi AL, Dodge J, et al. Endosomal-sorting complexes required for transport (ESCRT) pathway-dependent endosomal traffic regulates the localization of active Src at focal adhesions. *Proc Natl Acad Sci U S A* 2010; 107:16107-12; PMID:20805499; <http://dx.doi.org/10.1073/pnas.1009471107>
20. Yarar D, Waterman-Storer CM, Schmid SL. A dynamic actin cytoskeleton functions at multiple stages of clathrin-mediated endocytosis. *Mol Biol Cell* 2005; 16:964-75; PMID:15601897; <http://dx.doi.org/10.1091/mbc.E04-09-0774>
21. Collins A, Warrington A, Taylor KA, Svitkina T. Structural organization of the actin cytoskeleton at sites of clathrin-mediated endocytosis. *Curr Biol* 2011; 21:1167-75; PMID:21723126; <http://dx.doi.org/10.1016/j.cub.2011.05.048>
22. Dai J, Sheetz MP. Regulation of endocytosis, exocytosis, and shape by membrane tension. *Cold Spring Harb Symp Quant Biol* 1995; 60:567-71; PMID: 8824429; <http://dx.doi.org/10.1101/SQB.1995.060.01.060>
23. Upadhyaya A, Sheetz MP. Tension in tubulovesicular networks of Golgi and endoplasmic reticulum membranes. *Biophys J* 2004; 86:2923-8; PMID:15111408; [http://dx.doi.org/10.1016/S0006-3495\(04\)74343-X](http://dx.doi.org/10.1016/S0006-3495(04)74343-X)
24. Liu J, Sun Y, Oster GF, Drubin DG. Mechanochemical crosstalk during endocytic vesicle formation. *Curr Opin Cell Biol* 2010; 22:36-43; PMID:20022735; <http://dx.doi.org/10.1016/j.ceb.2009.11.009>
25. Duleh SN, Welch MD. WASH and the Arp2/3 complex regulate endosome shape and trafficking. *Cytoskeleton (Hoboken)* 2010; 67:193-206; PMID: 20175130
26. Zech T, Calaminus SD, Caswell P, Spence HJ, Carnell M, Insall RH, et al. The Arp2/3 activator WASH regulates  $\alpha 5 \beta 1$ -integrin-mediated invasive migration. *J Cell Sci* 2011; 124:3753-9; PMID:22114305; <http://dx.doi.org/10.1242/jcs.080986>
27. Huotari J, Helenius A. Endosome maturation. *EMBO J* 2011; 30:3481-500; PMID:21878991; <http://dx.doi.org/10.1038/emboj.2011.286>
28. Apodaca G. Endocytic traffic in polarized epithelial cells: role of the actin and microtubule cytoskeleton. *Traffic* 2001; 2:149-59; PMID:11260520; <http://dx.doi.org/10.1034/j.1600-0854.2001.020301.x>
29. Bloom GS, Goldstein LS. Cruising along microtubule highways: how membranes move through the secretory pathway. *J Cell Biol* 1998; 140:1277-80; PMID:9508761; <http://dx.doi.org/10.1083/jcb.140.6.1277>
30. Taunton J, Rowning BA, Coughlin ML, Wu M, Moon RT, Mitchison TJ, et al. Actin-dependent propulsion of endosomes and lysosomes by recruitment of N-WASP. *J Cell Biol* 2000; 148:519-30; PMID: 10662777; <http://dx.doi.org/10.1083/jcb.148.3.519>
31. Rozelle AL, Machesky LM, Yamamoto M, Driessens MH, Insall RH, Roth MG, et al. Phosphatidylinositol 4,5-bisphosphate induces actin-based movement of raft-enriched vesicles through WASP-Arp2/3. *Curr Biol* 2000; 10:311-20; PMID:10744973; [http://dx.doi.org/10.1016/S0960-9822\(00\)00384-5](http://dx.doi.org/10.1016/S0960-9822(00)00384-5)
32. Schuh M. An actin-dependent mechanism for long-range vesicle transport. *Nat Cell Biol* 2011; 13:1431-6; PMID:21983562; <http://dx.doi.org/10.1038/ncb2353>
33. Carnell M, Zech T, Calaminus SD, Ura S, Hagedorn M, Johnson SA, et al. Actin polymerization driven by WASH causes V-ATPase retrieval and vesicle neutralization before exocytosis. *J Cell Biol* 2011; 193:831-9; PMID:21606208; <http://dx.doi.org/10.1083/jcb.201009119>
34. Seaman MN, Harbour ME, Tattersall D, Read E, Bright N. Membrane recruitment of the cargo-selective retromer subcomplex is catalysed by the small GTPase Rab7 and inhibited by the Rab-GAP TBC1D5. *J Cell Sci* 2009; 122:2371-82; PMID:19531583; <http://dx.doi.org/10.1242/jcs.048686>
35. Harbour ME, Breusegem SY, Antrobus R, Freeman C, Reid E, Seaman MN. The cargo-selective retromer complex is a recruiting hub for protein complexes that regulate endosomal tubule dynamics. *J Cell Sci* 2010; 123:3703-17; PMID:20923837; <http://dx.doi.org/10.1242/jcs.071472>
36. Harbour ME, Breusegem SY, Seaman MN. Recruitment of the endosomal WASH complex is mediated by the extended 'tail' of Fam21 binding to the retromer protein Vps35. *Biochem J* 2012; 442:209-20; PMID:22070227; <http://dx.doi.org/10.1042/BJ20111761>
37. Jia D, Gomez TS, Billadeau DD, Rosen MK. Multiple repeat elements within the FAM21 tail link the WASH actin regulatory complex to the retromer. *Mol Biol Cell* 2012; 23:2352-61; PMID:22513087; <http://dx.doi.org/10.1091/mbc.E11-12-1059>
38. Temkin P, Lauffer B, Jäger S, Cimermancic P, Krogan NJ, von Zastrow M. SNX27 mediates retromer tubule entry and endosome-to-plasma membrane trafficking of signalling receptors. *Nat Cell Biol* 2011; 13:715-21; PMID:21602791; <http://dx.doi.org/10.1038/ncb2252>
39. Campellone KG, Webb NJ, Znameroski EA, Welch MD. WHAMM is an Arp2/3 complex activator that binds microtubules and functions in ER to Golgi transport. *Cell* 2008; 134:148-61; PMID:18614018; <http://dx.doi.org/10.1016/j.cell.2008.05.032>
40. Fricke R, Gohl C, Dharmalingam E, Grevelhörster A, Zahedi B, Harden N, et al. Drosophila Cip4/Toca-1 integrates membrane trafficking and actin dynamics through WASP and SCAR/WAVE. *Curr Biol* 2009; 19:1429-37; PMID:19716703; <http://dx.doi.org/10.1016/j.cub.2009.07.058>
41. Duleh SN, Welch MD. WASH and the Arp2/3 complex regulate endosome shape and trafficking. *Cytoskeleton (Hoboken)* 2010; 67:193-206; PMID: 20175130
42. Caswell PT, Spence HJ, Parsons M, White DP, Clark K, Cheng KW, et al. Rab25 associates with alpha5-beta1 integrin to promote invasive migration in 3D microenvironments. *Dev Cell* 2007; 13:496-510; PMID:17925226; <http://dx.doi.org/10.1016/j.devcel.2007.08.012>
43. Caswell PT, Vadrevu S, Norman JC. Integrins: masters and slaves of endocytic transport. *Nat Rev Mol Cell Biol* 2009; 10:843-53; PMID:19904298; <http://dx.doi.org/10.1038/nrm2799>
44. Lobert VH, Brech A, Pedersen NM, Wesche J, Oppelt A, Malerød L, et al. Ubiquitination of alpha 5 beta 1 integrin controls fibroblast migration through lysosomal degradation of fibronectin-integrin complexes. *Dev Cell* 2010; 19:148-59; PMID:20643357; <http://dx.doi.org/10.1016/j.devcel.2010.06.010>
45. Dozynkiewicz MA, Jamieson NB, Macpherson I, Grindlay J, van den Berghe PV, von Thun A, et al. Rab25 and CLIC3 collaborate to promote integrin recycling from late endosomes/lysosomes and drive cancer progression. *Dev Cell* 2012; 22:131-45; PMID: 22197222; <http://dx.doi.org/10.1016/j.devcel.2011.11.008>
46. Murk JL, Stoorvogel W, Kleijmeer MJ, Geuze HJ. The plasticity of multivesicular bodies and the regulation of antigen presentation. *Semin Cell Dev Biol* 2002; 13:303-11; PMID:12243730; <http://dx.doi.org/10.1016/S1084952102000605>

47. Puthenveedu MA, Lauffer B, Temkin P, Vistein R, Carlton P, Thorn K, et al. Sequence-dependent sorting of recycling proteins by actin-stabilized endosomal microdomains. *Cell* 2010; 143:761-73; PMID: 21111236; <http://dx.doi.org/10.1016/j.cell.2010.10.003>
48. Collinet C, Stöter M, Bradshaw CR, Samusik N, Rink JC, Kenski D, et al. Systems survey of endocytosis by multiparametric image analysis. *Nature* 2010; 464: 243-9; PMID:20190736; <http://dx.doi.org/10.1038/nature08779>
49. Haglund K, Dikic I. The role of ubiquitylation in receptor endocytosis and endosomal sorting. *J Cell Sci* 2012; 125:265-75; PMID:22357968; <http://dx.doi.org/10.1242/jcs.091280>
50. Rodahl LM, Stuffers S, Lobert VH, Stenmark H. The role of ESCRT proteins in attenuation of cell signalling. *Biochem Soc Trans* 2009; 37:137-42; PMID:19143618; <http://dx.doi.org/10.1042/BST0370137>
51. Jovic M, Sharma M, Rahajeng J, Caplan S. The early endosome: a busy sorting station for proteins at the crossroads. *Histol Histopathol* 2010; 25:99-112; PMID:19924646
52. Gowrishankar K, Ghosh S, Saha S, C R, Mayor S, Rao M. Active remodeling of cortical actin regulates spatiotemporal organization of cell surface molecules. *Cell* 2012; 149:1353-67; PMID:22682254; <http://dx.doi.org/10.1016/j.cell.2012.05.008>