Mullins, N. et al. (2021) Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. *Nature Genetics*, 53(6), pp. 817-829.

(doi: [10.1038/s41588-021-00857-4](https://doi.org/10.1038/s41588-021-00857-4))

This is the Author Accepted Manuscript.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/227778/>

Deposited on: 7 January 2021

Genome-wide association study of over 40,000 bipolar disorder cases provides novel biological insights

Authors

Niamh Mullins^{1,2,236,*}, Andreas J. Forstner^{3,4,5,236}, Kevin S. O'Connell^{6,7,237}, Brandon Coombes^{8,237}, Jonathan R. I. Coleman^{9,10,237}, Zhen Qiao^{11,237}, Thomas D. Als^{12,13,14}, Tim B. Bigdeli^{15,16}, Sigrid Børte^{17,18,19}, Julien Bryois²⁰, Alexander W. Charney², Ole Kristian Drange^{21,22}, Michael J. Gandal²³, Saskia P. Hagenaars^{9,10}, Masashi Ikeda²⁴, Nolan Kamitaki^{25,26}, Minsoo Kim²³, Kristi Krebs²⁷, Georgia Panagiotaropoulou²⁸, Brian M. Schilder^{1,29,30,31}, Laura G. Sloofman¹, Stacy Steinberg³², Vassily Trubetskoy²⁸, Bendik S. Winsvold^{19,33}, Hong-Hee Won³⁴, Liliya Abramova³⁵, Kristina Adorjan^{36,37}, Esben Agerbo^{14,38,39}, Mariam Al Eissa⁴⁰, Diego Albani⁴¹, Ney Alliey-Rodriguez^{42,43}, Adebayo Anjorin⁴⁴, Verner Antilla⁴⁵, Anastasia Antoniou⁴⁶, Swapnil Awasthi²⁸, Ji Hyun Baek⁴⁷, Marie Bækvad-Hansen^{14,48}, Nicholas Bass⁴⁰, Michael Bauer⁴⁹, Eva C. Beins³, Sarah E. Bergen²⁰, Armin Birner⁵⁰, Carsten Bøcker Pedersen^{14,38,39}, Erlend Bøen⁵¹, Marco P. Boks⁵², Rosa Bosch^{53,54,55,56}, Murielle Brum⁵⁷, Ben M. Brumpton¹⁹, Nathalie Brunkhorst-Kanaan⁵⁷, Monika Budde³⁶, Jonas Bybjerg-Grauholm^{14,48}, William Byerley⁵⁸, Murray Cairns⁵⁹, Miquel Casas^{53,54,55,56}, Pablo Cervantes⁶⁰, Toni-Kim Clarke⁶¹, Cristiana Cruceanu^{60,62}, Alfredo Cuellar-Barboza^{63,64}, Julie Cunningham⁶⁵, David Curtis^{66,67}, Piotr M. Czerski⁶⁸, Anders M. Dale⁶⁹, Nina Dalkner⁵⁰, Friederike S. David³, Franziska Degenhardt^{3,70}, Srdjan Djurovic^{71,72}, Amanda L. Dobbyn^{1,2}, Athanassios Douzenis⁴⁶, Torbjørn Elvsåshagen^{18,73,74}, Valentina Escott-Price⁷⁵, I. Nicol Ferrier⁷⁶, Alessia Fiorentino⁴⁰, Tatiana M. Foroud⁷⁷, Liz Forty⁷⁵, Josef Frank⁷⁸, Oleksandr Frei^{6,18}, Nelson B. Freimer^{23,79}, Louise Frisén⁸⁰, Katrin Gade^{36,81}, Julie Garnham⁸², Joel Gelernter^{83,84,85}, Marianne Giørtz Pedersen^{14,38,39}, Ian R. Gizer⁸⁶, Scott D. Gordon⁸⁷, Katherine Gordon-Smith⁸⁸, Tiffany A. Greenwood⁸⁹, Jakob Grove^{12,13,14,90}, José Guzman-Parra⁹¹, Kyoosob Ha⁹², Magnus Haraldsson⁹³, Martin Hautzinger⁹⁴, Urs Heilbronner³⁶, Dennis Hellgren²⁰, Stefan Herms^{3,95,96}, Per Hoffmann^{3,95,96}, Peter A. Holmans⁷⁵, Laura Huckins^{1,2}, Stéphane Jamain^{97,98}, Jessica S. Johnson^{1,2}, Janos L. Kalman^{36,37,99}, Yoichiro Kamatani^{100,101}, James L. Kennedy^{102,103,104,105}, Sarah Kittel-Schneider^{57,106}, James A. Knowles^{107,108}, Manolis Kogevinas¹⁰⁹, Maria Koromina¹¹⁰, Thorsten M. Kranz⁵⁷, Henry R. Kranzler^{111,112}, Michiaki Kubo¹¹³, Ralph Kupka^{114,115,116}, Steven A. Kushner¹¹⁷, Catharina Lavebratt^{118,119}, Jacob Lawrence¹²⁰, Markus Leber¹²¹, Heon-Jeong Lee¹²², Phil H. Lee¹²³, Shawn E. Levy¹²⁴, Catrin Lewis⁷⁵, Calwing Liao^{125,126}, Susanne Lucae⁶², Martin Lundberg^{118,119}, Donald J. MacIntyre¹²⁷, Wolfgang Maier¹²⁸, Adam Maihofer⁸⁹, Dolores Malaspina^{1,2}, Eirini Maratou¹²⁹, Lina Martinsson⁸⁰, Manuel Mattheisen^{12,13,14,106,130}, Nathaniel W. McGregor¹³¹, Peter McGuffin⁹, James D. McKay¹³², Helena Medeiros¹⁰⁸, Sarah E. Medland⁸⁷, Vincent Millischer^{118,119}, Grant W. Montgomery¹¹, Jennifer L. Moran^{25,133}, Derek W. Morris¹³⁴, Thomas W. Mühleisen^{4,95}, Niamh O'Brien⁴⁰, Claire O'Donovan⁸², Loes M. Olde Loohuis^{23,79}, Lilijana Oruc¹³⁵, Sergi Papiol^{36,37}, Antonio F. Pardiñas⁷⁵, Amy Perry⁸⁸, Andrea Pfennig⁴⁹, Evgenia Porichi⁴⁶, James B. Potash¹³⁶, Digby Quested^{137,138}, Towfique Raj^{1,29,30,31}, Mark H. Rapaport¹³⁹, J. Raymond DePaulo¹³⁶, Eline J. Regeer¹⁴⁰, John P. Rice¹⁴¹, Fabio Rivas⁹¹, Margarita Rivera^{142,143}, Julian Roth¹⁰⁶, Panos Roussos^{1,2,29}, Douglas M. Ruderfer¹⁴⁴, Cristina Sánchez-Mora^{53,54,56,145}, Eva C. Schulte^{36,37}, Fanny Senner^{36,37}, Sally Sharp⁴⁰, Paul D. Shilling⁸⁹, Engilbert Sigurdsson^{93,146}, Lea Sirignano⁷⁸, Claire Slaney⁸², Olav B. Smeland^{6,7}, Daniel J. Smith¹⁴⁷, Janet L. Sobell¹⁴⁸, Christine Sørholm Hansen^{14,48}, Maria Soler Artigas^{53,54,56,145}, Anne T. Spijker¹⁴⁹, Dan J. Stein¹⁵⁰, John S. Strauss¹⁰², Beata Świątkowska¹⁵¹, Chikashi Terao¹⁰¹, Thorgeir E. Thorgeirsson³², Claudio Toma^{152,153,154}, Paul Tooney⁵⁹, Evangelia-Eirini Tsermpini¹¹⁰, Marquis P. Vawter¹⁵⁵, Helmut Vedder¹⁵⁶, James T. R. Walters⁷⁵, Stephanie H. Witt⁷⁸, Simon Xi¹⁵⁷, Wei Xu¹⁵⁸, Jessica Mei Kay Yang⁷⁵, Allan H. Young^{159,160}, Hannah Young¹, Peter P. Zandi¹³⁶, Hang Zhou^{83,84}, Lea Zillich⁷⁸, HUNT All-In Psychiatry¹⁶¹, Rolf Adolfsson¹⁶², Ingrid Agartz^{51,130,163}, Martin Alda^{82,164}, Lars Alfredsson¹⁶⁵, Gulja Babadjanova¹⁶⁶, Lena Backlund^{118,119}, Bernhard T. Baune^{167,168,169}, Frank Bellivier^{170,171}, Susanne Bengesser⁵⁰, Wade H. Berrettini¹⁷², Douglas H. R. Blackwood⁶¹, Michael Boehnke¹⁷³, Anders D. Børghlum^{14,174,175}, Gerome Breen^{9,10}, Vaughan J. Carr¹⁷⁶, Stanley Catts¹⁷⁷, Aiden Corvin¹⁷⁸, Nicholas Craddock⁷⁵, Udo Dannlowski¹⁶⁷, Dimitris Dikeos¹⁷⁹, Tõnu Esko^{26,27,180,181}, Bruno Etain^{170,171}, Panagiotis Ferentinos^{9,46}, Mark Frye⁶⁴, Janice M. Fullerton^{152,153}, Micha Gawlik¹⁰⁶, Elliot S. Gershon^{42,182}, Fernando Goes¹³⁶, Melissa J. Green^{152,176}, Maria Grigoriou-Serbanescu¹⁸³, Joanna Hauser⁶⁸, Frans Henskens⁵⁹, Jan Hillert⁸⁰, Kyung Sue Hong⁴⁷, David M. Hougaard^{14,48}, Christina M. Hultman²⁰, Kristian Hveem^{19,184}, Nakao Iwata²⁴, Assen V. Jablensky¹⁸⁵, Ian Jones⁷⁵, Lisa A. Jones⁸⁸, René S. Kahn^{2,52}, John R. Kelsoe⁸⁹, George Kirov⁷⁵, Mikael Landén^{20,186}, Marion Leboyer^{97,98,187}, Cathryn M. Lewis^{9,10,188}, Qingqin S. Li¹⁸⁹, Jolanta Lissowska¹⁹⁰, Christine Lochner¹⁹¹, Carmel Loughland⁵⁹, Nicholas G. Martin^{87,192}, Carol A. Mathews¹⁹³, Fermin Mayoral⁹¹, Susan L. McElroy¹⁹⁴, Andrew M. McIntosh^{127,195}, Francis J. McMahon¹⁹⁶, Ingrid Melle^{6,197}, Patricia Michie⁵⁹, Lili Milani²⁷, Philip B. Mitchell¹⁷⁶, Gunnar Morken^{21,198}, Ole Mors^{14,199}, Preben Bo Mortensen^{12,14,38,39}, Bryan Mowry¹⁷⁷, Bertram Müller-Myhsok^{62,200,201}, Richard M. Myers¹²⁴, Benjamin M. Neale^{25,45,180}, Caroline M. Nievergelt^{89,202}, Merete Nordentoft^{14,203}, Markus M. Nöthen³, Michael C. O'Donovan⁷⁵, Ketil J. Oedegaard^{204,205}, Tomas Olsson²⁰⁶, Michael J.

Owen⁷⁵, Sara A. Paciga²⁰⁷, Chris Pantelis²⁰⁸, Carlos Pato¹⁰⁸, Michele T. Pato¹⁰⁸, George P. Patrinos^{110,209,210}, Roy H. Perlis^{211,212}, Danielle Posthuma^{213,214}, Josep Antoni Ramos-Quiroga^{53,54,55,56}, Andreas Reif⁵⁷, Eva Z. Reininghaus⁵⁰, Marta Ribasés^{53,54,56,145}, Marcella Rietschel⁷⁸, Stephan Ripke^{25,28,45}, Guy A. Rouleau^{126,215}, Takeo Saito²⁴, Ulrich Schall⁵⁹, Martin Schalling^{118,119}, Peter R. Schofield^{152,153}, Thomas G. Schulze^{36,78,81,136,216}, Laura J. Scott¹⁷³, Rodney J. Scott⁵⁹, Alessandro Serretti²¹⁷, Cynthia Shannon Weickert^{152,176,218}, Jordan W. Smoller^{25,133,219}, Hreinn Stefansson³², Kari Stefansson^{32,220}, Eystein Stordal^{221,222}, Fabian Streit⁷⁸, Patrick F. Sullivan^{20,223,224}, Gustavo Turecki²²⁵, Arne E. Vaaler²²⁶, Eduard Vieta²²⁷, John B. Vincent¹⁰², Irwin D. Waldman²²⁸, Thomas W. Weickert^{152,176,218}, Thomas Werge^{14,229,230,231}, Naomi R. Wray^{11,232}, John-Anker Zwart^{18,19,33}, Joanna M. Biernacka^{8,64}, John I. Nurnberger²³³, Sven Cichon^{3,4,95,96}, Howard J. Edenberg^{77,234}, Eli A. Stahl^{1,2,180,238}, Andrew McQuillin^{40,238}, Arianna Di Florio^{75,224,238}, Roel A. Ophoff^{23,79,117,235,238}, Ole A. Andreassen^{6,7,238,*}

Abstract

Bipolar disorder (BD) is a heritable mental illness with complex etiology. We performed a genome-wide association study (GWAS) of 41,917 BD cases and 371,549 controls of European ancestry, which identified 64 associated genomic loci. BD risk alleles were enriched in genes in synaptic signaling pathways and brain-expressed genes, particularly those with high specificity of expression in neurons of the prefrontal cortex and hippocampus. Significant signal enrichment was found in genes encoding targets of antipsychotics, calcium channel blockers and antiepileptics. Integrating eQTL data implicated 15 genes robustly linked to BD via gene expression, including druggable genes such as *HTR6*, *MCHR1*, *DCLK3* and *FURIN*. This GWAS provides the best-powered BD polygenic scores to date, when applied in both European and diverse ancestry samples. Analyses of BD subtypes indicated high but imperfect genetic correlation between BD type I and II and identified additional associated loci. Together, these results advance our understanding of the biological etiology of BD, identify novel therapeutic leads and prioritize genes for functional follow-up studies.

Introduction

Bipolar disorder (BD) is a complex mental disorder characterized by recurrent episodes of (hypo)mania and depression. It is a common condition affecting an estimated 40 to 50 million people worldwide¹. This, combined with the typical onset in young adulthood, an often chronic course, and increased risk of suicide², make BD a major public health concern and a major cause of global disability¹. Clinically, BD is classified into two main subtypes: bipolar I disorder, in which manic episodes typically alternate with depressive episodes, and bipolar II disorder, characterized by the occurrence of at least one hypomanic and one depressive episode³. These subtypes have a lifetime prevalence of ~1% each in the population^{4,5}.

Family and molecular genetic studies provide convincing evidence that BD is a multifactorial disorder, with genetic and environmental factors contributing to its development⁶. On the basis of twin and family studies, the heritability of BD is estimated at 60-85%^{7,8}. Genome-wide association studies (GWAS)⁹⁻²³ have led to valuable insights into the genetic etiology of BD. The largest such study has been conducted by the Psychiatric Genomics Consortium (PGC), in which genome-wide SNP data from 29,764 BD patients and 169,118 controls were analyzed and 30 genome-wide significant loci were identified (PGC2)²⁴. SNP-based heritability (h^2_{SNP}) estimation using the same data, suggested that common genetic variants genome-wide explain ~20% of BD's phenotypic variance²⁴. Polygenic risk scores generated from the results of this study explained ~4% of phenotypic variance in independent samples. Across the genome, genetic associations with BD converged on specific biological pathways including regulation of insulin secretion^{25,26}, retrograde endocannabinoid signaling²⁴, glutamate receptor signaling²⁷ and calcium channel activity⁹.

Despite this considerable progress, only a fraction of the genetic etiology of BD has been identified and the specific biological mechanisms underlying the development of the disorder are still unknown. In the

present study, we report the results of the third GWAS meta-analysis of the PGC Bipolar Disorder Working Group, comprising 41,917 patients with BD and 371,549 controls. These results confirm and expand on many previously reported findings, identify novel therapeutic leads and prioritize genes for functional follow-up studies^{28,29}. Thus, our results further illuminate the biological etiology of BD.

Results

GWAS results

A GWAS meta-analysis was conducted of 57 BD cohorts collected in Europe, North America and Australia (Table S1), totaling 41,917 BD cases and 371,549 controls of European descent (Effective N = 101,962, see online methods). For 52 cohorts, individual-level genotype and phenotype data were shared with the PGC and cases met international consensus criteria (DSM-IV, ICD-9 or ICD-10) for lifetime BD, established using structured diagnostic interviews, clinician-administered checklists or medical record review. BD GWAS summary statistics were received for five external cohorts (iPSYCH³⁰, deCODE genetics³¹, Estonian Biobank³², Trøndelag Health Study (HUNT)³³ and UK Biobank³⁴), in which most cases were ascertained using ICD codes. The GWAS meta-analysis identified 64 independent loci associated with BD at genome-wide significance ($P < 5E-08$) (Figure 1, Table 1, Table S2). Using LD Score regression (LDSC)³⁵ the h^2_{SNP} of BD was estimated to be 18.6% (SE=0.008, $P=5.1E-132$) on the liability scale, assuming a BD population prevalence of 2%, and 15.6% (SE=0.006, $P=5.0E-132$) assuming a population prevalence of 1% (Table S3). The genomic inflation factor (λ_{GC}) was 1.38 and the LD Score regression (LDSC) intercept was 1.04 (SE=0.01, $P=2.5E-04$) (Supplementary Figure 1). While the intercept has frequently been used as an indicator of confounding from population stratification, it can rise above 1 with increased sample size and heritability. The attenuation ratio - (LDSC intercept - 1)/(mean of association chi-square statistics - 1) - which is not subject to these limitations, was 0.06 (SE=0.02), indicating that the majority of inflation of the GWAS test statistics was due to polygenicity^{35,36}. Of the 64 genome-wide significant loci, 33 are novel discoveries (ie. loci not overlapping with any locus previously reported as genome-wide significant for BD). Novel loci include the major histocompatibility complex (MHC) and loci previously reaching genome-wide significance for other psychiatric disorders, including 10 for schizophrenia, 4 for major depression and 3 for childhood-onset psychiatric disorders or problematic alcohol use (Table 1).

Figure 1: Manhattan plot of genome-wide association meta-analysis of 41,917 bipolar disorder cases and 371,549 controls

The x-axis shows genomic position (chromosomes 1-22 and X) and the y-axis shows statistical significance as $-\log_{10}(P \text{ value})$. The red line shows the genome-wide significance threshold ($P < 5 \times 10^{-8}$). SNPs in genome-wide significant loci are colored green for loci previously associated with bipolar disorder and yellow for novel associations from this study. The genes labeled are those prioritized by integrative eQTL analyses or notable genes in novel loci (*MHC*, *CACNB2*, *KCNB1*).

Table 1: Genome-wide significant loci for bipolar disorder from meta-analysis of 41,917 cases and 371,549 controls

Locus	CHR	BP	SNP	P	OR	SE	A1/A2	A1 freq in controls	Previous report ^A for BD (citation)	Name for novel locus ⁺	Previous report ^A for psychiatric disorders
1	1	61105668	rs2126180	1.6E-09	1.058	0.009	A/G	0.457		LINC01748	
2	1	163745389	rs10737496	7.2E-09	1.056	0.009	C/T	0.444		NUF2	CDG
3*	2	97416153	rs4619651	4.8E-11	1.068	0.010	G/A	0.670	LMAN2L (PGC2)		CDG
4	2	166152389	rs17183814	2.7E-08	1.108	0.019	G/A	0.924	SCN2A (PGC2)		
5	2	169481837	rs13417268	2.1E-08	1.064	0.011	C/G	0.758		CERS6	
6	2	193738336	rs2011302	4.3E-08	1.055	0.010	A/T	0.377		PCGEM1	CDG
7	2	194437889	rs2719164	4.9E-08	1.053	0.010	A/G	0.564	intergenic (PGC2)		CDG
8*	3	36856030	rs9834970	6.6E-19	1.087	0.009	C/T	0.481	TRANK1 (PGC2)		SCZ, CDG
9*	3	52626443	rs2336147	3.6E-13	1.070	0.009	T/C	0.498	ITIH1 (PGC2)		SCZ, CDG
10	3	70488788	rs115694474	2.4E-08	1.068	0.012	T/A	0.799		MDFIC2	
11	3	107757060	rs696366	4.5E-08	1.053	0.009	C/A	0.550	CD47 (PGC2)		
12*	4	123076007	rs112481526	1.9E-09	1.065	0.011	G/A	0.256		KIAA1109	MD
13*	5	7542911	rs28565152	2.0E-09	1.070	0.011	A/G	0.238	ADCY2 (PGC2)		
14*	5	78849505	rs6865469	1.7E-08	1.060	0.010	T/G	0.274		HOMER1	
15	5	80961069	rs6887473	8.8E-09	1.062	0.011	G/A	0.739	SSBP2 (PGC2)		
16*	5	137712121	rs10043984	3.7E-08	1.062	0.011	T/C	0.236		KDM3B	CDG
17	5	169289206	rs10866641	2.8E-11	1.065	0.009	T/C	0.575		DOCK2	
18*	6	26463575	rs13195402	5.8E-15	1.146	0.018	G/T	0.919		MHC	MD, SCZ, CDG, MOOD
19*	6	98565211	rs1487445	1.5E-15	1.078	0.009	T/C	0.487	POU3F2 (PGC2)		CDG
20	6	152793572	rs4331993	2.0E-08	1.056	0.010	A/T	0.382	SYNE1 (Green, 2013)		
21*	6	166995260	rs10455979	4.2E-09	1.057	0.010	G/C	0.500	RPS6KA2 (PGC2)		
22*	7	2020995	rs12668848	1.9E-09	1.059	0.010	G/A	0.575	MAD1L1 (Hou, 2016, Ikeda, 2017)		MD, SCZ, CDG
23*	7	11871787	rs113779084	1.4E-13	1.079	0.010	A/G	0.299	THSD7A (PGC2)		
24*	7	21492589	rs6954854	5.9E-10	1.060	0.009	G/A	0.425		SP4	
25	7	24647222	rs12672003	2.7E-09	1.096	0.016	G/A	0.113		MPP6	SCZ, CDG, MOOD
26	7	105043229	rs11764361	3.5E-09	1.063	0.010	A/G	0.668	SRPK2 (PGC2)		SCZ, ASD, CDG
27	7	131870597	rs6946056	3.7E-08	1.055	0.010	C/A	0.623		PLXNA4	
28	7	140676153	rs10255167	1.6E-08	1.068	0.012	A/G	0.778	MRPS33 (PGC2)		CDG
29*	8	9763581	rs62489493	2.6E-11	1.094	0.014	G/C	0.128		miR124-1	SCZ, ALC, ASD
30*	8	10226355	rs3088186	2.1E-08	1.058	0.010	T/C	0.287		MSRA	SCZ, ALC, ASD
31	8	34152492	rs2953928	6.3E-09	1.124	0.020	A/G	0.067		RP1-84O15.2 (lincRNA)	SCZ, ADHD, CDG
32*	8	144993377	rs6992333	1.6E-09	1.062	0.010	G/A	0.410		PLEC	
33	9	37090538	rs10973201	2.5E-08	1.101	0.017	C/T	0.110		ZCCHC7	MD, CDG, MOOD
34*	9	141066490	rs62581014	2.8E-08	1.067	0.012	T/C	0.366		TUBBP5	
35*	10	18751103	rs1998820	4.1E-08	1.087	0.015	T/A	0.886		CACNB2	SCZ, CDG
36*	10	62322034	rs10994415	1.1E-11	1.125	0.017	C/T	0.082	ANK3 (PGC2)		
37	10	64525135	rs10761661	4.7E-08	1.053	0.009	T/C	0.472		ADO	
38*	10	111648659	rs2273738	1.6E-11	1.096	0.014	T/C	0.135	ADD3 (Charney, 2017, PGC2)		
39*	11	61618608	rs174592	9.9E-14	1.074	0.010	G/A	0.395	FADS2 (PGC2)		MD, CDG, MOOD
40	11	64009879	rs4672	3.4E-09	1.107	0.017	A/G	0.083		FKBP2	
41*	11	65848738	rs475805	2.0E-09	1.070	0.011	A/G	0.767	PACS1 (PGC2)		
42*	11	66324583	rs678397	5.5E-09	1.056	0.009	T/C	0.457	PC (PGC1, PGC2)		
43*	11	70517927	rs12575685	1.2E-10	1.067	0.010	A/G	0.327	SHANK2 (PGC2)		MD
44	11	79092527	rs12289486	3.3E-08	1.086	0.015	T/C	0.115	ODZ4 (PGC1)		
45*	12	2348844	rs11062170	1.9E-15	1.081	0.010	C/G	0.333	CACNA1C (PGC2)		SCZ, CDG, MOOD
46	13	113869045	rs35306827	3.6E-09	1.068	0.011	G/A	0.775		CUL4A	
47	14	99719219	rs2693698	2.0E-08	1.055	0.009	G/A	0.551		BCL11B	SCZ, CDG
48*	15	38973793	rs35958438	3.8E-08	1.066	0.012	G/A	0.772		C15orf53	CDG
49*	15	42904904	rs4447398	2.6E-09	1.086	0.014	A/C	0.131	STARD9 (PGC2)		
50	15	83531774	rs62011709	1.4E-08	1.064	0.011	T/A	0.747		HOMER2	SCZ
51*	15	85149575	rs748455	5.0E-11	1.070	0.010	T/C	0.719	ZNF592 (PGC2)		SCZ, CDG
52	15	91426560	rs4702	3.5E-09	1.059	0.010	G/A	0.446		FURIN	SCZ, CDG
53	16	9230816	rs28455634	2.6E-10	1.065	0.010	G/A	0.620		C16orf72	
54	16	9926348	rs7199910	1.7E-08	1.057	0.010	G/T	0.312	GRIN2A (PGC2)		SCZ, CDG
55	16	89632725	rs12932628	6.7E-09	1.058	0.010	T/G	0.487		RPL13	
56	17	1835482	rs4790841	3.1E-08	1.075	0.013	T/C	0.151		RTN4RL1	
57	17	38129841	rs11870683	2.8E-08	1.059	0.010	T/A	0.650	ERBB2 (Hou, 2016)		
58	17	38220432	rs61554907	1.6E-08	1.091	0.015	T/G	0.124	ERBB2 (Hou, 2016)		
59*	17	42191893	rs228768	2.8E-10	1.067	0.010	G/T	0.294	HDAC5 (PGC2)		
60*	20	43682551	rs67712855	4.2E-11	1.070	0.010	T/G	0.687	STK4 (PGC2)		
61*	20	43944323	rs6032110	1.0E-09	1.059	0.009	A/G	0.512	WFDC12 (PGC2)		
62*	20	48033127	rs237460	4.3E-09	1.057	0.009	T/C	0.412		KCNB1	CDG
63	20	60865815	rs13044225	8.5E-09	1.056	0.010	G/A	0.440		OSBPL2	
64	22	41153879	rs5758064	2.0E-08	1.054	0.009	T/C	0.523		SLC25A17	MD, SCZ, CDG, MOOD

CHR, chromosome; BP, GRCh37 basepair position; SNP, single nucleotide polymorphism; OR, odds ratio; SE, standard error, A1, tested allele; A2, other allele; freq, frequency; BD, bipolar disorder; CDG, Cross-disorder GWAS of the Psychiatric Genomics Consortium; MD, major depression; SCZ, schizophrenia; MOOD, mood disorders; ASD, Autism Spectrum Disorder; ALC, Alcohol use disorder or problematic alcohol use; ADHD, attention deficit/hyperactivity disorder. *Locus overlaps with genome-wide significant locus for bipolar I disorder. ^APrevious report refers to previous association of a SNP in the locus with the psychiatric disorder at genome-wide significance. PGC1 = PMID 21926972, PGC2 = PMID 31043756, Hou, 2016 = PMID 27329760, Ikeda, 2017 = PMID:28115744, Green, 2013 = PMID 22565781. Charney, 2017 = PMID 28072414. ⁺Novel loci are named using the nearest gene to the index SNP.

Enrichment analyses

Genome-wide analyses using MAGMA³⁷ indicated significant enrichment of BD associations in 161 genes (Table S4) and 4 gene sets, related to synaptic signaling (Table S5). The BD association signal was enriched amongst genes expressed in different brain tissues (Table S6), especially genes with high specificity of gene expression in neurons (both excitatory and inhibitory) versus other cell types, within cortical and subcortical brain regions in mice (Supplementary Figure 2)³⁸. In human brain samples, signal enrichment was also observed in hippocampal pyramidal neurons and interneurons of the prefrontal cortex and hippocampus, compared with other cell types (Supplementary Figure 2).

In a gene-set analysis of the targets of individual drugs (from the Drug-Gene Interaction Database DGIdb v.2³⁹ and the Psychoactive Drug Screening Database Ki DB⁴⁰), no individual set was significantly enriched (Table S7). Grouping drugs according to their Anatomical Therapeutic Chemical (ATC) classes⁴¹, there was significant enrichment in the targets of four broad drug classes (Table S8): psycholeptics (drugs with a calming effect on behavior) (especially antipsychotics, hypnotics and sedatives, anxiolytics, and benzodiazepines), calcium channel blockers (especially those with mainly vascular effects and dihydropyridine derivatives), antiepileptics and drugs for functional gastrointestinal disorders (which include anticholinergics and serotonin receptor antagonists) (Table S8).

eQTL integrative analyses

A transcriptome-wide association study (TWAS) was conducted using FUSION⁴² and eQTL data from the PsychENCODE Consortium (1,321 brain samples)⁴³. BD-associated alleles significantly influenced expression of 77 genes in the brain (Table S9, Supplementary Figure 3). These genes encompassed 40 distinct regions. TWAS fine-mapping was performed using FOCUS⁴⁴ to model the correlation among the TWAS signals and prioritize the most likely causal gene(s) in each region. Within the 90%-credible set, FOCUS prioritised 22 genes with a posterior inclusion probability (PIP) > 0.9 (encompassing 20 distinct regions) and 32 genes with a PIP > 0.7 (29 distinct regions) (Table S10).

Summary data-based Mendelian randomization (SMR)^{45,46} was used to identify putative causal relationships between SNPs and BD via gene expression by integrating the BD GWAS results with brain eQTL summary statistics from the PsychENCODE⁴³ Consortium and blood eQTL summary statistics from the eQTLGen Consortium (31,684 whole blood samples)⁴⁷. The eQTLGen results represent the largest existing eQTL study and provide independent eQTL data. Of the 32 genes fine-mapped with PIP > 0.7, 15 were significantly associated with BD in the SMR analyses and passed the HEIDI (heterogeneity in dependent instruments) test^{45,46}, suggesting that their effect on BD is mediated via gene expression in the brain and/or blood (Table S11). The genes located in genome-wide significant loci are labeled in Figure 1. Other significant genes included *HTR6*, *DCLK3*, *HAPLN4* and *PACSLN2*.

MHC locus

Variants within and distal to the major histocompatibility complex (MHC) locus were associated with BD at genome-wide significance. The most highly associated SNP was rs13195402, 3.2 megabases distal to any *HLA* gene or the complement component 4 (*C4*) genes (Supplementary Figure 4). Imputation of *C4* alleles using SNP data uncovered no association between the five most common structural forms of the *C4A/C4B* locus (BS, AL, AL-BS, AL-BL, and AL-AL) and BD, either before or after conditioning on rs13195402 (Supplementary Figure 5). While genetically predicted *C4A* expression initially showed a weak association with BD, this association was non-significant after controlling for rs13195402 (Supplementary Figure 6).

Polygenic risk scoring

The performance of polygenic risk scores (PRS) based on these GWAS results was assessed by excluding cohorts in turn from the meta-analysis to create independent test samples. PRS explained ~4.57% of phenotypic variance in BD on the liability scale (at GWAS P value threshold (p_T) < 0.1, BD population prevalence 2%), based on the weighted mean R^2 across cohorts (Figure 2, Table S12). This corresponds to a weighted mean area under the curve (AUC) of 66%. Results per cohort and per wave of recruitment to the PGC are in Tables S12-S13 and Supplementary Figure 7. At p_T < 0.1, individuals in the top 10% of BD PRS had an odds ratio of 3.62 (95% CI 1.7-7.9) of being affected with the disorder compared with individuals in the middle decile (based on the weighted mean OR across PGC cohorts), and an odds ratio of 9.5 (95% CI 5.4-20) compared with individuals in the lowest decile. The generalizability of PRS from this meta-analysis was examined in several non-European cohorts. PRS explained up to 2.3% and 1.9% of variance in BD in two East Asian samples, and 1.2% and 0.4% in two admixed African American samples (Figure 2, Table S14). The variance explained by the PRS increased in every cohort with increasing sample size of the PGC BD European discovery sample (Supplementary Figure 8, Table S14).

Figure 2: Phenotypic variance in bipolar disorder explained by polygenic risk scores

Variance explained is presented on the liability scale, assuming a BD population prevalence of 2%. For European ancestry, the results shown are the weighted mean R^2 values across all cohorts analyzed, calculated weighted by the effective N per cohort. The numbers of cases and controls are shown under the barplot for each study. GWAS p_T - the color of the bars represents the P value threshold used to select SNPs from the discovery GWAS. GAIN-AA - Genetic Association Information Network African American cohort, AA-GPC - African American Genomic Psychiatry Cohort.

Genetic architecture of BD and other traits

The genome-wide genetic correlation (r_g) of BD with a range of diseases and traits was assessed on LD Hub⁴⁸. After correction for multiple testing, BD showed significant r_g with 16 traits among 255 tested from published GWAS (Table S15). Genetic correlation was positive with all psychiatric disorders assessed, particularly schizophrenia ($r_g = 0.68$) and major depression ($r_g=0.44$), and to a lesser degree anorexia, attention deficit/hyperactivity disorder and autism spectrum disorder ($r_g \approx 0.2$). We found evidence of positive r_g between BD and smoking initiation, cigarettes per day, problematic alcohol use and drinks per week (Figure 3). BD was also positively genetically correlated with measures of sleep quality (daytime sleepiness, insomnia, sleep duration) (Figure 3). Among 514 traits measured in the general population of the UK Biobank, there was significant r_g between BD and many psychiatric-relevant traits or symptoms, dissatisfaction with interpersonal relationships, poorer overall health rating and feelings of loneliness or isolation (Table S16).

Bivariate gaussian mixture models were applied to the GWAS summary statistics for BD and other complex traits using the MiXeR tool^{49,50} to estimate the number of variants influencing each trait that explain 90% of h_{SNP}^2 and their overlap between traits. MiXeR estimated that approximately 8.6 k (SE=0.2 k) variants influence BD, which is similar to the estimate for schizophrenia (9.7 k, SE=0.2 k) and somewhat lower than that for major depression (12.3 k, SE=0.6 k) (Table S17, Supplementary Figure 9). When considering the number of shared loci as a proportion of the total polygenicity of each trait, the vast majority of loci influencing BD were also estimated to influence major depression (97%) and schizophrenia (96%) (Table S17, Supplementary Figure 9). Interestingly, within these shared components, the variants that influenced both BD and SCZ had high concordance in direction of effect (80%, SE=2%), while the portion of concordant variants between BD and MDD was only 69% (SE=1%) (Table S17).

Genetic and causal relationships between BD and modifiable risk factors

Ten traits associated with BD from clinical and epidemiological studies were investigated in detail for genetic and potentially causal relationships with BD via LDSC³⁵, generalized summary statistics-based Mendelian randomization (GSMR)⁵¹ and bivariate gaussian mixture modeling⁴⁹. BD has been strongly linked with sleep disturbances⁵², alcohol use⁵³ and smoking⁵⁴, higher educational attainment^{55,56} and mood instability⁵⁷ (which is a defining characteristic of the disorder). Most of these traits had modest but significant genetic correlations with BD (r_g -0.05-0.35) (Figure 3). Examining the effects of these traits on BD via GSMR, smoking initiation was associated with BD, corresponding to an OR of 1.49 (95% CI 1.38-1.61) for developing the disorder ($P=1.74E-22$) (Figure 3). Testing the effect of BD on the traits, BD was significantly associated with reduced likelihood of being a morning person and increased number of drinks per week ($P<1.47E-03$) (Figure 3). Positive bi-directional relationships were identified between BD and longer sleep duration, problematic alcohol use, educational attainment (EA) and mood instability (Figure 3). Notably, the instrumental variables for mood instability were selected from a GWAS conducted in the general population, excluding individuals with psychiatric disorders⁵⁸. For all of the aforementioned BD-trait relationships, the effect size estimates from GSMR were consistent with those calculated using the inverse variance weighted regression method, and there was no evidence of bias from horizontal pleiotropy. Full MR results are in Tables S18-19. Bivariate gaussian mixture modeling using MiXeR, indicated large proportions of variants influencing both BD and all other traits tested, particularly educational attainment, where approximately 98% of variants influencing BD were estimated to also influence EA. While cigarettes per day was a trait of interest, MiXeR could not model these data due to low polygenicity and heritability, and the effect of cigarettes per day on BD was inconsistent between MR methods, suggesting a violation of MR assumptions (Tables S18-20).

Figure 3: Relationships between bipolar disorder and modifiable risk factors based on genetic correlations, generalized summary statistics-based Mendelian randomization and bivariate gaussian mixture modeling

Venn diagrams depict MiXeR results of the estimated number of influencing variants shared between BD and each trait of interest (grey), unique to BD (blue) and unique to the trait of interest (orange). The number of influencing variants and standard error are shown in thousands. The size of the circles reflects the polygenicity of each trait, with larger circles corresponding to greater polygenicity. The estimated genetic correlation (r_g) between BD and each trait of interest and standard error from LD Score regression is shown below the corresponding Venn diagram, with an accompanying scale (-1 to +1). The arrows above and below the Venn diagrams indicate the results of generalized summary statistics-based Mendelian randomization (GSMR) of BD on the trait of interest, and the trait of interest on BD, respectively. The GSMR effect size and standard error is shown inside the corresponding arrow. Solid arrows indicate a significant relationship between the exposure and the outcome, after correction for multiple comparisons ($P < 1.47E-03$) and dashed arrows indicate a non-significant relationship.

BD subtypes

We conducted GWAS meta-analyses of bipolar I disorder (BD I) (25,060 cases, 449,978 controls) and bipolar II disorder (BD II) (6,781 cases, 364,075 controls). The BD I analysis identified 44 genome-wide significant loci, 31 of which overlapped with genome-wide significant loci from the main BD GWAS (Table 1, Table S21). The remaining 13 genome-wide significant loci for BD I all had $P < 4.0E-05$ in the main BD GWAS. One genome-wide significant locus was identified in the GWAS meta-analysis of BD II and had a $P < 1.1E-04$ in the main GWAS of BD (Table S21). The h^2_{SNP} estimates on the liability scale for BD I and BD II

were 20.9% (SE=0.009, $P=1.0E-111$) and 11.6% (SE=0.01, $P = 3.9E-15$), respectively, assuming a 1% population prevalence of each subtype. These heritability values are significantly different from each other ($P=2.4E-25$, block jackknife). The genetic correlation between BD I and BD II was 0.85 (SE=0.05, $P = 2.88E-54$), which is significantly different from 1 ($P=1.6E-3$). The genetic correlation of BD I with schizophrenia ($r_g=0.66$, SE=0.02) was higher than that of BD II ($r_g=0.54$ SE=0.05), whereas major depression was more strongly genetically correlated with BD II ($r_g=0.66$, SE=0.05) than with BD I ($r_g=0.34$, SE=0.03) (Table S22).

Discussion

In a GWAS of 41,917 BD cases, we identify 64 associated genomic loci, 33 of which are novel discoveries. With a 1.5-fold increase in effective sample size compared with the PGC2 BD GWAS, this study more than doubled the number of associated loci, representing an inflection point in the rate of risk variant discovery. We observed consistent replication of known BD loci, including 28/30 loci from the PGC2 GWAS²⁴ and several implicated by other BD GWAS^{15,16,17}, including a study of East Asian cases⁵⁹.

The 33 novel loci discovered here encompass genes of expected biological relevance in BD, such as the ion channels *CACNB2* and *KCNB1*. Amongst the 64 BD loci, 17 have previously been implicated in GWAS of schizophrenia⁶⁰, and seven in GWAS of major depression⁶¹, representing the first overlap of genome-wide significant loci between the mood disorders. For the BD loci shared across disorders, 17/17 and 5/7 of the BD index SNPs had the same direction of effect on schizophrenia and major depression respectively (Table S23). Bivariate gaussian mixture modeling estimated that across the entire genome, almost all variants influencing BD also influence schizophrenia and major depression, albeit with variable effects⁶². SNPs in and around the MHC locus reached genome-wide significance for BD for the first time. However, unlike in schizophrenia, we found no influence of *C4* structural alleles or gene expression⁶³. Rather the association was driven by variation outside the classical MHC locus, with the index SNP (rs13195402) being a missense variant in *BTN2A1*, a brain-expressed gene⁶⁴ encoding a plasma membrane protein.

The genetic correlation of BD with other psychiatric disorders was consistent with previous reports^{65,66}. Our results also corroborate previous genetic and clinical evidence of associations between BD and sleep disturbances⁶⁷, problematic alcohol use⁶⁸ and smoking⁶⁹. While the genome-wide genetic correlations with these traits were modest (r_g -0.05-0.35), MiXeR estimated that for all traits, more than 55% of trait-influencing variants also influence BD (Figure 3). Taken together, these results point to shared biology as one possible explanation for the high prevalence of substance use in BD. However, excluding genetic variants associated with both traits, MR analyses suggested that smoking is also a putatively “causal” risk factor for BD, while BD has no effect on smoking, consistent with a previous report⁷⁰. [We use the word “causal” with caution here as we consider MR an exploratory analysis to identify potentially modifiable risk factors which warrant more detailed investigations to understand their complex relationship with BD.] In contrast, MR indicated that BD had bi-directional “causal” relationships with problematic alcohol use, longer sleep duration and mood instability. Insights into the relationship of such behavioral correlates with BD may have future impact on clinical decision making in the prophylaxis or management of the disorder. Higher educational attainment has previously been associated with BD in epidemiological studies^{55,56}, while lower educational attainment has been associated with schizophrenia and major depression^{71,72}. Here, educational attainment had a significant positive effect on risk of BD and vice versa. Interestingly, MiXeR estimated that almost all variants that influence BD also influence educational attainment. The substantial genetic overlap observed between BD and the other phenotypes suggests that many variants likely influence multiple phenotypes which may be differentiated by phenotype-specific effect size distributions among the shared influencing variants.

The integration of eQTL data with our GWAS results yielded 15 high-confidence genes for which there was converging evidence that their association with BD is mediated via gene expression. Amongst these were *HTR6*, a serotonin receptor targeted by antipsychotics and antidepressants⁷³ and *MCHR1* (melanin-concentrating hormone receptor 1), a target of the antipsychotic haloperidol⁷³. We note that for both of these genes, their top eQTLs have opposite directions of effect on gene expression in the brain and blood, possibly playing a role in the tissue-specific gene regulation influencing BD⁷⁴. BD was associated with decreased expression of *FURIN*, a gene with a neurodevelopmental function which has already been the subject of functional genomics experiments in neuronal cells, following its association with schizophrenia in GWAS⁷⁵. The top association in our GWAS was in the *TRANK1* locus on chromosome 3, which has previously been implicated in BD^{12,18,59}. Although BD-associated SNPs in this locus are known to regulate *TRANK1* expression⁷⁶, our eQTL analyses support a stronger but correlated regulation of *DCLK3*, located 87 kb upstream of *TRANK1*^{43,77}. Both *FURIN* and *DCLK3* also encode druggable proteins (although they are not targets for any current psychiatric medications)^{73,78}. These eQTL results provide promising BD candidate genes for functional follow-up experiments²⁹. While several of these are in genome-wide significant loci, many are not the closest gene to the index SNP, highlighting the value of probing underlying molecular mechanisms to prioritize the most likely causal genes in the loci.

GWAS signals were enriched in the gene targets of existing BD pharmacological agents, such as antipsychotics, mood stabilizers, and antiepileptics. However, enrichment was also found in the targets of calcium channel blockers, which are used to treat hypertension, and drugs for treating functional gastrointestinal disorders, many of which target cholinergic and serotonergic receptors (Table S8). Indeed, calcium channel antagonists have long been used in the treatment of BD, without becoming an established therapeutic approach and there is evidence that some antiepileptics have calcium channel-inhibiting effects^{79,80}. These results underscore the opportunity for repurposing some classes of drugs, particularly calcium channel antagonists, as potential BD treatments⁸¹. While there was no significant enrichment in the targets of any individual drug tested, this analysis may currently be limited by statistical power.

BD associations were enriched in gene sets involving neuronal parts and synaptic signaling. Neuronal and synaptic pathways have been described in cross-disorder GWAS of multiple psychiatric disorders including BD^{82–84}. Dysregulation of such pathways has also been suggested by previous functional and animal studies⁸⁵. Analysis of single-cell gene expression data revealed enrichment in genes with high specificity of gene expression in neurons (both excitatory and inhibitory), of many brain regions, in particular the cortex and hippocampus. These findings are similar to those reported in GWAS data of schizophrenia⁸⁶ and major depressive disorder³⁸.

PRS for BD explained on average 4.57% of phenotypic variance (liability scale) across European cohorts, although this varied in different waves of the BD GWAS, ranging from 6.6% in the PGC1 cohorts to 2.9% in the External biobank studies (Supplementary Figure 7, Table S12). These results are in line with the h^2_{SNP} of BD per wave, which ranged from 24.6% (SE=0.01) in PGC1 to 11.9% (SE=0.01) in External studies (Table S3). Some variability in h^2_{SNP} estimates may arise from the inclusion of cases from population biobanks ascertained using ICD codes, who may have more heterogeneous clinical presentations or less severe illness than BD patients ascertained via inpatient or outpatient psychiatric clinics. Across the waves of clinically ascertained samples within the PGC, h^2_{SNP} and the R^2 of PRS also varied, likely reflecting clinical and genetic heterogeneity in the type of BD cases ascertained; the PGC1 cohorts consisted mostly of BD I cases⁹, known to be the most heritable of the BD subtypes^{11,24}, while later waves included more individuals with BD II²⁴. Overall, the h^2_{SNP} of BD calculated from the meta-analysis summary statistics was 18% on the

liability scale, a decrease of ~2% compared with the PGC2 GWAS²⁴, which may be due to the addition of cohorts with lower h_{SNP}^2 estimates and heterogeneity between cohorts (Table S3). However, despite differences in h_{SNP}^2 and R^2 of PRS per wave, the genetic correlation of BD between all waves was high (weighted mean $r_g=0.94$, $SE=0.03$), supporting our rationale for combining cases with different BD subtypes or ascertainment to increase power for discovery of risk variants. In Europeans, individuals in the top 10% of PRS had an OR of 3.6 for BD, compared with individuals with average PRS (middle decile), which translates into a modest absolute lifetime risk of the disorder (7.2% based on PRS alone). While PRS are invaluable tools in research settings, the current BD PRS lack sufficient power to separate individuals into clinically meaningful risk categories, and therefore have no clinical utility at present^{87,88}. PRS from this European BD meta-analysis yield higher R^2 values in diverse ancestry samples than PRS based on any currently available BD GWAS within the same ancestry⁵⁹. However, performance still greatly lags behind that in Europeans, with ~2% variance explained in East Asian samples and substantially less in admixed African American samples, likely due to differences in allele frequencies and LD structures, consistent with previous studies^{89,90}. There is a pressing need for more and larger studies in other ancestry groups to ensure that any future clinical utility is broadly applicable. Exploiting the differences in LD structure between diverse ancestry samples will also assist in the fine-mapping of risk loci for BD.

Our analyses confirmed that BD is a highly polygenic disorder, with an estimated 8.6 k variants explaining 90% of its h_{SNP}^2 . Hence, many more SNPs than those identified here are expected to account for the common variant architecture underlying BD. This GWAS marks an inflection point in risk variant discovery and we expect that from this point forward, the addition of more samples will lead to a dramatic increase in genetic findings. Nevertheless, fewer genome-wide significant loci have been identified in BD than in a schizophrenia GWAS of comparable sample size⁶⁰. This may be due to the clinical and genetic heterogeneity that exists in BD.

Our GWAS of subtypes BD I and BD II identified additional associated loci. Consistent with previous findings²⁴, our analysis showed that the two subtypes were highly but imperfectly genetically correlated ($r_g=0.85$), and that BD I is more genetically correlated with schizophrenia, while BD II has stronger genetic correlation with major depression. The subtypes are sufficiently similar to justify joint analysis as BD, but are not identical in their genetic composition, and as such contribute to the genetic heterogeneity of BD⁹¹. We identified thirteen loci passing genome-wide significance for BD I, and one for BD II, which did not reach significance in the main BD GWAS, further illustrating the partially differing genetic composition of the two subtypes. Understanding the shared and distinct genetic components of BD subtypes and symptoms requires detailed phenotyping efforts in large cohorts, and is an important area for future psychiatric genetic research.

In summary, these new data advance our understanding of the biological etiology of BD and prioritize a set of candidate genes for functional follow-up experiments. Several lines of evidence converge on the involvement of calcium channel signaling, providing a promising avenue for future therapeutic development.

Methods

Sample description

The meta-analysis sample comprises 57 cohorts collected in Europe, North America and Australia, totaling 41,917 BD cases and 371,549 controls of European descent (Table S1). The total effective N, equivalent to an equal number of cases and controls in each cohort ($4*N_{cases}*N_{controls}/(N_{cases}+N_{controls})$), is 101,962. For 52 cohorts, individual-level genotype and phenotype data were shared with the PGC. Cohorts

have been added to the PGC in five waves (PGC1⁹, PGC2²⁴, PGC PsychChip, PGC3 and External Studies); all cohorts from previous PGC BD GWAS were included. The source and inclusion/exclusion criteria for cases and controls for each cohort, are described in the Supplementary Note. Cases were required to meet international consensus criteria (DSM-IV, ICD-9 or ICD-10) for a lifetime diagnosis of BD, established using structured diagnostic instruments from assessments by trained interviewers, clinician-administered checklists or medical record review. In most cohorts, controls were screened for the absence of lifetime psychiatric disorders and randomly selected from the population. For five cohorts (iPSYCH³⁰, deCODE genetics³¹, Estonian Biobank³², Trøndelag Health Study (HUNT)³³ and UK Biobank³⁴), GWAS summary statistics for BD were shared with the PGC. In these cohorts, BD cases were ascertained using ICD codes or self-report during a nurse interview, and the majority of controls were screened for the absence of psychiatric disorders via ICD codes. Follow-up analyses included four non-European BD case-control cohorts, two from East Asia (Japan⁵⁹ and Korea⁹²), and two admixed African American cohorts^{22,93}, providing a total of 5,847 cases and 65,588 controls. These BD cases were ascertained through psychiatric interviews (Supplementary Note).

Genotyping, quality control and imputation

PGC cohorts were genotyped following local protocols, after which standardized quality control, imputation and statistical analyses were performed centrally using RICOPII (Rapid Imputation for CONsortias PIpeLine)⁹⁴, separately for each cohort. Briefly, the quality control parameters for retaining SNPs and subjects were: SNP missingness < 0.05 (before sample removal), subject missingness < 0.02, autosomal heterozygosity deviation ($F_{het} < 0.2$), SNP missingness < 0.02 (after sample removal), difference in SNP missingness between cases and controls < 0.02, SNP Hardy-Weinberg equilibrium ($P > 10^{-10}$ in psychiatric cases and $P > 10^{-6}$ in controls). Relatedness was calculated across cohorts using identity by descent and one of each pair of related individuals ($\pi_{hat} > 0.2$) was excluded. Genotype imputation was performed using the pre-phasing/ imputation stepwise approach implemented in Eagle v2.3.5⁹⁵ and Minimac3⁹⁶ to the Haplotype Reference Consortium (HRC) reference panel v1.0⁹⁷. The five external cohorts were processed by the collaborating research teams using comparable procedures and imputed to the HRC or a custom reference panel as appropriate. Full details of the genotyping, quality control and imputation for each cohort are available in the Supplementary Note. Identical individuals between PGC cohorts and the Estonian Biobank and UK Biobank cohorts were detected using genotype-based checksums

(https://personal.broadinstitute.org/sripke/share_links/zpXkv8INxUg9bayDpLT0G4g58TMtjN_PGC_SCZ_w3.0718d.76) and removed from PGC cohorts.

Genome-wide association study

For PGC cohorts, GWAS were conducted within each cohort using an additive logistic regression model in PLINK v1.90⁹⁸. Ancestry informant principal components (PCs) were calculated within each cohort and included as covariates, as required. Association analyses of the X chromosome were conducted in males and females separately using the same procedures, with males coded as 0 or 2 for 0 or 1 copies of the reference allele. Results from males and females were then meta-analyzed within each cohort. For external cohorts, GWAS were conducted by the collaborating research teams using comparable procedures (Supplementary Note). To control test statistic inflation at SNPs with low minor allele frequency (MAF) in small cohorts, SNPs were retained only if cohort MAF was > 1% and minor allele count was > 10 in either cases or controls (whichever had smaller N). There was no evidence of stratification artifacts or uncontrolled inflation of test statistics in the results from any cohort (λ_{GC} 0.97-1.05)(Table S1). Meta-analysis of GWAS summary statistics was conducted using an inverse variance-weighted fixed effects model in METAL⁹⁹. A genome-wide significant locus was defined as the region around a SNP with $P < 5E-08$, with linkage disequilibrium (LD) $r^2 > 0.1$, within a 3000 kilobase (kb) window.

Overlap of loci with other psychiatric disorders

Genome-wide significant loci for BD were assessed for overlap with genome-wide significant loci for other psychiatric disorders, using the largest available GWAS results for major depression⁶¹, schizophrenia⁶⁰, attention deficit/hyperactivity disorder¹⁰⁰, post-traumatic stress disorder¹⁰¹, lifetime anxiety disorder¹⁰², Tourette's Syndrome¹⁰³, anorexia nervosa¹⁰⁴, alcohol use disorder or problematic alcohol use⁶⁸, autism spectrum disorder¹⁰⁵, mood disorders⁹¹ and the cross-disorder GWAS of the Psychiatric Genomics Consortium⁶⁶. The boundaries of the genome-wide significant loci were calculated in the original publications. Overlap of loci was calculated using bedtools¹⁰⁶.

Enrichment analyses

P values quantifying the degree of association of genes and gene sets with BD were calculated using MAGMA v1.08³⁷, implemented in FUMA v1.3.6a^{64,107}. MAGMA uses Brown's method to combine SNP P values, while accounting for LD. Gene-based tests were performed for 19,576 genes (Bonferroni-corrected P value threshold = 2.55E-06). A total of 11,858 curated gene sets including at least 10 genes from MSigDB V7.0 were tested for association with BD (Bonferroni-corrected P value threshold = 4.22E-06). Competitive gene-set tests were conducted correcting for gene size, variant density and LD within and between genes. Tissue-set enrichment analyses were also performed using MAGMA implemented in FUMA, to test for enrichment of association signal in genes expressed in 54 tissue types from GTEx V8 (Bonferroni-corrected P value threshold = 9.26E-04)^{64,107}.

For single-cell enrichment analyses, publicly available single-cell RNA-seq data were compiled from five studies of the adult human and mouse brain^{86,108–111}. Using a previously described method³⁸, a metric of gene expression specificity was calculated by dividing the expression of each gene in each cell type by the total expression of that gene in all cell types, leading to values ranging from 0 to 1 for each gene (0 meaning that the gene is not expressed in that cell type and 1 meaning that all of the expression of the gene is in that cell type). MAGMAv1.08³⁷ was used to test whether the 10% most specific genes for each cell type were associated with BD, including relevant covariates. The P value threshold for significance was $P < 9.1\text{E-}03$, representing a 5% false discovery rate (FDR) across datasets. Full details are in the Supplementary Note.

Further gene-set analyses were performed restricted to genes targeted by drugs, assessing individual drugs and grouping drugs with similar actions. This approach has been described previously⁴¹. Two groups of gene sets were defined. The first group comprised the targets of each drug in the Drug-Gene Interaction Database DGIdb v.2³⁹ and in the Psychoactive Drug Screening Database Ki DB⁴⁰, both downloaded in June 2016⁴¹. The second set were drug sets grouped according to Anatomical Therapeutic Chemical (ATC) classes⁴¹. All analyses were performed using competitive gene-set analyses in MAGMA v1.08³⁷. Multiple testing was controlled using a Bonferroni-corrected significance threshold of $P < 2.86\text{E-}05$ for drug-set analysis and $P < 3.40\text{E-}04$ for drug-class analysis, accounting for 1,746 drug-sets and 147 drug classes tested.

eQTL integrative analysis

A transcriptome-wide association study (TWAS) was conducted using the precomputed gene expression weights from PsychENCODE data (1,321 brain samples)⁴³, available online with the FUSION software⁴². For genes with significant *cis*-SNP heritability (13,435 genes), FUSION software was used to test whether SNPs influencing gene expression are also associated with BD (Bonferroni-corrected P value threshold < 3.72E-06). For regions including a TWAS significant gene, TWAS fine-mapping of the region was conducted using

FOCUS (fine-mapping of causal gene sets)⁴⁴. A posterior inclusion probability (PIP) was assigned to each gene for being causal for the observed TWAS association signal and used to generate the 90%-credible gene set for each region⁴⁴.

Summary data-based Mendelian randomization (SMR)^{45,46} was applied to further investigate putative causal relationships between SNPs and BD via gene expression. SMR was performed using eQTL summary statistics from the eQTLGen (31,684 blood samples)⁴⁷ and PsychENCODE⁴³ consortia. SMR analysis is limited to transcripts with at least one significant *cis*-eQTL ($P < 5E-08$) in each dataset (15,610 in eQTLGen; 10,871 in PsychENCODE). The Bonferroni-corrected significance threshold was $P < 3.20E-06$ and $P < 4.60E-06$ for eQTLGen and PsychENCODE respectively. The significance threshold for the HEIDI test (heterogeneity in dependent instruments) was $P_{\text{HEIDI}} \geq 0.01$ ⁴⁶. While the results of TWAS and SMR indicate an association between BD and gene expression, a non-significant HEIDI test additionally indicates either a direct causal role or a pleiotropic effect of the BD-associated SNPs on gene expression.

Complement component 4 (C4) imputation

To investigate the major histocompatibility complex (MHC; chr6:24-34 Mb on hg19), the alleles of complement component 4 genes (*C4A* and *C4B*) were imputed in 47 PGC cohorts for which individual-level genotype data were accessible, totaling 32,749 BD cases and 53,370 controls. The imputation reference panel comprised 2,530 reference haplotypes of MHC SNPs and *C4* alleles, generated using a sample of 1,265 individuals with whole-genome sequence data, from the Genomic Psychiatry cohort¹¹². Briefly, imputation of *C4* as a multi-allelic variant was performed using Beagle v4.1^{113,114}, using SNPs from the MHC region that were also in the haplotype reference panel. The output consisted of dosage estimates for each of the common *C4* structural haplotypes (e.g., AL-BS, AL-AL, etc.) for each individual. The five most common structural forms of the *C4A/C4B* locus (BS, AL, AL-BS, AL-BL, and AL-AL) could be inferred with reasonably high accuracy (generally $0.70 < r^2 < 1.00$). The imputed *C4* alleles were tested for association with BD in a joint logistic regression that included (i) terms for dosages of the five most common *C4* structural haplotypes (AL-BS, AL-BL, AL-AL, BS, and AL), (ii) rs13195402 genotype (top lead SNP in the MHC) and (iii) PCs as per the GWAS. The genetically regulated expression of *C4A* was predicted from the imputed *C4* alleles using a model previously described⁶³. Predicted *C4A* expression was tested for association with BD in a joint logistic regression that included (i) predicted *C4A* expression, (ii) rs13195402 genotype (top lead SNP in the MHC) and (iii) PCs as per the GWAS. Full details are in the Supplementary Note.

Polygenic risk scoring

PRS from our GWAS meta-analysis were tested for association with BD in individual cohorts, using a discovery GWAS where the target cohort was left out of the meta-analysis. Briefly, the GWAS results from each discovery GWAS were pruned for LD using the P value informed clumping method in PLINK v1.90⁹⁸ (r^2 0.1 within a 500 kb window) based on the LD structure of the HRC reference panel⁹⁷. Subsets of SNPs were selected from the results below nine increasingly liberal P value thresholds (p_T) (5E-08, 1E-04, 1E-03, 0.01, 0.05, 0.1, 0.2, 0.5, 1). Sets of alleles, weighted by their log odds ratios from the discovery GWAS, were summed into PRS for each individual in the target datasets, using PLINK v1.90 implemented via RICOPILI^{94,98}. PRS were tested for association with BD in the target dataset using logistic regression, covarying for PCs as per the GWAS in each cohort. PRS were tested in the external cohorts by the collaborating research teams using comparable procedures. The variance explained by the PRS (R^2) was converted to the liability scale to account for the proportion of cases in each target dataset, using a BD population prevalence of 2% and 1%¹¹⁵. The weighted average R^2 values were calculated using the effective N for each cohort. To assess cross-ancestry performance, PRS generated from the meta-analysis results were tested for association with BD using similar methods in a Japanese sample⁵⁹, a Korean

sample⁹² and two admixed African American samples, in which BD cases were also ascertained using international consensus criteria (DSM-IV)^{22,93}. Full details of the ascertainment, QC, imputation and analysis of these samples are in the Supplementary Note.

LD score regression

LD Score regression (LDSC)³⁵ was used to estimate the h_{SNP}^2 of BD from GWAS summary statistics. h_{SNP}^2 was converted to the liability scale, using a lifetime BD prevalence of 2% and 1%. LDSC bivariate genetic correlations attributable to genome-wide SNPs (r_g) were estimated with 255 human diseases and traits from published GWAS and 514 GWAS of phenotypes in the UK Biobank from LD Hub (<http://ldsc.broadinstitute.org>)⁴⁸. Adjusting for the number of traits tested, the Bonferroni-corrected P value thresholds were $P < 1.96E-04$ and $P < 9.73E-05$ respectively.

MiXeR

We applied bivariate gaussian mixture models to the GWAS summary statistics for BD and other complex traits, using MiXeR v1.3^{49,50}. MiXeR provides univariate estimates of the proportion of non-null SNPs (“polygenicity”) and the variance of effect sizes of non-null SNPs (“discoverability”) in each phenotype. In the cross-trait analysis, MiXeR models additive genetic effects as a mixture of four components, representing SNPs not influencing either trait, SNPs influencing only one of the two traits, and SNPs influencing both traits. These components are then plotted in Venn diagrams. After fitting parameters of the model, the dice coefficient, which represents the proportion of overlapping variants, is calculated. Further details are provided in the Supplementary Note.

Mendelian randomization

Seventeen traits associated with BD in clinical or epidemiological studies were selected for Mendelian randomization (MR) to dissect their relationship with BD (Supplementary Note). Bi-directional generalized summary statistics-based MR (GSMR)⁵¹ analyses were performed between BD and the traits of interest using GWAS summary statistics. The instrumental variables (IVs) were selected by a clumping procedure internal to the GSMR software with parameters: *--gwas-thresh 5e-8 --clump-r2 0.01*. Traits with less than 10 IVs available were excluded from the GSMR analyses to avoid conducting underpowered tests⁵¹, resulting in 10 traits tested (Bonferroni-corrected P value threshold $< 2.5E-03$). The HEIDI-outlier test (heterogeneity in dependent instruments) was applied to test for horizontal pleiotropy ($P_{HEIDI} < 0.01$)⁵¹. For comparison, the MR analyses were also performed using the inverse variance weighted regression method, implemented via the TwoSampleMR R package, using the IVs selected by GSMR^{116,117}. To further investigate horizontal pleiotropy, the MR Egger intercept test was conducted using the TwoSampleMR package^{116,117} and MR-PRESSO software was used to perform the Global Test and Distortion Test¹¹⁸.

BD subtypes

GWAS meta-analyses were conducted for BD I (25,060 cases, 449,978 controls from 55 cohorts, effective N = 64,802) and BD II (6,781 cases, 364,075 controls from 31 cohorts, effective N = 22,560) using the same procedures described for the main GWAS. BD subtypes were defined based on international consensus criteria (DSM-IV, ICD-9 or ICD-10), established using structured diagnostic instruments from assessments by trained interviewers, clinician-administered checklists or medical record review, or using ICD codes (Supplementary Note). LDSC³⁵ was used to estimate the h_{SNP}^2 of each subtype, and the genetic correlation between the subtypes. The difference between LDSC h_{SNP}^2 estimates for BD I and BD II was tested for deviation from 0 using the block jackknife¹¹⁹. The LDSC genetic correlation (r_g) was tested for difference from 1 by calculating a chi-square statistic corresponding to the estimated r_g , as $[(r_g - 1)/se]^2$.

Data availability

The PGC's policy is to make genome-wide summary results public. Summary statistics will be made available through the PGC website upon publication (<https://www.med.unc.edu/pgc/results-and-downloads>). Data are accessible with collaborative analysis proposals through the Bipolar Disorder Working Group of the PGC (<https://www.med.unc.edu/pgc/shared-methods/how-to/>). This study included some publicly available datasets accessed through dbGaP - PGC bundle phs001254.v1.p1.

Acknowledgements

We thank the participants who donated their time, life experiences and DNA to this research, and the clinical and scientific teams that worked with them. We are deeply indebted to the investigators who comprise the PGC. The PGC has received major funding from the US National Institute of Mental Health (PGC3: U01 MH109528; PGC2: U01 MH094421; PGC1: U01 MH085520). Statistical analyses were carried out on the NL Genetic Cluster Computer (<http://www.geneticcluster.org>) hosted by SURFsara and the Mount Sinai high performance computing cluster (<http://hpc.mssm.edu>), which is supported by the Office of Research Infrastructure of the National Institutes of Health under award numbers S10OD018522 and S10OD026880. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Full acknowledgements are included in the Supplementary Note.

Author contribution statement - to be added

Competing interests

T.E. Thorgeirsson, S. Steinberg, H. Stefansson and K. Stefansson are employed by deCODE Genetics/Amgen. Multiple additional authors work for pharmaceutical or biotechnology companies in a manner directly analogous to academic co-authors and collaborators. Full details of competing interests for all co-authors are included in the Supplementary Note.

Affiliations

¹Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ²Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³Institute of Human Genetics, University of Bonn, School of Medicine and University Hospital Bonn, Bonn, Germany. ⁴Institute of Neuroscience and Medicine (INM-1), Research Centre Jülich, Jülich, Germany. ⁵Centre for Human Genetics, University of Marburg, Marburg, Germany. ⁶Division of Mental Health and Addiction, Oslo University Hospital, Oslo, Norway. ⁷NORMENT, University of Oslo, Oslo, Norway. ⁸Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA. ⁹Social, Genetic and Developmental Psychiatry Centre, King's College London, London, UK. ¹⁰NIHR Maudsley BRC, King's College London, London, UK. ¹¹Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, Australia. ¹²iSEQ, Center for Integrative Sequencing, Aarhus University, Aarhus, Denmark. ¹³Department of Biomedicine - Human Genetics, Aarhus University, Aarhus, Denmark. ¹⁴iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, Denmark. ¹⁵Department of Psychiatry and Behavioral Sciences, SUNY Downstate Health Sciences University, Brooklyn, NY, USA. ¹⁶VA NY Harbor Healthcare System, Brooklyn, NY, USA. ¹⁷Research and Communication Unit for Musculoskeletal Health, Division of Clinical Neuroscience, Oslo University Hospital, Ullevål, Oslo, Norway. ¹⁸Institute of Clinical Medicine, University of Oslo, Oslo, Norway. ¹⁹K. G. Jebsen Center for Genetic Epidemiology, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway. ²⁰Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden. ²¹Department of Mental Health, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology (NTNU), Trondheim, Norway. ²²Department of Østmarka, Division of Mental Health Care, St. Olavs Hospital, Trondheim University Hospital, Trondheim, Norway. ²³Department of Psychiatry and Biobehavioral Science, Semel Institute, David Geffen School of Medicine, University of California, Los Angeles, Los Angeles, CA, USA. ²⁴Department of Psychiatry, Fujita Health University School of Medicine, Toyoake, Japan. ²⁵Stanley Center for

Psychiatric Research, Broad Institute, Cambridge, MA, USA. ²⁶Department of Genetics, Harvard Medical School, Boston, MA, USA. ²⁷Estonian Genome Center, Institute of Genomics, University of Tartu, Tartu, Estonia. ²⁸Department of Psychiatry and Psychotherapy, Charité - Universitätsmedizin, Berlin, Germany. ²⁹Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³⁰Ronald M. Loeb Center for Alzheimer's Disease, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³¹Estelle and Daniel Maggin Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, USA. ³²deCODE Genetics / Amgen, Reykjavik, Iceland. ³³Department of Research, Innovation and Education, Division of Clinical Neuroscience, Oslo University Hospital, Oslo, Norway. ³⁴Samsung Advanced Institute for Health Sciences and Technology (SAIHST), Sungkyunkwan University, Samsung Medical Center, Seoul, South Korea. ³⁵Russian Academy of Medical Sciences, Mental Health Research Center, Moscow, Russian Federation. ³⁶Institute of Psychiatric Phenomics and Genomics (IPPG), University Hospital, LMU Munich, Munich, Germany. ³⁷Department of Psychiatry and Psychotherapy, University Hospital, LMU Munich, Munich, Germany. ³⁸National Centre for Register-Based Research, Aarhus University, Aarhus, Denmark. ³⁹Centre for Integrated Register-based Research, Aarhus University, Aarhus, Denmark. ⁴⁰Division of Psychiatry, University College London, London, UK. ⁴¹Department of Neuroscience, Istituto Di Ricerche Farmacologiche Mario Negri IRCCS, Milano, Italy. ⁴²Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, USA. ⁴³Northwestern University, Chicago, IL, USA. ⁴⁴Psychiatry, Berkshire Healthcare NHS Foundation Trust, Bracknell, UK. ⁴⁵Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA. ⁴⁶National and Kapodistrian University of Athens, 2nd Department of Psychiatry, Attikon General Hospital, Athens, Greece. ⁴⁷Department of Psychiatry, Sungkyunkwan University School of Medicine, Samsung Medical Center, Seoul, South Korea. ⁴⁸Center for Neonatal Screening, Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark. ⁴⁹Department of Psychiatry and Psychotherapy, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany. ⁵⁰Medical University of Graz, Department of Psychiatry and Psychotherapeutic Medicine, Graz, Austria. ⁵¹Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway. ⁵²Psychiatry, Brain Center UMC Utrecht, Utrecht, The Netherlands. ⁵³Instituto de Salud Carlos III, Biomedical Network Research Centre on Mental Health (CIBERSAM), Madrid, Spain. ⁵⁴Department of Psychiatry, Hospital Universitari Vall d'Hebron, Barcelona, Spain. ⁵⁵Department of Psychiatry and Forensic Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain. ⁵⁶Psychiatric Genetics Unit, Group of Psychiatry Mental Health and Addictions, Vall d'Hebron Research Institut (VHIR), Universitat Autònoma de Barcelona, Barcelona, Spain. ⁵⁷Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital Frankfurt, Frankfurt am Main, Germany. ⁵⁸Psychiatry, University of California San Francisco, San Francisco, CA, USA. ⁵⁹University of Newcastle, Newcastle, NSW, Australia. ⁶⁰Department of Psychiatry, Mood Disorders Program, McGill University Health Center, Montreal, QC, Canada. ⁶¹Division of Psychiatry, University of Edinburgh, Edinburgh, UK. ⁶²Department of Translational Research in Psychiatry, Max Planck Institute of Psychiatry, Munich, Germany. ⁶³Department of Psychiatry, Universidad Autonoma de Nuevo Leon, Monterrey, Mexico. ⁶⁴Department of Psychiatry and Psychology, Mayo Clinic, Rochester, MN, USA. ⁶⁵Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN, USA. ⁶⁶Centre for Psychiatry, Queen Mary University of London, London, UK. ⁶⁷UCL Genetics Institute, University College London, London, UK. ⁶⁸Department of Psychiatry, Laboratory of Psychiatric Genetics, Poznan University of Medical Sciences, Poznan, Poland. ⁶⁹Center for Multimodal Imaging and Genetics, Departments of Neurosciences, Radiology, and Psychiatry, University of California, San Diego, CA, USA. ⁷⁰Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital Essen, University of Duisburg-Essen, Duisburg, Germany. ⁷¹Department of Medical Genetics, Oslo University Hospital Ullevål, Oslo, Norway. ⁷²NORMENT, Department of Clinical Science, University of Bergen, Bergen, Norway. ⁷³Department of Neurology, Oslo University Hospital, Oslo, Norway. ⁷⁴NORMENT, KG Jebsen Centre for Psychosis Research, Oslo University Hospital, Oslo, Norway. ⁷⁵Medical Research Council Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neurosciences, Cardiff University, Cardiff, UK. ⁷⁶Academic Psychiatry, Newcastle University, Newcastle upon Tyne, UK. ⁷⁷Department of Medical and Molecular Genetics, Indiana University, Indianapolis, IN, USA. ⁷⁸Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany. ⁷⁹Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior, Los Angeles, CA, USA. ⁸⁰Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. ⁸¹Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Göttingen, Germany. ⁸²Department of Psychiatry, Dalhousie University, Halifax, NS, Canada. ⁸³Department of Psychiatry, Yale School of Medicine, New Haven, CT, USA. ⁸⁴Veterans Affairs Connecticut

Healthcare System, West Haven, CT, USA. ⁸⁵Departments of Genetics and Neuroscience, Yale University School of Medicine, New Haven, CT, USA. ⁸⁶Department of Psychological Sciences, University of Missouri, Columbia, MO, USA. ⁸⁷Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia. ⁸⁸Psychological Medicine, University of Worcester, Worcester, UK. ⁸⁹Department of Psychiatry, University of California San Diego, La Jolla, CA, USA. ⁹⁰Bioinformatics Research Centre, Aarhus University, Aarhus, Denmark. ⁹¹Mental Health Department, University Regional Hospital, Biomedicine Institute (IBIMA), Málaga, Spain. ⁹²Department of Psychiatry, Seoul National University College of Medicine, Seoul, South Korea. ⁹³Landspítali University Hospital, Reykjavik, Iceland. ⁹⁴Department of Psychology, Eberhard Karls Universität Tübingen, Tübingen, Germany. ⁹⁵Department of Biomedicine, University of Basel, Basel, Switzerland. ⁹⁶Institute of Medical Genetics and Pathology, University Hospital Basel, Basel, Switzerland. ⁹⁷Neuropsychiatrie Translationnelle, Inserm U955, Créteil, France. ⁹⁸Faculté de Santé, Université Paris Est, Créteil, France. ⁹⁹International Max Planck Research School for Translational Psychiatry (IMPRS-TP), Munich, Germany. ¹⁰⁰Laboratory of Complex Trait Genomics, Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, The University of Tokyo, Tokyo, Japan. ¹⁰¹Laboratory for Statistical and Translational Genetics, RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. ¹⁰²Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, ON, Canada. ¹⁰³Neurogenetics Section, Centre for Addiction and Mental Health, Toronto, ON, Canada. ¹⁰⁴Department of Psychiatry, University of Toronto, Toronto, ON, Canada. ¹⁰⁵Institute of Medical Sciences, University of Toronto, Toronto, ON, Canada. ¹⁰⁶Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University Hospital Würzburg, Würzburg, Germany. ¹⁰⁷Cell Biology, SUNY Downstate Medical Center College of Medicine, Brooklyn, NY, USA. ¹⁰⁸Institute for Genomic Health, SUNY Downstate Medical Center College of Medicine, Brooklyn, NY, USA. ¹⁰⁹ISGlobal, Barcelona, Spain. ¹¹⁰University of Patras, School of Health Sciences, Department of Pharmacy, Laboratory of Pharmacogenomics and Individualized Therapy, Patras, Greece. ¹¹¹Mental Illness Research, Education and Clinical Center, Crescenz VAMC, Philadelphia, PA, USA. ¹¹²Center for Studies of Addiction, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA. ¹¹³RIKEN Center for Integrative Medical Sciences, Yokohama, Japan. ¹¹⁴Psychiatry, Altrecht, Utrecht, The Netherlands. ¹¹⁵Psychiatry, GGZ inGeest, Amsterdam, The Netherlands. ¹¹⁶Psychiatry, VU medisch centrum, Amsterdam, The Netherlands. ¹¹⁷Department of Psychiatry, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands. ¹¹⁸Department of Molecular Medicine and Surgery, Karolinska Institutet, Stockholm, Sweden. ¹¹⁹Center for Molecular Medicine, Karolinska University Hospital, Stockholm, Sweden. ¹²⁰Psychiatry, North East London NHS Foundation Trust, Ilford, UK. ¹²¹Clinic for Psychiatry and Psychotherapy, University Hospital Cologne, Cologne, Germany. ¹²²Department of Psychiatry, Korea University College of Medicine, Seoul, South Korea. ¹²³Psychiatric and Neurodevelopmental Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA. ¹²⁴HudsonAlpha Institute for Biotechnology, Huntsville, AL, USA. ¹²⁵Department of Human Genetics, McGill University, Montréal, QC, Canada. ¹²⁶Montreal Neurological Institute and Hospital, McGill University, Montréal, QC, Canada. ¹²⁷Division of Psychiatry, Centre for Clinical Brain Sciences, The University of Edinburgh, Edinburgh, UK. ¹²⁸Department of Psychiatry and Psychotherapy, University of Bonn, Bonn, Germany. ¹²⁹National and Kapodistrian University of Athens, Medical School, Clinical Biochemistry Laboratory, Attikon General Hospital, Athens, Greece. ¹³⁰Department of Clinical Neuroscience, Centre for Psychiatry Research, Karolinska Institutet, Stockholm, Sweden. ¹³¹Systems Genetics Working Group, Department of Genetics, Stellenbosch University, Stellenbosch, South Africa. ¹³²Genetic Cancer Susceptibility Group, International Agency for Research on Cancer, Lyon, France. ¹³³Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA. ¹³⁴Centre for Neuroimaging and Cognitive Genomics (NICOG), National University of Ireland Galway, Galway, Ireland. ¹³⁵Medical faculty, University Sarajevo School of Science and Technology, Sarajevo, Bosnia and Herzegovina. ¹³⁶Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD, USA. ¹³⁷Oxford Health NHS Foundation Trust, Warneford Hospital, Oxford, UK. ¹³⁸Department of Psychiatry, University of Oxford, Warneford Hospital, Oxford, UK. ¹³⁹Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, USA. ¹⁴⁰Outpatient Clinic for Bipolar Disorder, Altrecht, Utrecht, The Netherlands. ¹⁴¹Department of Psychiatry, Washington University in Saint Louis, Saint Louis, MO, USA. ¹⁴²Department of Biochemistry and Molecular Biology II, Faculty of Pharmacy, University of Granada, Spain. ¹⁴³Institute of Neurosciences, Biomedical Research Center (CIBM), University of Granada, Spain. ¹⁴⁴Medicine, Psychiatry, Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN, USA. ¹⁴⁵Department of Genetics, Microbiology and Statistics, Faculty of Biology, Universitat de Barcelona, Barcelona, Catalonia, Spain. ¹⁴⁶Faculty of Medicine, Department of Psychiatry, School of

Health Sciences, University of Iceland, Reykjavik, Iceland. ¹⁴⁷Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK. ¹⁴⁸Psychiatry and the Behavioral Sciences, University of Southern California, Los Angeles, CA, USA. ¹⁴⁹Mood Disorders, PsyQ, Rotterdam, The Netherlands. ¹⁵⁰SAMRC Unit on Risk and Resilience in Mental Disorders, Dept of Psychiatry and Neuroscience Institute, University of Cape Town, Cape Town, South Africa. ¹⁵¹Department of Environmental Epidemiology, Nofer Institute of Occupational Medicine, Lodz, Poland. ¹⁵²Neuroscience Research Australia, Sydney, NSW, Australia. ¹⁵³School of Medical Sciences, University of New South Wales, Sydney, NSW, Australia. ¹⁵⁴Centro de Biología Molecular Severo Ochoa, Universidad Autónoma de Madrid and CSIC, Madrid, Spain. ¹⁵⁵Department of Psychiatry and Human Behavior, School of Medicine, University of California, Irvine, CA, USA. ¹⁵⁶Psychiatry, Psychiatrisches Zentrum Nordbaden, Wiesloch, Germany. ¹⁵⁷Computational Sciences Center of Emphasis, Pfizer Global Research and Development, Cambridge, MA, USA. ¹⁵⁸Dalla Lana School of Public Health, University of Toronto, Toronto, ON, Canada. ¹⁵⁹Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK. ¹⁶⁰South London and Maudsley NHS Foundation Trust, Bethlem Royal Hospital, Monks Orchard Road, Beckenham, Kent, UK. ¹⁶¹A list of members and affiliations appears in the Supplementary Note. ¹⁶²Department of Clinical Sciences, Psychiatry, Umeå University Medical Faculty, Umeå, Sweden. ¹⁶³NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction, Institute of Clinical Medicine and Diakonhjemmet Hospital, University of Oslo, Oslo, Norway. ¹⁶⁴National Institute of Mental Health, Klecany, Czech Republic. ¹⁶⁵Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden. ¹⁶⁶Institute of Pulmonology, Russian State Medical University, Moscow, Russian Federation. ¹⁶⁷Department of Psychiatry, University of Münster, Münster, Germany. ¹⁶⁸Department of Psychiatry, Melbourne Medical School, The University of Melbourne, Melbourne, VIC, Australia. ¹⁶⁹The Florey Institute of Neuroscience and Mental Health, The University of Melbourne, Parkville, VIC, Australia. ¹⁷⁰Université de Paris, INSERM, Optimisation Thérapeutique en Neuropsychopharmacologie, UMRs-1144, Paris, France. ¹⁷¹APHP Nord, DMU Neurosciences, GHU Saint Louis-Lariboisière-Fernand Widal, Département de Psychiatrie et de Médecine Addictologique, Paris, France. ¹⁷²Psychiatry, University of Pennsylvania, Philadelphia, PA, USA. ¹⁷³Center for Statistical Genetics and Department of Biostatistics, University of Michigan, Ann Arbor, MI, USA. ¹⁷⁴Department of Biomedicine and the iSEQ Center, Aarhus University, Aarhus, Denmark. ¹⁷⁵Center for Genomics and Personalized Medicine, CGPM, Aarhus, Denmark. ¹⁷⁶School of Psychiatry, University of New South Wales, Sydney, NSW, Australia. ¹⁷⁷University of Queensland, Brisbane, QLD, Australia. ¹⁷⁸Neuropsychiatric Genetics Research Group, Dept of Psychiatry and Trinity Translational Medicine Institute, Trinity College Dublin, Dublin, Ireland. ¹⁷⁹National and Kapodistrian University of Athens, 1st Department of Psychiatry, Eginition Hospital, Athens, Greece. ¹⁸⁰Medical and Population Genetics, Broad Institute, Cambridge, MA, USA. ¹⁸¹Division of Endocrinology, Children's Hospital Boston, Boston, MA, USA. ¹⁸²Department of Human Genetics, University of Chicago, Chicago, IL, USA. ¹⁸³Biometric Psychiatric Genetics Research Unit, Alexandru Obregia Clinical Psychiatric Hospital, Bucharest, Romania. ¹⁸⁴HUNT Research Center, Department of Public Health and Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway. ¹⁸⁵University of Western Australia, Nedlands, WA, Australia. ¹⁸⁶Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, Sweden. ¹⁸⁷Department of Psychiatry and Addiction Medicine, Assistance Publique - Hôpitaux de Paris, Paris, France. ¹⁸⁸Department of Medical and Molecular Genetics, King's College London, London, UK. ¹⁸⁹Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Titusville, NJ, USA. ¹⁹⁰Cancer Epidemiology and Prevention, M. Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland. ¹⁹¹SA MRC Unit on Risk and Resilience in Mental Disorders, Dept of Psychiatry, Stellenbosch University, Stellenbosch, South Africa. ¹⁹²School of Psychology, The University of Queensland, Brisbane, QLD, Australia. ¹⁹³Department of Psychiatry and Genetics Institute, University of Florida, Gainesville, FL, USA. ¹⁹⁴Research Institute, Lindner Center of HOPE, Mason, OH, USA. ¹⁹⁵Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK. ¹⁹⁶Human Genetics Branch, Intramural Research Program, National Institute of Mental Health, Bethesda, MD, USA. ¹⁹⁷Division of Mental Health and Addiction, University of Oslo, Institute of Clinical Medicine, Oslo, Norway. ¹⁹⁸Psychiatry, St Olavs University Hospital, Trondheim, Norway. ¹⁹⁹Psychosis Research Unit, Aarhus University Hospital - Psychiatry, Risskov, Denmark. ²⁰⁰Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ²⁰¹University of Liverpool, Liverpool, UK. ²⁰²Research/Psychiatry, Veterans Affairs San Diego Healthcare System, San Diego, CA, USA. ²⁰³Mental Health Services in the Capital Region of Denmark, Mental Health Center Copenhagen, University of Copenhagen, Copenhagen, Denmark. ²⁰⁴Division of Psychiatry, Haukeland Universitetssjukehus, Bergen, Norway. ²⁰⁵Faculty of Medicine and Dentistry, University of Bergen, Bergen, Norway. ²⁰⁶Department of Clinical Neuroscience and Center for Molecular Medicine, Karolinska Institutet at Karolinska

University Hospital, Solna, Sweden. ²⁰⁷Human Genetics and Computational Biomedicine, Pfizer Global Research and Development, Groton, CT, USA. ²⁰⁸University of Melbourne, VIC, Australia. ²⁰⁹United Arab Emirates University, College of Medicine and Health Sciences, Department of Pathology, Al-Ain, United Arab Emirates. ²¹⁰United Arab Emirates University, Zayed Center of Health Sciences, Al-Ain, United Arab Emirates. ²¹¹Psychiatry, Harvard Medical School, Boston, MA, USA. ²¹²Division of Clinical Research, Massachusetts General Hospital, Boston, MA, USA. ²¹³Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands. ²¹⁴Department of Clinical Genetics, Amsterdam Neuroscience, Vrije Universiteit Medical Center, Amsterdam, The Netherlands. ²¹⁵Department of Neurology and Neurosurgery, McGill University, Faculty of Medicine, Montreal, QC, Canada. ²¹⁶Department of Psychiatry and Behavioral Sciences, SUNY Upstate Medical University, Syracuse, NY, USA. ²¹⁷Department of Biomedical and NeuroMotor Sciences, University of Bologna, Bologna, Italy. ²¹⁸Department of Neuroscience, SUNY Upstate Medical University, Syracuse, NY, USA. ²¹⁹Psychiatric and Neurodevelopmental Genetics Unit (PNGU), Massachusetts General Hospital, Boston, MA, USA. ²²⁰Faculty of Medicine, University of Iceland, Reykjavik, Iceland. ²²¹Department of Psychiatry, Hospital Namsos, Namsos, Norway. ²²²Department of Neuroscience, Norges Teknisk Naturvitenskapelige Universitet Fakultet for naturvitenskap og teknologi, Trondheim, Norway. ²²³Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA. ²²⁴Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA. ²²⁵Department of Psychiatry, McGill University, Montreal, QC, Canada. ²²⁶Dept of Psychiatry, Sankt Olavs Hospital Universitetssykehuset i Trondheim, Trondheim, Norway. ²²⁷Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS, CIBERSAM, Barcelona, Spain. ²²⁸Department of Psychology, Emory University, Atlanta, GA, USA. ²²⁹Institute of Biological Psychiatry, Mental Health Services, Copenhagen University Hospital, Copenhagen, Denmark. ²³⁰Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark. ²³¹Center for GeoGenetics, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark. ²³²Queensland Brain Institute, The University of Queensland, Brisbane, QLD, Australia. ²³³Psychiatry, Indiana University School of Medicine, Indianapolis, IN, USA. ²³⁴Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN, USA. ²³⁵Department of Human Genetics, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, USA. ²³⁶These authors contributed equally as first authors: Niamh Mullins, Andreas J. Forstner. ²³⁷These authors contributed equally as second authors: Kevin S. O'Connell, Brandon Coombes, Jonathan R. I. Coleman, Zhen Qiao. ²³⁸These authors jointly supervised this work: Eli A. Stahl, Andrew McQuillin, Arianna Di Florio, Roel A. Ophoff, Ole A. Andreassen. *Corresponding authors: Niamh Mullins, niamh.mullins@mssm.edu; Ole A. Andreassen, ole.andreassen@medisin.uio.no

References

1. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* **390**, 1211–1259 (2017).
2. Plans, L. *et al.* Association between completed suicide and bipolar disorder: A systematic review of the literature. *J. Affect. Disord.* **242**, 111–122 (2019).
3. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders 5th edn.* (American Psychiatric Association Publishing, 2013).

4. Merikangas, K. R. *et al.* Lifetime and 12-month prevalence of bipolar spectrum disorder in the National Comorbidity Survey replication. *Arch. Gen. Psychiatry* **64**, 543–552 (2007).
5. Merikangas, K. R. *et al.* Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Arch. Gen. Psychiatry* **68**, 241–251 (2011).
6. Craddock, N. & Sklar, P. Genetics of bipolar disorder. *Lancet* **381**, 1654–1662 (2013).
7. Song, J. *et al.* Bipolar disorder and its relation to major psychiatric disorders: a family-based study in the Swedish population. *Bipolar Disord.* **17**, 184–193 (2015).
8. Bienvenu, O. J., Davydow, D. S. & Kendler, K. S. Psychiatric ‘diseases’ versus behavioral disorders and degree of genetic influence. *Psychol. Med.* **41**, 33–40 (2011).
9. Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat. Genet.* **43**, 977–983 (2011).
10. Baum, A. E. *et al.* A genome-wide association study implicates diacylglycerol kinase eta (DGKH) and several other genes in the etiology of bipolar disorder. *Mol. Psychiatry* **13**, 197–207 (2008).
11. Charney, A. W. *et al.* Evidence for genetic heterogeneity between clinical subtypes of bipolar disorder. *Transl. Psychiatry* **7**, e993 (2017).
12. Chen, D. T. *et al.* Genome-wide association study meta-analysis of European and Asian-ancestry samples identifies three novel loci associated with bipolar disorder. *Mol. Psychiatry* **18**, 195–205 (2013).
13. Cichon, S. *et al.* Genome-wide association study identifies genetic variation in neurocan as a susceptibility factor for bipolar disorder. *Am. J. Hum. Genet.* **88**, 372–381 (2011).
14. Ferreira, M. A. R. *et al.* Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. *Nat. Genet.* **40**, 1056–1058 (2008).
15. Green, E. K. *et al.* Association at SYNE1 in both bipolar disorder and recurrent major depression.

- Mol. Psychiatry* **18**, 614–617 (2013).
16. Green, E. K. *et al.* Replication of bipolar disorder susceptibility alleles and identification of two novel genome-wide significant associations in a new bipolar disorder case-control sample. *Mol. Psychiatry* **18**, 1302–1307 (2013).
 17. Hou, L. *et al.* Genome-wide association study of 40,000 individuals identifies two novel loci associated with bipolar disorder. *Hum. Mol. Genet.* **25**, 3383–3394 (2016).
 18. Mühleisen, T. W. *et al.* Genome-wide association study reveals two new risk loci for bipolar disorder. *Nat. Commun.* **5**, 3339 (2014).
 19. Schulze, T. G. *et al.* Two variants in Ankyrin 3 (ANK3) are independent genetic risk factors for bipolar disorder. *Mol. Psychiatry* **14**, 487–491 (2009).
 20. Scott, L. J. *et al.* Genome-wide association and meta-analysis of bipolar disorder in individuals of European ancestry. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 7501–7506 (2009).
 21. Sklar, P. *et al.* Whole-genome association study of bipolar disorder. *Mol. Psychiatry* **13**, 558–569 (2008).
 22. Smith, E. N. *et al.* Genome-wide association study of bipolar disorder in European American and African American individuals. *Mol. Psychiatry* **14**, 755–763 (2009).
 23. Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* **447**, 661–678 (2007).
 24. Stahl, E. A. *et al.* Genome-wide association study identifies 30 loci associated with bipolar disorder. *Nat. Genet.* **51**, 793–803 (2019).
 25. Lee, S.-H., Zabolotny, J. M., Huang, H., Lee, H. & Kim, Y.-B. Insulin in the nervous system and the mind: Functions in metabolism, memory, and mood. *Mol Metab* **5**, 589–601 (2016).
 26. McIntyre, R. S. *et al.* A randomized, double-blind, controlled trial evaluating the effect of intranasal insulin on neurocognitive function in euthymic patients with bipolar disorder. *Bipolar Disord.* **14**,

- 697–706 (2012).
27. Nurnberger, J. I., Jr *et al.* Identification of pathways for bipolar disorder: a meta-analysis. *JAMA Psychiatry* **71**, 657–664 (2014).
 28. Gordovez, F. J. A. & McMahon, F. J. The genetics of bipolar disorder. *Mol. Psychiatry* **25**, 544–559 (2020).
 29. Zhang, C., Xiao, X., Li, T. & Li, M. Translational genomics and beyond in bipolar disorder. *Mol. Psychiatry* (2020) doi:10.1038/s41380-020-0782-9.
 30. Pedersen, C. B. *et al.* The iPSYCH2012 case-cohort sample: New directions for unravelling genetic and environmental architectures of severe mental disorders. *Preprint at bioRxiv* (2017) doi:10.1101/146670.
 31. Gudbjartsson, D. F. *et al.* Large-scale whole-genome sequencing of the Icelandic population. *Nat. Genet.* **47**, 435–444 (2015).
 32. Leitsalu, L. *et al.* Cohort Profile: Estonian Biobank of the Estonian Genome Center, University of Tartu. *Int. J. Epidemiol.* **44**, 1137–1147 (2015).
 33. Krokstad, S. *et al.* Cohort Profile: the HUNT Study, Norway. *Int. J. Epidemiol.* **42**, 968–977 (2013).
 34. Sudlow, C. *et al.* UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med.* **12**, e1001779 (2015).
 35. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature Genetics* vol. 47 291–295 (2015).
 36. Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A. P. & Price, A. L. Mixed-model association for biobank-scale datasets. *Nat. Genet.* **50**, 906–908 (2018).
 37. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput. Biol.* **11**, e1004219 (2015).
 38. Bryois, J. *et al.* Genetic identification of cell types underlying brain complex traits yields insights into

- the etiology of Parkinson's disease. *Nat. Genet.* **52**, 482–493 (2020).
39. Wagner, A. H. *et al.* DGIdb 2.0: mining clinically relevant drug-gene interactions. *Nucleic Acids Res.* **44**, D1036–44 (2016).
 40. Roth, B. L., Lopez, E., Patel, S. & Kroeze, W. K. The Multiplicity of Serotonin Receptors: Uselessly Diverse Molecules or an Embarrassment of Riches? *Neuroscientist* **6**, 252–262 (2000).
 41. Gaspar, H. A. & Breen, G. Drug enrichment and discovery from schizophrenia genome-wide association results: an analysis and visualisation approach. *Sci. Rep.* **7**, 12460 (2017).
 42. Gusev, A. *et al.* Integrative approaches for large-scale transcriptome-wide association studies. *Nat. Genet.* **48**, 245–252 (2016).
 43. Gandal, M. J. *et al.* Transcriptome-wide isoform-level dysregulation in ASD, schizophrenia, and bipolar disorder. *Science* **362**, (2018).
 44. Mancuso, N. *et al.* Probabilistic fine-mapping of transcriptome-wide association studies. *Nat. Genet.* **51**, 675–682 (2019).
 45. Zhu, Z. *et al.* Integration of summary data from GWAS and eQTL studies predicts complex trait gene targets. *Nat. Genet.* **48**, 481–487 (2016).
 46. Wu, Y. *et al.* Integrative analysis of omics summary data reveals putative mechanisms underlying complex traits. *Nat. Commun.* **9**, 918 (2018).
 47. Vösa, U. *et al.* Unraveling the polygenic architecture of complex traits using blood eQTL metaanalysis. *Preprint at bioRxiv* (2018) doi:10.1101/447367.
 48. Zheng, J. *et al.* LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis. *Bioinformatics* **33**, 272–279 (2017).
 49. Frei, O. *et al.* Bivariate causal mixture model quantifies polygenic overlap between complex traits beyond genetic correlation. *Nat. Commun.* **10**, 2417 (2019).

50. Holland, D. *et al.* Beyond SNP Heritability: Polygenicity and Discoverability of Phenotypes Estimated with a Univariate Gaussian Mixture Model. *PLoS Genet* **16**, e1008612 (2020).
51. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat. Commun.* **9**, 224 (2018).
52. Steardo, L., Jr *et al.* Sleep Disturbance in Bipolar Disorder: Neuroglia and Circadian Rhythms. *Front. Psychiatry* **10**, 501 (2019).
53. Hunt, G. E., Malhi, G. S., Cleary, M., Lai, H. M. X. & Sitharthan, T. Prevalence of comorbid bipolar and substance use disorders in clinical settings, 1990-2015: Systematic review and meta-analysis. *J. Affect. Disord.* **206**, 331–349 (2016).
54. Heffner, J. L., Strawn, J. R., DelBello, M. P., Strakowski, S. M. & Anthenelli, R. M. The co-occurrence of cigarette smoking and bipolar disorder: phenomenology and treatment considerations. *Bipolar Disord.* **13**, 439–453 (2011).
55. Vreeker, A. *et al.* High educational performance is a distinctive feature of bipolar disorder: a study on cognition in bipolar disorder, schizophrenia patients, relatives and controls. *Psychol. Med.* **46**, 807–818 (2016).
56. MacCabe, J. H. *et al.* Excellent school performance at age 16 and risk of adult bipolar disorder: national cohort study. *Br. J. Psychiatry* **196**, 109–115 (2010).
57. Broome, M. R., Saunders, K. E. A., Harrison, P. J. & Marwaha, S. Mood instability: significance, definition and measurement. *Br. J. Psychiatry* **207**, 283–285 (2015).
58. Ward, J. *et al.* The genomic basis of mood instability: identification of 46 loci in 363,705 UK Biobank participants, genetic correlation with psychiatric disorders, and association with gene expression and function. *Mol. Psychiatry* (2019) doi:10.1038/s41380-019-0439-8.
59. Ikeda, M. *et al.* A genome-wide association study identifies two novel susceptibility loci and trans population polygenicity associated with bipolar disorder. *Mol. Psychiatry* **23**, 639–647 (2018).

60. Pardiñas, A. F. *et al.* Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nat. Genet.* **50**, 381–389 (2018).
61. Howard, D. M. *et al.* Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat. Neurosci.* **22**, 343–352 (2019).
62. Smeland, O. B., Frei, O., Dale, A. M. & Andreassen, O. A. The polygenic architecture of schizophrenia - rethinking pathogenesis and nosology. *Nat. Rev. Neurol.* **16**, 366–379 (2020).
63. Sekar, A. *et al.* Schizophrenia risk from complex variation of complement component 4. *Nature* **530**, 177–183 (2016).
64. GTEx Consortium *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).
65. Brainstorm Consortium *et al.* Analysis of shared heritability in common disorders of the brain. *Science* **360**, (2018).
66. Cross-Disorder Group of the Psychiatric Genomics Consortium. Genomic Relationships, Novel Loci, and Pleiotropic Mechanisms across Eight Psychiatric Disorders. *Cell* **179**, 1469–1482.e11 (2019).
67. Lewis, K. J. S. *et al.* Comparison of Genetic Liability for Sleep Traits Among Individuals With Bipolar Disorder I or II and Control Participants. *JAMA Psychiatry* (2019) doi:10.1001/jamapsychiatry.2019.4079.
68. Zhou, H. *et al.* Genome-wide meta-analysis of problematic alcohol use in 435,563 individuals yields insights into biology and relationships with other traits. *Nat. Neurosci.* (2020) doi:10.1038/s41593-020-0643-5.
69. Okbay, A. *et al.* Genome-wide association study identifies 74 loci associated with educational attainment. *Nature* **533**, 539–542 (2016).
70. Vermeulen, J. M. *et al.* Smoking and the risk for bipolar disorder: evidence from a bidirectional Mendelian randomisation study. *Br. J. Psychiatry* 1–7 (2019).

71. Peyrot, W. J. *et al.* The association between lower educational attainment and depression owing to shared genetic effects? Results in ~25 000 subjects. *Molecular Psychiatry* vol. 20 735–743 (2015).
72. Swanson, C. L., Jr, Gur, R. C., Bilker, W., Petty, R. G. & Gur, R. E. Premorbid educational attainment in schizophrenia: association with symptoms, functioning, and neurobehavioral measures. *Biol. Psychiatry* **44**, 739–747 (1998).
73. Wishart, D. S. *et al.* DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res.* **46**, D1074–D1082 (2018).
74. Mizuno, A. & Okada, Y. Biological characterization of expression quantitative trait loci (eQTLs) showing tissue-specific opposite directional effects. *Eur. J. Hum. Genet.* **27**, 1745–1756 (2019).
75. Schrode, N. *et al.* Synergistic effects of common schizophrenia risk variants. *Nat. Genet.* **51**, 1475–1485 (2019).
76. Jiang, X. *et al.* Sodium valproate rescues expression of TRANK1 in iPSC-derived neural cells that carry a genetic variant associated with serious mental illness. *Mol. Psychiatry* **24**, 613–624 (2019).
77. Huckins, L. M. *et al.* Transcriptomic Imputation of Bipolar Disorder and Bipolar subtypes reveals 29 novel associated genes. *Preprint at bioRxiv* (2017) doi:10.1101/222786.
78. Finan, C. *et al.* The druggable genome and support for target identification and validation in drug development. *Sci. Transl. Med.* **9**, (2017).
79. von Wegerer, J., Hesslinger, B., Berger, M. & Walden, J. A calcium antagonistic effect of the new antiepileptic drug lamotrigine. *Eur. Neuropsychopharmacol.* **7**, 77–81 (1997).
80. Cipriani, A. *et al.* A systematic review of calcium channel antagonists in bipolar disorder and some considerations for their future development. *Mol. Psychiatry* **21**, 1324–1332 (2016).
81. Harrison, P. J., Tunbridge, E. M., Dolphin, A. C. & Hall, J. Voltage-gated calcium channel blockers for psychiatric disorders: genomic reappraisal. *Br. J. Psychiatry* **216**, 250–253 (2020).
82. Network and Pathway Analysis Subgroup of Psychiatric Genomics Consortium. Psychiatric genome-

- wide association study analyses implicate neuronal, immune and histone pathways. *Nat. Neurosci.* **18**, 199–209 (2015).
83. Forstner, A. J. *et al.* Identification of shared risk loci and pathways for bipolar disorder and schizophrenia. *PLoS One* **12**, e0171595 (2017).
84. Bipolar Disorder and Schizophrenia Working Group of the Psychiatric Genomics Consortium. Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell* **173**, 1705–1715.e16 (2018).
85. Lee, Y., Zhang, Y., Kim, S. & Han, K. Excitatory and inhibitory synaptic dysfunction in mania: an emerging hypothesis from animal model studies. *Exp. Mol. Med.* **50**, 12 (2018).
86. Skene, N. G. *et al.* Genetic identification of brain cell types underlying schizophrenia. *Nat. Genet.* **50**, 825–833 (2018).
87. Lewis, C. M. & Vassos, E. Polygenic risk scores: from research tools to clinical instruments. *Genome Med.* **12**, 44 (2020).
88. Torkamani, A., Wineinger, N. E. & Topol, E. J. The personal and clinical utility of polygenic risk scores. *Nat. Rev. Genet.* **19**, 581–590 (2018).
89. Duncan, L. *et al.* Analysis of polygenic risk score usage and performance in diverse human populations. *Nat. Commun.* **10**, 3328 (2019).
90. Martin, A. R. *et al.* Clinical use of current polygenic risk scores may exacerbate health disparities. *Nat. Genet.* **51**, 584–591 (2019).
91. Coleman, J. R. I. *et al.* The Genetics of the Mood Disorder Spectrum: Genome-wide Association Analyses of More Than 185,000 Cases and 439,000 Controls. *Biol. Psychiatry* (2019) doi:10.1016/j.biopsych.2019.10.015.
92. Moon, S. *et al.* The Korea Biobank Array: Design and Identification of Coding Variants Associated with Blood Biochemical Traits. *Sci. Rep.* **9**, 1382 (2019).

93. Bigdeli, T. B. *et al.* Contributions of common genetic variants to risk of schizophrenia among individuals of African and Latino ancestry. *Mol. Psychiatry* (2019) doi:10.1038/s41380-019-0517-y.
94. Lam, M. *et al.* RICOPILI: Rapid Imputation for COnsortias PIpeLIine. *Bioinformatics* (2019) doi:10.1093/bioinformatics/btz633.
95. Loh, P.-R. *et al.* Reference-based phasing using the Haplotype Reference Consortium panel. *Nat. Genet.* **48**, 1443–1448 (2016).
96. Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat. Genet.* **48**, 1284–1287 (2016).
97. McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nat. Genet.* **48**, 1279–1283 (2016).
98. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).
99. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).
100. Demontis, D. *et al.* Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder. *Nat. Genet.* **51**, 63–75 (2019).
101. Nievergelt, C. M. *et al.* International meta-analysis of PTSD genome-wide association studies identifies sex- and ancestry-specific genetic risk loci. *Nat. Commun.* **10**, 4558 (2019).
102. Purves, K. L. *et al.* A Major Role for Common Genetic Variation in Anxiety Disorders. *Mol Psychiatry* (2019) doi:10.1101/203844.
103. Yu, D. *et al.* Interrogating the Genetic Determinants of Tourette’s Syndrome and Other Tic Disorders Through Genome-Wide Association Studies. *Am. J. Psychiatry* **176**, 217–227 (2019).
104. Watson, H. J. *et al.* Genome-wide association study identifies eight risk loci and implicates metabolic/psychiatric origins for anorexia nervosa. *Nat. Genet.* **51**, 1207–1214 (2019).

105. Grove, J. *et al.* Identification of common genetic risk variants for autism spectrum disorder. *Nat. Genet.* **51**, 431–444 (2019).
106. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).
107. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).
108. Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* vol. 174 999–1014.e22 (2018).
109. Saunders, A. *et al.* Molecular Diversity and Specializations among the Cells of the Adult Mouse Brain. *Cell* **174**, 1015–1030.e16 (2018).
110. Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nature Methods* vol. 14 955–958 (2017).
111. Lake, B. B. *et al.* Integrative single-cell analysis of transcriptional and epigenetic states in the human adult brain. *Nat. Biotechnol.* **36**, 70–80 (2018).
112. Kamitaki, N. *et al.* Complement component 4 genes contribute sex-specific vulnerability in diverse illnesses. *Preprint at bioRxiv* (2019) doi:10.1101/761718.
113. Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am. J. Hum. Genet.* **81**, 1084–1097 (2007).
114. Browning, B. L. & Browning, S. R. Genotype Imputation with Millions of Reference Samples. *Am. J. Hum. Genet.* **98**, 116–126 (2016).
115. Lee, S. H., Goddard, M. E., Wray, N. R. & Visscher, P. M. A better coefficient of determination for genetic profile analysis. *Genet. Epidemiol.* **36**, 214–224 (2012).
116. Hemani, G., Tilling, K. & Davey Smith, G. Orienting the causal relationship between imprecisely

- measured traits using GWAS summary data. *PLoS Genet.* **13**, e1007081 (2017).
117. Hemani, G. *et al.* The MR-Base platform supports systematic causal inference across the human phenome. *Elife* **7**, (2018).
118. Verbanck, M., Chen, C.-Y., Neale, B. & Do, R. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat. Genet.* **50**, 693–698 (2018).
119. Hübel, C. *et al.* Genomics of body fat percentage may contribute to sex bias in anorexia nervosa. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* **180**, 428–438 (2019).