



University  
of Glasgow

McKernan, M. and McMillan, D.C. and Anderson, J.R. and Angerson, W.J. and Stuart, R.C. (2008) *The relationship between quality of life (EORTC QLQ-C30) and survival in patients with gastro-oesophageal cancer*. *British Journal of Cancer*, 98 (5). pp. 888-893. ISSN 0007-0920

<http://eprints.gla.ac.uk/5253/>

Deposited on: 26 March 2009

**The relationship between quality of life (EORTC QLQ-C30) and survival in patients with gastro-oesophageal cancer**

Margaret McKernan<sup>1</sup>, Donald C McMillan<sup>1</sup>, John R Anderson<sup>2</sup>, W.J. Angerson<sup>1</sup>, Robert C Stuart<sup>1</sup>

1. University Department of Surgery, Royal Infirmary, Glasgow, G31 2ER, UK
2. Department of Surgery, Southern General Hospital, Glasgow, G51, UK

Short title: Quality of life and survival in gastro-oesophageal cancer

Keywords: gastro-oesophageal cancer, stage, treatment, quality of life.

Correspondence to:

Dr Donald C McMillan

University Department of Surgery,

Royal Infirmary,

Glasgow G31 2ER, United Kingdom.

Tel No. 0141 211 5435

Fax No. 0141 552 3229

E-mail [d.c.mcmillan@clinmed.gla.ac.uk](mailto:d.c.mcmillan@clinmed.gla.ac.uk)

## **Abstract**

It remains unclear whether any aspect of quality of life has a role in predicting survival in an unselected cohort of patients with gastro-oesophageal cancer. Therefore the aim of the present study was to examine the relationship between quality of life (EORTC QLQ-C30), clinico-pathological characteristics and survival in patients with gastro-oesophageal cancer. Patients presenting with gastric or oesophageal cancer, staged using the UICC TNM classification and who were deemed operable/inoperable between November 1997 and December 2002 (n=152) participated in a quality of life study, using the EORTC QLQ-C30 core questionnaire. On univariate analysis, age (P<0.01), tumour length (P<0.0001), TNM stage (P<0.0001), weight loss (P<0.0001), dysphagia score (P<0.001), performance status (P<0.1) and treatment (P<0.0001) were significantly associated with cancer specific survival. EORTC QLQ-C30, physical functioning (P<0.0001), role functioning (P<0.001), cognitive functioning (P<0.01), social functioning (P<0.0001), global quality of life (P<0.0001), fatigue (P<0.0001), nausea/ vomiting (P<0.01), pain (P<0.001), dyspnoea (P<0.0001), appetite loss (P<0.0001) and constipation (P<0.05) were also significantly associated with cancer specific survival. On multivariate survival analysis, tumour stage (P<0.0001), treatment (P<0.001) and appetite loss (P<0.0001) were significant independent predictors of cancer specific survival. The present study highlights the importance of quality of life (EORTC QLQ-C30) measures, in particular appetite loss, as a prognostic factor in these patients.

## Introduction

Gastro-oesophageal cancer is the third commonest cause of cancer death in the UK. Each year, there are approximately 16,500 new cases and over 13,000 deaths attributable to the disease. Overall survival is poor with the majority of patients presenting with advanced, inoperable disease and less than 15% surviving 5 years (Cancerstats, 2004;). Although there have been improvements in survival following surgery (Ando et al., 2000; Hundahl et al., 2000; Hofstetter et al., 2002; von Rahden et al., 2004), for the majority of patients current treatment offers little in terms of improved survival. As a result quality of life in these patients is likely to be of considerable importance (Aaronson, Bullinger and Ahmedzai, 1988; Aaronson *et al.*, 1993).

The European Organisation for Research and Treatment of Cancer have developed and validated the EORTC-QLQ-C30 questionnaire designed to assess the quality of life of cancer patients (Aaronson *et al.*, 1993). Disease specific aspects of the questionnaire provide detailed information about the patients' perception of their health. Moreover, it has been reported that, in a few studies, the EORTC QLQ-C30 measurement of quality of life may have prognostic value in patients with gastro-oesophageal cancer (Conroy *et al.*, 2006).

Blazeby and co-workers (2001) reported that, in addition to age and TNM stage, physical function or emotional function had independent prognostic value in 92 patients with oesophageal cancer. However, treatment (whether or not the patient underwent surgery) was not included in the model (Blazeby *et al.*, 2001).

Fang and co-workers (2003) studied 110 patients with squamous oesophageal cancer and concluded that there was evidence to support the correlation of patient-reported QOL scores with survival; therefore, pre-treatment physical functioning might be a surrogate marker of an

unrecognised biological prognostic factor. Although performance status was significant on univariate analysis it was not significant on multivariate analysis, whereas physical functioning was significant (Fang *et al*, 2003).

In contrast, in a study of more than 1000 patients with inoperable gastro-oesophageal cancer, entering 3 randomised clinical trials, Chau and colleagues (2004) reported that no aspect of the QLQ-C30 had independent prognostic value when performance status was considered. However, physical function, role function and global quality of life were associated with survival on univariate analysis. There were no survival differences among patients with oesophageal or gastric cancer (Chau *et al*, 2004). However, this study was retrospective and included selected cohorts of patients.

Therefore, from the above it remains unclear whether any aspect of quality of life other than physical function has a role in predicting survival in an unselected cohort of patients with gastro-oesophageal cancer. The aim of the present study was to examine the relationship between quality of life (EORTC QLQ-C30), clinico-pathological characteristics and survival in patients with gastro-oesophageal cancer

## **PATIENTS AND METHODS**

### **Patients**

Patients presenting with adenocarcinoma or squamous carcinoma of the gastric or oesophageal tract at the Royal Infirmary and Southern General Hospital, Glasgow between November 1997 and December 2002 (n=152) participated in a quality of life study, using the EORTC QLQ-C30 core questionnaire.

The extent of tumour spread was recorded using the TNM 5th edition classification (Sobin and Wittekind, 1997). Tumours around the gastro-oesophageal junction were further classified according to tumour site, using the Siewert system; type 1 and 2 lesions of the gastro-oesophageal junction were designated as cancers of the oesophagus. Type 3 tumours of the cardia were designated as gastric cancers (Siewert and Stein, 1998). For gastric cancers, tumour node metastasis (TNM) stage I–III tumours were considered to be amenable to curative surgical resection. In the present study for oesophageal cancers, TNM stage I–III tumours, excluding T4, were deemed to be amenable to curative surgical resection.

Patients who had stage 1 and 2 disease but their performance status was poor were not deemed suitable for surgery and went forward for active palliative treatment or supportive care. There were 152 patients included in the study, 69 patients were suitable for surgery and 83 patients were inoperable.

The study was approved by the Research Ethics Committee of the Royal Infirmary and Southern General Hospital, Glasgow.

## Methods

Clinical and demographic variables were recorded at the patient's initial presentation and included age, sex, tumour type, site and length, TNM stage, ECOG performance status, weight loss, dysphagia and treatment.

Following diagnosis but prior to treatment the lead clinician approached patients as to whether they would participate in a study to examine their quality of life. If they gave informed consent they were given the EORTC QLQ-C30 questionnaire to complete.

Different aspects of quality of life were assessed using this cancer specific 30-item questionnaire, which has six functional scales (physical, role, emotional, cognitive, social, global health status) and several questions relating to a range of physical symptoms (Aaronson *et al*, 1993). Patients marked to what extent each statement applied to them. In the present study a number of patients were excluded because they were unlikely to understand the language, had brain metastases, delirium or confusion. Neither age nor performance status were considered when offering the patient questionnaire. Patients were not randomised. Few subjects were excluded (less than 10 patients) and therefore in those patients offered the questionnaire the bias was likely to be small.

## Statistics

Scoring algorithms have been produced by the EORTC Quality of Life Study Group. The sum of items in each category is added and the total divided by the number of questions in the category. A linear transformation is then undertaken to convert this to a percentage scale with a higher score representing a higher response level. Thus a high score for functional scale represents a high/healthy level of functioning. A high score for the global health status/quality of life represents a high quality of life. In contrast, a high score for the symptom scale represents a higher level of symptoms / problems (Aaronson *et al*, 1993).

Data are presented as the median and range. Survival was determined from time of biopsy proven diagnosis until time of death. Patients were followed up at their clinic or endoscopy appointments until time of death, if patients did not attend appointment, their general practitioner was contacted. Deaths up to April 2007 were included in the analysis. Univariate and multivariate survival analysis and calculation of hazard ratios (HR) were performed using a Cox regression model. Covariates that were significant on univariate analysis were included in the multivariate analysis. This was performed using a stepwise backward procedure to derive a final model of the variables that had a significant relationship with survival. To remove a variable from the model, the corresponding P-value had to be greater than 0.05. Comparison of the association between tumour site, TNM stage, treatment and the functional (physical, role, emotional, cognitive, social, global health status) and physical symptoms (fatigue, pain and appetite loss) scales of the EORTC-QLQ-C30 quality of life questionnaire was carried out using the  $X^2$ -test or Mann-Whitney U-test where appropriate. Analysis was performed using SPSS software (SPSS Inc, Chicago, IL, USA).

## Results

Patient characteristics and cancer specific survival analysis of patients with gastro-oesophageal cancer (n=152) are shown in Table 1. The minimum follow up period was 54 months or until date of death, the median follow up for survivors was 81 months, one patient was lost to follow up and one patient withdrew from the study. During this period 106 (70%) patients died from their disease and 14 (9%) died from co-morbid disease.

The relationship between clinico-pathological characteristics, quality of life and cancer specific survival in patients with gastro-oesophageal cancer is shown in Table 1 The majority of patients were over the age of 65 years (57%), male (68%) and had adenocarcinomas (84%). The majority of patients presented with weight loss (66%), had little or no dysphagia, and a near normal performance status (ECOG-ps, 71%). The majority of patients had EORTC QLQ-C30 function scores above 50 (physical functioning 100%, role functioning 65%, emotional functioning 74%, cognitive functioning 83%, social a functioning 79% and global quality of life 56%) and symptom scores below 50 (fatigue 69%, nausea/vomiting 85%, pain 86%, dyspnoea 79%, sleep disturbance 69%, appetite loss 64%, constipation 76%, diarrhoea 95% and financial difficulties 89%) and therefore had apparently normal quality of life (Table 1)

The relationship between clinico-pathological characteristics, quality of life and cancer specific survival in patients with gastro-oesophageal cancer is shown in Table 1. On univariate analysis, age (P<0.01), tumour length (P<0.0001), TNM stage (P<0.0001), weight loss (P<0.0001), dysphagia score (P<0.001), performance status (P <0.1) and treatment (P<0.0001) were significantly associated with cancer specific survival. EORTC QLQ-C30, physical functioning (P<0.0001), role functioning (P<0.001), cognitive functioning (P<0.1), social functioning (P<0.0001), global quality of life (P<0.0001), fatigue (P<0.0001), nausea/

vomiting ( $P<0.01$ ), pain ( $P<0.001$ ), dyspnoea ( $P<0.0001$ ), appetite loss ( $P<0.0001$ ) and constipation ( $P<0.05$ ) were also significantly associated with cancer specific survival.

On multivariate analysis, tumour stage ( $P<0.0001$ ), treatment ( $P<0.001$ ) and appetite loss ( $P<0.0001$ ) were significantly independent predictors of cancer specific survival. The relationship between appetite loss and cancer specific survival in patients with gastro-oesophageal cancer is shown in Figure 1.

In the present study C-reactive protein concentrations, at the time of quality of life assessment, were available in 94 patients (57 patients  $<10\text{mg/l}$ , 37 patients  $>10\text{mg/l}$ ) and were significantly associated with poorer cancer specific survival ( $P<0.0001$ ). Therefore we included C-reactive protein in addition to TNM stage, treatment and appetite loss in the multivariate survival model. TNM stage (HR 1.36, 95%CI 1.00-1.85,  $P=0.0532$ ), treatment (HR 3.72, 95%CI 1.73-7.99,  $P=0.0007$ ), appetite loss (HR 1.02, 95%CI 1.01-1.03,  $P<0.0001$ ) and C-reactive protein (HR 2.02, 95%CI 1.12-3.63,  $P=0.0187$ ) were independently associated with cancer specific survival.

The relationship between tumour site, clinico-pathological characteristics and quality of life in patients with gastro-oesophageal cancer is shown in Table 2. Compared with the gastric cancer patients, oesophageal cancer patients were older ( $P<0.01$ ), had more dysphagia ( $P<0.001$ ) and a poorer ECOG-ps ( $P<0.05$ ). In terms of quality of life, compared with the gastric cancer patients, oesophageal cancer patients had higher emotional functioning ( $P<0.01$ ), cognitive functioning ( $P<0.05$ ), less nausea and vomiting ( $P<0.05$ ).

The relationship between TNM stage and clinico-pathological and quality of life characteristics in patients with gastric-oesophageal cancer is shown in Table 3. With increasing TNM stage

patients had greater weight loss ( $P<0.01$ ) and were less likely to have had surgery ( $P<0.001$ ). In terms of quality of life, with increasing TNM stage there was poorer physical functioning ( $P<0.05$ ), emotional functioning ( $P<0.05$ ), social functioning ( $P<0.01$ ) and global quality of life ( $P<0.01$ ). In terms of symptoms, with increasing TNM stage there was more fatigue ( $P<0.01$ ), appetite loss ( $P<0.001$ ), dyspnoea ( $P<0.05$ ) and constipation ( $P<0.05$ ).

The relationship between appetite loss, clinico-pathological characteristics and quality of life in patients with gastric-oesophageal cancer is shown in Table 4. Increasing appetite loss was associated with greater tumour length ( $P<0.05$ ), TNM stage ( $P<0.001$ ) and the operability of the tumour ( $P<0.001$ ). Also, increasing appetite loss was associated with weight loss ( $P<0.001$ ) and dysphagia ( $P<0.001$ ). In terms of quality of life, increasing appetite loss was associated with poorer physical ( $P<0.001$ ), role ( $P<0.001$ ), emotional ( $P<0.01$ ), cognitive ( $P<0.01$ ), social ( $P<0.001$ ) and global quality of life ( $P<0.001$ ) functioning. In terms of symptoms, with increasing appetite loss there was more fatigue ( $P<0.01$ ), nausea and vomiting ( $P<0.001$ ), pain ( $P<0.001$ ), sleep disturbance ( $P<0.05$ ) and constipation ( $P<0.001$ ).

The relationship between systemic inflammatory response, as evidenced by elevated C-reactive protein, clinico-pathological and quality of life characteristics in patients with gastric-oesophageal cancer is shown in Table 5. An elevated C-reactive protein was associated with greater tumour length ( $P<0.01$ ), advanced TNM stage ( $P<0.01$ ) and the operability of the tumour ( $P<0.001$ ) and a poorer ECOG-ps ( $P<0.05$ ). In terms of quality of life, an elevated C-reactive protein was associated with poorer physical ( $P<0.01$ ), role ( $P<0.05$ ) and social ( $P<0.05$ ) functioning. In terms of symptoms, with an elevated C-reactive protein was associated with more fatigue ( $P<0.01$ ), pain ( $P<0.05$ ) and appetite loss ( $P<0.01$ ).

## Discussion

In the present study tumour site was not associated with major differences in EORTC QLQ-C30 quality of life function or symptom scores. However, there were major differences with increasing stage of disease. Nevertheless, the symptom score for appetite loss remained significant prognostic factor even after adjustment for TNM stage. Furthermore, when the relationship between appetite loss and the other quality of life function or symptom scores was examined it was clear that appetite loss was closely related to these measures. Taken together the results of the present study highlight the importance of appetite loss as a presenting symptom in patients with gastro-oesophageal cancer.

Few studies have examined the relationship between aspects of quality of life and survival in patients with gastro-oesophageal cancer. The results of the present study are consistent with the report of Fang and co-workers (2003) who reported that appetite loss was associated with poorer survival in 110 patients with oesophageal cancer. However, the association was much weaker than that of the present study and was not significant in multivariate analysis.

Furthermore, the follow-up period and the numbers of patients who died of their disease was not defined. Blazeby and colleagues (1995), in a smaller study of 59 patients with oesophageal cancer, also reported that appetite loss was associated with poorer survival (Blazeby *et al*, 1995)

The basis of the relationship between appetite loss and poorer cancer specific survival cannot be determined by the present cross sectional study. However, it was of interest that appetite loss was closely associated with nausea and vomiting, dysphagia and weight loss and therefore it may be that these symptoms result in appetite loss and the consequent loss of weight, which has long been recognised to impact on outcome (DeWys *et al*, 1980).

A number of workers have implicated the systemic inflammatory response in this process (Kotler, 2000; MacDonald, 2007). O’Gorman and co-workers (1998), in a cross sectional study, showed that in addition to appetite loss and weight loss, the systemic inflammatory response was an important factor in determining patients’ quality of life (EORTC QLQ-C30) in gastro-intestinal cancer patients (O’Gorman *et al*, 1998). Therefore, it is of interest that two recent studies have shown that the presence of a systemic inflammatory response, as evidenced by an elevated C-reactive protein, predicts survival in both operable (Crumley *et al*, 2006a) and inoperable (Crumley *et al*, 2006b) gastro-oesophageal cancer patients. In the present study C-reactive protein concentrations, at the time of quality of life assessment, were available in 94 (62%) patients. Consistent with previous work an elevated C-reactive protein concentration was associated with increased appetite loss and when included in the multivariate analysis, an elevated C-reactive protein concentration was independently associated with poorer cancer specific survival. However, even those patients without an elevated C-reactive protein concentration reported some appetite loss and the independent prognostic value of appetite loss remained, thus confirming the importance of appetite loss in the multifactorial nature of weight loss and poor outcome in these patients (MacDonald, 2007).

In summary, in patients with gastro-oesophageal cancer, routinely used prognostic factors are based predominantly on clinical and pathological findings. The present study highlights the importance of quality of life (EORTC QLQ-C30) measures, in particular appetite loss, as prognostic factors in these patients.

## References

Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, Filiberti A, Flechtner H, Fleishman SB, de Haes JC, et al. (1993) The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst.* 85:365-76.

Aaronson NK, Bullinger M, Ahmedzai S. (1988) A modular approach to quality-of-life assessment in cancer clinical trials. *Recent Results Cancer Res.* 111:231-49.

Ando N, Ozawa S, Kitagawa Y, Shinozawa Y, Kitajima M. (2000) Improvement in the results of surgical treatment of advanced squamous esophageal carcinoma during 15 consecutive years. *Annals of Surgery.* 232(2):225-32.

Blazeby JM, Brookes ST, Alderson D. (2001) The prognostic value of quality of life scores during treatment for oesophageal cancer. *Gut.* 49: 227-30.

Blazeby JM, Williams MH, Brookes ST, Alderson D, Farndon JR. (1995) Quality of life measurement in patients with oesophageal cancer. *Gut.* 37:505-8.

Cancer Research UK, Information Resource Centre 2004 and Welsh Cancer Intelligence

<http://info.cancerresearchuk.org/cancerstats/> Accessed June 2007.

Chau I, Norman AR, Cunningham D, Waters JS, Oates J, Ross PJ (2004). Multivariate prognostic factor analysis in locally advanced and metastatic esophago-gastric cancer--pooled analysis from three multicenter, randomized, controlled trials using individual patient data. *J Clin Oncol.* 22: 2395-403.

Conroy T, Marchal F, Blazeby JM. (2006) Quality of life in patients with oesophageal and gastric cancer: an overview. *Oncology.* 70:391-402.

Crumley AB, McMillan DC, McKernan M, Going JJ, Shearer CJ, Stuart RC. (2006a) An elevated C-reactive protein concentration, prior to surgery, predicts poor cancer-specific survival in patients undergoing resection for gastro-oesophageal cancer. *Br J Cancer.* 94:1568-71.

Crumley AB, McMillan DC, McKernan M, McDonald AC, Stuart RC. (2006b) Evaluation of an inflammation-based prognostic score in patients with inoperable gastro-oesophageal cancer. *Br J Cancer*. 94:637-41.

Dewys WD, Begg C, Lavin PT, Band PR, Bennett JM, Bertino JR, Cohen MH, Douglass HO Jr, Engstrom PF, Ezdinli EZ, Horton J, Johnson GJ, Moertel CG, Oken MM, Perlia C, Rosenbaum C, Silverstein MN, Skeel RT, Sponzo RW, Tormey DC. (1980) Prognostic effect of weight loss prior to chemotherapy in cancer patients. Eastern Cooperative Oncology Group. *Am J Med*. ;69:491-7.

Fang FM, Tsai WL, Chiu HC, Kuo WR, Hsiung CY. (2004) Quality of life as a survival predictor for esophageal squamous cell carcinoma treated with radiotherapy. *Int J Radiat Oncol Biol Phys*. 2004;58:1394-404.

Hofstetter W, Swisher SG, Correa AM, Hess K, Putnam JB Jr, Ajani JA, Dolormente M, Francisco R, Komaki RR, Lara A, Martin F, Rice DC, Sarabia AJ, Smythe WR, Vaporciyan AA, Walsh GL, Roth JA.(2002) Treatment outcomes of resected esophageal cancer. *Annals of Surgery*.236(3):376-84.

Hundahl SA, Phillips JL, Menck HR.(2000) The National Cancer Data Base Report on poor survival of U.S. gastric carcinoma patients treated with gastrectomy: Fifth Edition American Joint Committee on Cancer staging, proximal disease, and the "different disease" hypothesis. *Cancer*.88(4):921-32.

Kotler DP (2000). Cachexia. *Ann Intern Med*. 133:622-34.

MacDonald N (2007). Cancer cachexia and targeting chronic inflammation: a unified approach to cancer treatment and palliative/supportive care. *J Support Oncol*. 5:157-62.

O'Gorman P, McMillan DC, McArdle CS. (1988) Impact of weight loss, appetite, and the inflammatory response on quality of life in gastrointestinal cancer patients. *Nutr Cancer*. 32: 76-80.

Sobin LH, Wittekind CH (1997) International Union Against Cancer (UICC). TNM classification of malignant tumors. 5th edition. New York: John Wiley & Sons, Inc.

von Rahden BH, Stein HJ.(2004) Therapy of advanced esophageal malignancy. Current Opinion of Gastroenterology. 20(4):391-6.

Table 1. The relationship between clinico-pathological characteristics, quality of life and cancer specific survival in patients with gastro-oesophageal cancer (n=152)

	Patients (n=152)	Univariate analysis HR (95%CI)	P-value	Multivariate analysis HR (95%CI)	P-value
Age:(<65/65-74/≥75)	66/56/30	1.46 (1.13-1.88)	0.0033		
Sex:(male/female)	104/48	0.84 (0.55-1.30)	0.4377		
Tumour type:(adeno/squam)	127/25	1.40 (0.83-2.36)	0.2016		
Tumour site:(oesoph/gastric)	70/82	0.88 (0.60-1.29)	0.5163		
Tumour length:(<5/510/>10cm)	60/70/12	2.37 (1.71-3.27)	<0.0001		
TNM stage:(I/II/III/IV)	28/46/34/41	2.28 (1.84-2.83)	<0.0001	1.74 (1.36-2.23)	<0.0001
Weight loss:(yes/no)	101/51	3.08 (1.94-4.89)	<0.0001		
Dysphagia score:(1/2/3/4/5)	81/23/32/15/1	1.37 (1.57-1.62)	0.0003		
ECOG:(0-1/2/3-4)	108/38/6	1.60 (1.39-2.27)	0.0069		
Treatment:(operable/inoperable)	69/83	7.52 (4.68-12.07)	<0.0001	3.61 (2.00-6.53)	<0.0001
<b>EORTC QLQ-C30 (0-100)</b>					
Physical functioning	93 (66.7-100)	0.96 (0.94-0.98)	<0.0001		
Role functioning	66.7 (0-100)	0.99 (0.99-1.00)	0.0006		
Emotional functioning	66.7 (0-100)	1.00 (0.99-1.00)	0.1377		
Cognitive functioning	83.3 (0-100)	0.99 (0.98-0.99)	0.0045		
Social functioning	83.3 (0-100)	0.99 (0.98-1.00)	<0.0001		
Global quality of life	50 (0-100)	0.98 (0.97-0.99)	<0.0001		
Fatigue	33.3 (0-100)	1.02 (1.01-1.02)	<0.0001		
Nausea and vomiting	16.7 (0-100)	1.01 (1.00-1.01)	0.0050		
Pain	16.7 (0-100)	1.01 (1.01-1.02)	0.0002		
Dyspnoea	0 (0-100)	1.01 (1.00-1.02)	<0.0001		
Sleep disturbance	33.3 (0-100)	1.00 (0.99-1.01)	0.1558		
Appetite loss	33.3 (0-100)	1.02 (1.02-1.03)	<0.0001	1.02 (1.01-1.02)	<0.0001
Constipation	33.3 (0-100)	1.01 (1.00-1.01)	0.0261		
Diarrhoea	0 (0-100)	1.00 (0.99-1.01)	0.9586		
Financial difficulty	0 (0-100)	1.00 (0.99-1.01)	0.4626		

Table 2. The relationship between tumour site, clinico-pathological characteristics and quality of life in patients with gastro-oesophageal cancer (n=152)

	Gastric (n=82)	Oesophageal (n=70)	P-value
Age:(<65 yrs/ 65-74yrs/≥75 yrs)	41/29/12	25/27/18	0.0041
Sex:(male/female)	53/29	51/19	0.279
Type:(squam/adeno)	1/81	24/46	<0.001
Tumour length:(<5cm/5-10cm/>10cm)	33/33/7	27/37/5	0.724
Tumour stage:(I/II/III/IV)	22/13/18/28	6/33/16/13	0.528
Dysphagia score:(1/2/3/4/5)	64/9/8/1/0	17/14/24/1	<0.001
Weight loss:(yes/no)	53/29	48/22	0.610
ECOG:(0-1/2/3-4)	64/17/1	44/21/5	0.018
Treatment:(operable/inoperable)	37/45	32/38	0.942
EORTC QLQ-C30 (0-100)			
Physical functioning	93.3 (66.7-100)	93.3 (66.7-100)	0.733
Role functioning	66.7 (0-100)	66.7 (0-100)	0.923
Emotional functioning	66.7 (0-100)	83.3 (0-100)	0.007
Cognitive functioning	83.3 (0-100)	83.3 (0-100)	0.038
Social functioning	83.3 (0-100)	75 (0-100)	0.964
Global quality of life	50 (0-100)	50 (0-100)	0.284
Fatigue	33.3 (0-100)	22.2 (0-100)	0.077
Nausea and vomiting	16.7 (0-100)	0 (0-100)	0.036
Pain	16.7 (0-100)	16.7 (0-100)	0.716
Dyspnoea	33.3 (0-100)	0 (0-100)	0.123
Sleep disturbance	33.3 (0-100)	33.3 (0-100)	0.360
Appetite loss	33.3 (0-100)	33.3 (0-100)	0.624
Constipation	33.3 (0-100)	33.3 (0-100)	0.031
Diarrhoea	0 (0-100)	0 (0-100)	0.802
Financial difficulty	0 (0-100)	0 (0-66.7)	0.098

Table 3. The relationship between TNM stage and clinico-pathological characteristics and quality of life in patients with gastro-oesophageal cancer (n=152)

	TNM I (n=28)	TNM II (n=46)	TNM III (n=34)	TNM IV (n=41)	P-value
Age:(<65 yrs/ 65-74yrs/≥75 yrs)	15/12/1	23/21/2	18/9/7	18/13/10	0.482
Sex:(male/female)	17/11	33/13	20/14	31/10	0.387
Tumour type:(squam/adeno)	2/26	13/33	6/28	4/37	0.576
Tumour site:(oesophagus/gastric)	6/22	33/13	16/18	13/28	0.528
Tumour length:(<5cm/5-10cm/>10cm)	19/7/0	21/22/3	10/19/3	8/22/6	<0.001
Weight loss:(yes/no)	14/14	29/17	23/11	34/7	0.004
Dysphagia score:(1/2/3/4/5)	22/3/3/0/0	20/10/9/7/0	14/5/9/5/1	23/5/10/3/0	0.130
ECOG:(0-1/2/3-4)	22/6/0	33/11/2	27/6/1	24/15/2	0.099
Treatment:(operable/inoperable)	24/4	26/20	16/18	1/40	<0.001
<b>EORTC QLQ-C30 (0-100)</b>					
Physical functioning	93.3 (73.3-100)	100 (66.7-100)	100 (73.3-100)	86.7 (66.7-100)	0.023
Role functioning	66.7 (0-100)	66.7 (0-100)	66.7 (0-100)	50 (0-100)	0.058
Emotional functioning	66.7 (8.3-100)	75 (25-100)	83.3 (0-100)	58.3 (0-100)	0.042
Cognitive functioning	83.3 (50-100)	83.3 (33.3-100)	83.3 (16.7-100)	75 (0-100)	0.042
Social functioning	100 (33.3-100)	83.3 (0-100)	66.7 (0-100)	50 (0-100)	0.002
Global quality of life	66.7 (8.3-100)	66.7 (0-100)	50 (16.7-100)	41.7 (0-100)	0.001
Fatigue	27.8 (0-66.7)	22.2 (0-88.9)	33.3 (0-100)	55.6 (0-100)	0.002
Nausea and vomiting	16.7 (0-100)	0 (0-100)	16.7 (0-100)	16.7 (0-100)	0.553
Pain	16.7 (0-66.7)	16.7 (0-100)	33.3 (0-100)	16.7 (0-100)	0.098
Dyspnoea	16.7 (0-100)	0 (0-100)	0 (0-100)	33.3 (0-100)	0.014
Sleep disturbance	33.3 (0-100)	33.3 (0-100)	33.3 (0-100)	66.7 (0-100)	0.689
Appetite loss	0 (0-100)	33.3 (0-100)	33.3 (0-100)	66.7 (0-100)	<0.001
Constipation	33.3 (0-66.7)	16.7 (0-100)	0 (0-100)	33.3 (0-100)	0.013
Diarrhoea	0 (0-100)	0 (0-100)	0 (0-33.3)	0 (0-66.7)	0.601
Financial difficulty	0 (0-66.7)	0 (0-100)	0 (0-66.7)	0 (0-100)	0.306

Table 4. The relationship between appetite loss, clinico-pathological characteristics and quality of life in patients with gastro-oesophageal cancer (n=152)

	Not at all (n=55)	A little (n=43)	Quite a bit (n=26)	Very much (n=28)	P-value
Age:(<65 yrs/ 65-74yrs/≥75 yrs)	27/18/10	18/15/10	8/15/3	13/8/7	0.540
Sex:(male/female)	43/12	26/17	15/11	20/8	0.312
Tumour type:(squam/adeno)	7/48	5/38	7/19	6/22	0.138
Tumour site:(oesophagus/gastric)	25/30	17/26	15/11	13/15	0.603
Tumour length:(<5cm/5-10cm/ >10cm)	27/20/4	18/21/2	9/15/2	6/14/4	0.016
TNM stage:(I/II/III/IV)	16/17/13/8	8/17/9/9	3/8/7/8	1/4/5/16	<0.001
Weight loss:(yes/no)	26/29	27/16	22/4	26/2	<0.001
Dysphagia score:(1/2/3/4/5)	37/10/6/2/0	24/6/10/3/0	9/4/9/3/1	11/3/7/7/0	<0.001
ECOG:(0-1/2/3-4)	39/14/2	34/8/1	18/7/1	17/9/2	0.281
Treatment:(operable/inoperable)	35/20	20/23	11/15	3/25	<0.001
<b>EORTC QLQ-C30 (0-100)</b>					
Physical functioning	100 (73.3-100)	100 (73.3-100)	86.7 (66.7-100)	80 (66.7-100)	<0.001
Role functioning	100 (0-100)	66.7 (0-100)	58.3 (0-100)	33.3 (0-100)	<0.001
Emotional functioning	75 (73.3-100)	66.7 (0-100)	83.3 (8.3-100)	58.3 (0-96.7)	0.003
Cognitive functioning	83.3 (16.7-100)	83.3 (0-100)	83.3 (50-100)	66.7(0-100)	0.001
Social functioning	100 (0-100)	83.3 (0-100)	66.7 (0-100)	50 (0-100)	<0.001
Global quality of life	66.7 (16.7-100)	50 (0-100)	45.8(16.7-100)	29.1 (0-66.7)	<0.001
Fatigue	11.1 (0-88.9)	33.3 (0-83.2)	33.3 (0-100)	77.7 (22.2-100)	<0.001
Nausea and vomiting	0 (0-100)	16.7 (0-100)	16.7 (0-100)	41.7 (0-100)	<0.001
Pain	16.7 (0-100)	16.7 (0-83.3)	33.3 (0-100)	25 (0-100)	<0.001
Dyspnoea	0 (0-66.7)	0 (0-100)	0 (0-100)	50 (0-100)	<0.001
Sleep disturbance	0 (0-100)	33.3 (0-100)	33.3 (0-100)	33.3 (0-100)	0.044
Constipation	0 (0-100)	33.3 (0-100)	33.3 (0-100)	66.7 (0-100)	<0.001
Diarrhoea	0 (0-100)	0 (0-66.7)	0 (0-100)	0 (0-100)	0.512
Financial difficulty	0 (0-66.7)	0 (0-100)	0 (0-100)	0 (0-100)	0.296

Table 5: The relationship between systemic inflammatory response, as evidenced by elevated C-reactive protein, clinico-pathological and quality of life characteristics in patients with gastric-oesophageal cancer (n=94)

	CRP $\leq$ 10 (n=57)	CRP>10 (n=37)	P value
Age:(<65 yrs/65-74yrs/ $\geq$ 75yrs)	34/16/7	18/12/7	0.258
Sex:(male/female)	38/19	27/10	0.520
Tumour type:(adeno/squam)	48/9	30/7	0.695
Tumour site:(oesoph/gastric)	23/34	20/17	0.195
Tumour length:(<5cm/5-10cm/>10cm)	35/17/2	11/20/3	0.005
Tumour stage:(I/II/III/IV)	15/20/13/8	4/10/10/13	0.006
Weight loss:(yes/no)	31/16	27/10	0.072
Dysphagia score:(1/2/3/4/5)	29/14/11/3/0	18/6/10/2/1	0.390
ECOG:(0-1/2/3-4)	52/5/0	27/10/0	0.019
Treatment:(operable/inoperable)	40/17	8/29	<0.001
EORTC:(0-100)			
Physical functioning	100 (73-100)	86.7 (66.7-100)	0.001
Role functioning	66.7 (0-100)	66.7 (0-100)	0.040
Emotional functioning	66.7 (0-100)	70.8 (0-100)	0.343
Cognitive functioning	83.3 (16.7-100)	83.3 (33.3-100)	0.875
Social functioning	83.3 (0-100)	66.7 (0-100)	0.045
Global quality of life	66.7 (0-100)	50 (0-100)	0.068
Fatigue	33.3 (0-100)	44.4 (0-88.9)	0.003
Nausea and vomiting	16.7 (0-100)	16.7 (0-100)	0.152
Pain	16.7 (0-100)	16.7 (0-100)	0.040
Appetite loss	33.3 (0-100)	66.7 (0-100)	0.001
Dyspnoea	0 (0-100)	0 (0-100)	0.055
Sleep disturbance	33.3 (0-100)	33.3 (0-100)	0.518
Constipation	0 (0-100)	33.3 (0-100)	0.142
Diarrhoea	0 (0-66.7)	0 (0-100)	0.304
Financial difficulty	0 (0-100)	0 (0-100)	0.362




Figure 1. The relationship between appetite loss (None, A little, Quite a bit, Very much, from top to bottom) and cancer specific survival in patients with gastro-oesophageal cancer.