



University  
of Glasgow

Stipetic, L. H., Hamilton, G., Dalby, M. J., Davies, R. L., Meek, R.M. D., Ramage, G., Smith, D. G.E., and Burgess, K. E.V. (2015) Draft genome sequence of isolate staphylococcus aureus LHSKBclinical, isolated from an infected hip. *Genome Announcements*, 3(2), e00336-15.

Copyright © 2015 The Authors

This work is made available under the Creative Commons Attribution 3.0 License (CC BY 3.0)

Version: Published

<http://eprints.gla.ac.uk/104531/>

Deposited on: 25 May 2015

# Draft Genome Sequence of Isolate *Staphylococcus aureus* LHSKBClinical, Isolated from an Infected Hip

Laurence H. Stipetic,<sup>a,b</sup> Graham Hamilton,<sup>b</sup> Matthew J. Dalby,<sup>c</sup> Robert L. Davies,<sup>a</sup> R. M. Dominic Meek,<sup>e</sup> Gordon Ramage,<sup>a,d</sup> David G. E. Smith,<sup>a,f</sup> Karl E. V. Burgess<sup>a,b</sup>

Institute of Infection, Immunity and Inflammation, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom<sup>a</sup>; Glasgow Polyomics, University of Glasgow, Glasgow, United Kingdom<sup>b</sup>; Institute of Molecular Cell and Systems Biology, University of Glasgow, Glasgow, United Kingdom<sup>c</sup>; Glasgow Dental School, School of Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom<sup>d</sup>; Department of Orthopedics, Southern General Hospital, Glasgow, United Kingdom<sup>e</sup>; Moredun Research Institute, Edinburgh, United Kingdom<sup>f</sup>

**We report here the genome sequence of a clinical isolate of *Staphylococcus aureus* from an orthopedic infection. Phenotypically diverse *Staphylococcus aureus* strains are associated with orthopedic infections and subsequent implant failure, and some are highly resistant to antibiotics. This genome sequence will support further analyses of strains causing orthopedic infections.**

Received 9 March 2015 Accepted 19 March 2015 Published 30 April 2015

**Citation** Stipetic LH, Hamilton G, Dalby MJ, Davies RL, Meek RMD, Ramage G, Smith DGE, Burgess KEV. 2015. Draft genome sequence of isolate *Staphylococcus aureus* LHSKBClinical, isolated from an infected hip. *Genome Announc* 3(2):e00336-15. doi:10.1128/genomeA.00336-15.

**Copyright** © 2015 Stipetic et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/4.0/).

Address correspondence to Karl E. V. Burgess, [karl.burgess@glasgow.ac.uk](mailto:karl.burgess@glasgow.ac.uk).

Members of the genus *Staphylococcus* are primarily responsible for orthopedic infections after arthroplasty surgery (1–3) and subsequent implant failure (4). When isolated from orthopedic cases (2), *Staphylococcus aureus* is considered a primary medical concern due to the emergence of antimicrobial-resistant strains (5) and the ability of *S. aureus* to attach to surfaces and excrete an extracellular matrix to form biofilms (1, 6). Biofilm formation has been shown to exacerbate antibiotic resistance (7, 8), providing a protective environment for the indwelling bacteria. Biofilm cells display an altered gene expression profile (9) and metabolism in contrast to their planktonic counterparts (10–12). Differences occur in biofilm capability between *S. aureus* strains, with certain strains forming biofilms more readily than others (6). A number of genes have been found to be important in the formation of a biofilm by *S. aureus* (9).

*S. aureus* LHSKBClinical was isolated at the Southern General Hospital Microbiology laboratories (Glasgow, United Kingdom) from pus aspirated from the hip of a 75-year-old female. Speciation was confirmed on the basis of a positive Staph Xtra latex test (Pro-Lab Diagnostics) and growth on bioMérieux chromID agar (SAID). Genomic DNA was prepared from a brain-heart infusion (BHI) broth culture using the DNeasy blood and tissue kit (Qiagen) according to the manufacturer's instructions, utilizing the protocol for Gram-positive cell lysis (13). Sequencing was carried out using the Illumina MiSeq system, employing 300-bp paired-end sequencing. The reads were trimmed of Illumina adapter sequences and low-quality bases, and then *de novo* assembled and scaffolded using the CLC Genomics Workbench version 7.5.1 (Qiagen). Assembled scaffolds totaled 24, consisting of a total of 2,762,774 bp with an  $N_{50}$  of 587,246 bp and an A-T content of 67%, similar to other sequenced strains of *S. aureus* (14, 15). Assembled scaffolds were annotated using the Rapid Annotations using Subsystem Technology (RAST) resource (16, 17). The genome sequence was predicted to encode 2,566 proteins.

Interrogating <http://saureus.mlst.net> (18–20) showed that this

strain did not match any existing sequence type, and alignment of assembled scaffolds using NCBI BLAST searches (21) and Mauve genome analysis software (22) identified regions of heterogeneity compared against other previously sequenced *S. aureus* genomes.

Genes encoding *icaA* and *icaD*, polysaccharide intracellular adhesion proteins associated with *S. aureus* biofilm formation (9), were found. Furthermore, *sarA*, another gene implicated in biofilm formation by *S. aureus* (23, 24), was also found in the genome. Moreover, *fnbB* was found in the genome, perceived as a possible genetic marker for biofilm formation (25). Additionally, several genes encoding exotoxins or superantigens were found in the genome, including several associated with super-antigen-encoding *S. aureus* pathogenicity islands (SaPI), a virulence factor responsible for increased pathogenicity (26, 27).

The genome shows a novel sequence type for this isolate and provides a reference genome for *S. aureus* from orthopedic infections, both aspects of pertinence to further comparative studies of *S. aureus* pathogenicity.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [JZAL000000000](https://www.ncbi.nlm.nih.gov/nuccore/JZAL000000000). The version described in this paper is version [JZAL010000000](https://www.ncbi.nlm.nih.gov/nuccore/JZAL010000000).

## ACKNOWLEDGMENTS

This work was supported by the Doctoral Training Centre (DTC) in Technologies at the Interface between Engineering, the Physical Sciences and the Life Sciences, University of Glasgow, funded by the Engineering and Physical Sciences Research Council (EPSRC) (grant number EP/F500424/1) and the Biotechnology and Biological Sciences Research Council (BBSRC).

We thank Jim Guthrie of the Southern General Hospital Microbiology Laboratories for supplying the isolate and carrying out initial identification and Julie Galbraith, Next Generation Sequencing Department, Glasgow Polyomics, for processing and sequencing the sample.

## REFERENCES

- Esteban J, Molina-Manso D, Spiliopoulou I, Cordero-Ampuero J, Fernández-Roblas R, Foka A, Gómez-Barrena E. 2010. Biofilm development by clinical isolates of *Staphylococcus* spp. from retrieved orthopedic prostheses. *Acta Orthop* 81:674–679. <http://dx.doi.org/10.3109/17453674.2010.537810>.
- Dempsey KE, Riggio MP, Lennon A, Hannah VE, Ramage G, Allan D, Bagg J. 2007. Identification of bacteria on the surface of clinically infected and non-infected prosthetic hip joints removed during revision arthroplasties by 16S rRNA gene sequencing and by microbiological culture. *Arthritis Res Ther* 9:R46. <http://dx.doi.org/10.1186/ar2201>.
- Zimmerli W, Trampuz A, Ochsner PE. 2004. Prosthetic-joint infections. *N Engl J Med* 351:1645–1654. <http://dx.doi.org/10.1056/NEJMra040181>.
- National Health Service (NHS). 2014. Scottish Arthroplasty Project biennial report 2014. Information Services Division, Scotland.
- Enright MC, Robinson DA, Randle G, Feil EJ, Grundmann H, Spratt BG. 2002. The evolutionary history of methicillin-resistant *Staphylococcus aureus* (MRSA). *Proc Natl Acad Sci U S A* 99:7687–7692. <http://dx.doi.org/10.1073/pnas.122108599>.
- Smith K, Perez A, Ramage G, Lappin D, Gemmell CG, Lang S. 2008. Biofilm formation by Scottish clinical isolates of *Staphylococcus aureus*. *J Med Microbiol* 57:1018–1023. <http://dx.doi.org/10.1099/jmm.0.2008/000968-0>.
- Ramage G, Tunney MM, Patrick S, Gorman SP, Nixon JR. 2003. Formation of *Propionibacterium acnes* biofilms on orthopaedic biomaterials and their susceptibility to antimicrobials. *Biomaterials* 24:3221–3227. [http://dx.doi.org/10.1016/S0142-9612\(03\)00173-X](http://dx.doi.org/10.1016/S0142-9612(03)00173-X).
- Drenkard E, Ausubel FM. 2002. *Pseudomonas* biofilm formation and antibiotic resistance are linked to phenotypic variation. *Nature* 416:740–743. <http://dx.doi.org/10.1038/416740a>.
- Resch A, Rosenstein R, Nerz C, Götz F. 2005. Differential gene expression profiling of *Staphylococcus aureus* cultivated under biofilm and planktonic conditions. *Appl Environ Microbiol* 71:2663–2676. <http://dx.doi.org/10.1128/AEM.71.5.2663-2676.2005>.
- Ammons MC, Tripet BP, Carlson RP, Kirker KR, Gross MA, Stanisich JJ, Copié V. 2014. Quantitative NMR metabolite profiling of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus* discriminates between biofilm and planktonic phenotypes. *J Proteome Res* 13:2973–2985. <http://dx.doi.org/10.1021/pr500120c>.
- Gjersing EL, Herberg JL, Horn J, Schaldach CM, Maxwell RS. 2007. NMR metabolomics of planktonic and biofilm modes of growth in *Pseudomonas aeruginosa*. *Anal Chem* 79:8037–8045. <http://dx.doi.org/10.1021/ac070800t>.
- Zhang B, Powers R. 2012. Analysis of bacterial biofilms using NMR-based metabolomics. *Future Med Chem* 4:1273–1306. <http://dx.doi.org/10.4155/fmc.12.59>.
- Qiagen. 2006. DNeasy blood and tissue handbook: pretreatment of Gram-positive bacteria. Qiagen, Venlo, Netherlands.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23:673–679. <http://dx.doi.org/10.1093/bioinformatics/btm009>.
- Ramaraj T, Matyi SA, Sundararajan A, Lindquist IE, Devitt NP, Schilkey FD, Lamichhane-khadka R, Hoyt PR, Mudge J, Gustafson E. 2014. Draft genome sequences of vancomycin-susceptible *Staphylococcus aureus* related to heterogeneous vancomycin-intermediate *S. Aureus*. *Genome Announc* 2(5):e01033-14. <http://dx.doi.org/10.1128/genomeA.01033-14>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
- Enright MC, Day NP, Davies CE, Peacock SJ, Spratt BG. 2000. Multilocus sequence typing for characterization of methicillin-resistant and methicillin-susceptible clones of *Staphylococcus aureus*. *J Clin Microbiol* 38:1008–1015.
- Maiden MC, Bygraves JA, Feil E, Morelli G, Russell JE, Urwin R, Zhang Q, Zhou J, Zurth K, Caugant DA, Feavers IM, Achtman M, Spratt BG. 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc Natl Acad Sci U S A* 95:3140–3145. <http://dx.doi.org/10.1073/pnas.95.6.3140>.
- Spratt BG. 1999. Multilocus sequence typing: molecular typing of bacterial pathogens in an era of rapid DNA sequencing and the internet. *Curr Opin Microbiol* 2:312–316. [http://dx.doi.org/10.1016/S1369-5274\(99\)80054-X](http://dx.doi.org/10.1016/S1369-5274(99)80054-X).
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [http://dx.doi.org/10.1016/S0022-2836\(05\)80360-2](http://dx.doi.org/10.1016/S0022-2836(05)80360-2).
- Darling AC, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <http://dx.doi.org/10.1101/gr.2289704>.
- Beenken KE, Blevins JS, Smeltzer MS. 2003. Mutation of *sarA* in *Staphylococcus aureus* limits biofilm formation. *Infect Immun* 71:4206–4211. <http://dx.doi.org/10.1128/IAI.71.7.4206-4211.2003>.
- Beenken KE, Dunman PM, McAleese F, Macapagal D, Murphy E, Projan SJ, Blevins JS, Smeltzer MS. 2004. Global gene expression in *Staphylococcus aureus* biofilms. *J Bacteriol* 186:4665–4684. <http://dx.doi.org/10.1128/JB.186.14.4665-4684.2004>.
- Lim Y, Shin HJ, Kwon AS, Reu JH, Park G, Kim J. 2013. Predictive genetic risk markers for strong biofilm-forming *Staphylococcus aureus*: *fnbB* gene and SCC<sub>mec</sub> type III. *Diagn Microbiol Infect Dis* 76:539–541. <http://dx.doi.org/10.1016/j.diagmicrobio.2013.04.021>.
- Lindsay JA, Holden MT. 2004. *Staphylococcus aureus*: superbug, super genome? *Trends Microbiol* 12:378–385. <http://dx.doi.org/10.1016/j.tim.2004.06.004>.
- Novick RP. 2003. Mobile genetic elements and bacterial toxins: the superantigen-encoding pathogenicity islands of *Staphylococcus aureus*. *Plasmid* 49:93–105. [http://dx.doi.org/10.1016/S0147-619X\(02\)00157-9](http://dx.doi.org/10.1016/S0147-619X(02)00157-9).