

Antigen-based immunotherapy (AIT) for autoimmune and allergic disease

Megan KL MacLeod¹ and Stephen M Anderton²



Autoimmune and allergic diseases are major causes of morbidity. Antigen-based immunotherapy (AIT) is immunologically the most satisfying means of specifically targeting only those T cells driving disease, thereby inducing antigen-specific immune tolerance, with the lowest adverse risk profile. AIT is highly effective in rodent models of T cell-driven inflammation and is now in clinical trials. The range of approaches to applying AIT in the clinic prevents a consensus on the molecular basis for this form of tolerance. In particular, there has been a paucity of information on how pre-activated effector and memory T cells respond to AIT. New, advanced murine models of AIT are beginning to deliver such information at the cellular, biochemical, transcriptional and epigenetic levels.

Addresses

¹ University of Glasgow, Institute of Infection, Immunity and Inflammation, Sir Graeme Davies Building, 120 University Place, Glasgow G12 8TA, UK

² University of Edinburgh, MRC Centre for Inflammation Research, Queen's Medical Research Institute, 47 Little France Crescent, Edinburgh EH16 4TJ, UK

Corresponding author: MacLeod, Megan KL
(Megan.Macleod@glasgow.ac.uk)

Current Opinion in Pharmacology 2015, 23:11–16

This review comes from a themed issue on **Immunomodulation**

Edited by **Stephen M Anderton** and **Simon Fillatreau**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 22nd May 2015

<http://dx.doi.org/10.1016/j.coph.2015.05.003>

1471-4892/© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Autoimmune diseases and allergic conditions are caused by inappropriate immune responses that attack either self-tissues or innocuous environmental antigens, respectively. Most current treatments for these conditions target the symptoms of the diseases rather than offering cures. Many of these treatments, especially in the context of autoimmunity, lead to significant immunodeficiency that can expose the patient to severe, sometimes life-threatening, infections. Such disease-modifying drugs are usually given over a long period of time, often for the lifetime of the individual. By contrast, specifically removing or modifying only those immune cells that drive the allergic

or autoreactive immune response could lead to a permanent cessation of the disease — a cure.


The CD4⁺ T cells believed to initiate and drive many autoimmune and allergic conditions are activated through their cell surface T cell receptor (TCR) that specifically recognises the self-antigen or allergen. By targeting the TCR, AIT aims to silence the damaging T cells whilst leaving the rest of the immune system intact. Success in animal studies and historical allergy treatments in humans has led to clinical studies in both fields. Here we will consider recent examples of AIT in both animal and human studies and discuss the underlying immunological mechanisms.

Immune tolerance — a physiologic basis for immunotherapy

T cells gain their heterodimeric TCR through random gene re-arrangement during thymic development. This is essential to generate a broad repertoire of T cells that can respond to the variable array of pathogens the host will be exposed to over a lifetime. TCR gene re-arrangement also generates T cells with the potential to be activated by self-peptides. Many of these potentially destructive T cells are deleted during negative thymic selection [1]. This ‘central tolerance’ is not complete, however. In all individuals, some self-reactive T cells mature and migrate to the periphery where, if activated, they have the potential to cause autoimmunity. To prevent this, ‘peripheral tolerance’ mechanisms regulate T cell activation to both self and innocuous foreign antigens, such as the dietary and environmental antigens that can cause allergy.

A key mechanism by which peripheral tolerance is maintained is the presentation of antigen in non-inflammatory contexts. Full T cell activation requires two signals stemming from cell surface receptors delivered when the T cell interacts with an antigen presenting cell (APC). Signal 1 is provided by TCR binding to short antigenic peptides presented by MHC molecules. Signal 2 (costimulation) is delivered by a number of different receptor–ligand pairs, upregulated in inflammatory environments. Delivery of signal 1 alone leads to T cell tolerance either as a consequence of T cell death, modulation of the T cell's ability to respond to further activation (often called anergy), or the T cell gains regulatory function, capable of suppressing the activation of other immune cells (Figure 1) [2–4]. AIT aims to mimic these natural tolerogenic processes by delivery of the appropriate self- or allergen-derived peptide in the absence of inflammatory

Figure 1



T cell activation requires two signals. CD4⁺ T cells are tolerised if they encounter their antigen (signal 1) presented by an antigen presenting cell (APC) that has not been activated. Inflammation and/or infection cause APCs to upregulate costimulatory molecules that deliver signal 2 to T cells. T cells that receive both signals 1 and 2 become activated and differentiate into effector cells. In the context of infection, effector T cells co-ordinate the clearance of invading pathogens. If the T cells recognise allergen or self-antigens, the host can develop allergy or autoimmune disease respectively. If a T cell receives a tolerising TCR signal in advance of encountering its antigen presented by an activated APC, the T cell will not become fully activated – it is tolerant.

signals. This attractively simple concept belies a complex process which requires extensive knowledge of the target antigen; the location, phenotype and function of the 'disease-causing' T cells; and an ability to track these T cells to evaluate the success of any treatment.

Allergy

For over 100 years [5], immunotherapy has been practiced in allergic individuals. Initially involving crude allergen extracts, antigens are now carefully prepared to reduce adverse events. Allergen-derived peptides rather than whole antigens are often used as peptide immunotherapy (PIT), particularly as these can be selected not to cross link allergen-specific IgE, a process that can lead to enhanced rather than reduced allergic symptoms, including anaphylactic shock [6]. Increasing the dose of the allergen over a period of months to years is thought to modulate the allergen specific immune response including inducing a switch from an IgE dominated antibody response to IgG [7] and/or production of the regulatory cytokine interleukin (IL)-10 [8]. More recent advances in the ability to track allergen-specific T cells in the blood of patients undergoing immunotherapy suggest that AIT causes the deletion, rather than the modulation, of the disease causing T cells [9[•],10^{••}].

The route of delivery of the immunotherapy is likely to affect the dose of antigen that reaches the immune system. In Europe, the sublingual route is approved for allergy immunotherapy while the subcutaneous route is approved in the USA [5]. While the efficacies of these two routes have rarely been compared directly, a recent meta-analysis indicates that tablets designed to treat grass pollen allergy and subcutaneous delivery of pollen antigens can each reduce symptoms of rhinoconjunctivitis to a similar extent [11[•]]. Delivery of antigen can also be increased by either direct injection into lymph nodes with the aid of ultrasound or by binding the antigen to particular adjuvants such as aluminium hydroxide [7,12].

Autoimmune disease

In allergy, the inciting antigen is usually well-defined. This is far from the case for many autoimmune diseases. Considerable progress has been made in recent years in the identification of potential antigens in many autoimmune conditions [13–20]. However, pinning down the specificity of the T cells that cause these diseases is a complex process, further confused by the accumulation of different antigenic targets as the disease progresses, a process known as epitope spreading [21].

Despite these complexities, a number of human trials have taken place or are currently ongoing. Individuals considered at risk of diabetes have been given insulin or the pancreatic enzyme, glutamine decarboxylase, Gad-65, in an attempt to remove or modify the immune cells that cause pancreatic β -cell destruction (reviewed in [22]). These studies have found no delay in, or modification of, disease progression. Such trials, as Harrison *et al.* suggest, are unlikely to provide meaningful data without biomarkers to evaluate the impact of the therapy [22].

Similar approaches have been made to reduce disease progression in individuals with multiple sclerosis. Myelin peptides delivered alone, in the context of self-MHC molecules, or chemically fixed to patient cells have had a variety of effects from some reduction in relapses, no effect, or, in some cases, serious hypersensitivity (reviewed in [23]). Hypersensitivity responses were found in studies aiming to modulate the T cell response using peptides with a slightly altered sequence compared to the self peptide. Such altered peptide ligands (APLs) have different TCR-binding affinities and consequently provide altered signals for T cell activation leading to quantitative and/or qualitative changes in the T cell response. These effects are often specific for individual TCRs, which makes clinical translation from reductionist animal models to outbred human populations with more complex TCR repertoires inevitably unreliable.

Modulation of T cell responses has also been observed in rheumatoid arthritis (RA) patients given a peptide from dnaJ, a heat shock protein, which shares similarities with the RA 'shared epitope' [24,25]. Oral delivery of this peptide led to improved clinical responses and a shift from inflammatory to regulatory cytokine production. By contrast, intranasal delivery of a potential RA autoantigen, human cartilage glycoprotein-39, failed to provide any additional benefit compared to placebo [26]. More recent phase 1 studies in RA patients, and also in diabetic patients, have delivered the tolerising antigen using patient derived APCs. Dendritic cells (DCs), a key population of APCs, are differentiated from the patient's blood cells in the presence of tolerogenic signals, incubated with antigens before transfer back into the patient [27,28]. Such approaches ensure that the antigen is targeted to APCs that are likely to interact with the disease-causing T cells and can ameliorate disease in animal models by either reducing T cell effector responses and/or increasing immune regulation [29,30]. Likewise, antigen can be targeted to DCs by binding it to nanoparticles. Such approaches have been successfully used in animal models of allergy and autoimmunity [31,32].

As in allergy trials, the ability to track the target T cell population will be crucial to evaluate the success, and investigate the mechanisms of, disease modification. Recent studies using MHC class II tetramers containing

RA-associated peptides show that, while healthy controls and RA patients have similar numbers of RA-relevant T cells, the T cell populations from patients are more likely to have a memory T cell phenotype [33]. Memory T cells are generated following immune responses and may, as discussed below, present a more complex challenge to immunotherapy.


Mechanisms of action

Many animal studies have investigated the mechanisms of tolerance induction following AIT (Figure 2 and reviewed in [34]). Recent full genome transcriptional analysis of T cells exposed to a dose escalation of peptide suggest that, as in some of the human trials discussed above, AIT shifts T cells towards production of the regulatory cytokine, IL-10 [35]. In other experimental scenarios, a single high dose of AIT can drive T cells to apoptosis [36].

Many of these studies have, however, investigated the mechanisms of AIT in the context of naïve T cells. Allergen or autoreactive T cells either involved in ongoing or previous disease will be effector or memory T cells. These previously activated T cells show responses that are distinct from those of naïve T cells. Critically, they can be activated by lower levels of antigen and do not necessarily require costimulatory signals to make an effector response [37]. These differences suggest that effector and memory T cells will fail to be tolerised by AIT, instead potentially causing more immune-driven damage.

Our recent studies have investigated the consequences of exposing memory T cells to AIT. In a mouse allergy model, this response varied depending on the phenotype of the targeted memory cells with effector memory T cells undergoing a greater reduction in their subsequent responses than central memory T cells [38]. This might have consequences for when best to apply AIT in seasonal allergies. Effector memory T cells would be expected to be dominant during active allergen exposure, whereas immune memory during the 'off-season' might be maintained predominantly by central memory T cells. A good knowledge of the target cell population is, therefore, likely to be key to selecting the most effective treatment strategy. Similarly, while memory T cells proved to be initially resistant to AIT, they were altered such that further activation led to apoptosis [39]. These data suggest that memory cells retain information about their activation history which affects their subsequent responses. Potentially this cellular memory is retained via epigenetic changes to the cell's DNA. Indeed, we recently demonstrated that effector T cells silenced by AIT must express the cell surface protein, PD-1 [40]. Ligation of this inhibitory receptor serves to diminish TCR-driven T cell activation/function and its expression is usually tightly related to TCR ligation. AIT led to DNA

Figure 2



Mechanisms of action of AIT. AIT can alter the responses of allergen- or self-reactive T cells via a number of mechanisms. In some cases, the pathogenic T cells are deleted following recognition of antigen on tolerising APCs. Alternatively, these T cells can be modulated either differentiating into regulatory T cells that can suppress the immune response, or altering their effector response by decreasing inflammatory cytokine production or increasing expression of immunomodulatory molecules such as IL-10 or PD-1.

demethylation within the PD-1 gene promoter, enabling long-term expression of the receptor. These data begin to provide a molecular basis for the loss of pathogenic T cell function following AIT.

Future perspectives

From its demonstrated effectiveness in rodent models of autoimmune and allergic diseases, AIT has reached implementation in clinical trials. Key to translational success will be rigorous interrogation of molecular mechanisms, particularly in T effector and T memory cells. The use of peptide-MHC tetramer staining can identify epitope-specific T cell in a range of human diseases, most notably allergies where the inciting antigen is well characterized. This remains more of a challenge for autoimmune diseases with complex autoaggressive T cell repertoires. A variety of treatment regimes — routes, doses, delivery systems and proteins versus peptides — are being trialled. Reaching a consensus from the likely different outcomes will be challenging at the molecular level. It is necessary to specifically focus on those T cells with TCRs recognizing the therapeutic antigen. Low cell yields from patients will make large scale, unbiased, ‘omics’ analyses impractical. For this we must rely on more advanced mouse models that allow retrieval of sufficient material (tolerant T cells) to identify alterations in key genes/pathways that then can be interrogated more easily in human studies. These might provide useful biomarkers (in concert with tetramer staining) for confirming the

establishment of tolerance and, importantly, for signs that it is beginning to wane, prompting renewed immunotherapy.

Conflict of interest statement

Nothing declared.

Acknowledgements

MKLM is an Arthritis Research UK Career Development Fellow and recipient of a Marie Curie Career Integration Grant. Work in SMA's laboratory was supported by the UK Medical Research Council, the Wellcome Trust, the UK Multiple Sclerosis Society and Asthma Research UK.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

- Collado JA, Guitart C, Ciudad MT, Alvarez I, Jaraquemada D: **The repertoires of peptides presented by MHC-II in the thymus and in peripheral tissue: a clue for autoimmunity?** *Front Immunol* 2013, **4**:442.
- Bour-Jordan H, Esensten JH, Martinez-Llordella M, Penaranda C, Stumpf M, Bluestone JA: **Intrinsic and extrinsic control of peripheral T-cell tolerance by costimulatory molecules of the CD28/B7 family.** *Immunol Rev* 2011, **241**:180-205.
- Chappert P, Schwartz RH: **Induction of T cell anergy: integration of environmental cues and infectious tolerance.** *Curr Opin Immunol* 2010, **22**:552-559.
- Gamper CJ, Powell JD: **Genetic and biochemical regulation of CD4 T cell effector differentiation: insights from examination of T cell clonal anergy.** *Immunol Res* 2010, **47**:162-171.

5. Smarr CB, Bryce PJ, Miller SD: **Antigen-specific tolerance in immunotherapy of Th2-associated allergic diseases.** *Crit Rev Immunol* 2013, **33**:389-414.
 6. Smith CE, Eagar TN, Strominger JL, Miller SD: **Differential induction of IgE-mediated anaphylaxis after soluble vs. cell-bound tolerogenic peptide therapy of autoimmune encephalomyelitis.** *Proc Natl Acad Sci U S A* 2005, **102**:9595-9600.
 7. Francis JN, James LK, Paraskevopoulos G, Wong C, Calderon MA, Durham SR, Till SJ: **Grass pollen immunotherapy: IL-10 induction and suppression of late responses precedes IgG4 inhibitory antibody activity.** *J Allergy Clin Immunol* 2008, **121**:1120-1125.e1122.
 8. Wakkach A, Fournier N, Brun V, Breittmayer JP, Cottrez F, Groux H: **Characterization of dendritic cells that induce tolerance and T regulatory 1 cell differentiation in vivo.** *Immunity* 2003, **18**:605-617.
 9. Wambre E, DeLong JH, James EA, Torres-Chinn N, Pflutzer W, Mobs C, Durham SR, Till SJ, Robinson D, Kwok WW: **Specific immunotherapy modifies allergen-specific CD4(+) T-cell responses in an epitope-dependent manner.** *J Allergy Clin Immunol* 2014, **133**:872-879.e877.
- This study uses human MHC class II tetramers to demonstrate a preferential deletion of grass pollen specific T cells in allergic individuals following immunotherapy.
10. Zaleska A, Eiwegger T, Soyer O, van de Veen W, Rhyner C, Soyka MB, Bekpen C, Demiroz D, Treis A, Sollner S *et al.*: **Immune regulation by intralymphatic immunotherapy with modular allergen translocation MAT vaccine.** *Allergy* 2014, **69**:1162-1170.
- To increase the efficacy of specific immunotherapy, this study involved the injection of peptides from cat allergens directly into the lymph nodes of allergic individuals. This led to an increase in allergen specific regulatory T cells and a decrease in allergen specific effector cells.
11. Nelson H, Cartier S, Allen-Ramey F, Lawton S, Calderon MA: **Network meta-analysis shows commercialized subcutaneous and sublingual grass products have comparable efficacy.** *J Allergy Clin Immunol Pract* 2014.
- This important meta-analysis enables comparison of the effectiveness of different routes of delivery of allergen immunotherapy.
12. Senti G, Prinz Vavricka BM, Erdmann I, Diaz MI, Markus R, McCormack SJ, Simard JJ, Wuthrich B, Cramer R, Graf N *et al.*: **Intralymphatic allergen administration renders specific immunotherapy faster and safer: a randomized controlled trial.** *Proc Natl Acad Sci U S A* 2008, **105**:17908-17912.
 13. Tsark EC, Wang W, Teng YC, Arkfeld D, Dodge GR, Kovats S: **Differential MHC class II-mediated presentation of rheumatoid arthritis autoantigens by human dendritic cells and macrophages.** *J Immunol* 2002, **169**:6625-6633.
 14. Blass S, Union A, Raymackers J, Schumann F, Ungethum U, Muller-Steinbach S, De Keyser F, Engel JM, Burmester GR: **The stress protein BiP is overexpressed and is a major B and T cell target in rheumatoid arthritis.** *Arthritis Rheum* 2001, **44**:761-771.
 15. Law SC, Street S, Yu CH, Capini C, Ramnoruth S, Nel HJ, van Gorp E, Hyde C, Lau K, Pahau H *et al.*: **T-cell autoreactivity to citrullinated autoantigenic peptides in rheumatoid arthritis patients carrying HLA-DRB1 shared epitope alleles.** *Arthritis Res Ther* 2012, **14**:R118.
 16. Varrin-Doyer M, Shetty A, Spencer CM, Schulze-Topphoff U, Weber MS, Bernard CC, Forsthuber T, Cree BA, Slavin AJ, Zamvil SS: **MOG transmembrane and cytoplasmic domains contain highly stimulatory T-cell epitopes in MS.** *Neuro Immunol Neuroinflamm* 2014, **1**:e20.
 17. Wucherpfennig KW, Sette A, Southwood S, Oseroff C, Matsui M, Strominger JL, Hafler DA: **Structural requirements for binding of an immunodominant myelin basic protein peptide to DR2 isotypes and for its recognition by human T cell clones.** *J Exp Med* 1994, **179**:279-290.
 18. Elong Ngono A, Pette S, Salou M, Bahbouhi B, Soullillou JP, Brouard S, Laplaud DA: **Frequency of circulating autoreactive T cells committed to myelin determinants in relapsing-remitting multiple sclerosis patients.** *Clin Immunol* 2012, **144**:117-126.
 19. Yang J, Chow IT, Sosinowski T, Torres-Chinn N, Greenbaum CJ, James EA, Kappler JW, Davidson HW, Kwok WW: **Autoreactive T cells specific for insulin B:11-23 recognize a low-affinity peptide register in human subjects with autoimmune diabetes.** *Proc Natl Acad Sci U S A* 2014, **111**:14840-14845.
 20. Yang J, James EA, Sanda S, Greenbaum C, Kwok WW: **CD4⁺ T cells recognize diverse epitopes within GAD65: implications for repertoire development and diabetes monitoring.** *Immunology* 2013, **138**:269-279.
 21. Vanderlugt CL, Miller SD: **Epitope spreading in immune-mediated diseases: implications for immunotherapy.** *Nat Rev Immunol* 2002, **2**:85-95.
 22. Harrison LC, Wentworth JM, Zhang Y, Bandala-Sanchez E, Bohmer RM, Neale AM, Stone NL, Naselli G, Bosco JJ, Auyeung P *et al.*: **Antigen-based vaccination and prevention of type 1 diabetes.** *Curr Diab Rep* 2013, **13**:616-623.
 23. Lutterotti A, Martin R: **Antigen-specific tolerization approaches in multiple sclerosis.** *Expert Opin Investig Drugs* 2014, **23**:9-20.
 24. Koffeman EC, Genovese M, Amox D, Keogh E, Santana E, Matteson EL, Kavanaugh A, Molitor JA, Schiff MH, Posever JO *et al.*: **Epitope-specific immunotherapy of rheumatoid arthritis: clinical responsiveness occurs with immune deviation and relies on the expression of a cluster of molecules associated with T cell tolerance in a double-blind, placebo-controlled, pilot phase II trial.** *Arthritis Rheum* 2009, **60**:3207-3216.
 25. Gregersen PK, Silver J, Winchester RJ: **The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis.** *Arthritis Rheum* 1987, **30**:1205-1213.
 26. Landewe RB, Houbiers JG, Van den Bosch F, in't Hout J, Verschuere PC, Meijerink JH, van den Hoogen FH, Masek BA, Bruyn GA, Wouters JM *et al.*: **Intranasal administration of recombinant human cartilage glycoprotein-39 as a treatment for rheumatoid arthritis: a phase II, multicentre, double-blind, randomised, placebo-controlled, parallel-group, dose-finding trial.** *Ann Rheum Dis* 2010, **69**:1655-1659.
 27. Hilkens CM, Isaacs JD: **Tolerogenic dendritic cell therapy for rheumatoid arthritis: where are we now?** *Clin Exp Immunol* 2013, **172**:148-157.
 28. Giannoukakis N, Phillips B, Finegold D, Harnaha J, Trucco M: **Phase I (safety) study of autologous tolerogenic dendritic cells in type 1 diabetic patients.** *Diabetes Care* 2011, **34**:2026-2032.
 29. Martin E, Capini C, Duggan E, Lutzky VP, Stumbles P, Pettit AR, O'Sullivan B, Thomas R: **Antigen-specific suppression of established arthritis in mice by dendritic cells deficient in NF-kappaB.** *Arthritis Rheum* 2007, **56**:2255-2266.
 30. Stoop JN, Harry RA, von Delwig A, Isaacs JD, Robinson JH, Hilkens CM: **Therapeutic effect of tolerogenic dendritic cells in established collagen-induced arthritis is associated with a reduction in Th17 responses.** *Arthritis Rheum* 2010, **62**:3656-3665.
- This animal study demonstrates that nanoparticles are an effective delivery method for immunotherapy.
31. Maldonado RA, LaMothe RA, Ferrari JD, Zhang AH, Rossi RJ, Kolte PN, Griset AP, O'Neil C, Altretreuter DH, Browning E *et al.*: **Polymeric synthetic nanoparticles for the induction of antigen-specific immunological tolerance.** *Proc Natl Acad Sci U S A* 2015, **112**:E156-E165.
- See annotation to Ref. [30*].
32. Hunter Z, McCarthy DP, Yap WT, Harp CT, Getts DR, Shea LD, Miller SD: **A biodegradable nanoparticle platform for the induction of antigen-specific immune tolerance for treatment of autoimmune disease.** *ACS Nano* 2014, **8**:2148-2160.
- See annotation to Ref. [30*].
33. Snir O, Rieck M, Gebe JA, Yue BB, Rawlings CA, Nepom G, Malmstrom V, Buckner JH: **Identification and functional characterization of T cells reactive to citrullinated vimentin in HLA-DRB1*0401-positive humanized mice and rheumatoid arthritis patients.** *Arthritis Rheum* 2011, **63**:2873-2883.

34. Podojil JR, Miller SD: **Molecular mechanisms of T-cell receptor and costimulatory molecule ligation/blockade in autoimmune disease therapy.** *Immunol Rev* 2009, **229**:337-355.

35. Burton BR, Britton GJ, Fang H, Verhagen J, Smithers B, Sabatos-
 ●● Peyton CA, Carney LJ, Gough J, Strobel S, Wraith DC: **Sequential transcriptional changes dictate safe and effective antigen-specific immunotherapy.** *Nat Commun* 2014, **5**:4741.

Microarray technology is used to demonstrate that dose escalation of peptide immunotherapy progressively alters the transcriptome of targeted T cells.

36. Konkel JE, Frommer F, Leech MD, Yagita H, Waisman A, Anderton SM: **PD-1 signalling in CD4(+) T cells restrains their clonal expansion to an immunogenic stimulus, but is not critically required for peptide-induced tolerance.** *Immunology* 2010, **130**:92-102.

37. MacLeod MK, Clambey ET, Kappler JW, Marrack P: **CD4 memory T cells: what are they and what can they do?** *Semin Immunol* 2009, **21**:53-61.

38. Mackenzie KJ, Nowakowska DJ, Leech MD, McFarlane AJ,
 ●● Wilson C, Fitch PM, O'Connor RA, Howie SE, Schwarze J,

- Anderton SM: **Effector and central memory T helper 2 cells respond differently to peptide immunotherapy.** *Proc Natl Acad Sci U S A* 2014, **111**:E784-E793.

This study demonstrates that different memory T cell subsets are affected distinctly by immunotherapy indicating the importance of phenotyping the disease causing T cells.

39. David A, Crawford F, Garside P, Kappler JW, Marrack P,
 ●● MacLeod M: **Tolerance induction in memory CD4 T cells requires two rounds of antigen-specific activation.** *Proc Natl Acad Sci U S A* 2014, **111**:7735-7740.

By analyzing the effect of immunotherapy on memory T cells, this study demonstrates that memory T cells are more resistant to tolerance than naïve T cells.

40. McPherson RC, Konkel JE, Prendergast CT, Thomson JP,
 ●● Ottaviano R, Leech MD, Kay O, Zandee SE, Sweenie CH, Wraith DC *et al.*: **Epigenetic modification of the PD-1 (Pdc1) promoter in effector CD4(+) T cells tolerized by peptide immunotherapy.** *Elife* 2015, **4**.

This study demonstrates the importance of epigenetic modulation of the T cell inhibitory molecule, PD1, which is induced on effector T cells targeted by immunotherapy.