

Bakker, K. M., Martinez-Bakker, M. E., Helm, B., and Stevenson, T. J. (2016) Digital epidemiology reveals global childhood disease seasonality and the effects of immunization. Proceedings of the National Academy of Sciences of the United States of America, (doi:10.1073/pnas.1523941113).

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

http://eprints.gla.ac.uk/118563/

Deposited on: 01 June 2016

Enlighten – Research publications by members of the University of Glasgow http://eprints.gla.ac.uk

# Digital Epidemiology Reveals Global Childhood Disease Seasonality and the Effects of Immunization

Kevin M Bakker<sup>a,b</sup>, Micaela Martinez-Bakker<sup>c</sup>, Barbara Helm<sup>d</sup> and Tyler J Stevenson<sup>e,1</sup>

<sup>a</sup>Department of Ecology and Evolutionary Biology and

<sup>b</sup>Center for the Study of Complex Systems, University of Michigan, Ann Arbor, MI 48109, USA <sup>c</sup>Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, 08544, USA

<sup>d</sup>Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Graham Kerr Building, Glasgow G12 8QQ, United Kingdom

<sup>e</sup>Institute for Biological and Environmental Sciences, University of Aberdeen, Aberdeen, United Kingdom, AB24 2TZ

April 20, 2016

#### Abstract

Public health surveillance systems are important for tracking disease dynamics. In recent years, social and real-time digital data sources have provided new means of studying disease transmission. Such affordable and accessible data have the potential to offer new insights into disease epidemiology at national and international scales. We used the extensive information repository Google Trends to examine the digital epidemiology of a common childhood disease, chicken pox, caused by varicella zoster virus (VZV), over an eleven-year period. We (1) report robust seasonal information seeking behavior for chicken pox using Google data from 36 countries, (2) validate Google data using clinical chicken pox cases, (3) demonstrate that Google data can be used to identify recurrent seasonal outbreaks and forecast their magnitude and seasonal timing, and (4) reveal that VZV immunization significantly dampened seasonal cycles in information seeking behavior. Our findings provide strong evidence that VZV transmission is seasonal and that seasonal peaks show remarkable latitudinal variation. We attribute the dampened seasonal cycles in chicken pox information seeking behavior to VZV vaccine-induced reduction of seasonal transmission. These data and the methodological approaches provide a novel way to track the global burden of childhood disease, and illustrate population-level effects of immunization. The global latitudinal patterns in outbreak seasonality could direct future studies of environmental and physiological drivers of disease transmission.

#### Significance

Disease surveillance systems largely focus on infectious diseases with high mortality, while benign diseases often go unreported. Using chicken pox as an example, we demonstrate that internet queries can be used as a proxy for disease incidence when reporting is lacking. We established that Google Trends accurately reflected clinical cases in countries with surveillance, and thus, population-level dynamics of chicken pox incidence. Then we discovered robust seasonal variation in query behavior, with a striking latitudinal gradient on a global scale. Third, we showed that real-time data-mining of queries could forecast the timing and magnitude of outbreaks. Finally, our analyses revealed that

countries with government-mandated vaccination programs have significantly reduced seasonality of queries, indicating vaccination efforts mitigated chicken pox outbreaks.

#### Introduction

Childhood infectious diseases continue to be a major global problem, and surveillance is needed to inform strategies for the prevention and mitigation of disease transmission. Our ability to characterize the global picture of childhood diseases is limited, as detailed epidemiological data are generally nonexistent or inaccessible across much of the world. Available data suggest that recurrent outbreaks of acute infectious diseases peak within a relatively consistent, but disease-specific seasonal window, which differs geographically [1, 2, 3, 4, 5]. Geographic variation in disease transmission is poorly understood, suggesting substantial knowledge gains from methods that can expand global epidemiological surveillance. Seasonal variations in host-pathogen interactions are common in nature [6]. In humans, the immune system undergoes substantial seasonal changes in gene expression, which is inverted between European locations and Oceana [7]. The regulation of seasonal changes in both disease incidence and immune defense is known to interact with environmental factors such as annual changes in day length, humidity and ambient temperature [8]. Accordingly, quantification of global spatio-temporal patterns of disease incidence can help to disentangle environmental, demographic, and physiological drivers of infectious disease transmission. Furthermore, the recognition of the regional timing of outbreaks would establish the groundwork for anticipating clinical cases, and when applicable, initiating public health interventions.

Since childhood disease outbreaks are often explosive and short-lived [9], temporally rich (i.e., weekly or monthly) data are needed to understand their dynamics. Similarly, in order to establish the recurrent nature of outbreaks that occur at annual or multi-annual frequencies, long-term data are needed. Thus, ideal disease incidence data have both high temporal resolution and breadth (i.e., frequent observations over many years). Over the past decade, the internet has become a significant health resource for the general public and health professionals [10, 11]. Internet query platforms, such as Google Trends, have provided powerful and accessible resources for identifying outbreaks and for implementing intervention strategies [12, 13, 14]. Research on infectious disease information seeking behaviour has demonstrated that internet queries can complement traditional surveillance by providing a rapid and efficient means of obtaining large epidemiological datasets [13, 15, 16, 17, 18]. For example, epidemiological information contained within Google Trends has been used in the study of rotavirus, norovirus, and influenza [14, 15, 17, 18]. These tools offer substantial promise for the global monitoring of diseases in countries that lack clinical surveillance but have sufficient internet coverage to allow for surveillance via digital epidemiology.

Here we focused on one common childhood disease, chicken pox, as a study system because it would allow us to validate internet query data using clinical data from the small number of geographically distinct countries that report cases, and to address the impact of VZV vaccination on outbreaks. Chicken pox—a highly contagious disease caused by VZV—has low mortality but exceptionally high morbidity, with most unvaccinated children infected by age 15 in developed countries [19, 20]. The burden of VZV extends beyond chicken pox, because a VZV infection causes fluid filled blisters, which eventually burst, creating the opportunity for infection from various invasive bacterial pathogens (e.g., Group A streptococcal infections) [21]. Chicken pox is not included in the World Health Organizations (WHO) global monitoring system for vaccine preventable diseases [22], meaning there are few countries which report clinical cases. In the United States (US), a country that immunizes against VZV; chicken pox was historically a notifiable disease. A lapse in national surveillance 1981–2001 compromised the ability of researchers to examine the long-term disease dynamics and the impact of immunization. [3, 23, 24]. Although the clear symptomatology of chicken pox makes the disease readily observable at the individual level, the lack of reporting makes VZV transmission dynamics cryptic at the population-level and obscures



**Figure 1: Left Figure:** Global seasonality of chicken pox outbreaks measured using Google Trends as a proxy for chicken pox dynamics. Countries are organized by geographic region and latitude. Latitudinal variation in seasonal chicken pox information seeking behavior was observed for countries with wavelet confirmed significant seasonality. The seasonality was estimated by fitting a General Additive Model (GAM) to the detrended Google data from each country. GAM values using week number as the predictive variable for Google data are shown in the heatmap and correspond to the GAM curves to the right. **Right Figure:** Data processing and regional GAM values (Top row) Data processing steps. (Top row, left panel) Detrended Google data for Italy. (Top row, middle panel) Box-and-whisker plot of Google data for Italy: 1st-to-3rd quartiles in solid color with whiskers representing 95% confidence intervals. All other panels represent GAM values using week number as the predictive variable for Google data in each country. European countries include Finland, Sweden, Denmark, Ireland, Netherlands, Poland, UK, Hungary, France, Romania, Italy, Spain, and Portugal. Asian countries include Vietnam, India, Thailand, and the Philippines. Americas include Mexico (with peak in week 10), Colombia, Brazil, and Argentina.

its spatio-temporal patterns.

The VZV vaccine is on the WHO list of essential medicines, which specifies the most important medicines needed for basic health systems [25], and is available as either stand-alone VZV or as the measles, mumps, rubella, and varicella vaccine (MMRV). However, the US, Germany, Canada, Uruguay, Australia and regions of Spain and Italy are among the few locations that have included VZV vaccination in their childhood immunization schedules for multiple years [26, 27, 28, 29, 30]. Short term surveillance studies in select locations of the US have demonstrated that moderate levels of vaccine coverage were able to dramatically reduce chicken pox incidence [31, 32], partially through the effect of herd immunity [33]. However, the effects of VZV vaccination on morbidity and mortality remain poorly understood [29] because global chicken pox report rates are low (e.g., the US rate is estimated to range from < 0.1% to 20% [34]). Therefore, the public health community is faced with a scarcity of chicken pox data to inform VZV vaccination policy. In certain locations (e.g., Madrid), VZV immunization has ceased, possibly due to the lack of information regarding the effects of immunization that can be used to assess health gains and economic feasibility. Clearly, there remains a lack of research, especially in countries that have recently introduced the VZV vaccine into the childhood immunization schedule [29, 35].

In this study, we took advantage of the extensive data available in Google Trends to study the global seasonal transmission of chicken pox. We (1) data mined chicken pox information seeking behavior

using language-specific Google queries of "chicken pox" from 36 countries spanning 5 continents over an eleven-year period (See SI Appendix, Table S3), and characterized the seasonality of outbreaks. We then (2) validated Google Trends data using detailed time series of clinical cases from 5 countries within 4 continents. We (3) confirmed that statistical models have a profound ability to forecast outbreaks in unvaccinated and vaccinated populations. Finally, we (4) verified the impact of VZV vaccination on the seasonality of chicken pox outbreaks.

# Seasonality of Chicken Pox Information Seeking & Validation

We detected significant seasonality of the Google Trends data [36] in 27 out of the 34 countries for which weekly data were available. Each geographic region in our study displayed distinct seasonal patterns of information seeking behaviour (Fig 1). A strong latitudinal pattern was clearly discernible (Fig 1), with peaks early in the year in the Northern Hemisphere and later in the Southern Hemisphere, corresponding to springtime outbreaks worldwide. These spring peaks agree with the seasonal timing found in historical datasets and previous studies of chicken pox [1, 3, 37, 38, 39] (See SI Appendix, Fig S8). European countries were mostly unimodal, with a peak in March–May, but some had an additional, smaller, peak in late December. Several countries in the Southern hemisphere, including Australia, New Zealand, South Africa, and those located in South America, had a single discernible peak at various times of the year. China and Japan had bimodal peaks, which occurred March–May and December–January, balanced by deep summer troughs occurring July–August. Other Asian countries had a single peak that occurred February–March punctuated by a relatively shallow trough.

To validate Google Trends as a reasonable proxy for chicken pox dynamics, we compared clinical data to Google Trends data from five countries. We found that in the three countries lacking VZV vaccination— Mexico, Thailand, and Estonia—chicken pox information seeking was significantly correlated with reported cases of chicken pox, with  $R^2 = 0.70$ , 0.81, and 0.65 respectively (Fig 2). The correlation was reduced, but still significant, in Australia ( $R^2 = 0.26$ ), which implemented nationwide immunization in 2005. In the US, which has actively vaccinated since 1995, the association between information seeking and reported chicken pox cases was low ( $R^2 = 0.018$ ). To better understand the patterns in these countries, we investigated the context of the language-specific Google searches of "chicken pox". We compared the context of top searches containing "chicken pox" to determine whether they were searches for disease, symptoms, or treatment (i.e., indicative of disease in the household/community) or vaccination (i.e., not necessarily related to disease incidence).

The proportion of searches related to disease—rather than chicken pox vaccination or other search contexts—differed between focal countries. In Mexico and Thailand, which do not vaccinate, we classified the relative frequency of the top searches indicating disease to be 0.82 and 0.80 (See SI Appendix, Fig S7). In both Australia and the US, where the VZV vaccine is required, vaccination and other search contexts had a higher relative frequency, while disease indicators had a frequency of 0.71 and 0.66, respectively. Interestingly, in the US, the second highest query involved chicken pox vaccination (See SI Appendix, Table S2). Estonia had low search volume and the top searches were not available.

These data indicate the value of adding further layers to search terms. By doing so, we identified two patterns in information seeking: i) the data reflected seasonal dynamics in most countries (Mexico, Thailand, Estonia, and Australia); and ii) in the US, where vaccination has long been introduced, the data reflected a shift in the motivation of information seeking. These findings suggest that Google Trends reflect chicken pox dynamics more closely in countries that do not vaccinate versus countries that do vaccinate, and highlight the dynamic nature of information seeking. This likely explains why chicken pox information seeking in Mexico, Thailand, Estonia, and Australia, but not in the US, strongly reflected reported cases.



**Figure 2:** Relationship between chicken pox cases and information seeking. (Left column) Time series of reported chicken pox cases and information seeking behaviour for chicken pox (i.e., Google Trends data) in Mexico, the US, Thailand, Australia, and Estonia. Google data were detrended to remove long-term trends and focus on seasonal variation in information seeking. (Right column) Relationship between reported cases of chicken pox and chicken pox information seeking when both were available, with applicable  $R^2$  and p values. Chicken pox case data from Mexico and the US were weekly, whereas chicken pox case data from Thailand, Australia, and Estonia were monthly.

# Forecasting Outbreaks using Google Trends

To determine whether the information seeking behaviour observed in Google data, T, was able to forecast chicken pox outbreak magnitude and timing, we built and fitted eight different statistical models to forecast chicken pox case data. We evaluated the epidemiological information contained in Google Trends by comparing these Google Trends models with a seasonal null model that did not incorporate Google data (See SI Appendix,Table S1, Fig S1, S2). Three models that included the Google data fit better than the null model, however we focused on the model with best fit, and refer to it as the Google model hereafter. The null model lacked information seeking in the force of infection parameter, which we defined as the monthly per capita rate at which children age 0–14 years are infected. In order to estimate the number of symptomatic VZV infections per month, we multiplied the force of infection with an estimate of the population aged 0–14 years (SI, Eq 1). Both the Google and null models were fitted to the case data from a VZV-unvaccinated population (Thailand), which showed robust seasonality, and a VZV-vaccinated population (Australia), which exhibited reduced seasonality. To estimate the number of symptomatic VZV infections each month,  $I_t$ , we used Google Trends data from the previous two months,  $T_{t-1}$  and  $T_{t-2}$ , where t is time in monthly time steps. The chicken pox process model tracked the force of infection.

$$\lambda_t = \left[\beta_1 \cos\left(\frac{2\pi}{12}(t+\omega)\right) \mathsf{T}_{t-1} + \beta_2 |\mathsf{T}_{t-1} - \mathsf{T}_{t-2}| + \beta_3\right] \epsilon_t.$$
(1)

The model also contained environmental stochasticity,  $\epsilon_t$ , which was drawn from a gamma distribution with a mean of 1 and variance  $\theta$ . We estimated 6 parameters for the model: the mean and the phase of the seasonality ( $\beta_1$  and  $\omega$ ), a parameter scaling the Google Trends data ( $\beta_2$ ), the baseline force of infection ( $\beta_3$ ), the process noise dispersion parameter ( $\theta$ ), and the reporting dispersion parameter ( $\tau$ ) of a normal distribution, with a mean of 1, from which case reports were drawn. The parameters were estimated using maximum likelihood by iterated particle filtering (MIF) in the R-package pomp [40, 41].

The Google Trends model fit the case data (See SI Appendix, Fig S1) and performed better than the null model (which estimated 5 parameters) for both Australia and Thailand; null model AICs were > 28 units above Google Trends model AICs in both locations. Since each model was seasonally forced, all models captured the seasonal timing of outbreaks. However, the Google Trends model was able to predict the interannual variation in outbreak size (Fig 3), while the null model could not (See SI Appendix, Fig S1).

### The Signature of Varicella Zoster Immunization

We investigated the signature of VZV immunization by examining Google Trends data in countries that actively immunize and those that do not. Seasonality of information seeking behaviour was much stronger in countries lacking active immunization programs than in countries that include the VZV vaccine as part of the childhood immunization schedule (Fig 3). Germany, which made the VZV vaccine mandatory in July 2004 [27, 42], had weakening seasonality in information seeking until 2009, when a second VZV booster dose was added to the immunization schedule, drastically reducing information seeking seasonality (See SI Appendix, Fig S5, S6). In Australia, where the VZV vaccine was publicly funded in November 2005 [43], the amplitude of information seeking was severely dampened by the end of 2007. In the US, immunization began in 1995 [31] while Canada required the vaccination starting in 2000 [44]. In these two countries, where the VZV vaccine introduction predated Google Trends data, little seasonality was observed in the Google Trends data (Fig 3, See SI Appendix, S9). In Spain and Italy, where only a few regions or municipalities require VZV immunization [29], minimal change was observed in the Google Trends



Figure 3: Left figure Forecasting chicken pox cases using Google Trends. (Top) Forecasting model schematic, Google Trends data from months t - 2 and t - 1 are used to predict chicken pox cases in month t. (Bottom left) Observed and predicted chicken pox cases in Australia (active immunization) and Thailand (no immunization) from 10000 simulations of the fitted models parameterized with the maximum likelihood estimates; overpredicted (green hash marks) and underpredicted (red hash marks) regions are indicated. (Bottom right) Model predicted cases versus observed chicken pox cases along the dotted 1-to-1 line. Right figure Detrended chicken pox information seeking in relation to immunization. Data are weekly; x-axes indicate time, and y-axes are the detrended Google data (same scale for all panels). Countries with universal (national) immunization in red, countries with select (regional or municipal) immunization in blue, and countries lacking any mandatory immunization in black. Panels 1-2: the UK and Brazil, two countries with no immunization. Panels 3-4: Spain and Italy, two countries with no universal (national) immunization, but with select regional or municipal immunization. Vertical lines identify the implementation (blue for select, red for national) or termination (black) of immunization efforts. Cities and regions on these panels indicate where these efforts were focused. Panels 5-6: Australia and Germany, two countries that implemented national immunization since 2004. Australia had the vaccine since 2001, but nationwide immunization was not funded by the government until November 2005. Germany required a single dose for every child in July 2004, provided nationalized payment in 2007, and required a second dose in 2009. Panel 7: the US, which has had national immunization since 1995, required a booster dose in 2006.

dynamics following immunization implementation. However, in Spain there was a reduction in search amplitude when immunization efforts were at their maximum, likely indicating a reduction in searches following immunization, similar to Germany (See SI Appendix, Fig S6).

Since information seeking behavior strongly correlates with seasonal outbreaks of chicken pox (Fig 2), the loss of information seeking seasonality in countries that immunize can signal the loss of recurrent seasonal chicken pox outbreaks, indicating outbreak mitigation driven by VZV immunization. We suggest that if (i) disease transmission is seasonal—as it is for chicken pox and other childhood diseases—and (ii) vaccination reduces disease transmission, then the impact of immunization can be measured as the reduction in seasonal outbreak amplitude. This is because vaccination will strongly diminish the transmission rate during the high-transmission season. In the case of chicken pox, the reduction of seasonality in information seeking is likely due to diminished outbreak seasonality seen in clinical data (Fig 2, See SI Appendix, S8) and the subsequent shift of information seeking from disease queries to vaccination queries.

## Discussion

In this study, we used digital epidemiology of chicken pox to (1) reveal previously unreported seasonal outbreaks on a global scale, which displayed robust latitudinal dependence, (2) confirm the reliability of the Google data against known clinical cases, (3) forecast the size of annual outbreaks, and (4) uncover the population-level effects of routine VZV immunization. The lack of contemporary reporting, due to the benign nature of infections and the increased use of immunization, have made it difficult to decipher modern VZV global epidemiology. Here, we established that information seeking behaviour can be applied to reveal the underlying epidemiology of a childhood disease, chicken pox.

Our analyses reveal profound, global patterns of seasonality in chicken pox transmission dynamics. These seasonal patterns are spatially structured: we have demonstrated a latitudinal pattern in the timing of outbreaks, with inverted phases between the southern and northern hemisphere and an apex in the spring. Evidence of the underlying biological basis for seasonality in chicken pox transmission remains an open question. There is a significant latitudinal shift (i.e., near 6 months) in chicken pox outbreak timing from the northern to southern hemisphere, which suggests an influence of environmental, biological, and/or behavioral drivers that vary with latitude, such as seasonal immunity, environmental factors, and/or school terms. Interestingly, Google Trends also revealed seasonal variation in croup, fifth disease, and hand, foot, and mouth disease; each of these childhood diseases exhibited a unique annual peak with little overlap in their seasonal window of outbreak occurrence. The lack of synchrony among childhood diseases likely indicates that school terms and holidays are not the primary drivers determining outbreak timing (See SI Appendix, Fig S3). We speculate that seasonal information seeking behaviour linked to childhood illnesses reflects biologically based seasonality of host-pathogen interactions [6, 7]. Our present data open new possibilities for extensive global analyses, which could disentangle contributions of different seasonal drivers to a broad range of infectious diseases. There is a pressing need for such knowledge as global seasonality is becoming rapidly modified and disrupted through human action, with potentially far-reaching implications for infectious disease transmission [45].

By taking advantage of freely available, real-time, internet search query data, we were able to validate information seeking behaviour as an appropriate proxy for otherwise cryptic chicken pox outbreaks and use those data to forecast outbreaks one month in advance. Our modeling approach, which incorporated Google Trends and the knowledge of spring peaks, was able to better forecast outbreaks than models that ignored Google Trends. While this was particularly clear for Thailand, which does not immunize against VZV, it also held for Australia, a country that vaccinates. These results suggest that information seeking can be used for rapid forecasting, when the reporting of clinical cases is unavailable or too slow.

Comparisons of Google Trends data with the reported cases in countries that lacked VZV immunization revealed a significant positive relationship (70%, 81%, and 65% of variation in Google Trends explained by variation in reported cases). However, the relationship significantly decreased in countries that included VZV vaccination in their childhood immunization schedule and displayed either no seasonality or low-amplitude seasonal cycles (e.g., the US, 1.8% and Australia, 26%). Interestingly, in Italy and Spain, where the VZV vaccine was only required in specific regions or municipalities of the country, no change in seasonal information seeking behaviour was detected in the face of vaccination, implying that widespread immunization is necessary to mitigate seasonal cycles of disease and information seeking. These findings, particularly from the highly vaccinated countries in our data—the US and Australia—indicate that immunization programs diminish seasonal information seeking behaviour and likely represent decreased seasonality of outbreaks.

Studies of disease transmission at the global level, and the success of interventions, are limited by data availability. Disease surveillance is a major obstacle in the global effort to improve public health, and is made difficult by under-reporting, language barriers, the logistics of data acquisition, and the time required for data curation. We demonstrated that seasonal variation in information seeking reflected disease dynamics, and as such, we were able to reveal global patterns of outbreaks and their mitigation via immunization efforts. Thus, digital epidemiology is an easily accessible tool that can be used to complement traditional disease surveillance, and in certain instances, may be the only readily available data source for studying seasonal transmission of non-notifiable diseases. We focused on chicken pox and its dynamics to demonstrate the strength of digital epidemiology for studying childhood diseases at the population level, because VZV is endemic worldwide and the global landscape of VZV vaccination is rapidly changing. Unfortunately, there is still a geographic imbalance of data sources: the vast majority of digital epidemiology data are derived from temperate regions with high internet coverage. However, because many childhood diseases remain non-notifiable throughout the developing world, digital epidemiology provides a valuable approach for identifying recurrent outbreaks when clinical data are lacking. It remains an open challenge to extend the reach of digital epidemiology to study other benign and malignant diseases with under-reported outbreaks and to identify spatio-temporal patterns, where knowledge about the drivers of disease dynamics are most urgently needed.

#### Materials and Methods

Google Trends data for 'chicken pox' were downloaded, and examined for seasonality. These data were then compared against reported cases of chicken pox in countries where case reports were available. We then constructed and tested multiple statistical models to determine whether Google Trends data could forecast chicken pox seasonality. Finally, we examined the effect of national immunization campaigns on the seasonality and amplitude of Google searches. Further methodological descriptions are included in the supporting information. This study was done with freely available, de-indentified, pre-existing data, thus no consent was required.

#### Acknowledgements

We would like to thank Fernando Gonzalez-Dominguez and Gilberto Vaughan for providing the chicken pox case reports from Mexico, and the Estonia Health Board, Department of Communicable Disease Surveillance and Control, for Estonian chicken pox case reports. KB would like to thank Mercedes Pascual, her lab, and Marisa Eisenberg for helpful comments. Jesus Cantu (research assistant, Princeton University) translated and categorized chicken pox searches from Mexico, Thailand, Australia, and the US.

# References

- 1. London, W. P & Yorke, J. A. (1973) Recurrent Outbreaks of Measles, Chickenpox and Mumps. I. Seasonal Variation in Contact Rates. *American Journal of Epidemiology* **98**, 453–468.
- Metcalf, C. J. E, Bjornstad, O. N, Grenfell, B. T, & Andreasen, V. (2009) Seasonality and Comparative Dynamics of Six Childhood Infections in Pre-vaccination Copenhagen. *Proceedings of the Royal Society B: Biological Sciences* 276, 4111–4118.
- 3. van Panhuis, W. G, Grefenstette, J, Jung, S. Y, Chok, N. S, Cross, A, Eng, H, Lee, B. Y, Zadorozhny, V, Brown, S, Cummings, D, & Burke, D. S. (2013) Contagious diseases in the United States from 1888 to the present. *New England Journal of Medicine* **369(22)**, 2152–2158.
- 4. Altizer, S, Dobson, A, Hosseini, P, Hudson, P, Pascual, M, & Rohani, P. (2006) Seasonality and the Dynamics of Infectious Diseases. *Ecology letters* **9**, 467–84.
- 5. Grassly, N & Fraser, C. (2006) Seasonal Infectious Disease Epidemiology. *Proceedings of the Royal Society B: Biological Sciences* 273, 2541–50.
- 6. Martinez-Bakker, M & Helm, B. (2015) The influence of biological rhythms on host–parasite interactions. *Trends in Ecology & Evolution*.
- Dopico, X. C, Evangelou, M, Ferreira, R. C, Guo, H, Pekalski, M. L, Smyth, D. J, Cooper, N, Burren, O. S, Fulford, A. J, Hennig, B. J, et al. (2015) Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nature communications* 6.
- 8. Stevenson, T. J & Prendergast, B. J. (2015) Photoperiodic time measurement and seasonal immunological plasticity. *Frontiers in Neuroendocrinology* **37**, 76–88.
- 9. Keeling, M & Rohani, P. (2008) *Modeling infectious diseases in humans and animals*. (Princeton University Press).
- 10. Higgins, O, Sixsmith, J, Barry, M, & Domegan, C. (2011) A literature review on health information seeking behaviour on the web: a health consumer and health professional perspective. *ECDC Technical Report, Stockholm.*
- 11. Brownstein, J, Freifeld, C, & Madoff, L. (2009) Digital disease detection—harnessing the web for public health surveillance. *New England Journal of Medicine* **360(21)**, 2153–2157.
- 12. Bryden, J, Funk, S, & Jansen, V. A. (2013) Word usage mirrors community structure in the online social network Twitter. *EPJ Data Science* **1**, 1–9.
- Salathé, M, Bengtsson, L, Bodnar, T. J, Brewer, D. D, Brownstein, J. S, Buckee, C, Campbell, E. M, Cattuto, C, Khandelwal, S, Mabry, P. L, & Vespignani, A. (2012) Digital epidemiology. *PLoS Computational Biology* 8, 1–5.
- 14. Hulth, A, Rydevik, G, Linde, A, & Montgomery, J. (2009) Web queries as a source for syndromic surveillance. *PloS one* **4(2)**, e4378.
- 15. Shaman, J & Karspeck, A. (2012) Forecasting seasonal outbreaks of influenza. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 20425–30.
- 16. Ginsberg, J, Mohebbi, M. H, Patel, R. S, Brammer, L, Smolinski, M. S, & Brilliant, L. (2009) Detecting influenza epidemics using search engine query data. *Nature* **457**, 1012–1014.

- 17. Desai, R, Lopman, B. A, Shimshoni, Y, Harris, J. P, Patel, M. M, & Parashar, U. (2012) Use of internet search data to monitor impact of rotavirus vaccination in the united states. *Clinical Infectious Diseases* cis121.
- Desai, R, Hall, A, Lopman, B, Shimshoni, Y, Rennick, M, Efron, N, Matias, Y, Patel, M, & Parashar, U. (2012) Norovirus disease surveillance using google internet query share data. *Clinical Infectious Diseases* 55(8), e75–e78.
- 19. Sengupta, N & Brewer, J. (2009) A global perspective of the epidemiology and burden of varicellazoster virus. *Current Pediatric Reviews* **5.4**, 207–228.
- 20. WHO. (2014) Background Paper on Varicella Vaccine: SAGE Working Group on Varicella and Herpes Zoster Vaccines.
- Vugia, D, Peterson, C, Meyers, H, Kim, K, Arrieta, A, Schlievert, P, Kaplan, E, Werner, S, & Mascola.,
  L. (1996) Invasive group a streptococcal infections in children with varicella in southern california. *The Pediatric infectious disease journal* **15(2)**, 146–150.
- 22. WHO. (2015) Immunization surveillance, assessment and monitoring/Vaccine-preventable diseases. http://apps.who.int/immunization\_monitoring/diseases/en/. Accessed Sept 15, 2015.
- 23. CDC. (2012) Prevention and control of influenza with vaccines: Recommendations of the Advisory Committee on Immunization Practices (ACIP) United States 2012–13 season. *Morbidity and Mortality Weekly Report August 17, 2012* **61**, 605–8.
- 24. CDC. (2015) Conducting Varicella Surveillance. http://www.cdc.gov/chickenpox/hcp/conductingsurveillance.html. Accessed Sept 15, 2015.
- 25. WHO. (2013) WHO Model List of Essential Medicines. http://www.who.int/medicines/publications/essentialmedicines/en/. Accessed Sept 15, 2015.
- 26. PHAC. (2015) Varicella (Chickenpox). http://www.phac-aspc.gc.ca/im/vpd-mev/varicella-eng.php. Ac-cessed Sept 15, 2015.
- 27. Koch-Institut, R. (2006) Empfehlungen der Stündigen Impfkommission (STIKO) am Robert Koch-InstitutStand. *Epidemiol Bulletin* **30**, 235–254.
- 28. Siedler, A & Arndt, U. (2008) Varicella sentinel in Germany shows success of vaccination. 26th Annual Meeting of the European Society for Paediatric Infectious Diseases (ESPID).
- 29. ECDC. (2014) Varicella vaccine in the European Union. ECDC. Stockholm.
- Quian, J, Rüttimann, R, Romero, C, Dall Orso, P, Cerisola, A, Breuer, T, Greenberg, M, & Verstraeten, T. (2008) Impact of universal varicella vaccination on 1-year-olds in uruguay: 1997–2005. Archives of Disease in Childhood 93, 845–850.
- Seward, J. F, Watson, B. M, Peterson, C. L, Mascola, L, Pelosi, J. W, Zhang, J. X, Maupin, T. J, Goldman, G. S, Tabony, L. J, Brodovicz, K. G, Jumaan, A. O, & Wharton, M. (2002) Varicella disease after introduction of varicella vaccine in the United States, 1995–2000. *Journal of the American Medical Association* 287, 606–611.
- Baxter, R, Tran, T. N, Ray, P, Lewis, E, Fireman, B, Black, S, Shinefield, H. R, Coplan, P. M, & Saddier, P. (2014) Impact of vaccination on the epidemiology of varicella: 1995–2009. *Pediatrics* 134, 24–30.

- 33. Clements, D, Zaref, J. I, Bland, C. L, Walter, E. B, & Coplan, P. M. (2001) Partial uptake of varicella vaccine and the epidemiological effect on varicella disease in 11 day-care centers in North Carolina. *Archives of Pediatrics & Adolescent Medicine* **4**, 455–461.
- 34. CDC. (1999) Evaluation of varicella reporting to the National Notifiable Disease Surveillance System– United States, 1972–1997. *Morbidity and mortality weekly report January 29, 1999* **3**.
- 35. CDC. (2015) Infectious Diseases Related to Travel. http://wwwnc.cdc.gov/travel/yellowbook/2016/infectiousdiseases-related-to-travel/varicella-chickenpox. Accessed Sept 9, 2015.
- 36. Google. (2015) Google Trends. https://www.google.com/trends/.
- 37. Health-Department. (2002) Surveillance of viral pathogens in Australia: varicella. *Communicable Diseases Intelligence* **26**.
- 38. Seward, J, Galil, K, & Wharton, M. (2000) *Epidemiology of varicella*. (Cambridge University Press), pp. 187–205.
- Vergara-Castañeda, A, Escobar-Gutiérrez, A, Ruiz-Tovar, K, Sotelo, J, Ordoñez, G, Cruz-Rivera, M. Y, Fonseca-Coronado, S, Martinez-Guarneros, A, Carpio-Pedroza, J. C, & Vaughan, G. (2012) Epidemiology of varicella in mexico. *Journal of Clinical Virology* 55, 51–57.
- 40. King, A. A, Nguyen, D, & Ionides, E. L. (2015) Statistical inference for partially observed Markov processes via the R package pomp. *Journal of Statistical Software* **arXiv preprint arXiv:1509.00503**.
- 41. King, A. A, Ionides, E. L, Bretó, C. M, Ellner, S. P, Ferrari, M. J, Kendall, B. E, Lavine, M, Nguyen, D, Reuman, D. C, Wearing, H, & Wood, S. N. (2015) *pomp: Statistical Inference for Partially Observed Markov Processes*, (http://kingaa.github.io/pomp). R package, version 1.2.1.1.
- 42. Rasch, G & Hellenbrand, W. (2004) Germany adds varicella vaccine to the national vaccination programme. *Eurosurveillance* **8**.
- 43. Carville, K. S, Riddell, M. A, & Kelly, H. A. (2010) A decline in varicella but an uncertain impact on zoster following varicella vaccination in Victoria, Australia. *Vaccine* **28**, 2532–2538.
- 44. Sweet, L, Gallant, P, Morris, M, & Halperin, S. A. (2003) Canada's first universal varicella immunization program: Lessons from Prince Edward Island. *The Canadian Journal of Infectious Diseases (Journal Canadien des Maladies Infectieuses)* **14**, 41–44.
- Stevenson, T, Visser, M, Arnold, W, Barrett, P, Biello, S, Dawson, A, Denlinger, D, Dominoni, D, Ebling, F, Elton, S, et al. (2015) Disrupted seasonal biology impacts health, food security and ecosystems. *Proc. R. Soc. B* 282, 20151453.