

# Genome Sequence of the Parainfluenza Virus 5 Strain That Persistently Infects AGS Cells

Elizabeth Wignall-Fleming,<sup>a,b</sup> Dan F. Young,<sup>b</sup> Steve Goodbourn,<sup>c</sup> Andrew J. Davison,<sup>a</sup> Richard E. Randall<sup>b</sup>

MRC – University of Glasgow Centre for Virus Research, Glasgow, United Kingdom<sup>a</sup>; Centre for Biomolecular Sciences, School of Biology, University of St. Andrews, St. Andrews, United Kingdom<sup>b</sup>; Division of Basic Medical Sciences, St. George's, University of London, London, United Kingdom<sup>c</sup>

**We have sequenced the parainfluenza virus 5 strain that persistently infects the commonly used AGS human cell line without causing cytopathology. This virus is most closely related to human strains, indicating that it may have originated from biopsy material or from laboratory contamination during generation of the cell line.**

Received 17 May 2016 Accepted 25 May 2016 Published 21 July 2016

**Citation** Wignall-Fleming E, Young DF, Goodbourn S, Davison AJ, Randall RE. 2016. Genome sequence of the parainfluenza virus 5 strain that persistently infects AGS cells. *Genome Announc* 4(4):e00653-16. doi:10.1128/genomeA.00653-16.

**Copyright** © 2016 Wignall-Fleming et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Richard E. Randall, rer@st-and.ac.uk.

Parainfluenza virus 5 (PIV5; genus *Rubulavirus*, subfamily *Paramyxovirinae*, family *Paramyxoviridae* [1]) possesses a single-stranded, negative-sense RNA genome containing seven genes that encode eight proteins (2). This virus has been isolated from numerous species, including humans, monkeys, pigs, cattle, and dogs (3). PIV5 is not known to cause acute disease in humans but has been associated with kennel cough in dogs and acute respiratory symptoms in pigs and calves. It can also cause unrecognized persistent infections of tissue culture cells and is likely to establish persistent infections *in vivo* (4).

The AGS cell line was derived from a human gastric adenocarcinoma (5) and has been used commonly in biomedical research, but it is persistently infected with a strain of PIV5 (4). To determine the genetic characteristics of this strain (PIV5-AGS), RNA was isolated by TRIzol (Invitrogen) extraction of AGS cells (ATCC CRL-1739) and of A549 cells (ATCC CCL-185) infected with the strain isolated from AGS cells. Libraries were prepared using a TruSeq stranded mRNA kit (Illumina) and sequenced using a MiSeq platform (Illumina). The data, which consisted of 2,973,486 and 2,874,556 paired-end reads (301 nucleotides [nt]) for virus from AGS cells and PIV5-AGS-infected A549 cells, respectively, were aligned initially to the PIV5-W3 genome sequence (accession no. JQ743318) (6) using Bowtie2 version 2.2.8 (7), and the alignments were visualized using Tablet version 1.13.08.05 (8). Final corrections to the consensus sequences were then made on the basis of fresh alignments. In each case, examination of the RNA strands from which reads were generated revealed that genomic RNA had copurified with viral mRNA, presumably as a result of intermolecular hybridization, and thus it was possible to determine the genome sequences (15,346 nt). The percentages of the aligned reads were 3.11 and 1.94 for virus from AGS cells and PIV5-AGS-infected A549 cells, respectively. The consensus sequences were almost identical, differing only in the nucleotides at positions 6910 and 7445, which were polymorphic, consisting of mixtures of G and A residues in both genomes. This observation demonstrated that the virus in persistently infected cells is not defective; the possibility of defectiveness had been suggested pre-

viously from the observation that AGS cells do not exhibit viral cytopathology (4).

As in several other PIV5 strains (6), the SH gene is unlikely to be functional in PIV5-AGS, because it contains a 1-nt deletion. This deletion is compensated by a 1-nt insertion in the noncoding region between the HN and L genes, the genome thereby conforming to the rule of six (9). Phylogenetic analysis of the PIV5-AGS sequence in comparison with the sequences of 17 other strains available from GenBank showed that PIV5-AGS is most closely related to human strains. Although this finding does not explain the origin of PIV5 in the AGS cell line, it suggests that infection from a human was involved, either because virus was present in the biopsy material or because the cell line was contaminated in the laboratory during its generation.

**Nucleotide sequence accession number.** The PIV5-AGS genome sequence present in persistently infected AGS cells has been deposited in GenBank under the accession no. [KX060176](https://www.ncbi.nlm.nih.gov/nuclot/KX060176).

## ACKNOWLEDGMENTS

The University of St. Andrews and the University of Glasgow are charities registered in Scotland (SC013532 and SC004401, respectively). This study was funded by grants from the Wellcome Trust (101788/Z/13/Z) to Richard E. Randall, the Wellcome Trust (101792/Z/13/Z) to Steve Goodbourn, and the Medical Research Council (MRC) (G0801822) to the MRC – University of Glasgow Centre for Virus Research.

## FUNDING INFORMATION

This work, including the efforts of Richard E. Randall, was funded by Wellcome Trust (101788/Z/13/Z). This work, including the efforts of Steve Goodbourn, was funded by Wellcome Trust (101792/Z/13/Z).

## REFERENCES

- Wang L-F, Collins PL, Fouchier RAM, Kurath G, Lamb RA, Randall RE, Rima BK. 2012. Family *Paramyxoviridae*, p 672–685. In King AMQ, Adams MJ, Carstens EB, Lefkowitz EJ (ed), *Virus taxonomy*, ninth report of the International Committee on Taxonomy of Viruses. Elsevier Academic, London, United Kingdom.
- Parks GD, Manuse MJ, Johnson JB. 2011. The parainfluenza virus simian

- virus. In Samal SK (ed), The biology of paramyxoviruses, vol 5. Caister Academic Press, Norfolk, United Kingdom.
3. Chatziandreou N, Stock N, Young D, Andrejeva J, Hagmaier K, McGeoch DJ, Randall RE. 2004. Relationships and host range of human, canine, simian and porcine isolates of simian virus, 5 (parainfluenza virus 5). *J Gen Virol* 85:3007–3016.
  4. Young DF, Carlos TS, Hagmaier K, Fan L, Randall RE. 2007. AGS and other tissue culture cells can unknowingly be persistently infected with PIV5; a virus that blocks interferon signalling by degrading STAT1. *Virology* 365:238–240. <http://dx.doi.org/10.1016/j.virol.2007.03.061>.
  5. Barranco SC, Townsend CM, Jr, Casartelli C, Macik BG, Burger NL, Boerwinkle WR, Gourley WK. 1983. Establishment and characterization of an *in vitro* model system for human adenocarcinoma of the stomach. *Cancer Res* 43:1703–1709.
  6. Rima BK, Gatherer D, Young DF, Norsted H, Randall RE, Davison AJ. 2014. Stability of the parainfluenza virus 5 genome revealed by deep sequencing of strains isolated from different hosts and following passage in cell culture. *J Virol* 88:3826–3836. <http://dx.doi.org/10.1128/JVI.03351-13>.
  7. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* 9:357–359. <http://dx.doi.org/10.1038/nmeth.1923>.
  8. Milne I, Bayer M, Cardle L, Shaw P, Stephen G, Wright F, Marshall D. 2010. Tablet—next generation sequence assembly visualization. *Bioinformatics* 26:401–402. <http://dx.doi.org/10.1093/bioinformatics/btp666>.
  9. Calain P, Roux L. 1993. The rule of six, a basic feature for efficient replication of Sendai virus defective interfering RNA. *J Virol* 67:4822–4830.