



Costantini, D., Smith, S., Killen, S. S., Nielsen, J., and Steffensen, J. F. (2017) The Greenland shark: A new challenge for the oxidative stress theory of ageing? *Comparative Biochemistry and Physiology. Part A: Molecular and Integrative Physiology*, 203, pp. 227-232.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/130329/>

Deposited on: 31 March 2017

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

1 **The Greenland shark: a new challenge**
2 **for the oxidative stress theory of ageing?**

3
4
5 David Costantini^{a,b,c*}, Shona Smith^b, Shaun S. Killen^b,
6 Julius Nielsen^d, John F. Steffensen^d

7
8
9 ^a Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk,
10 Belgium

11 ^b Institute for Biodiversity, Animal Health and Comparative Medicine, University of
12 Glasgow, Glasgow G12 8QQ, UK

13 ^c Current Address: Department of Evolutionary Ecology, Leibniz Institute for Zoo and
14 Wildlife Research, Alfred-Kowalke-Str. 17, 10315 Berlin, Germany

15 ^d Marine Biological Section, Department of Biology, University of Copenhagen,
16 Strandpromenaden 5, DK-3000 Helsingør, Denmark

17
18
19 * email: davidcostantini@libero.it
20

21 ABSTRACT

22 The free radical theory of ageing predicts that long-lived species should be more
23 resistant to oxidative damage than short-lived species. Although many studies support
24 this theory, recent studies found notable exceptions that challenge the generality of this
25 theory. In this study, we have analysed the oxidative status of the Greenland shark
26 (*Somniosus microcephalus*), which has recently been found as the longest living
27 vertebrate animal known to science with a lifespan of at least 272 years. As compared to
28 other species, the Greenland shark had body mass-corrected values of muscle
29 glutathione peroxidase and red blood cells protein carbonyls (metric of protein oxidative
30 damage) above 75 percentile and below 25 percentile, respectively. None of the
31 biochemical metrics of oxidative status measured in either skeletal muscle or red blood
32 cells were correlated with maximum lifespan of species. We propose that the values of
33 metrics of oxidative status we measured might be linked to ecological features (e.g.,
34 adaptation to cold waters and deep dives) of this shark species rather to its lifespan.

35

36 *Keywords:* ageing, lifespan, oxidative stress, sharks, vertebrates

37

38 **1. Introduction**

39 Some species live for hundreds of years; the lifetime of some other species is limited to
40 only a few hours. This surprising variation in longevity has intrigued biologists for
41 decades. A key issue that has generated an overwhelming body of research is the
42 underlying cause of senescence and of this dramatic variation in maximum lifespan. In
43 1956, Denham Harman suggested that the production of free radicals might be
44 responsible for cell senescence because free radicals can damage vital biomolecules like
45 proteins, lipids and nucleic acids, causing cell senescence. The free radical theory of
46 ageing has since been refined and a number of iterations have been presented, including
47 the mitochondrial theory of ageing (Harman, 1972) and the oxidative stress hypothesis
48 of ageing (Yu and Yang, 1996).

49 One commonly used method of unraveling the factors linked to species
50 longevity is to compare a given biochemical metric of oxidative status across species
51 exhibiting variation in maximum lifespan. In this context, many studies concluded that
52 long-lived species are more resistant to oxidative damage (e.g., cell composition with
53 biomolecules less sensitive to oxidation) than short-lived species and that long-lived
54 species have lower antioxidant levels than short-lived species because they produce less
55 free radicals (Hulbert et al., 2007; Pamplona and Costantini, 2011). As with all
56 apparently general patterns in biology, there are important exceptions that challenge the
57 generality of the theory. Contrary to the predictions of the free radical theory of ageing,
58 naked mole rats (*Heterocephalus glaber*), a long-living rodent, do not have
59 exceptionally high levels of antioxidant protection (Andziak et al., 2005) and have
60 elevated damage to DNA, lipids and proteins (Andziak et al., 2006) as compared to
61 similar-sized mice. Another example is the olm (*Proteus anguinus*), a small cave

62 salamander that has evolved an extreme life history strategy with a predicted maximum
63 lifespan of over 100 years. Surprisingly, neither its basal metabolism nor oxidative
64 damage and activities of antioxidant enzymes explain why this species sits as an outlier
65 in the relationship between amphibian size and longevity (Issartel et al., 2009; Voituron
66 et al., 2011). These and other experimental findings led some authors to challenge the
67 oxidative stress theory of ageing (Speakman and Selman, 2011).

68 The Greenland shark (*Somniosus microcephalus*) is one of the world's largest
69 predatory sharks (maximum body length > 5 meters), mainly distributed in Arctic and
70 subarctic regions of the North Atlantic (Bigelow and Schroeder, 1948; Nielsen et al.,
71 2014). Surprisingly little is known about the general biology of Greenland shark
72 (Hansen 1952, 1963), but a recent study based on radiocarbon dating of the sharks eye
73 lens found lifespan of at least 272 years and that it has a very long phase before sexual
74 maturity (around 130 years until maturity for females, Nielsen et al., 2016), which
75 makes the Greenland shark the longest living vertebrate animal known to science. The
76 Greenland shark is clearly suitable for evaluating whether its oxidative status explains
77 its lifespan or is rather a consequence of its ecology (e.g., exposure to repeated episodes
78 of hypoxia associated with deep water dives). We have therefore measured two metrics
79 of oxidative status in both blood and muscle of Greenland sharks and compared their
80 levels to those of species with different maximum lifespans.

81

82 **2. Material and methods**

83 *2.1. Tissue collection from sharks*

84 Samples from non-sexually mature Greenland sharks (n = 11 individuals, 2 males and 9
85 females; Table 1) were collected in 2012 in Ammassalik Fjord, south-eastern Greenland

86 from the Danish research vessel Dana. Sharks were caught on bottom long lines at
87 depths between 300 and 500 m. Immediately after capture, a sample of blood (ca. 100
88 ml) was collected, centrifuged for 3 minutes at 10,000 rpm and stored at -18°C while on
89 the field. Samples of skeletal muscle (white muscle) were collected from the dorsal side
90 above the gills and stored at -18°C while on the field. All samples were then stored at -
91 80°C. Body mass of each individual shark was measured on a digital scale except for
92 four sharks. For these sharks body mass was calculated according to Nielsen et al.
93 (2014). Animals were either released with tags or euthanized for research purposes.
94 Sampling of sharks was carried out in accordance with laws and regulations and with
95 authorization from the Government of Greenland (Ministry of Fisheries, Hunting &
96 Agriculture, document number 935119).

97

98 *2.2. Laboratory analyses*

99 Samples of skeletal muscle were homogenised in ice cold PBS (supplemented with 20%
100 (v/v) glycerol and 0.2 mM phenylmethylsulfonyl fluoride as an inhibitor of proteases)
101 using a pestle. Samples were then sonicated for 10 minutes and then centrifuged for 10
102 minutes at 10,000 rpm. The supernatant was split into different aliquots, which were
103 stored at -80 °C for later analyses of protein carbonyls (metric of protein oxidative
104 damage) and activity of glutathione peroxidase, two metrics that were found to be
105 linked to species lifespan (Pamplona and Costantini, 2011; Halliwell and Gutteridge,
106 2015). Protein carbonyls (PCs) were measured according to Levine et al. (1990).
107 Carbonyls (C=O) are introduced into proteins from free radicals or via reactions with
108 lipid peroxidation products (malondialdehyde and hydroxynonenal) or carbohydrates;
109 protein carbonylation is mostly irreversible (Halliwell and Gutteridge, 2015). Nucleic

110 acids were removed by adding 1 volume of a 10% solution of streptomycin sulfonate to
111 9 volumes of sample. PCs were then derivatised to 2,4-dinitrophenylhydrazone by
112 reaction with 2,4-dinitro-phenylhydrazine (DNPH). The Ransel assay (RANDOX
113 Laboratories, Crumlin, UK) was used to quantify the activity of glutathione peroxidase
114 (GPX). Glutathione peroxidase uses the reduced form of glutathione to reduce
115 peroxides and hydroperoxides to water and alcohols, respectively (Halliwell and
116 Gutteridge, 2007). The Ransel assay is based on the method of Paglia and Valentine
117 (1967) using cumene hydroperoxide as a substrate. Measures of protein carbonyls and
118 GPX were standardised by expressing the concentrations per mg of proteins as
119 measured by the Bradford protein assay (Bio-Rad Laboratories, Hercules, USA) using a
120 standard curve of bovine serum albumin. Hence, protein carbonyls were expressed as
121 nmol/mg proteins, while GPX was expressed as Units/mg proteins.

122

123 *2.3. Data collection from literature*

124 A literature search for studies in which protein carbonyls and glutathione peroxidase
125 (GPX) were analysed using our same protocols, respectively, was conducted on both
126 Web of Science and Scopus using a combination of keywords (“protein carbonyls”,
127 “GPX”, “muscle”, “red blood cells”). Data are reported in Table 2.

128 Data on body mass and maximum lifespan were collected from
129 genomics.senescence.info/species/ or fishbase.org/ when not reported in the articles
130 selected for this study. Longevity of *Hypophthalmichthys molitrix* and of *Hoplias*
131 *malabaricus* were obtained, respectively, from
132 animaldiversity.ummz.umich.edu/accounts/Hypophthalmichthys_molitrix/ and Novaes
133 and Carvalho (2011).

134

135 *2.4. Statistical analyses*

136 All statistical analyses were performed using SPSS version 22 (Chicago, USA). Values
137 of biochemical metrics, body mass and maximum lifespan (Table 2) were transformed
138 as $\log(x)$ or $\log(x + 1)$ before analyses in order to improve normality of distribution.
139 Similar results were obtained including untransformed variables in all models (data not
140 shown). To describe the distribution of PCs and GPX across species while controlling
141 for multiple sources of variation, we extracted residuals of each biochemical measure
142 (included in the models as dependent variables) from Linear Mixed Models (LMMs)
143 including ‘study’ as a random factor (because some articles analysed multiple species
144 for muscle PCs and GPX and for red blood cell GPX) and two independent fixed
145 factors: 1) origin of the animals (i.e. wild or cultured; humans were included in the
146 group of artificially selected strains); and 2) taxonomic class (fish, amphibians, reptiles,
147 birds or mammals). Mean adult body mass of each species was included as a covariate
148 to control for any allometric effects. Given that body mass data were obtained from
149 different sources, we opted to run additional LMMs where body mass was not included
150 in the models in order to control for any bias. In order to assess whether species that live
151 longer have lower production of PCs and lower levels of GPX in both skeletal muscle
152 and red blood cells, we calculated correlations between residuals of biochemical metrics
153 and residuals of maximum lifespan obtained from a linear regression of log-transformed
154 maximum lifespan onto log-transformed body mass. Given the moderate sample size,
155 we present results from Spearman test. Note that results were similar where Pearson test
156 was used. In preliminary analyses, we also used LMMs to test the covariation between a
157 given biochemical metric and maximum lifespan while controlling for potential sources

158 of variation, like taxonomic class or body mass. Given that results were similar to those
159 of bivariate correlations (data not shown), we opted to present only data from
160 correlations for easy of presentation.

161

162 **3. Results**

163 Compared to other species included in this analysis, the Greenland shark has
164 particularly high values of PCs and GPX in muscle (both above 75 percentile; Table 3).
165 When among species differences in body mass are corrected for, compared to other
166 species, the Greenland shark has high muscle GPX (above 75 percentile) and low values
167 of red blood cell PCs (below 25 percentile; Table 3). In contrast, values of muscle PCs
168 now sit in the middle of the whole variation of our sample. Activity of GPX in red
169 blood cells sits in the middle of the whole variation of our sample irrespective of
170 whether body mass is controlled for or is not.

171 None of the biochemical metrics of oxidative status were related to species
172 maximum lifespan (Spearman correlation coefficients: range -0.18 to 0.18, $p \geq 0.30$),
173 regardless of whether body mass was or was not included in the models (Fig. 1).

174

175 **4. Discussion**

176 Results of our comparative study indicated that the oxidative status of Greenland shark
177 appears to differ from that of other species. This is particularly evident from body mass-
178 corrected values of muscle GPX and red blood cells PCs, which were above 75
179 percentile and below 25 percentile, respectively. These characteristics of the Greenland
180 shark do not seem, however, to be the reason for its long lifespan as compared to other
181 species. In fact, none of the metrics of oxidative status in either skeletal muscle or red

182 blood cells were correlated with species maximum lifespan. A question then is why
183 Greenland sharks have high muscle GPX and low red blood cell PCs. The Greenland
184 shark inhabits cold waters, mostly in the range -1.8 to 10°C (Stokesbury et al., 2005).
185 Moreover, data from 25 years of surveys show that Greenland sharks are usually caught
186 at 400–700 m, but were found at all depths between 100 and 1,200 m (Nielsen et al.,
187 2014). Hence, the oxidative status of the Greenland shark might have been shaped to
188 some degree by adaptation to a life in the cold Arctic waters and to deep dives. For
189 example, many species that are faced with repeated episodes of hypoxia/reoxygenation
190 (as typical of diving animals) tend to have high basal levels of antioxidant defences in
191 order to protect them against high free radical production that occurs during
192 reoxygenation (Costantini, 2014). This might be one reason that explains why
193 Greenland sharks have high GPX (an antioxidant enzyme) as compared to the other
194 species included in this study.

195 Estimates of maximum lifespan include both the pre- and post-reproductive
196 phases. Hence, maximum lifespan of long-lived species may be inflated with the
197 addition of a period of life over which individuals are under a waning force of selection
198 (Reznick et al., 2006). Conversely, evolution would have shaped antioxidant
199 mechanisms to impact on the organism only in the period of life where selection is
200 stronger. This is particularly relevant for the Greenland shark. It has in fact a very long
201 phase before sexual maturity (Nielsen et al., 2016) during which selection would be
202 expected to be particularly long and sustained. Hence, individuals poorly protected
203 against oxidative stress might be rapidly eliminated by natural selection. All Greenland
204 shark individuals included in this study were not sexually mature. However, growth
205 does not appear to be a reason for the high GPX because sharks have an indeterminate

206 growth, i.e., they keep growing all their life long. Also many studies found juveniles to
207 have lower GPX than adults (Costantini, 2014). Further studies will clearly be needed to
208 unravel the reasons for the high activity of GPX in skeletal muscle and the low level of
209 protein damage in red blood cells. Although both protein carbonyls and GPX were
210 previously found to be linked to species lifespan (Pamplona and Costantini, 2011;
211 Halliwell and Gutteridge, 2015), it will be important to collect data on other metrics of
212 oxidative status (e.g., DNA damage and repair) that might reveal routes through which
213 Greenland sharks pave the way to their exceptional longevity. Species-based approaches
214 may, however, hide within-species variation in the association between oxidative status
215 and longevity that has been found in several vertebrate species (Costantini, 2014). Thus,
216 a proper understanding of the oxidative stress theory of ageing will require much further
217 study, in particular a combination of comparative species- and individual-based
218 approaches tailored to elucidate the association between oxidative status and longevity,
219 and whether the nature of the association is species-specific rather than general.

220 In conclusion, our study showed that levels of GPX and oxidative damage to
221 proteins (protein carbonyls) in muscle and red blood cells were not associated with
222 species lifespan, suggesting that these two metrics of oxidative status do not explain the
223 exceptional longevity of the Greenland shark. The high and low values of muscle GPX
224 and red blood cells PCs, respectively, we found in the Greenland shark as compared to
225 other vertebrate species might be explained by other ecological features of this shark
226 species, such as the adaptation to cold waters and deep dives.

227

228 **Competing interests**

229 The authors declare no competing financial interests.

230

231 **Funding**

232 Support from the Danish Center for Marine Research financed the research cruise with
233 R/V Dana to Ammasalik, Greenland in 2012. Additional support from Save Our Seas
234 Foundation, the Carlsberg Foundation and the Danish Research Council is gratefully
235 acknowledged.

236

237 **Acknowledgements**

238 We thank two anonymous reviewers for providing comments that helped us in the
239 interpretation and presentation of results.

240

241 **References**

242 Abele, D., Roecken, D., Graeve, M., Buck, B.H., 2007. Body growth, mitochondrial
243 enzymatic capacities and aspects of the antioxidant system and redox balance under
244 calorie restriction in young turbot (*Scophthalmus maximus*, L.) *Aquacult. Res.*, 38,
245 467-477.

246 Andziak, B., O'Connor, T.P., Buffenstein, R., 2005. Antioxidants do not explain the
247 disparate longevity between mice and the longest-living rodent, the naked mole-rat.
248 *Mech. Ageing Dev.* 126, 1206–1212.

249 Andziak, B., O'Connor, T.P., Qi, W.B., Dewaal, E.M., Pierce, E.M., Chaudhuri, A.R.,
250 Van Remmen, H., Buffenstein, R., 2006. High oxidative damage levels in the
251 longest-living rodent, the naked mole-rat. *Aging Cell* 5, 463–71.

252 Bigelow, H., Schroeder, W., 1948. Sharks. In *Fishes of the Western North Atlantic, Part*
253 *1* (Tee-Van, J., Breder, C. M., Hildebrand, S. F., Parr, A. E. & Schroeder, W. C.,
254 eds), pp. 59–546 Yale, CT: Yale University, Sears Foundation for Marine Research.

255 Costantini, D., 2014. Oxidative stress and hormesis in evolutionary ecology and
256 physiology. Springer-Verlag, Berlin Heidelberg.

257 Espín Luján, S., 2013. Biomonitoring of persistent environmental pollutants using
258 feathers and assessment of sublethal effects using oxidative stress biomarkers in
259 wildbirds. PhD thesis, University of Murcia, Spain.

260 Feidantsis, K., Anestis, A., Michaelidis, B., 2013. Seasonal variations of anti-/apoptotic
261 and antioxidant proteins in the heart and gastrocnemius muscle of the water frog
262 *Pelophylax ridibundus*. *Cryobiology* 67, 175-183.

263 Fisk, A.T., Tittlemier, S.A., Pranscke, J.L., Norstrom, R.J., 2002. Using anthropogenic
264 contaminants and stable isotopes to assess the feeding ecology of Greenland sharks.
265 *Ecology* 83, 2162-2172.

266 García-Medina, S., Nunez-Betancourt, J.A., Garcia-Medina, A.L., Galar-Martinez, M.,
267 Neri-Cruz, N., Islas-Flores, H., Gomez-Olivan, L.M., 2013. The relationship of
268 cytotoxic and genotoxic damage with blood aluminum levels and oxidative stress
269 induced by this metal in common carp (*Cyprinus carpio*) erythrocytes. *Ecotoxicol.*
270 *Environm. Safety* 96, 191-197.

271 Grim, J.M., Simonik, E.A., Semones, M.C., Kuhn, D.E., Crockett, E.I., 2013. The
272 glutathione-dependent system of antioxidant defense is not modulated by
273 temperature acclimation in muscle tissues from striped bass, *Morone saxatilis*.
274 *Comp. Biochem. Physiol. Part A* 164, 383–390.

275 Halliwell, B.H., Gutteridge, J.M.C. 2015. *Free Radicals in Biology and Medicine*, 4th
276 edn. Oxford University Press, Oxford.

277 Hansen, P.M., 1952. Tidsskiftet Grønland volume 5: 182-188

278 Hansen, P.M., 1963. Tagging experiments with the Greenland shark (*Somniosus*
279 *microcephalus* (Block and Schneider)) in Subarea 1. International Commission for
280 the Northwest Atlantic Fisheries. Special Publication 4, 172-175.

281 Harman, D., 1956. Aging: a theory based on free radical and radiation chemistry. *J.*
282 *Gerontol. A* 11, 298–300.

283 Hulbert, A.J., Pamplona, R., Buffenstein, R., Buttemer, W.A., 2007. Life and death:
284 metabolic rate, membrane composition, and life span of animals. *Physiol. Rev.* 87,
285 1175–1213.

286 Issartel, J., Hervant, F., de Fraipont, M., Clobert, J., Voituron, Y., 2009. High anoxia
287 tolerance in the subterranean salamander *Proteus anguinus* without oxidative stress
288 nor activation of antioxidant defenses during reoxygenation. *J. Comp. Physiol. B*
289 179, 543–551.

290 Kaczor, J.J., Hall, J.E., Payne, E., Tarnopolsky, M.A., 2007. Low intensity training
291 decreases markers of oxidative stress in skeletal muscle of mdx mice. *Free Rad. Biol.*
292 *Medic.* 43, 145-154.

293 Kubrak, O.I., Rovenko, B.M., Husak, V.V., Storey, J.M., Storey, K.B., Lushchak, V.I.,
294 2012. Nickel induces hyperglycemia and glycogenolysis and affects the antioxidant
295 system in liver and white muscle of goldfish *Carassius auratus* L. *Ecotoxicol.*
296 *Environ. Safety* 80, 231–237.

297 Levine, R.L., Garland, D., Oliver, C.N., Amici, A., Climent, I., Lenz, A.G., Ahn, B.W.,
298 Shaltiel, S., Stadtman, E.R., 1990. Determination of carbonyl content in oxidatively
299 modified proteins. *Methods Enzymol.* 186, 464-478.

300 Linsak, Z., Linsak, D.T., Spiric, Z., Srebocan, E., Glad, M., Milin, C., 2013. Effects of
301 mercury on glutathione and glutathione-dependent enzymes in hares (*Lepus*
302 *europaeus* Pallas). *J. Environ. Sci. Health A* 48, 1325-1332,

303 Luqman, S., Rizvi, S.I., 2006. Protection of lipid peroxidation and carbonyl formation
304 in proteins by capsaicin in human erythrocytes subjected to oxidative stress.
305 *Phytother. Res.* 20, 303-306.

306 MacNeil, M.A., McMeans, B.C., Hussey, N.E., Vecsei, P., Svavarsson, J., Kovacs,
307 K.M., Lydersen, C., Ramsey, M., Fisk, A.T., 2012. Biology of the Greenland shark
308 *Somniosus microcephalus*. *J. Fish Biol.* 80, 991-1018.

309 Maity, S., Nag, N., Chatterjee, S., Adhikari, S., Mazumder, S., 2013. Bilirubin clearance
310 and antioxidant activities^[11] of ethanol extract of *Phyllanthus amarus* root^[12] in
311 phenylhydrazine-induced neonatal jaundice in mice. *J. Physiol. Biochem.* 69, 467–
312 476.

313 Marasco, V., Spencer, K.A., Robinson, J., Herzyk, P., Costantini, D., 2013.
314 Developmental post-natal stress can alter the effects of pre-natal stress on the adult
315 redox balance. *Gener. Comp. Endocrinol.* 191, 239–246.

316 Martinez-Haro, M., Green, A.J., Mateo, R., 2011. Effects of lead exposure on oxidative
317 stress biomarkers and plasma biochemistry in waterbirds in the field. *Environ. Res.*
318 111, 530-538.

319 Monteiro, D.A., Rantin, F.T., Kalinin, A.L., 2013. Dietary intake of inorganic mercury:
320 bioaccumulation and oxidative stress parameters in the neotropical fish *Hoplias*
321 *malabaricus*. *Ecotoxicol.* 22, 446-456.

322 Montgomery, M.K., Hulbert, A.J., Buttemer, W.A., 2011. The long life of birds: the rat-
323 pigeon comparison revisited. *PLoS ONE* 6, e24138.

324 Montgomery, M.K., Buttemer, W.A., Hulbert, A.J., 2012. Does the oxidative stress
325 theory of aging explain longevity differences in birds? II. Antioxidant systems and
326 oxidative damage. *Exp. Geront.* 47, 211-222.

327 Nielsen, J., Hedeholm, R.B., Simon, M., Steffensen, J.F., 2014. Distribution and feeding
328 ecology of the Greenland shark (*Somniosus microcephalus*) in Greenland waters.
329 *Pol. Biol.* 37, 37-46.

330 Nielsen, J., Hedeholm, R.B., Heinemeier, J., Bushnell, P.G., Christiansen, J.S., Olsen,
331 J., Bronk Ramsey, C., Brill, R.W., Simon, M., Steffensen, K.F., Steffensen, J.F.,
332 2016. Eye lens radiocarbon reveals centuries of longevity in Greenland shark
333 (*Somniosus microcephalus*). *Science* 353, 702-704.

334 Novaes, J.L., Carvalho, E.D., 2011. Population structure and stock assessment of
335 *Hoplias malabaricus* (Characiformes: Erythrinidae) caught by artisanal fishermen in
336 river-reservoir transition area in Brazil. *Rev. Biol. Trop.* 59, 71-83.

337 Paglia, D.E., Valentine, W.N., 1967. Studies on the quantitative and qualitative
338 characterization of erythrocyte glutathione peroxidase. *J. Lab. Clin. Med.* 70, 158-
339 169.

340 Pamplona, R., Costantini, D., 2011. Molecular and structural antioxidant defences
341 against oxidative stress in animals. *Am. J. Physiol.* 301, R843-R863.

342 Pansarasa, O., Bertorelli, L., Vecchiet, J., Felzani, G., Marzatico, F., 1999. Age-
343 dependent changes of antioxidant activities and markers of free radical damage in
344 human skeletal muscle. *Free Rad. Biol. Med.* 27, 617-622.

345 Passi, S., Ricci, R., Cataudella, S., Ferrante, I., De Simone, F., Rastrelli, L., 2004. Fatty
346 acid pattern, oxidation product development, and antioxidant loss in muscle tissue of
347 rainbow trout and *Dicentrarchus labrax* during growth. *J. Agric. Food. Chem.* 52,
348 2587–2592.

349 Qujeq, D., Habibinudeh, M., Daylmatoli, H., Rezvani, T., 2005. Malondialdehyde and
350 carbonyl contents in the erythrocytes of streptozotocin-induced diabetic rats. *Int. J.*
351 *Diabetes Metabol.* 13, 96-98.

352 Radi, A.A.R., Hai, D.Q., Matkovics, B., Gabrielak, T., 1985. Comparative antioxidant
353 enzyme study in freshwater fish with different types of feeding behaviour. *Comp.*
354 *Biochem. Physiol. Part C* 81, 395-399.

355 Rey, B., Halsey, L.G., Domazon, V., Rouanet, J.-L., Roussel, D., Handrich, Y., Butler,
356 P.J., Duchamp, C., 2008. Long-term fasting decreases mitochondrial avian UCP-
357 mediated oxygen consumption in hypometabolic king penguins. *Am. J. Physiol.* 295,
358 92–100.

359 Reznick, D., Bryant, M., Holmes, D., 2006. The evolution of senescence and post-
360 reproductive lifespan in guppies (*Poecilia reticulata*). *PLoS Biol.* 4, e7.

361 Sanz, A., Furne, M., Trenzado, C.E., 2012. Study of oxidative state, as a marker of
362 welfare, on gilthead sea bream, *Sparus aurata*, subjected to handling stress. *J. World*
363 *Aquacul. Soc.* 43, 707-715.

364 Smith, S., 2016. The role of oxidative stress as an underlying mechanism in life-history
365 trade-offs. PhD Thesis, University of Glasgow, Glasgow, Scotland.

366 Speakman, J.R., Selman, C., 2011. The free-radical damage theory: Accumulating
367 evidence against a simple link of oxidative stress to ageing and lifespan. *Bioessays*
368 33, 255-259.

369 Stokesbury, M.J.W., Harvey-Clark, C., Gallant, J., Block, B.A., Myers, R.A., 2005.
370 Movement and environmental preferences of Greenland sharks (*Somniosus*
371 *microcephalus*) electronically tagged in the St. Lawrence Estuary, Canada. *Mar.*
372 *Biol.* 148, 159–165.

373 Voituron, Y., Servais, S., Romestaing, C., Douki, T., Barre, H., 2006. Oxi-dative DNA
374 damage and antioxidant defenses in the European common lizard (*Lacerta vivipara*)
375 in supercooled and frozen states. *Cryobiology* 52, 74–82.

376 Voituron, Y., de Fraipont, M., Issartel, J., Guillaume, O., Clobert, J., 2011. Extreme
377 lifespan of the human fish (*Proteus anguinus*): a challenge for ageing mechanisms.
378 *Biol. Lett.* 7, 105-107.

379 Yu, B.P., Yang, R., 1996. Critical evaluation of the free radical theory of aging. A
380 proposal for the oxidative stress hypothesis. *Ann. N. Y. Acad. Sci.* 786, 1–11.

381 Williams, C.A., Kronfeld, D.S., Hess, T.M., Saker, K.E., Waldron, J.N., Crandell,
382 K.M., Hoffman, R.M., Harris, P.A., 2004. Antioxidant supplementation and
383 subsequent oxidative stress of horses during an 80-km endurance race. *J. Anim. Sci.*
384 82, 588-594.

385 Willmore, W.G., Storey, K.B., 1997. Antioxidant systems and anoxia tolerance in a
386 freshwater turtle *Trachemys scripta elegans*. *Mol. Cell. Biochem.* 170, 177–185.

387 Woods, A.K., Storey, K.B., 2005. Effects of hibernation on multicatalytic proteinase
388 complex in thirteen-lined ground squirrels, *Spermophilus tridecemlineatus*. *Mol.*
389 *Cell. Biochem.* 271, 205–213.

390

391 Table 1. Descriptive statistics of Greenland sharks included in our study. X = the
 392 biological matrix collected from a single specimen. RBCs = red blood cells; PCs =
 393 protein carbonyls (expressed as nmol/mg proteins); GPX = glutathione peroxidase
 394 (expressed as Units/mg proteins). * For these sharks body mass was calculated
 395 according to Nielsen et al. (2014). Mean total length (TL) was 327 cm (SD = 38; n =
 396 11) and mean body mass (BM) was 343 kg (SD = 123; n = 11).

397

398

Sex	TL (cm)	BM (kg)	RBC PCs	Muscle PCs	RBC GPX	Muscle GPX	RBCs	Muscle
F	346	416	7.0		0.008		×	
F	306	246	3.1	10.5	0.007	0.091	×	×
F	264	168	1.9	4.5	0.006	0.099	×	×
F	386	560	7.0	3.1	0.004	0.118	×	×
F	365	430		1.8		0.059		×
F	336	338	3.3	3.4	0.007	0.070	×	×
F	355	435*	6.9		0.013		×	
F	351	452	2.8	6.3	0.011	0.075	×	×
F	302	262*	7.8		0.007		×	
M	290	231*	3.3	9.2	0.006	0.142	×	×
M	291	234*	3.6		0.006		×	

399

400

401 Table 2. Summary information of species and articles included in the comparative
 402 work. MLS = maximum lifespan; rbc = red blood cells; PCs = protein carbonyls
 403 (expressed as nmol/mg proteins); GPX = glutathione peroxidase (expressed as Units/mg
 404 proteins).

Class	Scientific name	Origin	Body mass (g)	MLS (years)	PCs muscle	PCs rbc	GPX muscle	GPX rbc	Article
Amphibians	<i>Pelophylax ridibundus</i>	Wild	39.5	11			0.0186		Feidantsis et al., 2013
Amphibians	<i>Proteus anguinus</i>	Captivity	10.6	102			0.0041		Issartel et al., 2009
Amphibians	<i>Calotriton asper</i>	Captivity	4.3	26			0.0041		Issartel et al., 2009
Birds	<i>Taeniopygia guttata</i>	Captivity	15.1	12		7.594			Costantini, unpublished data
Birds	<i>Bubo bubo</i>	Wild	2686	68				0.6110	Espín Luján, 2013
Birds	<i>Gyps fulvus</i>	Wild	7436	41.4				0.4490	Espín Luján, 2013
Birds	<i>Coturnix japonica</i>	Captivity	235.4	6		9.608		0.0072	Marasco et al., 2013
Birds	<i>Fulica atra</i>	Wild	734	20.6				0.7240	Martinez-Haro et al., 2011
Birds	<i>Aythya ferina</i>	Wild	823	23.2				0.2640	Martinez-Haro et al., 2011
Birds	<i>Anas platyrhynchos</i>	Wild	1048	29.1				0.3080	Martinez-Haro et al., 2011
Birds	<i>Gallinula chloropus</i>	Wild	392	18.6				0.5800	Martinez-Haro et al., 2011
Birds	<i>Columba livia</i>	Captivity	417	35	0.380				Montgomery et al., 2011
Birds	<i>Melopsittacus undulatus</i>	Captivity	26	21	0.295				Montgomery et al., 2012
Birds	<i>Nymphicus hollandicus</i>	Captivity	83.2	35	0.823				Montgomery et al., 2012
Birds	<i>Coturnix japonica</i>	Captivity	235.4	6	0.884				Montgomery et al., 2012
Birds	<i>Coturnix chinensis</i>	Captivity	45.4	5	0.819				Montgomery et al., 2012
Birds	<i>Agapornis</i> sp.	Captivity	48.9	16.1	1.528				Montgomery et al., 2012
Birds	<i>Aptenodytes patagonicus</i>	Wild	14000	26			0.0081		Rey et al., 2008
Birds	<i>Cyanistes caeruleus</i>	Wild	11.08	14.6		3.824		0.1257	Smith, 2016
Fish	<i>Scophthalmus maximus</i>	Captivity	113.3	26	6.288				Abele et al., 2007
Fish	<i>Cyprinus carpio</i>	Captivity	10600	47		11.030			García-Medina et al., 2013
Fish	<i>Morone saxatilis</i>	Captivity	31350	30			0.0105		Grim et al., 2013
Fish	<i>Carassius auratus</i>	Captivity	52.5	41	3.370				Kubrak et al., 2012
Fish	<i>Hoplias malabaricus</i>	Wild	246.32	9.3	3.000				Monteiro et al., 2013
Fish	<i>Oncorhynchus mykiss</i>	Captivity	663	11	0.260				Passi et al., 2004

Fish	Dicentrarchus labrax	Wild	1217	15	0.240				Passi et al., 2004
Fish	Somniosus microcephalus	Wild	343000	272	5.540	4.675	0.0936	0.0074	Present study
Fish	Barbus barbus	Captivity	12000	15			0.0190		Radi et al., 1985
Fish	Cyprinus carpio	Captivity	10600	47			0.0050		Radi et al., 1985
Fish	Carassius carassius	Captivity	3000	10			0.0700		Radi et al., 1985
Fish	Ctenopharyngodon idella	Captivity	1450	21			0.0019		Radi et al., 1985
Fish	Hypophthalmichthys molitrix	Captivity	450	20			0.0250		Radi et al., 1985
Fish	Sparus aurata	Captivity	9460	11			0.0023		Sanz et al., 2012
Mammals	Mus musculus	Captivity	20.5	4	0.995				Kaczor et al., 2007
Mammals	Lepus europaeus	Wild	4175	10.7			0.2100		Linsak et al., 2013
Mammals	Homo sapiens	Captivity	62000	122.5		4.670			Luqman and Rizvi, 2006
Mammals	Mus musculus	Captivity	17.6	4		3.150			Maity et al., 2013
Mammals	Rattus norvegicus	Captivity	343	3.8	0.696				Montgomery et al., 2011
Mammals	Homo sapiens	Captivity	62000	122.5	3.444				Pansarasa et al., 1999
Mammals	Rattus norvegicus	Captivity	240	3.8		1.100			Qujeq et al., 2005
Mammals	Equus caballus	Captivity	421000	57				0.0469	Williams et al., 2004
Mammals	Spermophilus tridecemlineatus	Captivity	230	7.9	0.567				Woods and Storey, 2005
Reptiles	Lacerta vivipara	Wild	2.94	11			0.0006		Voituron et al., 2006
Reptiles	Trachemys scripta	Captivity	240	41.3			0.0130		Willmore and Storey, 1997

405

406

407 Table 3. Descriptive statistics of biochemical metrics of oxidative status included in this
 408 study. Values of Greenland shark (GS) are also reported separately for comparison with
 409 the whole variation of our sample. N = number of vertebrate species data were collected
 410 from; SD = standard deviation; rbc = red blood cells; PCs = protein carbonyls
 411 (expressed as nmol/mg proteins); GPX = glutathione peroxidase (expressed as Units/mg
 412 proteins); res = residuals; NoBM = no correction for body mass; BMcorr = correction
 413 for body mass.

414

415

	PC s mu scl	PCs rbc	GPX muscle	GPX rbc	resPC muscle NoBM	resPC muscle BMcorr	resPCr bcs NoBM	resPCr bcs BMcorr	resGP Xmuscle NoBM	resGP Xmuscle BMcorr	resGP Xrbc NoBM	resGP Xrbc BMcorr
N	16	8	15	10	16	16	8	8	15	15	10	10
Mean	1.8	5.7	0.032	0.312	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
SD	1.9	3.4	0.056	0.267	0.058	0.054	0.121	0.107	0.009	0.008	0.054	0.044
Min	0.2	1.1	0.001	0.007	-0.149	-0.118	-0.242	-0.221	-0.010	-0.010	-0.102	-0.056
Max	6.3	11.0	0.210	0.724	0.141	0.152	0.189	0.131	0.019	0.021	0.083	0.081
Perc. 25	0.4	3.3	0.004	0.037	-0.019	-0.018	-0.040	-0.039	-0.008	-0.006	-0.040	-0.039
Perc. 50	0.9	4.7	0.011	0.286	-0.001	0.000	0.009	0.018	0.000	0.000	0.000	0.000
Perc. 75	3.3	9.1	0.025	0.588	0.027	0.014	0.050	0.078	0.003	0.004	0.047	0.035
GS	5.5	4.7	0.094	0.007	0.031	0.007	-0.009	-0.040	0.015	0.010	0.000	0.000

416

417


418

419

420 **Figure captions**

421 Figure 1. Scatterplots of the residuals of maximum lifespan (corrected for body mass in
422 all panels) against those of metrics of oxidative status (see text). None of the
423 biochemical metrics of oxidative status were significantly correlated to maximum
424 lifespan, regardless of whether body mass was or was not included in the models. The
425 Greenland shark is shown as a blue dot in each graph.

426



427

428

Figure 1