

Small RNAs and extracellular vesicles in filarial nematodes: From nematode development to diagnostics

J. F. Quintana¹ | S. A. Babayan² | A. H. Buck¹

¹Institute of Immunology and Infection Research and Centre for Immunity, Infection & Evolution, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

²Institute of Biodiversity Animal Health and Comparative Medicine, University of Glasgow, Glasgow, UK

Correspondence

Amy H. Buck, Ashworth Laboratories, Edinburgh, UK.
Email: a.buck@ed.ac.uk

Funding information

WT Pathfinder Award, Grant/Award Number: 201083/Z/16/Z; Institute for Medical Microbiology, Immunology and Parasitology, University Hospital of Bonn, Germany; Bill and Melinda Gates Foundation

Summary

Parasitic nematodes have evolved sophisticated mechanisms to communicate with their hosts in order to survive and successfully establish an infection. The transfer of RNA within extracellular vesicles (EVs) has recently been described as a mechanism that could contribute to this communication in filarial nematodes. It has been shown that these EVs are loaded with several types of RNAs, including microRNAs, leading to the hypothesis that parasites could actively use these molecules to manipulate host gene expression and to the exciting prospect that these pathways could result in new diagnostic and therapeutic strategies. Here, we review the literature on the diverse RNAi pathways that operate in nematodes and more specifically our current knowledge of extracellular RNA (exRNA) and EVs derived from filarial nematodes in vitro and within their hosts. We further detail some of the issues and questions related to the capacity of RNA-mediated communication to function in parasite–host interactions and the ability of exRNA to enable us to distinguish and detect different nematode parasites in their hosts.

KEYWORDS

diagnostics, exosomes, extracellular vesicles, filarial nematodes, host–pathogen, microRNAs

1 | INTRODUCTION

Filarial nematodes, the causative agents of some of the most prevalent poverty-related diseases, are tissue-dwelling nematodes that are transmitted by blood-feeding arthropods to terrestrial vertebrate hosts, from amphibians to mammals. For those nematodes infecting humans, their distribution is confined to tropical and subtropical regions, therefore representing a matter of public health in developing countries.¹ Latest estimations of the World Health Organization suggest that over 120 million people are infected by filarial parasites, causing considerable morbidity despite long-term chemotherapy-based control programmes.^{1,2} The clinical manifestations (e.g. lymphoedema, hypertrophy of the skin and blindness) are rarely associated with high mortality rates, but their chronicity and morbidity impose a tremendous socio-economic burden on these countries.

From a host–pathogen standpoint, filarial nematodes are fascinating organisms for their ability to persist in their hosts for long periods, surviving and reproducing for over a decade in some cases.³ This can be attributed to a repertoire of adaptations and strategies that the parasites employ, including secretion of factors with immunomodulatory properties.^{4,5} Furthermore, the complex life cycles and ecological interactions of parasitic nematodes make them an interesting object of study with regard to moulting, growth and survival in challenging environments, such as those encountered upon infection of the definitive host. However, many mechanistic and molecular aspects associated with the biology of these parasitic nematodes have not been fully elucidated. RNA interference (RNAi) pathways have been shown to play important roles in the free-living nematode *Caenorhabditis elegans*, including regulation of developmental timing, genome defence and adaptation to the environment.^{6–8} Here, we describe the current understanding of how different RNAi pathways operate in filarial

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



nematodes, making use of comparisons with studies in *C. elegans* in which many mechanistic aspects of RNAi were discovered.^{9,10}

In the last 8 years, it has been shown that the small RNAs involved in RNAi within cells are also found extracellularly. Their association with extracellular vesicles (EVs) in parasite infections may implicate them as novel players in the transmission of information between the parasites and their hosts.¹¹ We will describe recent evidence of extracellular RNAs derived from filarial nematodes, their potential use for diagnostics and current challenges and outstanding questions in the field.

2 | RNAI PATHWAYS IN FILARIAL NEMATODES

Three primary RNAi pathways have been characterized in animals: the microRNA (miRNA) pathway, the endo/exo-small interfering RNA (endo/exo-siRNA) pathway and the P-element-induced wimpy testis (PIWI)-interacting RNA (piRNA) pathway.^{9,12} These pathways are distinguished by the origin and identity of the small RNA guide and target, as well as the properties of the Argonaute (AGO) protein to which they bind. In general, AGOs have two main functions: (1) recognizing and binding small RNA and (2) mediating the interaction with other proteins required for small RNA loading, association with targeted RNAs, gene silencing activity and/or subcellular localization.^{8,13} From a structural standpoint, they are generally ~90–100-kDa monomeric proteins containing several domains a PAZ domain involved in 3'-end recognition and binding of the small RNA, a MID domain that binds the 5' end of the small RNA and a PIWI domain, that in some cases includes an RNaseH-like activity that can carry out endonucleolytic cleavage ("slicing") of the targets.^{13,14} The ancestral AGOs that bind to miRNAs are called ALG1 and ALG2 (AGO-like gene). The piRNA pathway is thought primarily to operate in genome defence through targeting transposable elements, mediated by the PIWI clade of AGOs. Homologs to these proteins are not present in clade III nematodes; the phylogenetic classification proposed by Blaxter et al.¹⁵ is used throughout this manuscript. Rather, it is thought instead that other AGOs and small RNA classes could be involved in genome defence in this clade.¹⁶ Indeed, a remarkable feature of nematodes is their extended AGOs (27 identified in *C. elegans*^{8,13,17}), reflecting the diversity of RNAi pathways that can operate in these animals. The majority of the AGOs in *C. elegans* belong to the WAGO (worm-specific AGO) clade, and many members of this clade are expected to be found in filarial nematodes.^{8,13} From studies in *C. elegans*, the WAGOs are thought to bind to a class of secondary siRNAs that can act through a range of mechanisms including chromosome segregation and epigenetic modifications^{18–20} and can mediate transgenerational inheritance.²¹ A more extensive description of different structural, functional and mechanistic aspects of AGO proteins is provided in recent reviews.^{8,13,22}

3 | BIOGENESIS OF MIRNAS

The microRNA pathway is one of the best characterized RNAi pathways in nematodes.²³ These molecules, first described in *C. elegans*

over two decades ago, are encoded within the genome as stem-loop structures that undergo a series of maturation events to produce the short RNA guide. In nematodes, as in other animals, miRNAs can either derive from within intragenic sequences (generally within the introns) or from independent, intergenic transcriptional units.²⁴ These transcripts, termed the primary miRNAs (pri-miRNAs), are mostly derived from the activity of RNA polymerase II (Figure 1). Some miRNAs are clustered together in discrete genomic regions suggesting coordinated expression.¹⁰


Once transcribed, miRNA biogenesis involves a series of maturation events starting with cleavage by the microprocessor complex in the nucleus.^{10,25} The microprocessor is composed of the RNase III endonuclease DROSHA and DCRG8, among other scaffold proteins, and cleaves the pri-miRNA to produce a shorter hairpin (pre-miRNA) with a 5' monophosphate and a ~2-nt overhang at the 3' end (Figure 1). The pre-miRNA is then actively transported to the cytoplasm by Ran-GTP protein and members of the exportin family (predominantly EXP-5). Once in the cytoplasm, the pre-miRNA is recognized by a second RNase III endonuclease called DICER that catalyses cleavage of the hairpin to produce a double-stranded duplex approximately 22 nt in length, where both 3' ends display a ~2-nt overhang.^{10,25} One strand of this miRNA duplex is then incorporated into the RNA-induced silencing complex (RISC) through association with the AGO protein (Figure 1). The miRNA then guides RISC to target messenger RNAs to elicit inhibition of translation, accelerated mRNA de-adenylation and/or endonucleolytic cleavage of the mRNA, depending on the degree of complementarity between the miRNA and its target.^{10,25} In animals, miRNAs generally are not perfectly complementary to their targets and recognition is dominated by the "seed" site defined as nucleotides 2–7 in the 5' end of the miRNA.

4 | MIRNA DISCOVERY AND EVOLUTION IN FILARIAL NEMATODES

A number of studies have now documented miRNAs in filarial nematodes as well as the related clade III nematodes *Ascaris suum* and *Ascaris lumbricoides*^{26–28} (Table 1). Poole et al.²⁹ first reported miRNAs in the filarial nematode *Brugia malayi* (Table 1) using bioinformatic predications as well as classical cloning from mixed life stages: adult males, gravid adult females and microfilariae (Mf). The authors identified 32 miRNAs including families well conserved in nematodes. A subsequent report by Winter et al.³⁰ identified miRNAs in the genome of *B. pahangi*, reporting a total of 132 miRNA loci that encode 104 unique mature sequences, including 29 of the 32 miRNAs previously discovered in *B. malayi*. Winter et al. carried out a side-by-side comparison of miRNAs sequenced from the clade V gastrointestinal parasite *Haemonchus contortus* and the clade III filarial parasite *B. pahangi* and were able to show that most of the miRNAs in each organism were not conserved in the other. Some of the newly evolved miRNAs were highly abundant and/or showed stage-specific expression.

Beyond the studies with *Brugia spp.*, miRNAs have also been identified in the dog heartworm parasite *Dirofilaria immitis*.³¹ Here, a total

FIGURE 1 Simplified schematic of biogenesis and potential export pathways of microRNAs. miRNAs are generally produced from primary miRNA transcripts that are processed by the microprocessor in the nucleus and exported to the cytoplasm where they are further processed by Dicer to produce a 22-nt duplex RNA. One strand of the duplex (the mature miRNA) is loaded onto an Argonaute (AGO) protein and guides the RISC complex to mediate control of gene expression by translational repression or accelerated miRNA decay (A). The microRNA can also be exported out of the cell, either in association with AGO or in another form (the uncertainty is depicted with a question mark), through directly fusing with components of the plasma membrane into extracellular vesicles (EVs) termed microvesicles (B), or through incorporation into the exosomal biogenesis pathway into multivesicular bodies (MVBs) (C)



of 1063 miRNA candidates were identified by sequence alignment of mixed adult libraries against the miRBase repository,³² corresponding to 808 miRNA families.³¹ While the large number of miRNAs reported here could reflect an expanded miRNA repertoire in this parasite, it also highlights the fact that different studies use different criteria for assigning a small RNA sequence as a miRNA. In the study by Winter et al.³⁰, for example, the authors used both miReap and miRDeep prediction programs, but then further filtered the results manually with the requirement that both arms of the hairpin must be present in their data sets. All of these factors, along with the depth of sequencing that is carried out, will affect the number and identity of miRNAs identified in different nematode species, in addition to the quality of the genomes available. This becomes an issue when trying to examine acquisitions and losses, as well as species specificity of miRNAs for use in diagnostic applications (detailed further below).

While it is tempting to speculate that the evolution of miRNAs in filarial nematodes relates to parasitism, it should also be noted that a study comparing miRNAs in the free-living nematode *Pristionchus pacificus* to the *Caenorhabditis* spp. (clade V nematodes) also showed that the majority of miRNAs were not conserved.³³ Likewise, another

study examining miRNAs in nematodes spanning clades I-V showed that at least 20% of *C. elegans* miRNAs were conserved. This work also demonstrated that homology inversely correlated with phylogenetic distance for both free-living and parasitic nematodes.¹⁶ Consequently, it seems likely that different miRNAs follow diverse evolutionary trajectories linked to various aspects of nematode biology in both free-living and parasitic organisms. It is still challenging to pinpoint correlations between specific behavioural and physiological adaptations and the fluidity at which miRNA families are lost or gained. Gene duplication and "arm switching" (a process that leads to a switch in the arm from which the functional mature miRNA is derived) have been proposed as common mechanisms for the evolution of miRNAs and expansion of some miRNA family members.^{30,34}

5 | FUNCTIONAL IMPLICATIONS OF STAGE-SPECIFIC EXPRESSION OF MIRNAS

A number of observations suggest that discrete miRNA subsets might be important regulators of processes in a particular life stage of filarial

**TABLE 1** miRNA identification in clade III nematodes

Clade III nematode	Life stage(s)	Method	Depth of coverage	miRNA diversity	Reference
Filarial nematodes					
<i>Brugia malayi</i>	Mixed AM, gAF & Mf	RNA 5' ligation independent protocol + RT-PCR/Capillary Sequencing	503 inserts cloned	32 miRNAs	[29]
<i>Brugia pahangi</i>	iL3s & mixed AM & gAF	Small RNA library prep kit/Illumina sequencing platform	13 million reads for <i>B. pahangi</i> iL3s/~13 million reads for <i>B. pahangi</i> mixed adults	125 precursor sequences that produce 99 mature miRNAs and 81 unique star sequences	[30]
<i>Brugia malayi</i>	AM, gAF & Mf	RNA 5' ligation-dependent protocol + RT-PCR/Illumina sequencing platform	3.5-3.7 million reads in adult stages and Mf (8.9-10.5 million reads in Mf libraries with alternative treatments)	129 precursor sequences that produce 145 mature miRNAs	[36]
<i>Dirofilaria immitis</i>	Mixed AM & gAF	RNA di-tagging + RT-PCR/Solexa sequencing platform	9.8 million reads	1063 miRNA candidates	[31]
Ascaris genus					
<i>Ascaris suum</i>	Embryonic stages & early development	RNA di-tagging approach+ RT-PCR/Illumina sequencing platform	6-46 million reads	97 miRNAs grouped into 59 <i>Ascaris</i> seed families	[28]
<i>Ascaris suum</i>	AM & gAF	RNA di-tagged approach/Solexa sequencing platform	11.7 million reads for each life stage	494 and 505 miRNA candidates in gAF and AM, respectively	[26]
<i>Ascaris suum</i> , <i>Ascaris lumbricoides</i>	gAF	Illumina small RNA library prep kit/Solexa sequencing platform	14.7 and 9.8 million reads in <i>A. lumbricoides</i> and <i>A. suum</i> libraries, respectively	494 miRNA candidates in <i>A. suum</i> ; 171 miRNA candidates in <i>A. lumbricoides</i>	[27]

iL3s = infective L3s, AM = adult male, gAF = gravid adult female, Mf = microfilariae.

nematodes. For example, many of the miRNAs identified in *B. malayi* showed stage-specific expression, including miR-2b and let-7 which are abundant in adult stages.³⁰ Interestingly, in a further study, Winter et al.³⁵ demonstrated that one particular member of the let-7 family, miR-5364, is the most upregulated miRNA in the infective L3s (iL3s) during the vector-to-host transition (~12X compared to vector-derived L3s), as soon as 24 hours post-infection. Analysis of the pre-miRNA sequence suggests that this let-7 family member is present in all clade III nematodes for which there is sequence information but is absent in other nematodes, including the clade V nematodes *C. elegans*, *Heligmosomoides polygyrus* and *H. contortus*.

It has also been reported that miR-71 was one of the most abundant miRNAs in small RNA libraries prepared from total RNA from mixed adult worms in *D. immitis*.³¹ A later study in *B. malayi* showed that miR-71 represents ~27% of the total miRNA reads identified in Mf data sets and is 3-5X more enriched in Mf than adult worms.³⁶ It is possible that some of miR-71 detected in the study with *D. immitis* could potentially originate from Mf found in gravid female worms and not from adult worms per se, although this is still unclear. A recent report demonstrated the functional activity of miR-71 in developmentally competent *B. malayi* embryos using a luciferase reporter assay, concluding that miR-71 can act as a post-transcriptional repressor of mRNA targets in this life stage.³⁷ In *C. elegans*, miR-71 regulates longevity and life span,³⁸ where it is upregulated in L1 diapause and dauer larvae but not particularly in other life stages.³⁹

It has been shown that filarial nematodes adjust their developmental schedule and fecundity in response to host-derived immunological factors.⁴⁰ This is indicative of different developmental trajectories depending upon environmental signals, a phenomenon referred to as phenotypic plasticity.⁴¹⁻⁴⁴ miRNAs, as well as other RNAi pathways, have been shown to control developmental choices and life-history traits in post-dauer *C. elegans*, which shares behavioural and physiological traits with infective L3 larvae in parasitic nematodes.^{41,43,45-47} Therefore, it is likely that the same mechanisms operate in filarial nematodes to control development and fertility in response to immunological cues from the host. A comparative analysis evaluating the RNAi landscape throughout filarial development in different environmental contexts will help to clarify whether such molecular "switches" (discrete small RNA populations) could be the drivers or modulators of such morphological and developmental choices.

6 | ENDOGENOUS SMALL INTERFERING RNA (ENDO-SIRNA) PATHWAYS IN FILARIAL NEMATODES

Most commonly, RNAi pathways in parasitic nematodes are discussed in relation to the ability to trigger an RNAi-mediated gene silencing response upon stimulation with exogenous (or environmental) double-stranded RNAs (exo-dsRNAs). This requires uptake of

TABLE 2 Extracellular filarial-derived miRNAs reported in vitro and in vivo

Clade III nematode	Host	Sample type	Depth of coverage (Total parasite-specific RNA reads)	miRNA diversity	Reference
In vitro					
<i>Brugia malayi</i> (iL3s + adult males and females)	iL3s derived from mosquito/Adult worms obtained from NIAID-NIH/FR3	Excretion/Secretion (ES) products	11 139 <i>B. malayi</i> reads in extracellular vesicles (EV's) (2% of total reads)/1 519 403 <i>B. malayi</i> reads in iL3s (50% of total reads)	52 miRNAs detected in iL3s-derived EVs	[69]
In vivo					
<i>L. sigmodontis</i>	BALB/c mice	Serum (d60p.i.–Patent infection)	1188 <i>L. sigmodontis</i> reads (1.5% of total reads)	16 <i>L. sigmodontis</i> miRNAs in mouse serum	[68]
<i>Dirofilaria immitis</i>	Dog	Plasma	>338,694 <i>D. immitis</i> reads	245 <i>D. immitis</i> miRNAs in dog plasma	[78]
<i>Onchocerca volvulus</i>	Human	Serum	>46 <i>O. volvulus</i> reads	21 <i>O. volvulus</i> miRNAs in Human serum	
<i>Loa loa</i>	Baboon	Plasma	Unknown	22 <i>L. loa</i> miRNAs in baboon plasma	[79]
<i>Onchocerca ochengi</i>	Cattle	Plasma	Unknown	10 <i>O. ochengi</i> miRNAs in bovine serum	
<i>Onchocerca ochengi</i>	Cattle	Nodular fluid	157 633 <i>O. ochengi</i> reads (1.1% of total reads)	62 <i>Onchocerca</i> miRNAs in onchocercoma fluids	[80]
<i>Onchocerca volvulus</i>	Humans	Serum/plasma	108 323 and 355 397 <i>O. volvulus</i> reads in two separate libraries (1.1 and 1.5% of total reads)	6 <i>Onchocerca</i> miRNAs in human serum/plasma	

double-stranded RNA (dsRNA), processing this into primary siRNAs, amplification involving an RNA-dependent RNA polymerase to produce the secondary siRNAs and ability to spread the signal, (reviewed in ^{48–51}). In many nematodes, the absence of the dsRNA import protein SID-1 is thought to explain the lack of efficient RNAi carried out experimentally.⁵² However, endogenous pathways are expected to exist in these organisms where siRNAs are generated by a variety of mechanisms and these can have a variety of functions.^{50,53} In *C. elegans*, two major categories of endo-siRNAs have been identified 22G-RNAs and 26G-RNAs, both displaying a strong bias for guanine at the 5' end. The endogenous 26G-RNAs are normally produced from mature mRNA transcripts by the action of the RNA-dependent RNA polymerase (RdRP) RRF-3.⁵⁴ These 26G-RNA precursors act as triggers for the production of a second class of 22G-RNAs that are synthesized de novo by RdRPs.⁵⁴ Other triggers such as piRNAs can also induce the de novo synthesis of 22G-siRNAs.^{53,54} The function of the secondary siRNAs is dictated by the association with different types of AGO proteins. These have been shown to have multiple roles in *C. elegans* including chromosome segregation,¹⁸ genome defence, surveillance and integrity,¹³ as well as transgenerational epigenetic inheritance.²¹

In *Ascaris*, it was shown that 26G-RNAs as well as 22G-RNAs were predominantly detected in the germline through to 128-cell embryos.²⁸ The majority of these endo-siRNAs mapped to a broad spectrum of coding genes in an antisense fashion.²⁸ On the other hand, a total of 40 repeat-associated siRNAs were identified in adult stages of *B. malayi*.²⁹ Similarly, several sense and antisense siRNAs were

detected in the small RNA data from iL3s and mixed adult stages in *B. pahangi*, with at least eight sequences derived from repetitive elements.³⁰ A closer examination revealed that these sequences were mainly associated with retrotransposons and mapped to nonannotated repeats. Interestingly, a phylumwide survey suggested that in clade III nematodes, the 22G-RNAs preferentially target antisense to predicted repetitive elements and have been proposed as a mechanism to control transposon activity in the absence of piRNAs.¹⁶ Beyond genome defence, it is possible that endo-siRNAs might be involved in a wide range of biological processes in nematodes, including sophisticated (and potentially novel) gene silencing mechanisms as well as epigenetic regulation. Our understanding of these phenomena will be greatly enhanced with further studies of the post-transcriptional regulatory networks of different life stages across this clade.

7 | EXTRACELLULAR VESICLES AND EXTRACELLULAR RNA IN FILARIAL NEMATODES

It is now recognized that RNA molecules can also operate beyond the limits of the cell. One key feature of extracellular RNA (exRNA) is its remarkable stability in hostile environments such as human biofluids. Several studies have demonstrated that the stabilization of exRNA can occur through direct association with protein and lipid partners such as AGO complexes or LDH particles or encapsulation within EVs,



reviewed in.⁵⁵⁻⁵⁷ EVs and exRNAs have been found in excretion/secretion (ES) products from a diverse range of parasites, from microbes to nematodes (reviewed in¹¹). This has been suggested as an active exchange of genetic material that can mediate communication between organisms of the same species, or even between evolutionarily distant organisms.^{58,59}

Most of the literature detailing exRNA in helminths focuses on their encapsulation within EVs although the origins of these are not all well documented (Figure 1). EVs that pellet upon ultracentrifugation can derive from the endocytic pathway (termed exosomes) or from budding off the plasma membrane (often termed microvesicles), and these can be difficult to distinguish by their sizes: exosomes are generally 40-100 nm and microvesicles can range from 100 to 1000 nm. These can also be difficult to distinguish based on their protein content; for example, recent research with mammalian EVs has demonstrated that proteins previously referred to as “canonical exosomal markers” (MHC I and II, flotillins, actin or heat-shock proteins 70, among others^{56,57}) can be detected in other classes of EV. The authors further showed that even within small EVs of the same density and size, there were multiple categories that could be distinguished by displaying different combinations of protein markers.⁶⁰ It seems likely that such heterogeneity exists in parasite EVs, an area that remains largely unexplored, which could be key to understanding the diversity of their functions.¹¹

Initial reports in the trematodes *Echinostoma caproni* and *Fasciola hepatica* suggested that EVs (30-100 nm) could derive from tegumental structures and could be a mechanism for transferring material to host cells.^{61,62} EVs with similar sizes have also been characterized in the human pathogenic trematodes *Schistosoma mansoni*⁶³ and *Schistosoma japonicum*,⁶⁴ the carcinogenic liver fluke *Opisthorchis viverrini*,⁶⁵ the clade V gastrointestinal nematode *Teladorsagia circumcincta*⁶⁶ and the clade I whipworm *Trichuris suis*.⁶⁷ In our own work, we showed that the clade V gastrointestinal parasitic nematode *H. polygyrus* secretes EVs that are enriched in proteins known to be abundant in the intestinal tissue of the parasite as well proteins associated with exosome biogenesis (e.g. Alix).⁶⁸ In the context of filarial infections, a recent report focusing on *B. malayi* showed that both iL3s and gravid adult females secreted EVs in vitro.⁶⁹ The EVs detected in excretion/secretion (ES) products from iL3s were described as homogeneous, based on size, ranging between 50 and 120 nm. Proteomic analysis of the iL3s revealed an enrichment for several proteins previously termed exosome markers, including HSP70 and Rab-1.⁶⁹

Nematode-derived miRNAs were identified in both *H. polygyrus* and *B. malayi* EVs with some overlap in those that were found including miR-71 and members of the let-7 and miR-100 families. Both studies in *H. polygyrus*⁶⁸ and *B. malayi*⁶⁹ showed that the secreted RNA population is distinct from the RNA isolated from the total worm. Similar observations were made when comparing the RNA from EVs and worms in *F. hepatica*.⁷⁰ While this suggests distinct miRNAs are secreted, it does not inform on whether this subset is selectively exported from the cell from which it derives. Some mechanisms for selective sorting of miRNAs into EVs have been described in mammalian systems, involving RNA-binding proteins.^{71,72} The presence of ribosomal proteins in the EVs secreted by *B. malayi* iL3s was also noted,⁶⁹

although it is not known whether these were associated with rRNA fragments that were also found. It is unclear whether or how different RNA processing pathways converge with EV biogenesis and secretion. In some systems, components of the RISC complex have been detected in EVs or shown to comigrate with endosomal MVB fractions in density gradients.⁷³ Interestingly, one AGO protein was also identified in both vesicle and vesicle-depleted fractions from *H. polygyrus* in vitro,⁶⁸ although the mechanistic aspects associated with secretion of AGO proteins in nematodes or others parasites are unknown.

8 | REGULATION AND PLASTICITY OF EV SECRETION?

The population of EVs detected in ES products seems to be variable between life stages, with reduced content observed in *B. malayi* gravid adult females compared to iL3s.⁶⁹ Interestingly, the EV release rate from iL3s was reduced by ~two fold between 24 and 72 hours (estimated as the amount of particles released by parasite over time), and this is thought to be associated with worm viability in the culture conditions tested. Indeed, one outstanding question in the field is whether exRNA and EVs could derive from dead or moribund worms. It is intriguing to think of vesicle secretion as a regulated mechanism involving specialized tissues and/or organs in the nematode, whereby release could occur in response to environmental conditions, vector-to-host transition, activation of specific receptors stimulated by host hormones, etc. However, there is very little evidence at present to support this, in part because of the youth of this field. It was proposed by Zamanian et al.⁶⁹ that the release of EVs could be a phenotype restricted to larval stages and might be involved in invasion during the onset of infection by modulating immune responses in the host. Given the precedent for immune modulation by parasite EVs,^{11,68,74} it does seem likely that their release would be subject to control, possibly by both parasite and host. It is also possible that properties of the EVs and their cargo can change throughout filarial development and in response to particular environmental challenges and/or cues. In the context of filarial parasites, for example, iL3s could release exRNA-loaded EVs aiming to ensure successful migration through an active interaction with cells at the site of the infection and/or evasion of early innate immune cells. Similarly, gravid adult females could release exRNAs involved in downregulation of immune response against Mf, thus ensuring their survival. Although exciting, the hypothesis of “plasticity” in the exRNA signals and EVs secreted by different life stages in filarial nematodes remains unstudied.

9 | EXTRACELLULAR SMALL RNAS AS BIOMARKERS FOR FILARIASIS—TOWARDS DIAGNOSTIC APPLICATIONS

One potential application of these parasite-derived exRNAs is in the area of biomarkers for helminthiasis. This is based on a key observation that parasite-derived exRNAs can be detected in biofluids from

their hosts as demonstrated by small RNA sequencing and qRT-PCR. This was first documented in schistosomiasis,⁷⁵⁻⁷⁷ but has also been examined by multiple groups in the context of filarial infections⁷⁸⁻⁸⁰ (Table 2). In an initial report, Tritten et al.⁷⁸ documented a total of 245 miRNA candidates of potential nematode origin in the plasma of dogs infected with the heartworm *D. immitis* based on sequencing. In a subsequent report, they documented a total of 22 unique sequences derived from *L. loa* in human serum and 10 sequences derived from *Onchocerca ochengi* in infected cattle serum.⁷⁹ We have also identified 62 *O. ochengi*-derived miRNAs in onchocercomata fluid⁸⁰ and found a total of 16 *Litomosoides sigmodontis*-derived miRNAs in the serum of mice during the patent stage of the infection.⁶⁸ Common to all of these studies was the presence of extracellular miR-71 and miR-100 family members. Further comparisons are challenging due to differences in the technical methods and analysis reported from the studies, which use different cut-offs for defining a candidate miRNA and are carried out at different depths of coverage. Identification of

nematode-derived small RNAs in host fluids is also challenging given the dominance of host-derived sequences in these samples, and it may be appropriate to remove all sequences that could derive from the host prior to assignment of these as parasitic in origin.

While it might be expected that all nematode parasites can or do release exRNA and EVs, there are a number of factors that will influence the ability to detect these molecules in different host fluids. It is logical that the localization of the parasite within the host dictates the presence of parasite-derived exRNAs in different biofluids. In line with this, nematode miRNAs could be identified in the serum of mice infected with *L. sigmodontis*, but not in serum from mice infected with *H. polygyrus* (which resides in the small intestine) in a side-by-side comparison.⁶⁸ The close contact between some filarial nematodes and the lymphatic system (*Wuchereria bancrofti*, *Loa loa*, *Brugia* spp.) could account for a widespread distribution of secreted parasite products into the bloodstream, such that they are readily detectable in serum and plasma (and perhaps urine) (Figure 2). On the other hand,




FIGURE 2 Proposed routes of extracellular vesicles (EV) secretion in vivo in lymphatic filariasis. Depiction of filarial nematodes (for example *Brugia* spp.) residing within a lymph node. A hypothesis is that the lack of a nodular structure (as observed in infection from *Onchocerca* spp) might facilitate the accessibility of EVs into the circulation. It is also unclear whether the detection of EVs and small RNAs is exclusively associated with viable worms or can be also derived from moribund or dead worms

the presence of a nodular structure in onchocerciasis may impose a physical barrier for the trafficking of locally secreted parasite products to the bloodstream. However, the presence of a vascularized system surrounding the onchocercomata nodule may be seen as a "window" for dissemination of such products⁸¹ (Figure 3). The mechanisms by which parasite EVs and exRNAs can bypass physical barriers (e.g. those imposed by the nodular structure in *Onchocerca* spp.) and reach the bloodstream are not well understood yet but could help inform to what extent these molecules can be effectively used as biomarkers for different filarial infections.

The potential species specificity of some miRNAs makes them attractive candidates as diagnostics where co-infection is an issue, for example in distinguishing *Onchocerca volvulus* and *L. loa* in co-endemic communities. A pan-filarial small RNA-based biomarker could also be useful as a point-of-care diagnostic test aiming to monitor populations subjected to mass drug administration (MDA) or for elimination programs. Studies conducted in biofluids from several filarial infections suggest that, for instance, miR-71 can be used as a biomarker for filarial infection.⁷⁸⁻⁸⁰ However, it is expected that several technological approaches will be considered to improve not only the platforms

currently available for exRNA detection (reviewed in^{82,83}) but also the way in which these technologies can be transferred in a field-friendly manner. Advancing inexpensive technologies and streamlined purification protocols will certainly increase the likelihood of adopting small RNA-based biomarkers in the field.

10 | FINAL CONSIDERATIONS & OUTSTANDING QUESTIONS

The field of EVs and small RNAs in parasitic nematodes is in its infancy and rapidly growing alongside efforts to exploit these in therapeutic and diagnostic applications. From a biological perspective, several outstanding questions should be addressed to drive this field forward. It is still unknown whether the secretion of exRNA-loaded EVs is developmentally regulated in parasitic nematodes, whether there are mechanisms to sort and package different small RNAs into EVs and whether all the different types of exRNAs reported so far in ES products play a role in parasite-to-host communication. If there are mechanisms in place in the parasitic nematodes




FIGURE 3 Proposed routes of extracellular vesicle (EV) secretion in vivo in Onchocerciasis. Depiction of a nodule-forming species member of the *Onchocerca* genus (e.g. *Onchocerca volvulus* or *Onchocerca ochengi*) residing within a nodular structure termed an onchocercoma. It is not yet clear whether or how the nodular structure imposes a physical barrier for dissemination of small RNA-loaded EVs into the bloodstream. As in Figure 2, it is unclear whether the detection of EVs and small RNAs is exclusively associated with viable worms or can be also derived from moribund or dead worms

to control EV secretion and dictate the cargo that is exported, then it is plausible that these mechanisms might have evolved as an additional axis of adaptation of the parasites to regulate their hosts' immune system.⁸⁴

If this is the case, several aspects need to be considered. First, if the parasites effectively use EVs and small RNAs as a mechanism for invasion, colonization and immune evasion, then one possibility is that these functions are specifically compartmentalized within the parasites. We therefore should expect that certain tissues or organs be directly involved in their production and secretion/excretion, for example those with glandular functions such as the pharynx or cells producing ES products. Building from this idea, we could also propose that the profile and exRNA content of EVs, as well as the diversity of exRNAs, will be different between life stages, as a result of their development. For example, gravid adult females could actively secrete a plethora of exRNAs and EVs, which, together with other soluble proteins in the ES products, aim to maintain a downregulated immune status in the host in order to aid in survival not only of the adults but of the Mf as well, a life stage that is particularly less complex. This interplay with the host could begin early in the infection and would be maintained throughout. The idea of "maternally mediated" Mf survival is exciting as it offers a new possibility for treatment and intervention. However, the ideas proposed so far remain speculative and will require further analysis.

Although less explored in helminth infection, exRNAs and EVs could also have functions as mediators of parasite-to-parasite cross-talk. Studies in the protozoan *Plasmodium falciparum* showed that EVs can be involved in communication between parasite populations as well as with the host.⁸⁵ Other organisms such as fungi and bacteria, which are typically found in the normal microbiota in the host, also secrete EVs. EVs may not only be involved in communication with the vertebrate host but also could be relevant for the establishment of relevant ecological interactions between pathogens that might co-exist in natural conditions, for example in situations where multiple infections occur at the same time.⁸⁶ For example, it remains possible that exRNAs and EVs could be involved in *a*) intraspecies communication, for example chemoattractant derived from female worms to increase adult male motility or fertility, maternally derived prosurvival signals to increase Mf survival or *b*) interspecies communication, for example *Wolbachia*-derived EVs that confer nutritional advantages to the filarial host,⁸⁷⁻⁸⁹ modulation of the host's immune response by filarial-derived EVs.⁶⁹

Although still far off, further work in this area could contribute to the development of novel technological applications to diagnose and control filarial infections (reviewed in⁹⁰). Basic research in this area also offers new scenarios to understand the real complexity that exists in the interaction between organisms at a molecular level. This is useful for understanding how parasitic nematodes can manipulate the gene expression machinery in their host for their own benefit, or the molecular basis of the mutualistic interactions between endosymbionts, for example *Wolbachia* and their nematode hosts. An exciting and intriguing avenue is the possibility of merging genome editing and functional genomic tools (reviewed in^{91,92}) to engineer specific EVs

cargos. These "tailored" EVs could be used as vehicles to further our understanding on how multiple organisms use these extracellular systems to transfer information and to maintain a dialogue with their surroundings. Towards this goal, further work will be required to improve the genetic manipulation toolkit currently available for filarial nematodes and to advance the basic research on EV and exRNA secretion and function in these parasites.

ACKNOWLEDGEMENTS

We thank our collaborators on filarial nematode projects for many helpful discussions and in particular Ken Pfarr, Achim Hoeruf, Ben Makepeace, Mark Blaxter, David Taylor and Coralie Martin. We also thank Ben Makepeace, Ken Pfarr and Paul Dickinson for comments on the manuscript. The studies on filarial nematode diagnostics are supported by a WT Pathfinder Award (201083/Z/16/Z) as well as previous funding from the Institute for Medical Microbiology, Immunology and Parasitology, University Hospital of Bonn, Germany, as part of a consortium grant funded by the Bill and Melinda Gates Foundation. Basic research in AB's laboratory on EVs and exRNA is supported by a WTRCDF (097394/Z/11/Z) and HFSP Young Investigator Award (RGY0069).

DISCLOSURES

None.

REFERENCES

1. Taylor MJ, Hoerauf A, Bockarie M. Lymphatic filariasis and onchocerciasis. *Lancet*. 2010;376:1175–1185.
2. Crump A, Morel CM, Omura S. The onchocerciasis chronicle: from the beginning to the end? *Trends Parasitol*. 2012;28:280–288.
3. Gems D. Longevity and ageing in parasitic and free-living nematodes. *Biogerontology*. 2000;1:289–307.
4. Maizels RM, McSorley HJ. Regulation of the host immune system by helminth parasites. *J Allergy Clin Immunol*. 2016;138:666–675.
5. Robinson MW, Donnelly S, Dalton JP. Helminth defence molecules-immunomodulators designed by parasites!. *Front Microbiol*. 2013;4:1–4.
6. Decottignies A. Endogenous RNAi and adaptation to environment in *C. elegans*. *Worm*. 2012;1:129–133.
7. Ghildiyal M, Zamore PD. Small silencing RNAs: an expanding universe. *Nat Rev Genet*. 2009;10:94–108.
8. Youngman EM, Claycomb JM. From early lessons to new frontiers: the worm as a treasure trove of small RNA biology. *Front Genet*. 2014;5:1–13.
9. Hoogstrate SW, Volkens RJ, Sterken MG, Kammenga JE, Snoek LB. Nematode endogenous small RNA pathways. *Worm*. 2014;3:e28234.
10. Kim VN, Han J, Siomi MC. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol*. 2009;10:126–139.
11. Coakley G, Maizels RM, Buck AH. Exosomes and other extracellular vesicles: the new communicators in parasite infections. *Trends Parasitol*. 2015;31:477–489.
12. Weick E-M, Miska E. piRNAs: from biogenesis to function. *Development*. 2014;141:3458–3471.
13. Buck AH, Blaxter M. Functional diversification of Argonautes in nematodes: an expanding universe. *Biochem Soc Trans*. 2013;41:881–886.



14. Hutvagner G, Simard MJ. Argonaute proteins: key players in RNA silencing. *Nat Rev Mol Cell Biol.* 2008;9:22–32.
15. Blaxter ML, De Ley P, Garey JR, et al. A molecular evolutionary framework for the phylum Nematoda. *Nature.* 1998;392:71–75.
16. Sarkies P, Selkirk ME, Jones JT, Blok V, et al. Ancient and novel small RNA pathways compensate for the loss of piRNAs in multiple independent nematode lineages. *PLoS Biol.* 2015;13:1–20.
17. Yigit E, Batista PJ, Bei Y, et al. Analysis of the *C. elegans* Argonaute family reveals that distinct Argonautes act sequentially during RNAi. *Cell.* 2006;127:747–757.
18. Wedeles CJ, Wu MZ, Claycomb JM. A multitasking Argonaute: exploring the many facets of *C. elegans* CSR-1. *Chromosome Res.* 2013;21:573–586.
19. Tu S, Wu MZ, Wang J, Cutter AD, Weng Z, Claycomb JM. Comparative functional characterization of the CSR-1 22G-RNA pathway in *Caenorhabditis* nematodes. *Nucleic Acids Res.* 2015;43:208–224.
20. Gu W, Shirayama M, Conte D, et al. Distinct Argonaute-mediated 22G-RNA pathways direct genome surveillance in the *C. elegans* germline. *Mol Cell.* 2009;36:231–244.
21. Klosin A, Lehner B. Mechanisms, timescales and principles of transgenerational epigenetic inheritance in animals. *Curr Opin Genet Dev.* 2016;36:41–49.
22. Meister G. Argonaute proteins: functional insights and emerging roles. *Nat Rev Genet.* 2013;14:447–459.
23. Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell.* 2009;136:215–233.
24. Grishok A. RNAi mechanisms in *Caenorhabditis elegans*. *FEBS Lett.* 2005;579:5932–5939.
25. Winter J, Jung S, Keller S, Gregory RI, Diederichs S. Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nat Cell Biol.* 2009;11:228–234.
26. Xu MJ, Fu JH, Nisbet AJ, et al. Comparative profiling of microRNAs in male and female adults of *Ascaris suum*. *Parasitol Res.* 2013;112:1189–1195.
27. Shao C-C, Xu M-J, Alasaad S, et al. Comparative analysis of microRNA profiles between adult *Ascaris lumbricoides* and *Ascaris suum*. *BMC Vet Res.* 2014;10:99.
28. Wang J, Czech B, Crunk A, et al. Deep small RNA sequencing from the nematode *Ascaris* reveals conservation, functional diversification, and novel developmental profiles. *Genome Res.* 2011;21:1462–1477.
29. Poole CB, Davis PJ, Jin J, McReynolds LA. Cloning and bioinformatic identification of small RNAs in the filarial nematode, *Brugia malayi*. *Mol Biochem Parasitol.* 2010;169:87–94.
30. Winter A, Weir W, Hunt M, et al. Diversity in parasitic nematode genomes: the microRNAs of *Brugia pahangi* and *Haemonchus contortus* are largely novel. *BMC Genom.* 2012;13:4.
31. Fu Y, Lan J, Wu X, et al. Identification of *Dirofilaria immitis* miRNA using illumina deep sequencing. *Vet Res.* 2013;44:1–11.
32. Kozomara A, Griffiths-Jones S. MiRBase: annotating high confidence microRNAs using deep sequencing data. *Nucleic Acids Res.* 2014;42:68–73.
33. De Wit E, Linsen SEV, Cuppen E, Berezikov E. Repertoire and evolution of miRNA genes in four divergent nematode species. *Genome Res.* 2009;19:2064–2074.
34. Griffiths-Jones S, Hui JHL, Marco A, Ronshaugen M. MicroRNA evolution by arm switching. *EMBO Rep.* 2011;12:172–177.
35. Winter AD, Gillan V, Maitland K, et al. A novel member of the let-7 microRNA family is associated with developmental transitions in filarial nematode parasites. *BMC Genom.* 2015;16:331.
36. Poole CB, Gu W, Kumar S, et al. Diversity and expression of microRNAs in the filarial parasite, *Brugia malayi*. *PLoS ONE.* 2014;9:5.
37. Liu C, Voronin D, Poole CB, et al. Functional analysis of microRNA activity in *Brugia malayi*. *Int J Parasitol.* 2015;45:579–583.
38. Boulias K, Horvitz HR. The *C. elegans* microRNA mir-71 acts in neurons to promote germline-mediated longevity through regulation of DAF-16/FOXO. *Cell Metab.* 2012;15:439–450.
39. Karp X, Hammell M, Ow MC, et al. Effect of life history on microRNA expression during *C. elegans* development. Effect of life history on microRNA expression during *C. elegans* development. *RNA.* 2011;17:639–651.
40. Babayan SA, Read AF, Lawrence RA, Bain O, Allen JE. Filarial parasites develop faster and reproduce earlier in response to host immune effectors that determine filarial life expectancy. *PLoS Biol.* 2010;8:e1000525.
41. Kochin BF, Bull JJ, Antia R. Parasite evolution and life history theory. *PLoS Biol.* 2010;8:10–13.
42. Schlichting CD. Origins of differentiation via phenotypic plasticity. *Evol Dev.* 2003;5:98–105.
43. Schlichting CD, Smith H. Phenotypic plasticity: linking molecular mechanisms with evolutionary outcomes. *Evol Ecol.* 2002;16:189–211.
44. Viney M, Cable J. Macroparasite life histories. *Curr Biol.* 2011;21:R767–R774.
45. Fielenbach N, Antebi A. *C. elegans* dauer formation and the molecular basis of plasticity. *Genes Dev.* 2008;22:2149–2165.
46. Davies SJ, McKerrow JH. Developmental plasticity in schistosomes and other helminths. *Int J Parasitol.* 2003;33:1277–1284.
47. Hall SE, Chirn G-W, Lau NC, Sengupta P. RNAi pathways contribute to developmental history-dependent phenotypic plasticity in *C. elegans*. *RNA.* 2013;19:306–319.
48. Maule AG, McVeigh P, Dalzell JJ, Atkinson L, Mousley A, Marks NJ. An eye on RNAi in nematode parasites. *Trends Parasitol.* 2011;27:505–513.
49. Piatek MJ, Werner A. Endogenous siRNAs: regulators of internal affairs. *Biochem Soc Trans.* 2014;42:1174–1179.
50. Hoogstrate S, Volkers R. Nematode endogenous small RNA pathways. *Worm.* 2014;3:e28234.
51. Britton C, Winter AD, Marks ND, et al. Application of small RNA technology for improved control of parasitic helminths. *Vet Parasitol.* 2015;212:47–53.
52. Dalzell JJ, McVeigh P, Warnock ND, et al. RNAi effector diversity in nematodes. *PLoS Negl Trop Dis.* 2011;5:e1176.
53. Sarkies P, Miska E. Small RNAs break out: the molecular cell biology of mobile small RNAs. *Nat Rev Mol Cell Biol.* 2014;15:525–535.
54. Billi AC. Endogenous RNAi pathways in *C. elegans*. *WormBook (10) WormBook* 2014;7:1–49.
55. Hoy AM, Buck AH. Extracellular small RNAs: what, where, why? *Biochem Soc Trans.* 2012;40:886–890.
56. Mittelbrunn M, Sánchez-Madrid F. Intercellular communication: diverse structures for exchange of genetic information. *Nat Rev Mol Cell Biol.* 2012;13:328–335.
57. Turchinovich A, Samatov TR, Tonevitsky AG, Burwinkel B. Circulating miRNAs: cell-cell communication function? *Front Genet.* 2013;4:1–10.
58. Sarkies P, Miska E. Is there social RNA? *Science.* 2013;341:467–468.
59. Knip M, Constantin ME, Thordal-Christensen H. Trans-kingdom cross-talk: small RNAs on the move. *PLoS Genet.* 2014;10:e1004602.
60. Kowal J, Arras G, Colombo M, et al. Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci USA.* 2016;113:E968–E977.
61. Marcilla A, Trellis M, Cortes A, et al. Extracellular vesicles from parasitic helminths contain specific excretory/secretory proteins and are internalized in intestinal host cells. *PLoS ONE.* 2012;7:e45974.
62. Cwiklinski K, de la Torre-Escudero E, Trellis M, et al. The extracellular vesicles of the helminth pathogen, *Fasciola hepatica*: biogenesis pathways and cargo molecules involved in parasite pathogenesis. *Mol Cell Proteomics.* 2015;14:3258–3273.
63. Nowacki FC, Swain MT, Klychnikov OI, et al. Protein and small non-coding RNA-enriched extracellular vesicles are released by the pathogenic blood fluke *Schistosoma mansoni*. *J Extracell Vesicles.* 2015;4:28665.

64. Wang L, Li Z, Shen J, et al. Exosome-like vesicles derived by *Schistosoma japonicum* adult worms mediates M1 type immune-activity of macrophage. *Parasitol Res.* 2015;114:1865–1873.
65. Chaiyadet S, Sotillo J, Smout M, et al. Carcinogenic liver fluke secretes extracellular vesicles that promote cholangiocytes to adopt a tumorigenic phenotype. *J Infect Dis.* 2015;212:1636–1645.
66. Tzelos T, Matthews JB, Buck AH, et al. A preliminary proteomic characterisation of extracellular vesicles released by the ovine parasitic nematode, *Teladorsagia circumcincta*. *Vet Parasitol.* 2016;221:84–92.
67. Hansen EP, Kringel H, Williams AR, Nejsum P. Secretion of RNA-containing extracellular vesicles by the porcine whipworm, *Trichuris suis*. *J Parasitol.* 2015;101:336–340.
68. Buck AH, Coakley G, Simbari F, et al. Exosomes secreted by a nematode parasite transfer small RNAs to mammalian cells and regulate genes of the innate immune system. *Nat Commun.* 2014;5:1–11.
69. Zamanian M, Fraser LM, Agbedanu PN, et al. Release of small RNA-containing exosome-like vesicles from the human filarial parasite *Brugia malayi*. *PLoS Negl Trop Dis.* 2015;9:1–23.
70. Fromm B, Trelis M, Hackenberg M, Cantalapiedra F, Bernal D, Marcilla A. The revised microRNA complement of *Fasciola hepatica* reveals a plethora of overlooked microRNAs and evidence for enrichment of immuno-regulatory microRNAs in extracellular vesicles. *Int J Parasitol.* 2015;45:697–702.
71. Villarroya-Beltri C, Gutiérrez-Vázquez C, Sánchez-Cabo F, et al. Sumoylated hnRNPA2B1 controls the sorting of miRNAs into exosomes through binding to specific motifs. *Nat Commun.* 2013;4:2980.
72. Mukherjee K, Ghoshal B, Ghosh S, et al. Reversible HuR-microRNA binding controls extracellular export of miR-122 and augments stress response. *EMBO Rep.* 2016;17:11841203.
73. Gibbings DJ, Ciaudo C, Erhardt M, Voinnet O. Multivesicular bodies associate with components of miRNA effector complexes and modulate miRNA activity. *Nat Cell Biol.* 2009;11:1143–1149.
74. Montaner S, Galiano A, Trelis M, et al. The role of extracellular vesicles in modulating the host immune response during parasitic infections. *Front Immunol.* 2014;5:1–8.
75. Hoy AM, Lundie RJ, Ivens A, et al. Parasite-derived microRNAs in host serum as novel biomarkers of helminth infection. *PLoS Negl Trop Dis.* 2014;8:e2701.
76. Cai P, Gobert GN, You H, Duke M, McManus DP. Circulating miRNAs: potential novel biomarkers for hepatopathology progression and diagnosis of Schistosomiasis japonica in two murine models. *PLoS Negl Trop Dis.* 2015;9:1–18.
77. Cheng G, Luo R, Hu C, Cao J, Jin Y. Deep sequencing-based identification of pathogen-specific microRNAs in the plasma of rabbits infected with *Schistosoma japonicum*. *Parasitology.* 2013;140:1751–1761.
78. Tritten L, Burkman E, Moorhead A, et al. Detection of circulating parasite-derived MicroRNAs in filarial infections. *PLoS Negl Trop Dis.* 2014;8:e2971.
79. Tritten L, O'Neill M, Nutting C, et al. *Loa loa* and *Onchocerca ochengi* miRNAs detected in host circulation. *Mol Biochem Parasitol.* 2014;198:14–17.
80. Quintana JF, Makepeace BL, Babayan SA, et al. Extracellular *Onchocerca*-derived small RNAs in host nodules and blood. *Parasit Vectors.* 2015;8:58.
81. Attout T, Hoerauf A, Dénécé G, et al. Lymphatic vascularisation and involvement of Lyve-1 + macrophages in the human *Onchocerca* nodule. *PLoS ONE.* 2009;4:e8234.
82. Pritchard CC, Cheng HH, Tewari M. MicroRNA profiling: approaches and considerations. *Nat Rev Genet.* 2012;13:358–369.
83. Alhassan A, Li Z, Poole CB, Carlow CKS. Expanding the MDx toolbox for filarial diagnosis and surveillance. *Trends Parasitol.* 2015;31:391–400.
84. Twu O, Johnson PJ. Parasite extracellular vesicles: mediators of intercellular communication. *PLoS Pathog.* 2014;10:8–10.
85. Mantel PY, Hoang AN, Goldowitz I, et al. Malaria-infected erythrocyte-derived microvesicles mediate cellular communication within the parasite population and with the host immune system. *Cell Host Microbe.* 2013;13:521–534.
86. Barteneva NS, Maltsev N, Vorobjev IA. Microvesicles and intercellular communication in the context of parasitism. *Front Cell Infect Microbiol.* 2013;3:49.
87. Fischer K, Beatty WL, Jiang D, Weil GJ, Fischer PU. Tissue and stage-specific distribution of *Wolbachia* in *Brugia malayi*. *PLoS Negl Trop Dis.* 2011;5:e1174.
88. McNulty SN, Fischer K, Curtis KC, Weil GJ, Brattig NW, Fischer PU. Localization of *Wolbachia*-like gene transcripts and peptides in adult *Onchocerca flexuosa* worms indicates tissue specific expression. *Parasit Vectors.* 2013;6:2.
89. Voronin D, Bachu S, Shlossman M, Unnasch TR, Ghedin E, Lustigman S. Glucose and glycogen metabolism in *Brugia malayi* is associated with *wolbachia* symbiont fitness. *PLoS ONE.* 2016;11:1–18.
90. Tritten L, Geary TG. MicroRNAs of filarial nematodes: a new frontier in host-pathogen interactions. In: Ana LL, Francisco JE, eds. *Non-coding RNAs Inter-kingdom Communication*. Switzerland: Springer International Publishing; 2016: 207–223.
91. Zamanian M, Andersen EC. Prospects and challenges of CRISPR/Cas genome editing for the study and control of neglected vector-borne nematode diseases. *FEBS J.* 2016;283:3204–3221.
92. Ward JD. Rendering the intractable more tractable: tools from *Caenorhabditis elegans* ripe for import into parasitic nematodes. *Genetics.* 2015;201:1279–1294.

How to cite this article: Quintana JF, Babayan SA, Buck AH. Small RNAs and extracellular vesicles in filarial nematodes: From nematode development to diagnostics. *Parasite Immunology.* 2017;39:e12395. <https://doi.org/10.1111/pim.12395>