

Liu, D. J. et al. (2017) Exome-wide association study of plasma lipids in >300,000 individuals. *Nature Genetics*, 49, pp. 1758-1766. (doi:10.1038/ng.3977)

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

http://eprints.gla.ac.uk/150909

Deposited on: 01 November 2017

 $Enlighten-Research \ publications \ by \ members \ of \ the \ University \ of \ Glasgow \ \underline{http://eprints.gla.ac.uk}$ 

Title: Exome-wide association study of plasma lipids in >300,000 individuals

Authors: Dajiang J. Liu<sup>1†</sup>, Gina M Peloso<sup>2,3†</sup>, Haojie Yu<sup>4†</sup>, Adam Butterworth<sup>5,6†</sup>, Xiao Wang<sup>7†</sup>, Anubha Mahajan<sup>8†</sup>, Danish Saleheen<sup>5,9,10†</sup>, Connor Emdin<sup>3,11†</sup>, Dewar Alam<sup>12</sup>, Alexessander Couto Alves<sup>13</sup>, Philippe Amouyel<sup>14</sup>, Emanuele di Angelantonio<sup>5,6</sup>, Dominique Arveiler<sup>15</sup>, Themistocles L. Assimes<sup>16,17</sup>, Paul L. Auer<sup>18</sup>, Usman Baber<sup>19</sup>, Christie M. Ballantyne<sup>20</sup>, Lia E. Bang<sup>21</sup>, Marianne Benn<sup>22,23</sup>, Joshua C. Bis<sup>24</sup>, Michael Boehnke<sup>25</sup>, Eric Boerwinkle<sup>26</sup>, Jette Bork-Jensen<sup>27</sup>, Erwin P. Bottinger<sup>28</sup>, Ivan Brandslund<sup>29,30</sup>, Morris Brown<sup>31</sup>, Fabio Busonero<sup>32</sup>, Mark J Caulfield<sup>33,34</sup>, John C Chambers<sup>35-37</sup>, Daniel I. Chasman<sup>38,39</sup>, Y. Eugene Chen<sup>40</sup>, Yii-Der Ida Chen<sup>41</sup>, Rajiv Chowdhury<sup>5</sup>, Cramer Christensen<sup>42</sup>, Audrey Y. Chu<sup>38,43</sup>, John M Connell<sup>44</sup>, Francesco Cucca<sup>32,45</sup>, L. Adrienne Cupples<sup>2,43</sup>, Scott M. Damrauer<sup>46,47</sup>, Gail Davies<sup>48,49</sup>, Ian J Deary<sup>48,49</sup>, George Dedoussis<sup>50</sup>, Joshua C. Denny<sup>51,52</sup>, Anna Dominiczak<sup>53</sup>, Marie-Pierre Dubé<sup>54-56</sup>, Tapani Ebeling<sup>57</sup>, Gudny Eiriksdottir<sup>58</sup>, Tõnu Esko<sup>3,59</sup>, Aliki-Eleni Farmaki<sup>50</sup>, Mary F Feitosa<sup>60</sup>, Marco Ferrario<sup>61</sup>, Jean Ferrieres<sup>62</sup>, Ian Ford<sup>63</sup>, Myriam Fornage<sup>64</sup>, Paul W. Franks<sup>65-67</sup>, Timothy M. Frayling<sup>68</sup>, Ruth Frikke-Schmidt<sup>69,70</sup>, Lars Fritsche<sup>25</sup>, Philippe Frossard<sup>10</sup>, Valentin Fuster<sup>19</sup>, Santhi K. Ganesh<sup>40,71</sup>, Wei Gao<sup>72</sup>, Melissa E. Garcia<sup>73</sup>, Christian Gieger<sup>74-76</sup>, Franco Giulianini<sup>38</sup>, Mark O. Goodarzi<sup>77,78</sup>, Harald Grallert<sup>74-76</sup>, Niels Grarup<sup>27</sup>, Leif Groop<sup>79</sup>, Megan L. Grove<sup>26</sup>, Vilmundur Gudnason<sup>58,80</sup>, Torben Hansen<sup>27,81</sup>, Tamara B. Harris<sup>73</sup>, Caroline Hayward<sup>82</sup>, Joel N. Hirschhorn<sup>3,83</sup>, Oddgeir L. Holmen<sup>84,85</sup>, Jennifer Huffman<sup>82</sup>, Yong Huo<sup>86</sup>, Kristian Hveem<sup>84</sup>, Sehrish Jabeen<sup>10</sup>, Anne U Jackson<sup>25</sup>, Johanna Jakobsdottir<sup>58,80</sup>, Marjo-Riitta Jarvelin<sup>13</sup>, Gorm B Jensen<sup>87</sup>, Marit E. Jørgensen<sup>88,89</sup>, J. Wouter Jukema<sup>90,91</sup>, Johanne M. Justesen<sup>27</sup>, Pia R. Kamstrup<sup>22,23</sup>, Stavroula Kanoni<sup>92</sup>, Fredrik Karpe<sup>93,94</sup>, Frank Kee<sup>95</sup>, Amit V. Khera<sup>3,11</sup>, Derek Klarin<sup>3,11,46</sup>, Heikki A. Koistinen<sup>96-98</sup>, Jaspal S Kooner<sup>36,37,99</sup> Charles L. Kooperberg<sup>100</sup>, Kari Kuulasmaa<sup>96</sup>, Johanna Kuusisto<sup>101</sup>, Markku Laakso<sup>101</sup>, Timo Lakka<sup>102,103</sup>, Claudia Langenberg<sup>104</sup>, Anne Langsted<sup>22,23</sup>, Lenore J. Launer<sup>73</sup>, Torsten Lauritzen<sup>105</sup>, David CM Liewald<sup>48,49</sup>, Li An Lin<sup>64</sup>, Allan Linneberg<sup>106-108</sup>, Ruth J.F. Loos<sup>28,109</sup>, Yingchang Lu<sup>28</sup>, Xiangfeng Lu<sup>40,110</sup>, Reedik Mägi<sup>59</sup>, Anders Malarstig<sup>111,112</sup>, Ani Manichaikul<sup>113</sup>, Alisa K. Manning<sup>3,11,114</sup>, Pekka Mäntyselkä<sup>115</sup>, Eirini Marouli<sup>92</sup>, Nicholas GD Masca<sup>116,117</sup>, Andrea Maschio<sup>32</sup>, James B. Meigs<sup>3,114,118</sup>, Olle Melander<sup>119</sup>, Andres Metspalu<sup>59</sup>, Andrew P Morris<sup>8,120</sup>, Alanna C. Morrison<sup>26</sup>, Antonella Mulas<sup>32</sup>, Martina Müller-Nurasyid<sup>75,121,122</sup>, Patricia B. Munroe<sup>33,123</sup>, Matt J Neville<sup>93</sup>, Jonas Nielsen<sup>40</sup>, Sune F Nielsen<sup>22,23</sup>, Børge G Nordestgaard<sup>22,23</sup>, Jose M. Ordovas<sup>124-126</sup>, Roxana Mehran<sup>19</sup>, Christoper J. O'Donnell<sup>46,127</sup>, Marju Orho-Melander<sup>119</sup>, Cliona M. Molony<sup>128</sup>, Pieter Muntendam<sup>129</sup>, Sandosh Padmanabhan<sup>53</sup>, Colin NA Palmer<sup>44</sup>, Dorota Pasko<sup>68</sup>, Aniruddh P. Patel<sup>3,11,127,130</sup>, Oluf Pedersen<sup>27</sup>, Markus Perola<sup>96,131</sup>, Annette Peters<sup>74,75,122</sup>, Charlotta Pisinger<sup>108</sup>, Giorgio Pistis<sup>32</sup>, Ozren Polasek<sup>132,133</sup>, Neil Poulter<sup>134</sup>, Bruce M. Psaty<sup>24,135,136</sup>, Daniel J. Rader<sup>47,137</sup>, Asif Rasheed<sup>10</sup>, Rainer Raumaraa<sup>103,138</sup>, Dermot Reilly<sup>128</sup>, Alex P. Reiner<sup>100,139</sup>, Frida Renström<sup>65,140</sup>, Stephen S Rich<sup>113</sup>, Paul M Ridker<sup>38</sup>, John D. Rioux<sup>54</sup>, Neil R Robertson<sup>8,93</sup>, Dan M. Roden<sup>52</sup>, Jerome I. Rotter<sup>41</sup>, Igor Rudan<sup>133</sup>, Veikko Salomaa<sup>96</sup>, Nilesh J Samani<sup>116,117</sup>, Serena Sanna<sup>32</sup>, Naveed Sattar<sup>53</sup>, Ellen M. Schmidt<sup>141</sup>, Robert A. Scott<sup>104</sup>, Peter Sever<sup>134</sup>, Raquel S. Sevilla<sup>142</sup>, Christian M. Shaffer<sup>52</sup>, Xueling Sim<sup>25,143</sup>, Suthesh Sivapalaratnam<sup>144</sup>, Kerrin S Small<sup>145</sup>, Albert V. Smith<sup>58,80</sup>, Blair H Smith<sup>146,147</sup>, Sangeetha Somayajula<sup>148</sup>, Lorraine Southam<sup>8,149</sup>, Timothy D Spector<sup>145</sup>, Elizabeth K. Speliotes<sup>141,150</sup>, John M Starr<sup>48,151</sup>, Kathleen E Stirrups<sup>92,152</sup>, Nathan Stitziel<sup>153,154</sup>, Konstantin Strauch<sup>75,155</sup>, Heather M Stringham<sup>25</sup>, Praveen Surendran<sup>5</sup>, Hayato Tada<sup>156</sup>, Alan R. Tall<sup>157</sup>, Hua Tang<sup>158</sup>, Jean-Claude Tardif<sup>54,56</sup>, Kent D Taylor<sup>159</sup>, Stella Trompet<sup>90,160</sup>, Philip S. Tsao<sup>16,17</sup>, Jaakko Tuomilehto<sup>161-164</sup>, Anne Tybjaerg-Hansen<sup>69,70</sup>, Natalie R van Zuydam<sup>8,44</sup>, Anette Varbo<sup>22,23</sup>, Tibor V Varga<sup>65</sup>, Jarmo Virtamo<sup>96</sup>, Melanie Waldenberger<sup>75,76</sup>, Nan Wang<sup>157</sup>, Nick J. Wareham<sup>104</sup>, Helen R Warren<sup>33</sup>, Peter E. Weeke<sup>52,165</sup>, Joshua Weinstock<sup>25</sup>, Jennifer Wessel<sup>166,167</sup>, James G. Wilson<sup>168</sup>, Peter W. F. Wilson<sup>169,170</sup>, Ming Xu<sup>171</sup>, Hanieh Yaghootkar<sup>68</sup>, Robin Young<sup>5</sup>, Eleftheria Zeggini<sup>149</sup>, He Zhang<sup>40</sup>, Neil S. Zheng<sup>172</sup>, Weihua Zhang<sup>35</sup>, Yan Zhang<sup>86</sup>, Wei Zhou<sup>141</sup>, Yanhua Zhou<sup>2</sup>, Magdalena Zoledziewska<sup>32</sup>, Charge Diabetes Working Group, The EPIC-InterAct consortium, EPIC-CVD Consortium, GOLD Consortium, VA Million Veteran Program, Joanna MM Howson<sup>5†</sup>, John Danesh<sup>5,6,149†</sup>, Mark I McCarthy<sup>8,93,94†</sup>, Chad Cowan<sup>4,173†</sup>, Goncalo Abecasis<sup>25†</sup>, Panos Deloukas<sup>92,174†</sup>, Kiran Musunuru<sup>7†</sup>, Cristen J. Willer<sup>40,71,141†\*</sup>, Sekar Kathiresan<sup>3,11,127,130†\*</sup>

# Affiliations

- 1. Department of Public Health Sciences, Institute of Personalized Medicine, Penn State College of Medicine, Hershey, PA, USA.
- 2. Department of Biostatistics, Boston University School of Public Health, Boston, MA, USA.
- 3. Program in Medical and Population Genetics, Broad Institute, Cambridge, MA, USA.
- 4. Department of Stem Cell and Regenerative Biology, Harvard Stem Cell Institute, Harvard University, Cambridge, MA 02138, USA.
- 5. Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK.
- 6. The National Institute for Health Research Blood and Transplant Unit (NIHR BTRU) in Donor Health and Genomics at the University of Cambridge, Cambridge, UK.
- 7. Cardiovascular Institute, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104 USA.
- 8. Wellcome Trust Centre for Human Genetics, University of Oxford, Oxford, UK.
- 9. Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, PA, USA.
- 10. Center for Non-Communicable Diseases, Karachi, Pakistan.
- 11. Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA.
- 12. ICDDR, B, Mohakhali, Dhaka 1212, Bangladesh.
- 13. Imperial College London, London, UK.
- 14. Univ. Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U1167 RID-AGE Risk factors and molecular determinants of aging-related diseases, F-59000 Lille, France.
- 15. Department of Epidemiology and Public Health, EA 3430, University of Strasbourg, Strasbourg, F- 67085, France.
- 16. VA Palo Alto Health Care System, Palo Alto, CA, USA.
- 17. Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.
- 18. Zilber School of Public Health, University of Wisconsin-Milwaukee, Milwaukee WI, USA.
- 19. Cardiovascular Institute, Mount Sinai Medical Center, Icahn School of Medicine, Mount Sinai, New York, NY, USA.
- 20. Department of Medicine, Baylor College of Medicine, Houston, TX, USA.
- 21. Department of Cardiology, Rigshospitalet, Copenhagen University Hospital, Copenhagen, Denmark.
- 22. Department of Clinical Biochemistry and The Copenhagen General Population Study, Herlev and Gentofte Hospital, Copenhagen University Hospital, Denmark.
- 23. Faculty of Health and Medical Sciences, University of Denmark, Denmark.
- 24. Cardiovascular Health Research Unit, Department of Medicine, University of Washington, Seattle, WA, USA.
- 25. Center for Statistical Genetics, Department of Biostatistics, University of Michigan School of Public Health, Ann Arbor, Michigan, USA.
- 26. Human Genetics Center, School of Public Health, The University of Texas School Health Science Center at Houston, Houston, Texas.
- 27. The Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
- 28. The Charles Bronfman Institute for Personalized Medicine, Ichan School of Medicine at Mount Sinai, New York, NY 10029, USA.
- 29. Department of Clinical Biochemistry, Lillebaelt Hospital, Vejle, Denmark.
- 30. Institute of Regional Health Research, University of Southern Denmark, Odense, Denmark.
- 31. Clinical Pharmacology Unit, University of Cambridge, Addenbrookes Hospital, Hills Road, Cambridge CB2 2QQ, UK.

- 32. Istituto di Ricerca Genetica e Biomedica, Consiglio Nazionale delle Ricerche (CNR), Monserrato, Cagliari, Italy.
- 33. Clinical Pharmacology, William Harvey Research Institute, Barts and The London, Queen Mary University of London, Charterhouse Square, London, EC1M 6BQ, UK.
- 34. The Barts Heart Centre, William Harvey Research Institute, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, UK.
- 35. Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, Norfolk Place, London W2 1PG, UK.
- 36. Department of Cardiology, Ealing Hospital NHS Trust, Uxbridge Road, Southall, Middlesex UB1 3HW, UK.
- 37. Imperial College Healthcare NHS Trust, London, UK.
- 38. Division of Preventive Medicine, Boston, MA, USA.
- 39. Harvard Medical School, Boston, MA, USA.
- 40. Department of Internal Medicine, Division of Cardiovascular Medicine, University of Michigan, Ann Arbor, Michigan 48109, USA.
- 41. The Institute for Translational Genomics and Population Sciences, LABioMed at Harbor-UCLA Medical Center, Departments of Pediatrics and Medicine, Los Angeles, CA, USA.
- 42. Medical Department, Lillebaelt Hospital, Vejle, Denmark.
- 43. NHLBI Framingham Heart Study, Framingham, MA, USA.
- 44. Medical Research Institute, University of Dundee, Ninewells Hospital and Medical School, Dundee, UK.
- 45. Dipartimento di Scienze Biomediche, Universita' degli Studi di Sassari, Sassari, Italy.
- 46. Massachusetts Veterans Epidemiology Research and Information Center (MAVERIC), VA Boston Healthcare System, Boston, MA, USA.
- 47. Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.
- 48. Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh, UK.
- 49. Department of Psychology, University of Edinburgh, Edinburgh, UK.
- 50. Department of Nutrition and Dietetics, School of Health Science and Education, Harokopio University, Athens, 17671, Greece.
- 51. Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN, USA.
- 52. Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA.
- 53. British Heart Foundation Glasgow Cardiovascular Research Centre, Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK.
- 54. Montreal Heart Institute, Montreal, Quebec, Canada.
- 55. Université de Montréal Beaulieu-Saucier Pharmacogenomics Center, Montreal, Quebec, Canada.
- 56. Université de Montréal, Montreal, Quebec, Canada.
- 57. Department of Medicine, Oulu University Hospital and University of Oulu, Oulu, Finland.
- 58. The Icelandic Heart Association, Kopavogur, Iceland.
- 59. Estonian Genome Center, University of Tartu, Tartu, Estonia.
- 60. Division of Statistical Genomics, Department of Genetics, Washington University School of Medicine, St. Louis, MO, USA.
- 61. Research Centre in Epidemiology and Preventive Medicine EPIMED, Department of Clinical and Experimental Medicine, University of Insubria, Varese, Italy.
- 62. Department of Epidemiology, UMR 1027- INSERM, Toulouse University-CHU Toulouse, Toulouse, France.
- 63. University of Glasgow, Glasgow, UK.

- 64. Institute of Molecular Medicine, the University of Texas Health Science Center at Houston, Houston, TX, USA.
- 65. Department of Clinical Sciences, Genetic and Molecular Epidemiology Unit, Lund University, Malmö, Sweden.
- 66. Department of Public Health & Clinical Medicine, Umeå University, Umeå, Sweden.
- 67. Department of Nutrition, Harvard T. H. Chan School of Public Health, Boston, MA, USA.
- 68. Genetics of Complex Traits, University of Exeter Medical School, University of Exeter, Exeter EX2 5DW, UK.
- 69. Department of Clinical Biochemistry, Rigshospitalet, Copenhagen, Denmark.
- 70. Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
- 71. Department of Human Genetics, University of Michigan, Ann Arbor, Michigan 48109, USA.
- 72. Department of Cardiology, Peking University Third Hospital, Key Laboratory of Cardiovascular Molecular Biology and Regulatory Peptides, Ministry of Health, Beijing 100191, China.
- 73. Laboratory of Epidemiology and Population Sciences, National Institute on Aging, Bethesda, MD 20892, USA.
- 74. German Center for Diabetes Research (DZD e.V.), Neuherberg, Germany.
- 75. Institute of Genetic Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany.
- 76. Research Unit of Molecular Epidemiology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany.
- 77. Department of Medicine and Department of Biomedical Sciences, Cedars-Sinai Medical Center, Los Angeles, California 90048, USA.
- 78. Division of Endocrinology, Diabetes and Metabolism, Cedars-Sinai Medical Center, Los Angeles, California 90048, USA.
- 79. Department of Clinical Sciences, Diabetes and Endocrinology, Clinical Research Centre, Lund University, Malmö, Sweden.
- 80. The University of Iceland, Reykjavik, Iceland.
- 81. Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark.
- 82. Medical Research Council Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, UK.
- 83. Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston Children's Hospital, Boston, MA, USA.
- 84. Department of Public Health and General Practice, HUNT Research Centre, Norwegian University of Science and Technology, 7600 Levanger, Norway.
- 85. St Olav Hospital, Trondheim University Hospital, 7030 Trondheim, Norway.
- 86. Department of Cardiology, Peking University First Hospital, Beijing 100034, China.
- 87. The Copenhagen City Heart Study, Frederiksberg Hospital, Denmark.
- 88. Steno Diabetes Center, Gentofte, Denmark.
- 89. National Institute of Public Health, Southern Denmark University, Denmark.
- 90. Department of Cardiology, Leiden University Medical Center, Leiden, The Netherlands.
- 91. The Interuniversity Cardiology Institute of the Netherlands, Utrecht, The Netherlands.
- 92. William Harvey Research Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK.
- 93. Oxford Centre for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford, Oxford, UK.
- 94. Oxford NIHR Biomedical Research Centre, Oxford University Hospitals Trust, Oxford, UK.
- 95. Director, UKCRC Centre of Excellence for Public Health, Queens University, Belfast, Northern Ireland.
- 96. Department of Health, National Institute for Health and Welfare, FI-00271, Helsinki, Finland.
- 97. Department of Medicine, and Abdominal Center: Endocrinology, Helsinki University Central Hospital, Helsinki, Finland.

- 98. Minerva Foundation Institute for Medical Research, Helsinki, Finland.
- 99. National Heart and Lung Institute, Imperial College London, Hammersmith Hospital Campus, London, UK.
- 100. Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle WA, USA.
- 101. Institute of Clinical Medicine, Internal Medicine, University of Eastern Finland and Kuopio University Hospital, 70210 Kuopio, Finland.
- 102. Department of Physiology, Institute of Biomedicine, University of Eastern Finland, Kuopio Campus, Kuopio, Finland.
- 103. Kuopio Research Institute of Exercise Medicine, Kuopio, Finland.
- 104. MRC Epidemiology Unit, Institute of Metabolic Science, University of Cambridge School of Clinical Medicine, Cambridge, UK.
- 105. Department of Public Health, Section of General Practice, University of Aarhus, Aarhus, Denmark.
- 106. Department of Clinical Experimental Research, Rigshospitalet, Glostrup, Denmark.
- 107. Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.
- 108. Research Center for Prevention and Health, Capital Region of Denmark, Copenhagen, Denmark.
- 109. The Mindich Child Health and Development Institute, Ichan School of Medicine at Mount Sinai, New York, NY 10029, USA.
- 110. State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, National Center for Cardiovascular Diseases, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, China.
- 111. Cardiovascular Genetics and Genomics Group, Cardiovascular Medicine Unit, Department of Medicine, Solna, Karolinska Institutet, Stockholm, Sweden.
- 112. Pharmatherapeutics Clinical Research, Pfizer Worldwide R&D, Sollentuna, Sweden.
- 113. Center for Public Health Genomics, University of Virginia, Charlottesville, VA USA.
- 114. Department of Medicine, Massachusetts General Hospital, Boston, MA, USA.
- 115. Unit of Primary Health Care, University of Eastern Finland and Kuopio University Hospital, Kuopio, Finland.
- 116. Department of Cardiovascular Sciences, University of Leicester, UK.
- 117. NIHR Leicester Cardiovascular Biomedical Research Unit, Glenfield Hospital, Leicester, UK.
- 118. Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA, USA.
- 119. Department of Clinical Sciences, University Hospital Malmo Clinical Research Center, Lund University, Malmo, Sweden.
- 120. Department of Biostatistics, University of Liverpool, Liverpool, UK.
- 121. Department of Medicine, Ludwig-Maximilians-University, Munich, Germany.
- 122. DZHK German Centre for Cardiovascular Research, Munich, Germany.
- 123. NIHR Barts Cardiovascular Biomedical Research Unit, Queen Mary University of London, London, UK.
- 124. Department of Cardiovascular Epidemiology and Population Genetics, National Center for Cardiovascular Investigation, Madrid 28049, Spain.
- 125. IMDEA-Alimentacion, Madrid 28049, Spain.
- 126. Nutrition and Genomics Laboratory, Jean Mayer-USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA 02111, USA.
- 127. Department of Medicine, Harvard Medical School, Boston, MA, USA.
- 128. Genetics, Merck Sharp & Dohme Corp., Kenilworth, New Jersey, USA.
- 129. BG Medicine Inc, Waltham, MA.
- 130. Cardiovascular Research Center, Massachusetts General Hospital, Boston, MA, USA.
- 131. Institute of Molecular Medicine FIMM, University of Helsinki, Finland.
- 132. Faculty of Medicine, University of Split, Split, Croatia.
- 133. Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK.

- 134. International Centre for Circulatory Health, Imperial College London, W2 1PG, UK.
- 135. Group Health Research Institute, Group Health Cooperative, Seattle, WA.
- 136. Departments of Epidemiology and Health Services, University of Washington, Seattle, WA, USA.
- 137. Department of Genetics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.
- 138. Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, Kuopio, Finland.
- 139. Department of Epidemiology, University of Washington, Seattle WA, USA.
- 140. Department of Biobank Research, Umeå University, Umeå, Sweden.
- 141. Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, Michigan 48109, USA.
- 142. Imaging, Merck Sharp & Dohme Corp., Kenilworth, New Jersey, USA.
- 143. Saw Swee Hock School of Public Health, National University of Singapore, Singapore, 117549, Singapore.
- 144. Department of Vascular Medicine, Academic Medical Center, University of Amsterdam, Amsterdam, NL.
- 145. Department of Twin Research and Genetic Epidemiology, King's College London, London, UK.
- 146. Division of Population Health Sciences, Ninewells Hospital and Medical School, University of Dundee, Dundee, Scotland.
- 147. Generation Scotland, Centre for Genomic and Experimental Medicine, University of Edinburgh, Edinburgh, UK.
- 148. Scientific Informatics, Merck Sharp & Dohme Corp., Kenilworth, New Jersey, USA.
- 149. Wellcome Trust Sanger Institute, Genome Campus, Hinxton, UK.
- 150. Department of Internal Medicine, Division of Gastroenterology, University of Michigan, Ann Arbor, Michigan 48109, USA.
- 151. Alzheimer Scotland Dementia Research Centre, University of Edinburgh, Edinburgh, UK.
- 152. Department of Haematology, University of Cambridge, Cambridge, UK.
- 153. Cardiovascular Division, Departments of Medicine and Genetics, Washington University School of Medicine, St. Louis, MO, USA.
- 154. The McDonnell Genome Institute, Washington University School of Medicine, St. Louis, MO, USA.
- 155. Institute of Medical Informatics, Biometry and Epidemiology, Chair of Genetic Epidemiology, Ludwig-Maximilians-Universität, Munich, Germany.
- 156. Division of Cardiovascular Medicine, Kanazawa University Graduate School of Medicine, Kanazawa, Japan.
- 157. Division of Molecular Medicine, Department of Medicine, Columbia University, New York, NY, USA.
- 158. Department of Genetics, Stanford University School of Medicine, Stanford CA 94305, USA.
- 159. Los Angeles Biomedical Research Institute at Harbor, UCLA, Los Angeles, CA, USA.
- 160. Department of Gerontology and Geriatrics, Leiden University Medical Center, Leiden, the Netherlands.
- 161. Chronic Disease Prevention Unit, National Institute for Health and Welfare, 00271 Helsinki, Finland.
- 162. Dasman Diabetes Institute, Dasman 15462, Kuwait.
- 163. Centre for Vascular Prevention, Danube-University Krems, 3500 Krems, Austria.
- 164. Saudi Diabetes Research Group, King Abdulaziz University, Fahd Medical Research Center, Jeddah 21589, Saudi Arabia.
- 165. The Heart Centre, Department of Cardiology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark.
- 166. Department of Epidemiology, Indiana University Fairbanks School of Public Health, Indianapolis, Indiana, USA.

- 167. Department of Medicine, Indiana University School of Medicine, Indianapolis, Indiana, USA.
- 168. Department of Physiology and Biophysics, University of Mississippi Medical Center, Jackson, MS, USA.
- 169. Atlanta VA Medical Center, Decatur, GA, USA.
- 170. Emory Clinical Cardiovascular Research Institute, Atlanta, GA, USA.
- 171. Department of Cardiology, Institute of Vascular Medicine, Peking University Third Hospital, Key Laboratory of Molecular Cardiovascular Sciences, Ministry of Education, Beijing 100191, China.
- 172. Yale University, New Haven, CT, USA.
- 173. Center for Regenerative Medicine, Massachusetts General Hospital, Boston, MA 02114, USA.
- 174. Princess Al-Jawhara Al-Brahim Centre of Excellence in Research of Hereditary Disorders (PACER-HD), King Abdulaziz University, Jeddah 21589, Saudi Arabia.

<sup>†</sup>These authors contributed equally.

\*Correspondence to:

Cristen Willer, cristen@umich.edu

Sekar Kathiresan, SKATHIRESAN1@mgh.harvard.edu

Plasma lipid levels are modifiable risk factors for atherosclerotic cardiovascular disease. In >300,000 participants, we screened DNA sequence variants on an exome-focused array and identified 444 independent variants in 250 loci significantly associated with total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), and/or triglycerides (TG). At two loci (JAK2 and A1CF), experimental analysis in mice revealed TC and TG changes, respectively, consistent with the human data. We utilized 444 mapped variants to address several questions of clinical relevance and found the following: (1) beta-thalassemia trait carriers displayed lower total cholesterol and were protected from coronary artery disease; (2) multiple CETP variants each consistently associated with not only higher HDL-C but also elevated risk for age-related macular degeneration; (3) only some mechanisms of lowering LDL-C seemed to increase risk for type 2 diabetes; and (4) at TG genes involved in hepatic production of TG-rich lipoproteins (e.g., TM6SF2, PNPLA3), the plasma TG-lowering allele tracked with higher liver fat, higher risk for type 2 diabetes, and lower risk for coronary artery disease whereas at TG genes involed in peripheral lipolysis (e.g., LPL, ANGPTL4), the plasma TG-lowering allele had no effect on liver fat but lowered risks for both type 2 diabetes and coronary artery disease.

# Main Text

Association studies testing common DNA sequence variation have uncovered 175 genetic loci that affect lipid levels in the population<sup>1-7</sup>, with most variants residing in the non-coding portion of the genome. These findings have informed biology of lipoproteins and elucidated the causal roles of lipid levels on cardiovascular disease<sup>8-11</sup>. Here, we build on these previous efforts to: 1) perform an exome-wide association screen for plasma lipids in >300,000 individuals; 2) evaluate discovered alleles experimentally in mouse models; and 3) test the inter-relationship of mapped lipid variants with four diseases – coronary artery disease (CAD), age-related macular degeneration (AMD), fatty liver, and type 2 diabetes.

We tested the association of genotypes from the HumanExome BeadChip (i.e., exome array) with lipid levels in each of 73 studies encompassing >300,000 participants (Supplementary Material, Tables S1-S3) across several ancestries (European, African, South Asian, Hispanic, and other). The maximal sample sizes from each ancestry were 237,050 for European, 16,935 for African, 37,613 for South Asian, and 5,082 for Hispanic or other. A companion manuscript describes results for 47,532 East Asian participants<sup>12</sup>. A total of 242,289 variants were analyzed after quality control (Supplementary Methods). About one-third of the genotyped variants are non-synonymous with minor allele frequency (MAF) < 0.1% (Table S4).

Single-variant association statistics and inter-marker linkage disequilibrium information summarized across 1 megabase sliding windows were generated from each cohort using RAREMETALWORKER or RVTESTS<sup>13,14</sup> software. Meta-analyses of genetic associations were performed using the R-package rareMETALS (version 6.0). In addition to testing each variant individually for association, we also performed gene-level association tests, which aggregate multiple rare coding variants in each of 16,798 protein-coding genes. Genomic control values for meta-analysis results were between 1.09 and 1.14 for all four lipid traits (Fig. S1), suggesting that population structure in our analysis is well-controlled<sup>3,15</sup>. Genomic control values scale with sample size<sup>16</sup> and as such, our genomic control values are in line with previous reports including similar sample sizes<sup>15</sup>.

In single-variant association analyses, we identified 1,445 variants associated at  $P < 2.1 \times 10^{-7}$  (Bonferroni correction of 242,289 variants analyzed) (Fig. S2; full association results are available at http://csg.sph.umich.edu/abecasis/public/lipids2017/). Of these, 75 were 'novel' (i.e., located in previously unreported loci): 35 of these were protein-altering variants and 40 were non-coding variants (**Table 1**, Table S5 & S6). These 75 new loci are each located at least 1 megabase from previously reported genome-wide association study (GWAS) signals (Table S7). The lead variant was common (MAF > 5%) in 61 of these 75 loci. When analyzed separately by ancestry, European ancestry participants provided the most significant associations for the 75 novel variants, except for two noncoding variants associated with LDL (rs201148465 and rs147032017) which were driven by the South Asian participants (Table S8). Gene-level association analyses revealed an additional five genes where the signal was driven by multiple rare variants ( $P < 4.2 \times 10^{-7}$ , Bonferroni correction threshold for performing 5 tests on ~20,000 genes, Table S9).

We tested the new single variant association findings for replication in an independent set of up to 286,268 participants from three studies – Nord-Trøndelag Health Study<sup>17</sup>, (HUNT; max n = 62,168), Michigan Genomics Initiative (MGI; <u>www.michigangenomics.org</u>; max n = 6,411) and the Million Veteran Program<sup>18</sup> (MVP; max n = 218,117). Of the novel primary trait associations, 73/73 associations were directionally consistent (Table S10); two SNPs were not available for replication (rs201148465, rs75862065). Furthermore, we were able to replicate the associations of 66/73 (90%) at an alpha of 0.05.

At any given genetic locus, multiple coding and non-coding variants may contribute to plasma lipid levels in an independent fashion. We quantified this phenomenon by iteratively performing association analyses conditional on the top variants at all lipid loci (175 previously identified and 75 newly identified). With this procedure, we identified 444 variants independently associated with one or more of the four lipid traits in 250 loci (Fig. S3; Table S11

and Table S12). Collectively, these 444 variants explain 9%-12% of the overall phenotypic variance for plasma lipid levels; previously mapped variants explain about 5%-9% of the variance in the current dataset.

The identification of lipid-associated coding variants using the exome array may help refine association signals at previously identified GWAS loci. We were able to evaluate this possibility in 131 of the 175 previously reported GWAS loci where the index variant or a proxy was available on the exome array, and associated with lipids levels with  $P < 2.1 \times 10^{-7}$  (Table S13 and Table S14). For example, rs11136341, an intronic SNP close to the *PLEC* gene reported to be associated with LDL-C, was the original lead SNP in its GWAS locus ( $P = 2 \times 10^{-13}$ ). In the current study, a protein-altering variant in *PARP10* is the top variant in the same locus (rs11136343; Leu395Pro;  $P = 7 \times 10^{-26}$ ). After conditioning on *PARP10* Leu395Pro, the evidence for rs11136341 diminished (P = 0.02); in contrast, *PARP10* Leu395Pro remained significant ( $P = 9 \times 10^{-13}$ ) after conditioning on rs11136341. *PARP10* has been shown to affect the hepatic secretion of apolipoprotein B (apoB) in human hepatocytes<sup>19</sup>; these results prioritize *PARP10* as a causal gene at this locus.

Experimental analysis of discovered mutations in model systems is a powerful approach to validate human association findings. This approach holds particular appeal in following up the results of our exome array analysis, in which hundreds of protein-altering variants were robustly associated with circulating lipids. We prioritized two coding mutations for experimental analysis: *JAK2* (Janus Kinase 2) p.Val617Phe and *A1CF* (APOBEC1 complementation factor) p.Gly398Ser.

*JAK2* p.Val617Phe is a recurrent somatic mutation arising in hematopoietic stem cells which can lead to myeloproliferative disorders or clonal hematopoiesis of indeterminate potential<sup>20-24</sup>. We recently showed that carriage of p.Val617Phe increases with age and confers higher risk for CAD<sup>25</sup>. Surprisingly, the 617Phe allele which increases risk for CAD is associated with *lower* LDL-C. Mice knocked in for *Jak2* p.Val617Phe were created as reported previously<sup>26</sup>. Hypercholesterolemia-prone mice that were engrafted with bone marrow obtained from *Jak2* p.Val617Phe transgenic mice displayed lower total cholesterol than did mice that had received control bone marrow (Fig. S4). These data are consistent with our human genetic observations. The mechanism by which *JAK2* p.Val617Phe leads to lower plasma TC/LDL-C but higher risk for CAD requires further study.

Another strong new association to emerge from genetic analyses was between *A1CF* p.Gly398Ser and TG [0.10-standard deviation (SD) increase in TG per copy of alternate allele,  $P = 4 \times 10^{-11}$ ]; this rare variant (MAF = 0.7%) was also associated with increased circulating TC ( $P = 4 \times 10^{-7}$ ) and nominally associated with increased risk of CAD (OR 1.12; P = 0.02). *A1CF* encodes APOBEC1 complementation factor, an RNA-binding protein which facilitates the RNA-editing action of APOBEC1 on the *APOB* transcript<sup>27,28</sup>. We performed CRISPR-Cas9 deletion, rescue, and knock-in experiments to assess whether *A1CF* p.Gly398Ser is a causal mutation that alters TG metabolism.

CRISPR-Cas9-induced deletion of *A1CF* led to 72% and 65% reduction in secreted APOB100 compared to control cells in Huh7 and HepG2 human hepatoma cells, respectively (**Fig. 1A-1C;** Fig. S5). These findings are consistent with previous studies in rat primary hepatocytes that also showed significantly decreased apoB secretion after RNAi-based depletion of  $A1CF^{29}$ . Additionally, cellular APOB100 levels were significantly reduced in A1CF-deficient cells (Fig. S5B and S5C). A subsequent "rescue" experiment involving overexpression of wild-type or p.Gly398Ser A1CF in Huh7 cells with or without endogenous A1CF expression confirmed that significantly more APOB100 secretion was noted in cell lines expressing A1CF p.Gly398Ser (**Fig. 1D**).

We sought to further validate the *A1CF* gene and the p.Gly398Ser variant through the use of CRISPR-Cas9 to generate knock-in mice. Using a guide RNA targeting *A1cf* exon 9, the site of the codon for p.Gly398, and a 162-nucleotide single-strand DNA oligonucleotide repair template containing the p.Gly398Ser variant as well as extra synonymous changes to prevent recleavage by CRISPR-Cas9, we generated mice of the C57BL/6J inbred background with an *A1cf* Gly398Ser allele (hereafter referred to as KI) (Fig. S6A, S6B). We bred the KI allele to homozygosity and found that KI/KI mice were viable and healthy. We compared wild-type and KI/KI colony mates (n = 9, 8) with respect to TG levels (Fig. S6C). We found that KI/KI mice had 46% increased TG compared to wild-type mice (P = 0.05). In sum, these results indicate that *A1CF* is a causal gene for TG in humans and that the p.Gly398Ser variant is a causal mutation, with possible relevance to CAD.

Next, we used the 444 identified DNA sequence variants to address four questions relevant to clinical medicine, focusing on one question for each of the four traits: TC, HDL-C, LDL-C, and TG. First, a rare null mutation in the beta-globin gene (*HBB*; c.92+1G>A, rs33971440) associated with lower total cholesterol (Table S15), raising the question of the relationship between beta-thalassemia and risk for CAD. Approximately 80 to 90 million individuals worldwide are estimated to carry a heterozygous loss-of-function *HBB* mutation, termed beta-thalassemia trait<sup>30</sup>. Observational epidemiologic studies several decades ago showed that beta-thalassemia trait associates with lower blood cholesterol level<sup>31,32</sup>. We find that *HBB* c.92+1G>A is not only associated with lower hematocrit but also with a 17 mg/dl decrease in LDL-C (-17 CI -23, -11;  $P = 2.7 \times 10^{-8}$ ) and a 21 mg/dl decrease in TC (-21 CI -27, -14;  $P = 8.9 \times 10^{-11}$ ) (Fig. S7). In an analysis of 31,156 CAD cases and 65,787 controls, carriers of loss-offunction variants in *HBB* were protected against CAD (odds ratio for CAD, 0.70; 95% CI 0.54, 0.90; P = 0.005, Fig. S8).

Second, DNA sequence variants in the *CETP* gene which associate with higher HDL-C also correlate with higher risk for age-related macular degeneration (AMD), a leading cause of blindness around the world<sup>33-37</sup>; here, we ask if *any way* of increasing *plasma* HDL-C will predictably lead to increased AMD risk. Across 168 independent HDL-C variants at 109 distinct loci with MAF > 1%, we tested the association of each independent HDL-C variant with AMD risk. The effect size of variant on HDL-C was positively correlated with its effect on AMD risk (correlation in effect sizes, r = 0.41,  $P = 4.4 \times 10^{-8}$ ; Table S16, Fig. S9). However, this effect was driven by the 10 independent HDL-C associated variants in the *CETP* gene (heterogeneity across the different HDL-C-raising mechanisms ( $\tau^2 = 0.91$ ,  $P_{het} = 1.8 \times 10^{-15}$ ) (Table S17). A one-SD

increase in HDL-C via a genetic risk score of the 10 *CETP* variants robustly associated with increased AMD risk (OR = 2.01, 95% CI = 1.88-2.14, 16,144 AMD cases, 17,832 controls,  $P = 3.4 \times 10^{-97}$ ). When these 10 *CETP* variants were removed, there was no longer a relationship between genetically-altered HDL-C and AMD risk (P = 0.17). These genetic results suggest that outside of the *CETP* locus, there is not a predictable relationship between plasma HDL-C and risk for AMD.

Third, will lowering LDL-C with lipid-modifying medicines always increase risk for type 2 diabetes? This question is motivated by the fact that in randomized controlled trials, statin therapy increases risk for type 2 diabetes<sup>26,27</sup> and recent reports of *PCSK9* variants associating with higher risk for type 2 diabetes<sup>38-40</sup>. We confirmed the association of *PCSK9* p.Arg46Leu (R46L) with risk for type 2 diabetes among 222,877 participants (Table S18). We found that the 46Leu allele associated with lower LDL-C confers a 13% increased risk for type 2 diabetes (OR 1.13; 95% CI 1.06 – 1.20; P=6.96 x 10<sup>-5</sup>) (Fig. S10). In addition, across 113 independent LDL-C variants at 90 distinct loci, we compared each variant's effect on LDL-C with its subsequent effect on risk for type 2 diabetes. Across the 113 variants, there is a weak inverse correlation between a variant's effects on LDL-C and type 2 diabetes (r=-0.21, p=0.025); however, there is evidence for heterogeneity in this relationship ( $\tau 2 = 0.50$ ,  $P_{het} = 2.5 \times 10^{-9}$ ). Five LDL-C lowering genetic mechanisms had the most compelling evidence for association with higher risk for type 2 diabetes (TM6SF2 p.Glu167Lys, APOE chr19:4510002, HNF4A p.Thr136Ile, *PNPLA3* p.Ile148Met, and *GCKR* p.Leu446Pro) ( $P < 4.0 \times 10^{-4}$  for each, Bonferroni correction threshold for performing tests at 113 variants, Table S19; Fig. S11). These results suggest that only some ways of lowering LDL-C are likely to increase risk for type 2 diabetes.

Finally, two key processes – hepatic production and peripheral lipolysis – contribute to the blood level of TG. We asked how genes involved in hepatic production of TG-rich lipoproteins (*PNPLA3*, *TM6SF2*) differed from lipolysis pathway genes (*LPL*, *ANGPTL4*) in their impact on related metabolic traits - blood lipids, fatty liver, type 2 diabetes, and CAD (**Table 2**). The alternative alleles at *PNPLA3* p.Ile148Met, *TM6SF2* p.Glu167Lys, *LPL* 

p.Ser474Ter, and *ANGTPL4* p.Glu40Lys all associated with lower blood triglycerides and reduced risk for CAD. However, the blood TG-lowering alleles at *PNPLA3* and *TM6SF6* led to more fatty liver and higher risk for type diabetes. In contrast, the blood triglyceride-lowering alleles at *LPL* and *ANGPTL4* were neutral with respect to fatty liver and led to lower risk for type 2 diabetes. We confirmed the *LPL* observation using a phenome-wide association study approach in the UK Biobank, a population-based cohort study (Table S20). In UK Biobank, a one-SD decrease in TG mediated by *LPL* variants reduced risks for *both* type 2 diabetes and CAD (**Fig. 2**).

In summary, combining large-scale human genetic analysis with experimental evidence, we demonstrate: (1) 444 independent coding and non-coding variants at 250 loci as associated with plasma lipids; (2) the use of mouse models and genome editing to pinpoint causal genes and protein-altering variants, as with example of *A1CF* p.Gly398Ser; and (3) that *LPL* activation can be expected to lower triglycerides and reduce risks for *both* CAD and type 2 diabetes without increasing liver fat and thus be advantageous for patients with metabolic risk factors.

### **ONLINE METHODS**

#### **Study samples and phenotypes**

Seventy-three studies contributed association results for exome chip genotypes and plasma lipid levels. The outcomes were fasting lipid values in mg/dL [TC, HDL-C, LDL-C, TG] from the baseline, or earlier exam with fasting measures. If a study only had non-fasting levels then it contributed only to the TC and HDL-C analyses. LDL-C and TG analyses were only performed on fasting lipid values. Lipid lowering therapy with statins was not routinely used prior to the publication of the 4S study in 1994 which demonstrated the clinical benefit of statin therapy. Therefore, for data collected before 1994, no lipid medication adjustment was applied. For data collected after 1994, we adjusted the TC values for individuals on lipid medication by replacing their total cholesterol values by TC/0.8; this adjustment estimates the effect of statins on TC values. No adjustment was made on HDL-C or TG. LDL-C was calculated using the Friedewald equation for those with TG < 400 mg/dl (LDL-C = TC – HDL-C – (TG/5)). If TC was modified as described above for medication use after 1994, then modified TC was used in this formula. If only measured LDL-C was available in a study, we used LDL/0.7 for those on lipid-lowering medication when data were collected after 1994. TG values were natural log transformed. For each phenotype, residuals were obtained after accounting for age, age<sup>2</sup>, sex, principal components (as needed by each study, up to four), and inverse normal transform residuals were created for analysis. For studies ascertained on CAD case/control status, the two groups were modeled as separate studies.

### Genotyping and quality control

All studies assayed the Illumina or Affymetrix Human Exome array v1 or v1.1. Genotypes were determined from Zcall<sup>41</sup> or joint calling<sup>42</sup>. Individual studies performed the following quality control: call rate, heterozygosity, gender discordance, GWAS discordance (if GWAS data available), fingerprint concordance, if available, and PCA outliers.

# **Association analyses**

Each contributing cohort analyzed the ancestries within their cohorts separately and studies collected on case/control status analyzed cases separately from the controls. We performed both single variant and gene-level association tests. In the association analysis, we obtain residuals after controlling for sex, age, age<sup>2</sup> and up to 4 principal components as covariates. Studies that had related samples analyzed the association using linear mixed models with relatedness estimated from genome-wide SNPs or from pedigrees.

From each study, we collected single variant score statistics and their covariance matrix for variants in sliding windows across the genome. Summary association test statistics were generated using RAREMETALWORKER or RVTESTS. Using summary association statistics collected from each study, we performed meta-analysis of single variant association tests using the Mantel-Haenszel test and constructed burden, SKAT and variable threshold tests using the approach by Liu et al<sup>14</sup>. For burden and SKAT, we used minor allele frequency thresholds of 1% and 5% and for VT, we applied minor allele frequency threshold of 5%. In the SKAT test, variants are weighted according to their minor allele frequencies, using the beta kernel  $\beta$ (1,25).

Using covariance matrices between single variant association statistics, we were also able to perform conditional association analyses centrally, which distinguishes genuine signals from "shadows" of known loci. Details of the methods can be found in Liu et al<sup>14</sup>.

We centrally performed quality control for the data. We aligned study reported reference and alternative alleles with alleles reported in the NHLBI Exome Sequencing Project<sup>43</sup> and remove mis-labelled variant sites that can be strand ambiguous. For variant sites in each study, we removed variants that had call rate < 0.9 or had Hardy Weinberg *P* values <  $1 \times 10^{-7}$ . Finally, as additional checks, we visually inspected for each study the scatter plot of variant allele frequency against frequencies from ethnicity-matched populations in the 1000 Genomes Project<sup>44</sup>, and made sure that the strand and allele labels were well calibrated between studies.

Single variant associations with  $P < 2.1 \times 10^{-7}$  (0.05/242,289 variants analyzed) and genebased associations with  $P < 4.2 \times 10^{-07}$  (0.05/[20,000 genes \* 6 tests]) were considered significant. Novel loci were defined as being not within 1 megabase of a known lipid GWAS SNP. Additionally, linkage disequilibrium information was used to determine independent SNPs where a locus extended beyond 1 megabase. All novel loci reported in this manuscript are > 1 megabase from any previously reported locus and independent ( $r^2 < 0.2$  was required for variants within 3 megabases).

### Sequential forward selection

To identify independently associated variants for each known and newly identified locus, we performed sequential forward selection. We initialized the set of independently associated variants (denoted by  $\Phi$ ), starting with the top association signal in the locus. For each iteration, conditioning on variants in  $\Phi$ , we performed conditional association analyses for all remaining variants. If the top association signal after the conditional analysis remained significant, we added the top variant to the set  $\Phi$ , and then repeated the conditional association analysis. If the top variant after the conditional analysis was no longer significant, we stopped and reported

variants in the set  $\Phi$  as the final set of independent variants for that locus. We used the same single variant significance threshold (P < 2.1 x 10<sup>-7</sup>) to determine statistical significance with the sequential forward selection results (Fig. S3).

### Annotation

Sequence variants were annotated according to refSeq version 1.9, using the SEQMINER software (version 5.7)<sup>45</sup>. Transcript level annotations were obtained and prioritized. When multiple transcript level annotations were available, they were prioritized according to their functionality and deleteriousness. To implement gene-level association tests, the annotation with the highest priority was used (along with other filtering criteria such as minor allele frequencies) to determine the set of variants that are included.

#### Heritability and proportion of variance explained estimates

We estimated the proportion of variance explained by the set of 444 independently associated variants. The joint effects of variants in a locus were approximated by  $\hat{\beta}_{JOINT} = \mathbf{V}_{META}^{-1} \vec{U}_{META}$ , where  $\vec{U}_{META}$  is the single variant score statistics and  $\mathbf{V}_{META}^{-1}$  is the covariance matrix between them. The covariance between single variant genetic effects was approximated by the inverse of the variance-covariance matrix of score statistics, i.e.  $\mathbf{V}_{META}^{-1}$ . The phenotypic variance explained by the independently associated variants in a locus is given by  $\hat{\beta}_{JOINT}^{T}$  cov(G) $\hat{\beta}_{JOINT}$ , where G is the genotypes of the analyzed variants.

#### **Refinement of genome-wide association signals**

We sought to quantify what proportion of GWAS loci might be due to a protein-altering variant and, therefore, directly identify a functional gene. We made the assumption that a proteinaltering variant is the most likely causal variant for each region if it is the top signal, explains the signal, or is independent of the original signal. To identify putative functional coding variants accounting for the effects at known lipid loci, we performed reciprocal conditional analyses to control for the effects of known lipid GWAS or coding variants within 500kb, as this was the maximum distance for variants within the covariance matrix. Loci where coding variants are the most significant signals were considered as "coding as top". Loci where the initial GWAS variants had conditional P > 0.01 were considered to be explained by the coding variants. Loci where the coding variants had conditional  $P < 2.1 \times 10^{-7}$  were considered to be independent of the initial GWAS signals.

# *JAK2* p.Val617Phe and plasma cholesterol in a mouse model

*Jak2* p.Val617Phe MxCre mice were created and reported previously<sup>26</sup>. Bone marrow cells from the WT or *JAK2* p.Val617Phe MxCre mice, both treated with poly I:C, were transplanted into irradiated  $Ldlr^{-/-}$  recipients. After four weeks of recovery, the  $Ldlr^{-/-}$  recipient mice were fed a Western diet (TD88137, Harlan Teklad) for 8 weeks. Plasma was collected and 250 microliter of polled plasma from 7 WT $\rightarrow$ Ldlr-/- or 7 *Jak2* Val617Phe $\rightarrow$ Ldlr-/- recipient was subjected to fast protein liquid chromatography on Sepharose CL-6B size exclusion column. Total cholesterol content in each fraction was assessed by Cholesterol E kit (Wako Diagnostics).

#### Validation of A1CF with CRISPR-Cas9 in human cells

To knock out *A1CF* in Huh7 and HepG2 human hepatoma cells, three CRISPRs (sgRNA sequences are as follows: 5'-GGATGCCCCCAACAACAATAAT-3'; 5'-

TTGGTTTTATCTGCAGCGCT-3' and 5'-ATGACTCTCATACTCCACGA-3') targeting exon 4 of the *A1CF* gene were constructed by using the lentiviral vector lentiGuide-Puro. Packaged viruses were used to transduce the cells expressing Cas9 for 16 hours. Subsequently, cells were cultured in the presence of 5  $\mu$ g/ml puromycin for five days before splitting for assays. Cells for APOB secretion assay were cultured for 18 hours in serum-free medium, then the amount of APOB100 in medium was measured using an ELISA kit (MABTECH) according to the manufacturer's instructions.

In a rescue experiment, to avoid cutting of the *A1CF* coding region on the recombinant plasmids by previously designed exon-targeting CRISPRs, four new CRISPRs targeting introns flanking exon 4 were applied to deplete endogenous *A1CF*. The sequences for those sgRNAs are as follows: 5'-GGTTGAAAATGTATGGCTGT-3'; 5'-TGATTGATGAGAATCCTTGT-3'; 5'-TTTTCTTTGCGGAGGCAGGC-3' and 5'-GCAGAGAAGTCTCATCTTTG-3'. The *A1CF* p.Gly398Ser variant was generated by using overlapping PCR and confirmed by Sanger sequencing. Both wild-type and the *A1CF* p.Gly398Ser variant were constructed into lentiviral plasmids, respectively. After transduction, cells were cultured for 48 hours in the presence of 100 ng/ml doxycycline to induce recombinant expression of A1CF or p.Gly398Ser variant before performing different assays.

# A1cf p.Gly390Ser knock-in mice

All procedures used for animal studies were approved by Harvard University's Faculty of Arts and Sciences Institutional Animal Care and Use Committee and were consistent with local, state, and federal regulations as applicable. Knock-in mice were generated using a guide RNA designed to target the orthologous site of the *A1CF* p.Gly390Ser variant. In vitro transcribed Cas9 mRNA (100 ng/µL; TriLink BioTechnologies) and guide RNA (50 ng/µL) were co-injected with 100 ng/µL single-strand DNA oligonucleotide (Integrated DNA Technologies): (5'-GTTTGATTCAGTTTAATGCCTTCTTACATTCCTAGAAATTTACATGAATGTCCCTGTA GGGGCTGCGGG

GGTGAGAGGTCTGGGCAGCCGTGGCTATTTGGCATACACAGGCCTGGGTCGAGGAT ACCACGTCAAAGGAGACAAGAGAGAGAAGACAAACTCTA-3') into the cytoplasm of fertilized oocytes from C57BL/6J mice. Genomic DNA samples from founder mice were screened for knock-in mutations by PCR and confirmed by Sanger sequencing. Positive mice were bred with C57BL/6J mice to generate wild-type and homozygous knock-in mice. Male colony mates at 12 weeks of age were used for lipid measurements. Blood samples were collected from the lateral tail vein following an overnight fast. Plasma triglyceride levels were measured using Infinity Triglycerides Reagent (Thermo Fisher) according to the manufacturers' instructions.

### Intersection of lipid association signals with AMD, CAD, and T2D

To estimate the association of loss-of-function variants in *HBB* with cholesterol levels, participants from the following two consortia were studied: the Global Lipids Genetics Consortium and the Myocardial Infarction Genetics Consortium (MIGen, 27,939 participants in 12 cohorts). A rare loss-of-function variant in *HBB* (c.92+1G>A, rs33971440) was genotyped in participants from the Global Lipids Genetics Consortium Exome consortium. This variant was pooled with sequence data for the *HBB* gene in MIGen, available in 19,434 participants with blood cholesterol measurements. The association of loss-of-function variants with cholesterol was estimated using linear regression with adjustment for age, sex and up to five principal components of ancestry. Estimates from genotyped and sequence data were pooled using inverse variance weighted fixed effects meta-analysis.

To estimate the association of loss-of-function variants in *HBB* with CAD, participants from the following two consortia were studied: the CARDIoGRAM Exome Consortium (69,087 participants from 20 studies) and MIGen (12384 CAD cases and 15547 controls from 12 studies). 69,086 individuals who were genotyped for the c.92+1G>A variant in CARDIoGRAM Exome were pooled with sequence data for *HBB* from 27,931 individuals in MIGen. The association of loss-of-function variants with CAD was estimated using logistic regression with adjustment for age, sex and up to five principal components of ancestry. Estimates were pooled using inverse variance weighted fixed effects meta-analysis. To estimate the association of loss of function variants in *HBB* with hemoglobin and hematocrit levels, estimates from an exome chip analysis of red blood cell traits (24 814 individuals) were used<sup>8</sup>.

For 168 variants independently and significantly associated with HDL-C and a MAF > 1%, we looked up the association evidence in 16,144 age-related macular degeneration cases and 17,832 controls with exome chip genotypes<sup>46</sup>.

For 132 independently and significantly associated LDL-C variants and MAF > 1%, we looked up the association evidence in: (1) up to 120,575 individuals with and without CAD and exome chip genotypes (42,335 cases and 78,240 controls)<sup>47</sup>; and (2) up to 69,870 individuals with and without type 2 diabetes. Only 113 of the 132 LDL variants were available in the type 2 diabetes results. We used a Bonferroni correction for 132 variants to determine significance of the results (alpha =  $4.0 \times 10^{-4}$ ).

### Association of PCSK9 R46L with type 2 diabetes

For evaluating the association of *PCSK9* R46L with risk of type 2 diabetes, we considered a total of 42,011 type 2 diabetes cases and 180,834 controls from 30 studies from populations of European ancestry (Table S17). The variant was directly genotyped in all studies using the Metabochip or the Exome array. Sample and variant quality control was performed within each study as described previously<sup>48-51</sup>. Within each study, the variant was tested for association with type 2 diabetes under an additive model after adjustment for study-specific covariates, including principal components to adjust for population structure. Association summary statistics for the variant for each study was corrected for residual population structure using the genomic control inflation factor as described previously<sup>48-50</sup>. We then combined association summary statistics for the variant across studies via fixed-effects inverse-variance weighted meta-analysis.

#### TG variants, lipids, fatty liver, type 2 diabetes, and CAD

Exome chip results for four variants (*LPL* p.Ser474Ter [rs328], *ANGPTL4* p.Glu40Lys [rs116843064], *PNPLA3* p.Ile148Met [rs738409], and *TM6SF2* p.Glu167Lys [rs58542926]) were obtained from the following sources:

1. lipids: current analysis

2. fatty liver: Between 2002 and 2005, 1,400 individuals from the Framingham Offspring Study and 2,011 individuals from third generation underwent multi-detector computed tomograms (CT) on which we evaluated liver attenuation as previously described<sup>52</sup>. We tested the association of TG variants with CT liver fat after inverse normal transformation. Covariates in the regression models included age, age<sup>2</sup>, and gender. A similar analysis was conducted in 3,293 participants of

European ancestry from BioImage study<sup>53</sup>. Association results for liver attenuation from the Framingham and BioImage studies were combined through fixed-effects inverse-variance weighted meta-analysis (Table S20).

3. type 2 diabetes: ExTexT2D Consortium<sup>54</sup>

4. CAD: published results from the Myocardial Infarction Genetics and CARDIoGRAM Exome Consortia study<sup>47</sup>.

**Fig. 1:** *A1CF* **p.Gly398Ser mutant leads to increased APOB100 secretion. A**, Western blot showing the depletion of endogenous A1CF levels via CRISPR/Cas9 system in both Huh7 and HepG2 cells. **B and C**, Lack of A1CF leads to reduced APOB100 secretion in Huh7 (B) and HepG2 (C) human hepatoma cells. **D**, Recombinantly overexpressed A1CF p.Gly398Ser variant led to significantly increased APOB100 secretion compared to A1CF or GFP control in both Huh7 wild-type and A1CF knockout cells (labeled as A1CF KO), respectively. The error bars show SD from experiments with biological replicates, N=6 for B, C and D. Statistically significant differences are marked (\*p<0.05, \*\*p<0.01).



Fig. 2: Association of genetically-lowered triglycerides by LPL variants with a range of phenotypes. Estimates were derived in UK Biobank using logistic regression, adjusting for age, sex, ten principal components of ancestry and an indicator variable for array type. Effect estimates are for a 1 standard deviation lower plasma triglycerides. Definitions for all outcomes are provided in Table S16.



	Cases	Controls	OR per SD	Lowe	r TG	P Value
Cardiovascular						
Coronary Heart Disease	4461	107877	+	0.78	[0.69; 0.88]	0.0001
Stroke	2066	110272		1.06	[0.90; 1.25]	0.51
Heart Fallure	599	111739		0.83	[0.60; 1.13]	0.24
Atrial Fibrillation	2208	110130	-	0.93	[0.79; 1.09]	0.35
Aortic Stenosis	193	112145	++++	0.63	[0.36; 1.13]	0.12
Peripheral Vascular disease	692	111646		0.85	[0.64; 1.14]	0.28
Venous Thromboembolism	3306	109032	+	1.00	[0.87; 1.14]	0.97
Gastrointestinal disease						
Inflammatory Bowel Disease	1021	111317	+	1.17	[0.93; 1.48]	0.17
Gastric Reflux	4881	107457	+	1.03	[0.92; 1.14]	0.64
Galistones	1831	110507		1.13	[0.95; 1.34]	0.17
Irritable Bowel Syndrome	2679	109659	-	0.93	[0.80; 1.08]	0.36
Endocrine						
Type 2 Diabetes	5741	112338	-	0.88	[0.79; 0.97]	0.014
Gestational Diabetes	56	112282	••• <b>†</b> •	0.73	[0.26; 2.08]	0.56
Hyperthyroidlusm	868	111470	+	1.13	[0.88; 1.45]	0.33
Hypothyroldism	5433	106905	÷	1.05	[0.95; 1.17]	0.35
Gout	1612	110726	+	1.00	[0.83; 1.21]	0.98
Renal/urological						
Enlarged Prostate	1573	110765	·	1.08	[0.89; 1.30]	0.45
Uterine Fibroids	1634	110704	+	1.00	[0.83; 1.20]	0.99
Neurological/psychiatric						
Migraine	3161	109177	+	1.02	[0.89; 1.17]	0.74
Depression	6667	105671	노	1.06	[0.97; 1.17]	0.20
Anxiety	1545	110793	-F	1.02	[0.84; 1.23]	0.84
Musculooskeletal						
Backpain	584	111754		1.03	[0.76; 1.40]	0.85
Jointpain	377	111961	_	0.99	[0.68; 1.46]	0.97
Osteoporosis	1740	110598	+	1.02	[0.85; 1.22]	0.85
Osteoarthritis	9693	102645	±	0.96	[0.89; 1.05]	0.38
Sciatica	1035	111303	+	1.00	[0.79; 1.27]	0.98
Prolapsed Disc	1856	110482	+	1.04	[0.87; 1.24]	0.65
Respiratory						
Asthma	13941	98397	÷	0.99	[0.93; 1.06]	0.84
COPD/Emphysema	2363	109975		0.93	[0.80; 1.09]	0.40
Pneumonia	1581	110757		1.05	[0.87; 1.26]	0.63
Hayfever	6263	106075	<b>声</b>	1.06	[0.96; 1.17]	0.24
Cancer						
Lung Cancer	115	112223	• • • • •	0.92	[0.45; 1.86]	0.81
Breast Cancer	2383	109955	- <del></del> ·	0.90	[0.77; 1.06]	0.20
Colorectal Cancer	616	111722	++++	1.16	[0.86; 1.56]	0.32
Skin Cancer	2482	109856	+	0.96	[0.82; 1.12]	0.59
Prostate Cancer	840	111498	-+	0.99	[0.76; 1.28]	0.92
Cervical Cancer	872	111466	- <b> -</b>	1.09	[0.85; 1.40]	0.49
Other Cancer	2409	109929		0.88	[0.75; 1.03]	0.10
Any Cancer	9530	102808	4	0.94	[0.87; 1.02]	0.17

0.5 0.751 1.5

Table 1: Protein-altering variants at novel loci associated with lipid levels									
Chromosome: position (hg19)	rs ID	Alleles (reference/ alternative)	Gene	Protein change	Ν	Frequency alternative allele	Trait	P value	Beta±SE
Total Cholestero	l								
2:101627925	rs1062062	C/T	TBC1D8	p.Gly954Arg	292898	0.12	TC	1×10-7	-0.021±0.0040
4:69343287	rs976002	A/G	TMPRSS11E	p.Tyr303Cys	293961	0.23	TC LDL-C	5×10 <sup>-20</sup> 3×10 <sup>-12</sup>	0.029±0.0031, 0.023±0.0033
4:155489608	rs6054	C/T	FGB	p.Pro206Leu	307997	0.0038	TC TG	5×10 <sup>-12</sup> 3×10 <sup>-11</sup>	0.14±0.021, 0.14±0.021
9:5073770	rs77375493	G/T	JAK2	p.Val617Phe	188412	0.0011	TC LDL-C	1×10 <sup>-11</sup> 2×10 <sup>-9</sup>	-0.32±0.047, -0.30±0.049
9:117166246	rs2274159	A/G	DFNB31	p.Val400Ala	319677	0.48	TC	2×10-7	0.013±0.0026
17.8216468	rs871841	T/C	ARHGEF15	n Leu277Pro	298725	0.52	TC	2×10 <sup>-8</sup>	$0.015\pm0.0026$
19.18304700	rs874628	A/G	MPV17L2	n Met72Val	319677	0.26	TC	2×10-7	$0.012 \pm 0.0020$ 0.015 \pm 0.0029
LDL Cholesterol	10071020	120		pinienziia	515011	0.20	10	2 10	0.010-0.0025
1.155106227	rs4745	A/T	EFNA1	n Asn137Val	291361	0.49	LDL-C	5×10 <sup>-8</sup>	-0.015±0.0027
4.187120211	rs13146272	C/A	CYP4V2	n Gln259Lvs	295826	0.62	LDL-C	1×10-7	-0.015+0.0027
5.176520243	rs351855	G/A	FGFR4	n Glv388Arg	233058	0.29	LDL-C	4×10 <sup>-8</sup>	$-0.018\pm0.0023$
9.139368953	rs3812594	G/A	SEC16A	n Arg1039Cvs	293723	0.24	LDL-C	2×10 <sup>-8</sup>	-0.018+0.0031
10.118397971	rs10885997	A/G	PNLIPRP2	n Gln387Arg	258146	0.41	LDL-C	9×10 <sup>-8</sup>	0.015+0.0029
10:124610027	rs1891110	G/A	FAM24B	n Pro2Leu	295826	0.55	LDL-C	8×10 <sup>-15</sup>	$0.013 \pm 0.0025$ 0.021 \pm 0.0026
10.12101002,	1010)1110	0,11		pirrozitou	2,0020	0.00	TC	$2 \times 10^{-13}$	$0.019\pm0.0025$
12.72179446	rs61754230	C/T	RAB21	n Ser224Phe	292762	0.015	LDL-C	1×10 <sup>-7</sup>	$0.057\pm0.011$
14:94844947	rs28929474	C/T	SERPINA1	p.Glu366Lys	290263	0.015	LDL-C TC	$4 \times 10^{-14}$ $6 \times 10^{-14}$	0.081±0.011, 0.078+0.010
17:26694861	rs704	G/A	VTN	p.Thr400Met	295826	0.49	LDL-C	6×10 <sup>-16</sup> 1×10 <sup>-8</sup>	0.021±0.0026, 0.015±0.0025
10.42584058	rs201596848	C/T	ZNE574	n Arg734Cvs	273744	0.0014		5×10 <sup>-12</sup>	$-0.255\pm0.037$
Triglycerides	13201370040	0/1	2111 37 4	p./ iig/ 54Cy5	213144	0.0014	LDL C	5.10	0.235-0.057
2·202122995	rs3769823	A/G	CASP8	n Lys14Arg	295956	0.69	TG	1×10-9	0.017+0.0028
5:131008194	rs26008	T/C	ENIP1	n Gln620Arg	305699	0.92	TG	5×10-9	-0.028+0.0048
10:52573772	rs41274050	C/T	AICF	p.Gly398Ser	299984	0.0072	TG	4×10 <sup>-11</sup>	0.10±0.015,
							TC	1×10-7	0.08±0.015
13:45970147	rs138358301	A/G	SLC25A30	p.Phe280Leu	301087	0.0035	TG	3×10-11	0.15±0.022
15:40751555	rs3803357	C/A	BAHD1	p.Gln298Lys	305699	0.55	TG	1×10-10	-0.017±0.0026
17:17409560	rs7946	C/T	PEMT	p.Val212Met	304420	0.67	TG	1×10-8	-0.016±0.0029
20:56140439	rs41302559	G/A	РСК1	p.Arg483Gln	299984	0.0021	TG	9×10-8	-0.154±0.029
22:17625915	rs35665085	G/A	CECR5	p.Thr149Met	302582	0.050	TG	5×10-8	0.032±0.0059
HDL Cholesterol									
2:272203	rs11553746	C/T	ACP1	p.Thr95Ile	313148	0.33	HDL-C	5×10-8	0.015±0.0027
2:54482553	rs17189743	G/A	TSPYL6	p.Arg246Cys	314415	0.029	HDL-C	2×10-7	0.040±0.0076
2:179309165	rs75862065	G/A	PRKRA	p.Pro116Leu	105490	0.29	HDL-C	2×10-7	0.026±0.0050
3:48229366	rs146179438	C/A	CDC25A	p.Gln25His	288306	0.020	HDL-C	3×10 <sup>-11</sup>	-0.063±0.0095
5:176637576	rs28932178	T/C	NSD1	p.Ser457Pro	310567	0.17	HDL-C	8×10-9	0.020±0.0035
11:64031241	rs35169799	C/T	PLCB3	p.Ser778Leu	314415	0.060	HDL-C TG	$4 \times 10^{-13},$ $3 \times 10^{-12}$	-0.039±0.0054, 0.038±0.0055
11:68703959	rs622082	A/G	IGHMBP2	p.Thr671Ala	316391	0.31	HDL-C	6×10 <sup>-10</sup>	-0.017±0.0028

16:4755108	rs78074706	G/A	ANKS3	p.Arg286Trp	315298	0.022	HDL-C	1×10-9	-0.053±0.0087
16:69385641	rs76116020	A/G	TMED6	p.Phe6Leu	310822	0.033	HDL-C	7×10-9	-0.041±0.0071
17:40257163	rs2074158	T/C	DHX58	p.Gln425Arg	244331	0.19	HDL-C	1×10-7	-0.020±0.0038

Tally liver, type 2 ul	abeles, and corolla	Ty aftery disease.		
Gene	LPL	ANGPTL4	PNPLA3	TM6SF2
Variant	p.Ser474Ter	p.Glu40Lys	p.Ile148Met	p.Glu167Lys
Effect Allele	Х	Κ	Met	Κ
Frequency	10%	2%	23%	7%
		Blood triglycerides	5	
Effect Direction	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$
Beta	-0.18	-0.27	-0.018	-0.12
(SE)	(0.004)	(0.010)	(0.003)	(0.005)
Р	$P < 1 \ge 10^{-323}$	$P = 4 \ge 10^{-175}$	$P = 4 \ge 10^{-9}$	$P = 4 \ge 10^{-125}$
		Blood LDL cholester	rol	
Effect Direction	-	$\downarrow$	$\downarrow$	$\downarrow$
Beta	0.013	-0.004	-0.018	-0.103
(SE)	(0.004)	(0.010)	(0.003)	(0.005)
Р	P = 0.005	P = 0.70	$P = 1 \ge 10^{-8}$	$P = 7 \ge 10^{-93}$
		Fatty liver		
Effect Direction	-	-	$\uparrow$	$\uparrow$
Beta*	0.026	0.112	-0.247	-0.250
(SE)	(0.031)	(0.068)	(0.022)	(0.036)
Р	P = 0.41	P = 0.10	$P = 2 \ge 10^{-30}$	$P = 5 \ge 10^{-12}$
		Type 2 diabetes		
Effect Direction	$\downarrow$	$\downarrow$	$\uparrow$	$\uparrow$
Beta	-0.049	-0.097	0.039	0.07
(SE)	(0.01)	(0.040)	(0.005)	(0.01)
Р	$P = 7 \ge 10^{-9}$	$P = 1 \ge 10^{-4}$	$P = 2 \ge 10^{-10}$	$P = 5 \ge 10^{-12}$
		Coronary artery dise	ase	
Effect Direction	$\downarrow$	$\downarrow$	$\downarrow$	$\downarrow$
Beta	-0.074	-0.160	-0.044	-0.047
(SE)	(0.014)	(0.026)	(0.008)	(0.013)
Р	$P = 4 \ge 10^{-7}$	$P = 2 \ge 10^{-10}$	$P = 4 \ge 10^{-8}$	$P = 3 \ge 10^{-4}$

**Table 2.** Impact of genes involved in hepatic production of triglyceride-rich lipoproteins (*PNPLA3*, *TM6SF2*) verus lipolysis pathway genes (*LPL*, *ANGPTL4*) on related metabolic traits - blood lipids, fatty liver, type 2 diabetes, and coronary artery disease.

\*A negative beta reflects liver attenuation on computed tomography which is indicative of *higher* liver fat Association results for lipids are derived from present study

Association results for type 2 diabetes are from<sup>54</sup>

Association results for coronary artery disease are from<sup>47</sup>

# References

- 1. Teslovich, T.M. *et al.* Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* **466**, 707-13 (2010).
- 2. Asselbergs, F.W. *et al.* Large-scale gene-centric meta-analysis across 32 studies identifies multiple lipid loci. *Am J Hum Genet* **91**, 823-38 (2012).
- 3. Global Lipids Genetics Consortium, *et al.* Discovery and refinement of loci associated with lipid levels. *Nat Genet* **45**, 1274-83 (2013).
- 4. Albrechtsen, A. *et al.* Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. *Diabetologia* **56**, 298-310 (2013).
- 5. Peloso, G.M. *et al.* Association of Low-Frequency and Rare Coding-Sequence Variants with Blood Lipids and Coronary Heart Disease in 56,000 Whites and Blacks. *American journal of human genetics* **94**, 223-232 (2014).
- 6. Surakka, I. *et al.* The impact of low-frequency and rare variants on lipid levels. *Nat Genet* **47**, 589-97 (2015).
- 7. Tang, C.S. *et al.* Exome-wide association analysis reveals novel coding sequence variants associated with lipid traits in Chinese. *Nat Commun* **6**, 10206 (2015).
- 8. Musunuru, K. *et al.* From noncoding variant to phenotype via SORT1 at the 1p13 cholesterol locus. *Nature* **466**, 714-9 (2010).
- 9. Burkhardt, R. *et al.* Trib1 is a lipid- and myocardial infarction-associated gene that regulates hepatic lipogenesis and VLDL production in mice. *J Clin Invest* **120**, 4410-4 (2010).
- 10. Voight, B.F. *et al.* Plasma HDL cholesterol and risk of myocardial infarction: a mendelian randomisation study. *Lancet* **380**, 572-80 (2012).
- 11. Do, R. *et al.* Common variants associated with plasma triglycerides and risk for coronary artery disease. *Nat Genet* **45**, 1345-52 (2013).
- 12. Lu, X. *et al.* Exome chip meta-analysis identifies novel loci and East Asian-specific coding variants contributing to lipid levels and coronary artery disease. *Nature Genetics* (2017).
- 13. Feng, S., Liu, D., Zhan, X., Wing, M.K. & Abecasis, G.R. RAREMETAL: fast and powerful meta-analysis for rare variants. *Bioinformatics* **30**, 2828-9 (2014).
- 14. Liu, D.J. *et al.* Meta-analysis of gene-level tests for rare variant association. *Nat Genet* **46**, 200-4 (2014).
- 15. Locke, A.E. *et al.* Genetic studies of body mass index yield new insights for obesity biology. *Nature* **518**, 197-206 (2015).
- 16. de Bakker, P.I. *et al.* Practical aspects of imputation-driven meta-analysis of genome-wide association studies. *Hum Mol Genet* **17**, R122-8 (2008).
- Holmen, O.L. *et al.* Systematic evaluation of coding variation identifies a candidate causal variant in TM6SF2 influencing total cholesterol and myocardial infarction risk. *Nat Genet* 46, 345-51 (2014).
- 18. Klarin, D. *et al.* Genetics of lipids in >300,000 participants in the Million Veteran Program. *Under Review*.
- 19. Shen, X. *et al.* Identification of genes affecting apolipoprotein B secretion following siRNA-mediated gene knockdown in primary human hepatocytes. *Atherosclerosis* **222**, 154-7 (2012).

- 20. Baxter, E.J. *et al.* Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* **365**, 1054-61 (2005).
- 21. James, C. *et al.* A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* **434**, 1144-8 (2005).
- 22. Kralovics, R. *et al.* A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* **352**, 1779-90 (2005).
- 23. Levine, R.L. *et al.* Activating mutation in the tyrosine kinase JAK2 in polycythemia vera, essential thrombocythemia, and myeloid metaplasia with myelofibrosis. *Cancer Cell* **7**, 387-97 (2005).
- 24. Jaiswal, S. *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* **371**, 2488-98 (2014).
- 25. Jaiswal, S. *et al.* Clonal Hematopoiesis and Risk of Atherosclerotic Cardiovascular Disease. *N Engl J Med* **377**, 111-121 (2017).
- Mullally, A. *et al.* Physiological Jak2V617F expression causes a lethal myeloproliferative neoplasm with differential effects on hematopoietic stem and progenitor cells. *Cancer Cell* 17, 584-96 (2010).
- 27. Lellek, H. *et al.* Purification and molecular cloning of a novel essential component of the apolipoprotein B mRNA editing enzyme-complex. *J Biol Chem* **275**, 19848-56 (2000).
- 28. Mehta, A., Kinter, M.T., Sherman, N.E. & Driscoll, D.M. Molecular cloning of apobec-1 complementation factor, a novel RNA-binding protein involved in the editing of apolipoprotein B mRNA. *Mol Cell Biol* **20**, 1846-54 (2000).
- 29. Galloway, C.A., Ashton, J., Sparks, J.D., Mooney, R.A. & Smith, H.C. Metabolic regulation of APOBEC-1 complementation factor trafficking in mouse models of obesity and its positive correlation with the expression of ApoB protein in hepatocytes. *Biochim Biophys Acta* **1802**, 976-85 (2010).
- 30. Galanello, R. & Origa, R. Beta-thalassemia. Orphanet J Rare Dis 5, 11 (2010).
- 31. Fessas, P., Stamatoyannopoulos, G. & Keys, A. Serum-cholesterol and thalassemia trait. *Lancet* **1**, 1182-3 (1963).
- 32. Sidore, C. *et al.* Genome sequencing elucidates Sardinian genetic architecture and augments association analyses for lipid and blood inflammatory markers. *Nat Genet* **47**, 1272-81 (2015).
- 33. Chen, W. *et al.* Genetic variants near TIMP3 and high-density lipoprotein-associated loci influence susceptibility to age-related macular degeneration. *Proc Natl Acad Sci U S A* **107**, 7401-6 (2010).
- 34. Neale, B.M. *et al.* Genome-wide association study of advanced age-related macular degeneration identifies a role of the hepatic lipase gene (LIPC). *Proc Natl Acad Sci U S A* **107**, 7395-400 (2010).
- 35. Wang, Y.F. *et al.* CETP/LPL/LIPC gene polymorphisms and susceptibility to age-related macular degeneration. *Sci Rep* **5**, 15711 (2015).
- 36. Cheng, C.Y. *et al.* New loci and coding variants confer risk for age-related macular degeneration in East Asians. *Nat Commun* **6**, 6063 (2015).
- 37. Momozawa, Y. *et al.* Low-frequency coding variants in CETP and CFB are associated with susceptibility of exudative age-related macular degeneration in the Japanese population. *Hum Mol Genet* **25**, 5027-5034 (2016).

- 38. Lotta, L.A. *et al.* Association Between Low-Density Lipoprotein Cholesterol-Lowering Genetic Variants and Risk of Type 2 Diabetes: A Meta-analysis. *JAMA* **316**, 1383-1391 (2016).
- 39. Schmidt, A.F. *et al.* PCSK9 genetic variants and risk of type 2 diabetes: a mendelian randomisation study. *Lancet Diabetes Endocrinol* (2016).
- 40. Ference, B.A. *et al.* Variation in PCSK9 and HMGCR and Risk of Cardiovascular Disease and Diabetes. *N Engl J Med* **375**, 2144-2153 (2016).
- 41. Goldstein, J.I. *et al.* zCall: a rare variant caller for array-based genotyping: genetics and population analysis. *Bioinformatics* **28**, 2543-5 (2012).
- 42. Grove, M.L. *et al.* Best Practices and Joint Calling of the HumanExome BeadChip: The CHARGE Consortium. *PLoS One* **8**, e68095 (2013).
- 43. Tennessen, J.A. *et al.* Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science* **337**, 64-9 (2012).
- 44. 1000 Genomes Project Consortium, *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56-65 (2012).
- 45. Zhan, X. & Liu, D.J. SEQMINER: An R-Package to Facilitate the Functional Interpretation of Sequence-Based Associations. *Genet Epidemiol* **39**, 619-23 (2015).
- 46. Fritsche, L.G. *et al.* A large genome-wide association study of age-related macular degeneration highlights contributions of rare and common variants. *Nat Genet* **48**, 134-43 (2016).
- 47. Myocardial Infarction Genetics & CARDIoGRAM exome consortia Investigators. Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. *N Engl J Med* **374**, 1134-44 (2016).
- 48. Morris, A.P. *et al.* Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. *Nat Genet* **44**, 981-90 (2012).
- 49. Gaulton, K.J. *et al.* Genetic fine mapping and genomic annotation defines causal mechanisms at type 2 diabetes susceptibility loci. *Nat Genet* **47**, 1415-25 (2015).
- 50. Fuchsberger, C. et al. The genetic architecture of type 2 diabetes. Nature 536, 41-7 (2016).
- 51. The UK Biobank. Genotyping and quality control of UK Biobank, a large-scale, extensively phenotyped prospective resource. (<u>http://www.ukbiobank.ac.uk/wp-content/uploads/2014/04/UKBiobank genotyping QC documentation-web.pdf</u>, 2015).
- 52. Speliotes, E.K. *et al.* Genome-wide association analysis identifies variants associated with nonalcoholic fatty liver disease that have distinct effects on metabolic traits. *PLoS Genet* **7**, e1001324 (2011).
- 53. Baber, U. *et al.* Prevalence, impact, and predictive value of detecting subclinical coronary and carotid atherosclerosis in asymptomatic adults: the BioImage study. *J Am Coll Cardiol* **65**, 1065-74 (2015).
- 54. Mahajan, A. *et al.* Refining The Accuracy Of Validated Target Identification Through Coding Variant Fine-Mapping In Type 2 Diabetes. *bioRxiv* (2017).