

Research

Open Access

Differentiation of adult-type Leydig cells occurs in gonadotrophin-deficient mice

PJ Baker¹, H Johnston¹, M Abel², HM Charlton² and PJ O'Shaughnessy*¹

Address: ¹Institute of Comparative Medicine, University of Glasgow, Veterinary School, Bearsden Rd, Glasgow G61 1QH, UK and ²Department of Human Anatomy, University of Oxford, South Parks Rd, Oxford OX1 3QX, UK

Email: PJ Baker - p.j.baker@vet.gla.ac.uk; H Johnston - h.johnston@vet.gla.ac.uk; M Abel - margaret.abel@human-anatomy.oxford.ac.uk; HM Charlton - harry.charlton@human-anatomy.oxford.ac.uk; PJ O'Shaughnessy* - p.j.oshaughnessy@vet.gla.ac.uk

* Corresponding author

Published: 5 February 2003

Received: 23 January 2003

Reproductive Biology and Endocrinology 2003, 1:4

Accepted: 5 February 2003

This article is available from: <http://www.RBEj.com/content/1/1/4>

© 2003 Baker et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

During mammalian testis development distinct generations of fetal and adult Leydig cells arise. Luteinising hormone (LH) is required for normal adult Leydig cell function and for the establishment of normal adult Leydig cell number but its role in the process of adult Leydig cell differentiation has remained uncertain. In this study we have examined adult Leydig cell differentiation in gonadotrophin-releasing hormone (GnRH)-null mice which are deficient in circulating gonadotrophins. Adult Leydig cell differentiation was assessed by measuring expression of mRNA species encoding four specific markers of adult Leydig cell differentiation in the mouse. Each of these markers (3 β -hydroxysteroid dehydrogenase type VI (3 β HSD VI), 17 β -hydroxysteroid dehydrogenase type III (17 β HSD III), prostaglandin D (PGD)-synthetase and oestrogen sulphotransferase (EST)) is expressed only in the adult Leydig cell lineage in the normal adult animal. Real-time PCR studies showed that all four markers are expressed in adult GnRH-null mice. Localisation of 3 β HSD VI and PGD-synthetase expression by *in situ* hybridisation confirmed that these genes are expressed in the interstitial tissue of the GnRH-null mouse. Treatment of animals with human chorionic gonadotrophin increased expression of 3 β HSD VI and 17 β HSD III within 12 hours further indicating that differentiated, but unstimulated cells already exist in the GnRH-null mouse. Thus, while previous studies have shown that LH is required for adult Leydig cell proliferation and activity, results from the present study show that adult Leydig cell differentiation will take place in animals deficient in LH.

Background

In mice, as in most mammalian species, two generations of Leydig cells arise during normal testis development. A "fetal" population which appears shortly after testis differentiation *in utero* and an "adult" population which arises after birth around day 7 [1–3]. Morphological and functional maturation of the adult Leydig cells is critically dependent on luteinising hormone (LH) stimulation and this is clearly seen in mice lacking either circulating LH or

the LH-receptor [4–7]. In these animals Leydig cell number fails to develop normally after birth and circulating androgen levels are low or undetectable [6–9]. In contrast, testicular androgen levels and Leydig cell number is normal during fetal development in animals lacking gonadotrophins [8,9] showing that the fetal population of Leydig cells does not require gonadotrophins during this period. Despite the clear role for LH in proliferation and maturation of adult Leydig cells in the postnatal testis, it

is not known whether this hormone is essential for the onset of adult Leydig cell differentiation. Early studies suggested that LH is required for the differentiation process [10–12] while more recent, immunohistochemical studies have reported that specific steroidogenic enzymes are detectable in rat adult Leydig cell precursors before LH-receptors are detectable [13]. This evidence would suggest that Leydig cell differentiation begins in the absence of LH-stimulation although it remains possible that low, undetectable levels of receptor are present prior to differentiation. Certainly, Leydig cell precursor cells express the LH-receptor gene prior to apparent differentiation although it is not clear whether transcripts are translated or produce functional protein [14–16].

It has been shown previously that in the mouse mRNA species encoding β 3-hydroxysteroid dehydrogenase type VI (β 3HSD VI), 17β -hydroxysteroid dehydrogenase type III (17β HSD III), prostaglandin (PG) D-synthetase, estrogen sulfotransferase (EST) and vascular cell adhesion molecule 1 (VCAM 1) are expressed only in the adult Leydig cell population and not in the fetal population [1,17–20]. It is possible, therefore, to use these markers to examine the role of LH in the initiation of adult Leydig cell differentiation using mice which lack gonadotrophin-stimulation. Expression of these markers in mice lacking gonadotrophins would be a clear demonstration that Leydig cell differentiation does not require LH stimulation. It would be anticipated, however, that even if adult Leydig cells exist in the absence of LH-stimulation the expression of these specific markers will be very low or non-existent since overall Leydig cell activity is determined by LH [21]. Treatment of gonadotrophin-deficient mice with hCG is likely, however, to lead to a rapid increase in expression of markers within 24 hours if adult Leydig cells already exist. If the cells have not differentiated then a more prolonged period of treatment is likely to be required to induce differentiation of Leydig cells and expression of markers. For these reasons the expression of adult Leydig cell markers has been examined in the gonadotrophin-deficient gonadotrophin-releasing hormone (GnRH)-null (*hpg*) mouse. This animal has an advantage over receptor-deficient models in that the gonads remain responsive to exogenous hormone treatment.

Materials and Methods

Animals and treatments

Normal and GnRH-null mice were maintained at the University of Glasgow and at the University of Oxford under UK Home Office regulations as previously described [8]. Adult animals (greater than 2 months) (four per group) were used without treatment or after sub-cutaneous injections of recombinant hCG (4IU/injection in saline) (Sero-pharmaceuticals Ltd, Feltham, Middlesex, UK). The injections were given every 12 hours and animals were

killed 2 hours after the final injection. The testes were frozen in liquid N₂ or fixed overnight in Bouin's fluid before being stored in 70% ethanol.

Reverse transcription and Real-time PCR

For quantification of the testicular content of specific mRNA species a real-time PCR approach was used which utilised the TaqMan PCR method following reverse transcription of the isolated RNA [22]. To allow specific mRNA levels to be expressed per testis and to control for the efficiency of RNA extraction, RNA degradation and the reverse transcription step an external standard was used [18]. The external standard was luciferase mRNA (Promega UK, Southampton, UK) and 5 ng was added to each testis at the start of the RNA extraction procedure. Testis RNA was extracted using Trizol (Life Technologies, Paisley, UK) and residual genomic DNA was removed by DNase treatment (*DNA-free*, Ambion Inc, supplied by AMS biotechnology, UK). The RNA was reverse transcribed using random hexamers and Moloney murine leukemia virus reverse transcriptase (Superscript II, Life Technologies, UK) as described previously [23,24].

Primers and probes for use in the TaqMan method have been described previously [19]. The real-time PCRs were carried out in a 25 μ l volume using a 96-well plate format. Components for real-time PCR were purchased from Applied Biosystems (Warrington, UK) apart from primers and probes which came from MWG Biotech (Milton Keynes, UK). Each PCR well contained reaction buffer, 5 mM MgCl₂, 200 μ M dNTPs (including dUTP), 300 nM each primer, 200 nM probe and 0.02 U/ μ l enzyme (Ampliq Gold). Reactions were carried out and fluorescence detected on a GeneAmp 5700 system (Applied Biosystems, Warrington, Cheshire, UK). For each sample a replicate was run omitting the reverse transcription step and a template negative control was run for each primer/probe combination. The quantity of each measured cDNA was then expressed relative to the internal standard luciferase cDNA in the same sample. This method allows direct comparison of expression levels per testis between different samples [18].

In situ hybridisation and histology

Testes from adult GnRH-null mice and GnRH-null mice treated with hCG were fixed in Bouin's and prepared for *in situ* hybridisation as described previously [1]. *In situ* probes for β 3HSD type VI and PGD-synthetase were produced as described previously [1,18]. Sense and antisense cRNA probes were prepared using T3 and T7 polymerase and were labelled with [³⁵S]dUTP. Tissue sections (7 μ m and at least four sections per animal) were hybridised overnight at 60°C with probe (100,000 c.p.m./ μ L) in hybridisation buffer (50% deionized formamide, 10% dextran sulphate, 50 mM dithiothreitol, 500 μ g/mL DNA

from calf thymus, 1 × Denhardt's solution, 20 mM Tris/HCl, pH8.0, 5 mM EDTA, 10 mM sodium phosphate, pH 6.8). Following hybridisation, slides were washed, dehydrated and allowed to air dry. Autoradiography was carried out using Ilford K5 emulsion (Ilford, Cheshire, UK) and the slides were stained with Mayer's haematoxylin and eosin (Merck Ltd, Leicestershire, UK).

To generate semi-thin (1 µm) sections for histology testes were fixed in 1% glutaraldehyde-4% paraformaldehyde, in phosphate buffer (0.1 M), pH 7.2 for 24 h at 4 °C and embedded in araldite. Cut sections were stained with toluidine blue.

Statistical analysis

To examine the effects of hCG injection on Leydig cell gene expression data from two experiments were normalised and combined. Differences between groups was analysed by single factor analysis of variance of log-transformed data. Differences between means was determined using the Newman-Keul post-test.

Results

Leydig cell morphology in adult GnRH-null mice

Leydig cells in adult GnRH-null mice were characterised by the presence of numerous large lipid droplets dispersed through the cell (Fig 1A). In normal adult mice the Leydig cells also frequently contained lipid droplets although these were generally smaller than in the GnRH-null mice (Fig 1B). In normal neonatal testes which contain only fetal-type Leydig cells there was more variation in Leydig cell morphology with lipid droplets present in many cells in characteristic clusters (Fig 1C). Some Leydig cells in the neonatal testis did not appear to contain lipid droplets although using semi-thin sections it is possible that they are present in the cell but not in the section shown.

Expression of adult Leydig cell markers in GnRH-null mice

A comparison of the expression of Leydig cell-specific mRNA species in GnRH-null mice and normal mice is shown in Table 1. Results are expressed relative to an external standard (luciferase) which was added to all testes at the RNA extraction phase and results are, therefore, comparable on a per testis basis [18]. The number of Leydig cells in the adult GnRH-null mouse testis is 10% of adult normal [9] and in the final column of Table 1 levels of mRNA are expressed relative to adult normal and corrected for cell number. Data from neonatal normal animals is included for control purposes since they contain only fetal-type Leydig cells [1,2]. The results show that, of those mRNA species tested, two (glutathione-S-transferase (GST) 5-5 and 3βHSD type I) are expressed at adult normal levels in the GnRH-null testis while the rest show a markedly reduced level of expression. There is, however, considerable variation with cytochrome P450 side chain

cleavage (P450sc), 3βHSD VI and PGD-synthetase present at less than 1% of normal while epoxide hydrolase (EH) is nearly 20% of normal. The four markers of adult Leydig cell differentiation which were measured in this study (3βHSD VI, PGD-synthetase, EST and 17βHSD III) were all detectable in the adult GnRH-null testis although expressions levels were amongst the lowest for all mRNA species measured. In most cases expression levels of mRNA species in adult GnRH-null animals was similar to, or higher than, that in neonatal normal animals. The exceptions were P450sc and renin. As expected 3βHSD VI and EST levels were undetectable in neonatal normal animals while the relatively high levels of PGD-synthetase and 17βHSD III in these animals is due expression in the seminiferous tubules [17,25].

Effect of hCG injections on marker expression in GnRH-null mice


Injection of adult GnRH-null mice with hCG increased the levels of all mRNA species tested with the exception of 3βHSD I which showed no change under the injection regime used (Fig 1). A single injection of hCG was sufficient to increase significantly levels of mRNA encoding P450c17, P450sc and steroidogenic acute regulatory (StAR) protein while two injections, spaced over 12 hours, increased expression of 3βHSD VI and 17βHSD III. Levels of EST, PGD-synthetase and relaxin-like factor (RLF) did not increase significantly until after four injections. The biggest change in expression was seen in 3βHSD VI which showed a 191-fold increase over control after 4 injections.

In situ hybridisation

In situ hybridisation was used to localise expression of markers of adult Leydig cell differentiation in the GnRH-null testis. A probe for 3βHSD VI showed specific localisation to the interstitial tissue in untreated GnRH-null mice (Fig 2A, 2B) with a clear increase in expression after hCG treatment (Fig 2C). Hybridisation of the 3βHSD VI probe tended to be in the central interstitial tissue rather than in the peritubular region (Fig 2B). A probe for PGD-synthetase also showed localisation to the interstitial tissue in untreated GnRH-null mice (Fig 2E) although there was no clear change in expression after hCG treatment (not shown).

Discussion

The initial studies which identified fetal and adult Leydig cells as belonging to different cell populations were based on morphological differences between the two cell types [26,27]. These differences are not particularly suitable, however, for differentiating between fetal and adult cells under conditions of hormonal or growth-factor deprivation since the morphology itself will be affected by the state of deprivation. This is clearly seen in the present study, and in previous studies [4,9,28], which show that

**Figure 1**

Light micrographs showing Leydig cell morphology in adult hpg testes (A), adult normal testes (B) and neonatal (day 7) testes (C). On micrographs A and C the bar represents 10 μm while on B it represent 20 μm .

Leydig cells in GnRH-null mice contain numerous large lipid droplets which are unlike either normal fetal or adult Leydig cells but are similar to Leydig cells with significantly reduced levels of steroidogenesis [29]. It is worth noting, however, that lipid droplet distribution in the GnRH-null adult Leydig cell is similar to that of the normal adult Leydig cell although the lipid droplets are larger and more frequent. Lipid droplet distribution in adult GnRH-null mice is unlike the characteristic clusters seen in fetal Leydig cells.

While morphological markers are clearly not ideal for identifying fetal and adult Leydig cells other studies have

now shown that there are fundamental differences in gene expression between the two Leydig cell populations which could be used to differentiate between the cell types. In initial studies it was shown that 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) is expressed in the adult Leydig cells of the rat but that there is no expression in fetal cells [30,31]. Our group and others have extended this to identify four mRNA species (3 β HSD VI, EST, PGD-synthetase and 17 β HSD III) which are expressed in the adult Leydig cells, but not the fetal Leydig cells, of the mouse [17–20]. So far, the only clearly identified differences in gene expression between the two populations are for genes expressed in the adult cell population but not in the




Figure 2

Expression of Leydig cell-specific mRNA species in testes of GnRH-null mice following injection of hCG. Levels of mRNA were measured by real-time PCR. Results show expression of each mRNA species following 1, 2 or 4 injections of hCG spaced 12 hours apart. The results for each mRNA species are expressed relative to control (untreated) GnRH-null animals. In each group the mean ± SEM of four animals is shown. Groups with different letter superscripts are significantly ($P < 0.05$) different.

Table 1:

Gene	Expression per testis (relative to Luciferase × 10 ³)			GnRH-null relative to adult normal (corrected for cell number) ²
	Adult GnRH-null	Adult normal	Neonatal normal ¹	
P450scc	0.23 ± 0.04	242 ± 35	2.2 ± 0.5	0.95%
P450c17	15.3 ± 1.8	3453 ± 1042	13.5 ± 3.5	4.4%
GST5-5	60.6 ± 5.6	677 ± 91	76 ± 15	89.5%
RLF	19.2 ± 2.1	2490 ± 611	5.3 ± 1.3	7.7%
3βHSD I	7.5 ± 1.7	51 ± 10	4.6 ± 1.8	147.0%
Renin	0.027 ± 0.012	21.4 ± 3.5	5.8 ± 0.9	1.3%
EH	18.5 ± 1.6	996 ± 112	8.0 ± 1.6	18.5%
StAR	2.08 ± 0.2	177 ± 30	4.0 ± 1.1	11.7%
3βHSD VI*	0.014 ± 0.005	124 ± 15	ND ³	0.11%
PGD-synthetase*	0.26 ± 0.06	843 ± 43	2.5 ± 1.1	0.31%
EST*	0.014 ± 0.009	9.0 ± 2.4	ND ³	1.5%
17βHSD III*	2.39 ± 0.21	1644 ± 192	140 ± 7	1.45%


* Genes expressed in adult Leydig cells but not fetal Leydig cells ¹ Neonatal animals were 5 days old. ² Expression in GnRH-null mice is expressed as a percentage of normal mice and corrected for reduced Leydig cell number in GnRH-null animals [4]. ³ND = not detectable.

fetal population [1,17–19]. Genes which are highly expressed in the fetal Leydig cell population, such as those encoding renin and thrombospondin 2, continue to be expressed in the adult testis, albeit at low levels [19,32,33]. This may be because adult Leydig cells express these genes at low levels or, more likely, because the fetal Leydig cell population persists in the adult testis [19,34–36]. Identification of genes specifically expressed in the adult Leydig cell population but not the fetal Leydig cell population does, however, allow us to determine whether adult Leydig cells develop in the absence of specific hormones or growth factors.

Normal differentiation of the adult Leydig cell population in the mouse begins around postnatal day 7 [1–3]. It has been shown previously that in the absence of LH there is a failure of normal proliferation of these cells and development of normal steroidogenic function does not occur [8,9]. Thus, there is no doubt that once differentiation has started adult Leydig cells require LH-stimulation but the role of LH at the onset of the initial differentiation phase has remained uncertain. Results from this study now show, however, that adult Leydig cells will differentiate in mice deficient in circulating LH. This conclusion is based primarily on the measured expression of specific adult Leydig cell markers in the testes of untreated GnRH-null mice. Expression of all four markers of adult Leydig cell differentiation was low but clearly detectable by real-time PCR in adult GnRH-null mice. Localisation, by *in situ* hybridisation, of two of these markers confirmed that expression was present in the untreated GnRH-null mouse testis and was localised to the interstitial tissue. For this

data to be accepted as clear evidence that adult Leydig cells differentiate despite a deficiency of LH depends on the veracity of the markers chosen to indicate adult Leydig cell development. Evidence that these markers are specific to adult Leydig cells comes from previous studies which have shown that during development in normal mice mRNA species encoding 3βHSD VI and EST are undetectable by real-time PCR [19] [and confirmed by current studies], *in situ* hybridisation [1], immunohistochemistry [20] or Northern blotting [20] before adult Leydig cell differentiation occurs. In addition, while both PDG-synthetase and 17βHSD III are expressed in the fetal/neonatal testis, this expression is only in the seminiferous tubules and stops prior to puberty at which time expression appears in the adult Leydig cell population [17,18]. Expression of PGD-synthetase in the current experiments was confirmed by *in situ* hybridisation to be localised to the interstitial tissue of the adult GnRH-null mouse testis. Thus, as far as can be determined, the markers used are specific for the adult Leydig cell population and indicate that these cells are present in the GnRH-null testis.

Comparison of expression patterns in adult GnRH-null testes with normal neonatal testes is also informative. If adult GnRH-null testes do not contain adult Leydig cells and have only fetal-type Leydig cells present it would be expected that the general pattern of mRNA expression would be that of a neonatal testis lacking gonadotrophin support. In fact, with the exceptions of P450scc and renin, expression levels are similar or higher in the adult GnRH-null testis. This is consistent, therefore, with differentia-

**Figure 3**

In situ hybridization studies showing 3 β -hydroxysteroid dehydrogenase (3 β HSD) type VI and prostaglandin D (PGD)-synthetase expression in adult control GnRH-null mice and GnRH-null mice following injection with hCG. Hybridisation of 3 β HSD VI anti-sense probe to control GnRH-null testis is shown in A) and at higher power in B). In C) hybridisation to testes from GnRH-null mice after 4 injections of hCG is shown. Hybridisation to a sense 3 β HSD VI probe is shown in D). Hybridisation of an anti-sense PGD-synthetase probe to testis from control GnRH-null mice is shown in E) while hybridization of a sense probe is shown in F).

tion of an adult population of cells in the adult GnRH-null testis.

The effects of hCG treatment on gene expression in the adult GnRH-null mouse testis also supports the hypothesis that adult Leydig cells are present in this animal. Expression of the markers 3 β HSD VI and 17 β HSD III was increased significantly after 2 injections of hCG within a 12 hour period. During normal testicular development there is a gap of at least four days between initial Leydig cell differentiation [around day 7 [2]] and first expression of 3 β HSD VI mRNA (around day 11 [1]) and a longer gap before the first significant rise in 17 β HSD III mRNA levels (after day 20 [19]). If only undifferentiated precursor cells were present in the GnRH-null testis it is likely that a more prolonged period of hCG treatment would be required before increased expression of 3 β HSD VI and 17 β HSD III would be seen. In contrast to 3 β HSD VI and 17 β HSD III expression of mRNA species encoding EST and PGD did not increase until after 48 hours of hCG treatment. This is consistent with previous data which has shown that these two genes exhibit a delayed pattern of expression during normal development [19].

In the GnRH-null mouse lack of pituitary stimulation by GnRH causes a marked reduction in pituitary synthesis and release of gonadotrophins [8]. This means that circulating levels of LH and FSH are very low but they are not, necessarily, abolished. Thus, the possibility cannot be excluded that there is low-level gonadotrophin-stimulation of Leydig cell precursors. It should be noted, however, that intratesticular testosterone levels in GnRH-null mice are undetectable and are lower than in LH-receptor-null mice [6,7]. Any circulating LH must, therefore, be at a minimal level and, as such, is unlikely to affect Leydig cell development.

If, as data described here would suggest, adult Leydig cells develop without LH-stimulation this begs the question as to what state of differentiation/development the cells reach. Morphological and functional development of the adult Leydig cell lineage has recently been reviewed by Mendis-Handagama & Ariyaratne [37]. Spindle-shaped, Leydig precursor cells are found in the peritubular region of the interstitial tissue and differentiate initially to progenitor cells in the same region. Further development to "newly formed" and "immature" adult Leydig cells is associated with rounding of the cells and movement to the central interstitial region [37]. The expression of adult Leydig cell genes in the unstimulated GnRH-null testis and the rapid response of some of these to hCG stimulation suggests that the cells have moved beyond initial steps of differentiation and may represent a form of unstimulated, immature adult cell. Further evidence that the cells have moved beyond the progenitor stage is provided

by the 3 β HSD VI *in situ* hybridisation studies which show that the hybridisation signal is localised primarily in the central interstitial tissue area rather than the peritubular region where the progenitor cells are located. It is also interesting to note that the total number of morphologically-recognisable Leydig cells (with no distinction drawn between fetal and adult cells) actually doubles over the pubertal period in the GnRH-null animal [9]. This would be consistent with morphological differentiation of adult Leydig cells during this period in GnRH-null mice and ties in with the functional data described above.

Despite apparent adult Leydig cell differentiation in GnRH-null mice the total number of Leydig cells in these animals remains markedly reduced at about 10% of control [9]. It is clear that development of the final number of adult Leydig cells is a combination of precursor differentiation and proliferation of the mature cell although the relative contribution of the two processes remains unclear [37]. If it is accepted, from the data presented here, that Leydig cell differentiation is largely an LH-independent event the marked reduction in Leydig cell number in the GnRH-null mouse would be consistent with a reduced role for cell differentiation and a more important role for proliferation in determining final cell number. A further implication of this model would be that while differentiation is an LH-independent event cell proliferation (including precursor/progenitor proliferation) is LH-dependent. This would be consistent with earlier studies showing that LH will induce Leydig cell proliferation in GnRH-null mice [28].

Results from this study suggest that adult Leydig cell differentiation does not require LH although the hormone is necessary for functional development of the cells and for the establishment of normal cell number. The initial stimulus for adult Leydig cell differentiation remains unknown although thyroid stimulating hormone appears to be required [30,38] along with the Sertoli cell products Dhh and PDGF-A [35,39]. Whatever other factor or factors are involved it appears likely that at least some will be of Sertoli cell origin since these cells are closely involved in regulation of the Leydig cells and are necessary for Leydig cell survival and function [40]

Acknowledgements

This study was supported by grants from the BBSRC and the Wellcome Trust. Technical assistance for this study came from Gary Jackson and Lynne Fleming.

References

1. Baker PJ, Sha JA, McBride MW, Peng L, Payne AH and O'Shaughnessy PJ **Expression of 3 β -hydroxysteroid dehydrogenase type I and VI isoforms in the mouse testis during development.** *Eur J Biochem* 1999, **260**:911-916
2. Nef S, Shipman T and Parada LF **A molecular basis for estrogen-induced cryptorchidism.** *Dev Biol* 2000, **224**:354-361

3. Vergouwen RPFA, Jacobs SGPM, Huiskamp R, Davids JAG and Derooij **Proliferative activity of gonocytes, sertoli cells and interstitial cells during testicular development in mice.** *J Reprod Fertil* 1991, **93**:233-243
4. Cattanach BM, Iddon CA, Charlton HM, Chiappa SA and Fink G **Gonadotrophin releasing hormone deficiency in a mutant mouse with hypogonadism.** *Nature* 1977, **269**:338-340
5. Kendall SK, Samuelson LC, Saunders TL, Wood RI and Camper SA **Targeted disruption of the pituitary glycoprotein hormone α -subunit produces hypogonadal and hypothyroid mice.** *Gene Dev* 1995, **9**:2007-2019
6. Zhang F-P, Poutanen M, Wilbertz J and Huhtaniemi I **Normal prenatal but arrested postnatal sexual development of luteinizing hormone receptor knockout (LuRKO) mice.** *Mol Endocrinol* 2001, **15**:172-183
7. Lei ZM, Mishra S, Zou W, Xu B, Foltz M, Li X and Rao CV **Targeted disruption of luteinizing hormone/human chorionic gonadotrophin receptor gene.** *Mol Endocrinol* 2001, **15**:184-200
8. O'Shaughnessy PJ, Baker P, Sohnius U, Haavisto A-M, Charlton HM and Huhtaniemi I **Fetal development of Leydig cell activity in the mouse is independent of pituitary gonadotroph function.** *Endocrinology* 1998, **139**:1141-1146
9. Baker PJ and O'Shaughnessy PJ **Role of gonadotrophins in regulating numbers of Leydig and Sertoli cells during fetal and postnatal development in mice.** *Reproduction* 2001, **122**:227-234
10. Hardy MP, Kelce WR, Kliefelder GR and Ewing LL **Differentiation of leydig cell precursors invitro – a role for androgen.** *Endocrinology* 1990, **127**:488-490
11. Huhtaniemi IT, Katikineni M and Catt KJ **Regulation of luteinizing-hormone-receptors and steroidogenesis in the neonatal rat testis.** *Endocrinology* 1981, **109**:588-595
12. Teerds KJ, Closset J, Rommerts FF, de Rooij DG, Stocco DM, Colenbrander B, Wensing CJ and Hennen G **Effects of pure FSH and LH preparations on the number and function of Leydig cells in immature hypophysectomized rats.** *J Endocrinol* 1989, **120**:97-106
13. Siril Ariyaratne HB, Mendis-Handagama SM, Hales DB and Mason JI **Studies of the onset of Leydig precursor cell differentiation in the prepubertal rat testis.** *Biol Reprod* 2000, **63**:165-171
14. Tena-Sempere M, Zhang FP and Huhtaniemi I **Persistent expression of a truncated form of the luteinizing hormone receptor messenger ribonucleic acid in the rat testis after selective Leydig cell destruction by ethylene dimethane sulfonate.** *Endocrinology* 1994, **135**:1018-1024
15. Veldhuizen-Tsoerkan MB, Ivell R and Teerds KJ **hCG-induced changes in LH/CG receptor mRNA transcript levels in the testis of adult hypophysectomized, ethane dimethyl sulphate-treated rats.** *Mol Cell Endocrinol* 1994, **105**:37-44
16. Tena-Sempere M, Rannikko A, Kero J, Zhang FP and Huhtaniemi IT **Molecular mechanisms of reappearance of luteinizing hormone receptor expression and function in rat testis after selective Leydig cell destruction by ethylene dimethane sulfonate.** *Endocrinology* 1997, **138**:3340-3348
17. O'Shaughnessy PJ, Baker PJ, Heikkila M, Vainio S and McMahon AP **Localization of 17 β -hydroxysteroid dehydrogenase/17-ketosteroid reductase isoform expression in the developing mouse testis – androstenedione is the major androgen secreted by fetal/neonatal leydig cells.** *Endocrinology* 2000, **141**:2631-2637
18. Baker PJ and O'Shaughnessy PJ **Expression of prostaglandin D synthetase during development in the mouse testis.** *Reproduction* 2001, **122**:553-559
19. O'Shaughnessy PJ, Willerton L and Baker PJ **Changes in Leydig cell gene expression during development in the mouse.** *Biol Reprod* 2002, **66**:966-975
20. Song WC, Qian Y, Sun X and Negishi M **Cellular localization and regulation of expression of testicular estrogen sulfotransferase.** *Endocrinology* 1997, **138**:5006-5012
21. O'Shaughnessy PJ **Steroidogenic enzyme-activity in the hypogonadal (hpg) mouse testis and effect of treatment with luteinizing-hormone.** *J Steroid Biochem Mol Biol* 1991, **39**:921-928
22. Bustin SA **Absolute quantification of mRNA using real-time reverse transcription polymerase chain reaction assays.** *J Mol Endocrinol* 2000, **25**:169-193
23. O'Shaughnessy PJ and Murphy L **Cytochrome P-450 17 α -hydroxylase protein and mRNA in the testis of the testicular feminized (Tfm) mouse.** *J Mol Endocrinol* 1993, **11**:77-82
24. O'Shaughnessy PJ, Marsh P and Dudley K **Follicle-stimulating hormone receptor mRNA in the mouse ovary during post-natal development in the normal mouse and in the adult hypogonadal [hpg] mouse: structure of alternate transcripts.** *Mol Cell Endocrinol* 1994, **101**:197-201
25. Baker PJ and O'Shaughnessy PJ **Expression of prostaglandin D synthetase during development in the mouse testis.** *Reproduction* 2001, **122**:553-559
26. Roosen-Runge EC and Anderson D **The development of the interstitial cells of the albino rat.** *Acta Anat* 1959, **37**:125-137
27. Lording DW and De Kretser DM **Comparative ultrastructural and histochemical studies of the interstitial cells of the rat testis during fetal and postnatal development.** *J Reprod Fertil* 1972, **29**:261-269
28. Scott IS, Charlton HM, Cox BS, Grocock CA, Sheffield JW and O'Shaughnessy PJ **Effect of LH injections on testicular steroidogenesis, cholesterol side-chain cleavage P450 messenger RNA content and leydig cell morphology in hypogonadal mice.** *J Endocrinol* 1990, **125**:131-138
29. Hasegawa T, Zhao L, Caron KM, Majdic G, Suzuki T, Shizawa S, Sasano H and Parker KL **Developmental roles of the steroidogenic acute regulatory protein (StAR) as revealed by StAR knockout mice.** *Mol Endocrinol* 2000, **14**:1462-1471
30. Mendis-Handagama SM, Ariyaratne HB, Teunissen van Manen KR and Haupt RL **Differentiation of adult Leydig cells in the neonatal rat testis is arrested by hypothyroidism.** *Biol Reprod* 1998, **59**:351-357
31. Phillips DM, Lakshmi V and Monder C **Corticosteroid 11 beta-dehydrogenase in rat testis.** *Endocrinology* 1989, **125**:209-216
32. Perera EM, Martin H, Seeherunvong T, Kos L, Hughes IA, Hawkins JR and Berkovitz GD **Tescalcin, a novel gene encoding a putative EF-hand Ca²⁺-binding protein, Col9a3, and renin are expressed in the mouse testis during the early stages of gonadal differentiation.** *Endocrinology* 2001, **142**:455-463
33. O'Shaughnessy PJ, Johnston H and Baker PJ **Failure of normal adult Leydig cell development in androgen-receptor-deficient mice.** *J Cell Sci* 2002, **115**:3491-3496
34. Kerr JB and Knell CM **The fate of fetal leydig-cells during the development of the fetal and postnatal rat testis.** *Development* 1988, **103**:535-544
35. Bitgood MJ, Shen L and McMahon AP **Sertoli cell signaling by Desert hedgehog regulates the male germline.** *Curr Biol* 1996, **6**:298-304
36. Ariyaratne HB and Chamindrani Mendis-Handagama S **Changes in the testis interstitium of Sprague Dawley rats from birth to sexual maturity.** *Biol Reprod* 2000, **62**:680-690
37. Mendis-Handagama SM and Ariyaratne HB **Differentiation of the adult Leydig cell population in the postnatal testis.** *Biol Reprod* 2001, **65**:660-671
38. Teerds KJ, de Rooij DG, de Jong FH and van Haaster LH **1998 Development of the adult-type Leydig cell population in the rat is affected by neonatal thyroid hormone levels.** *Biol Reprod* 1998, **59**:344-350
39. Gnessi L, Basciani S, Mariani S, Arizzi M, Spera G, Wang C, Bondjers C, Karlsson L and Betsholtz C **Leydig cell loss and spermatogenic arrest in platelet-derived growth factor (PDGF)-A-deficient mice.** *J Cell Biol* 2000, **149**:1019-1026
40. Russell LD, Warren J, Debeljuk L, Richardson LL, Mahar PL, Waymire KG, Amy SP, Ross AJ and MacGregor GR **Spermatogenesis in Bclw-deficient mice.** *Biol Reprod* 2001, **65**:318-332