



Litthauer, S., Chan, K. X. and Jones, M. (2018) 3'-phosphoadenosine 5'-phosphate accumulation delays the circadian system. *Plant Physiology*, 176(4), pp. 3120-3135.

There may be differences between this version and the published version. You are advised to consult the publisher's version: <https://doi.org/10.1104/pp.17.01611> if you wish to cite from it.

<http://eprints.gla.ac.uk/193419/>

Deposited on: 19 August 2019

Short Title: PAP Signaling Delays the Circadian System

Corresponding Author:

Dr. Matt Jones, School of Biological Sciences, University of Essex, Wivenhoe Park,
Colchester, Essex, CO4 3SQ, United Kingdom; +44 (0) 1206-874740 (ph)

Article Title:

3'-Phosphoadenosine 5'-Phosphate Accumulation Delays the Circadian System

Authors and Affiliations:

Suzanne Litthauer^a, Kai Xun Chan^b, and Matthew Alan Jones^a

^a School of Biological Sciences, University of Essex, Wivenhoe Park, Colchester, Essex,
CO4 3SQ, United Kingdom

^b Center for Plant Systems Biology, VIB, 9052 Ghent, Belgium

One Sentence Summary:

Accumulation of the stress-induced metabolite PAP delays the circadian system in
Arabidopsis thaliana.

List of Author Contributions: S.L. and M.A.J. performed the research and analysed data.
S.L., K.X.C., and M.A.J. designed the research and wrote the article.

Funding Information: This work was supported by the Leverhulme Trust (ECF-2012-358),
The Royal Society (grant no. RG130746), The Oppenheimer Memorial Trust (PhD
studentship to S.L.), The Gen Foundation (to S.L.), the University of Essex, and the
Australian Research Council Centre of Excellence in Plant Energy Biology (CE140100008).

Corresponding Author Email: matthew.jones@essex.ac.uk

Abstract

The circadian system optimizes cellular responses to stress, but the signaling pathways that convey the metabolic consequences of stress into this molecular timekeeping mechanism remain unclear. Redox-regulation of the SAL1 phosphatase during abiotic stress initiates a signaling pathway from chloroplast to nucleus by regulating the accumulation of a metabolite, 3'-phosphoadenosine 5'-phosphate (PAP). Consequently, PAP accumulates in response to redox stress and inhibits the activity of exoribonucleases (XRNs) in the nucleus and cytosol. We demonstrated that osmotic stress induces a lengthening of circadian period and that genetically inducing the SAL1-PAP-XRN pathway in plants lacking either SAL1 or XRNs similarly delays the circadian system under these conditions. Exogenous application of PAP was also sufficient to extend circadian period. Thus, SAL1-PAP-XRN signaling likely regulates circadian rhythms in response to redox stress. Our findings exemplify how two central processes in plants, molecular timekeeping and responses to abiotic stress, can be interlinked to regulate gene expression.

Introduction

The rotation of the Earth confers overt environmental rhythms upon species living on its surface. Both temperature and incident light change dramatically (yet predictably) over any given 24-hour cycle, and as a consequence there is a selective pressure for species to anticipate changes in environmental conditions (Hut and Beersma, 2011). This selective pressure has led to the evolution of the circadian clock, an endogenous biological oscillator that modulates biochemical and physiological activity to optimize behaviour within the prevailing environmental context (Millar, 2016).

The pervasive nature of the circadian system has encouraged the detailed description of the network underpinning these biological rhythms. Circadian rhythms are entrained to the local day/night cycle by regular changes in temperature and light (Jones, 2009; Hsu and Harmer, 2014). Phytochromes act to input red light-derived signals, while cryptochromes and the ZEITLUPE (ZTL) family are the predominant blue photoreceptors that influence circadian rhythms (Fankhauser and Staiger, 2002; Hsu and Harmer, 2014). Nuclear circadian rhythms within *Arabidopsis thaliana* consist of multiple, interconnected transcriptional feedback loops (Hsu and Harmer, 2014). PSEUDORESPONSE REGULATOR9 (PRR9) acts sequentially with PRR7, PRR5, and PRR1/TIMING OF CAB EXPRESSION 1 (TOC1) to repress expression of *CIRCADIAN CLOCK ASSOCIATED1* (CCA1) and *LATE ELONGATED HYPOCOTYL* (LHY) throughout the day (Farré and Kay, 2007; Nakamichi et al., 2010; Gendron et al., 2012; Huang et al., 2012). In turn, CCA1 and LHY (whose expression is induced by light at dawn) repress expression of *PRR9/7/5/TOC1* (Alabadi et al., 2001; Adams et al., 2015) and additional circadian genes such as *GIGANTEA* (Lu et al., 2012). Subsequently, a complex of proteins, including EARLY FLOWERING4 (ELF4), act to repress circadian gene expression during the early night (McWatters et al., 2007; Nusinow et al., 2011; Chow et al., 2012; Herrero et al., 2012). The oscillations generated by these feedback loops modulate many physiological processes including growth, photosynthesis, flowering time, and responses to biotic and abiotic stresses (Dong et al., 2011; Eriksson and Webb, 2011; Hsu and Harmer, 2014; Song et al., 2015; Jones, 2017).

We have been interested in the molecular interactions between plants' response to stress and the circadian system so as to improve survival. Damage induced by abiotic factors is typically first observed within the chloroplast and mitochondria, where perturbations in metabolism rapidly induce oxidative damage (Mittler et al., 2011). These perturbations are communicated from organelles to the nucleus *via* multiple retrograde signaling pathways that

adjust nuclear gene expression. However, the extent to which retrograde signals can regulate plant homeostasis, and by what mechanism(s), remain enigmatic (Chan et al., 2016). We therefore examined a candidate signaling pathway that could be responsible for coordinating nuclear circadian rhythms in response to metabolic stress in the chloroplast. SAL1 is a redox-sensitive phosphatase localized to the chloroplast and mitochondria (Chen et al., 2011; Estavillo et al., 2011; Chan et al., 2016). A recent model of SAL1-initiated signaling from chloroplast to nucleus has proposed that upon oxidative stress, redox-induced impairment of SAL1 activity leads to accumulation of its substrate 3'-phosphoadenosine 5'-phosphate (PAP), resulting in inhibition of 5'→3' exoribonuclease (XRN) activity and subsequent changes in expression of plastid redox-associated nuclear genes and abscisic acid (ABA) signaling (Dichtl et al., 1997; Mechold et al., 2006; Estavillo et al., 2011; Chan et al., 2016; Pornsiriwong et al., 2017). This model is supported by analysis of the transcriptomes of *xrn* and *sal1* mutants, with considerable overlap between the mis-regulated transcripts in each of these genotypes (Gy et al., 2007; Estavillo et al., 2011; Kurihara et al., 2012). In this study, we demonstrated that osmotic stress delays the circadian system and that constitutive activation of PAP signalling in *sal1* and *xrn* plants results in an extended circadian period. These data provide an additional mechanism through which the molecular clock, a cornerstone of plant function, can be coordinated with the metabolic status of a plant cell to guide molecular responses to environmental fluctuations.

Results

Whole-plant osmotic stress treatments lengthen circadian period and induce accumulation of PAP

Drought is a multifaceted stress that arises from limited water availability and is one of the primary abiotic stresses that limits crop yield (Verslues et al., 2006; Steduto et al., 2012). We therefore sought to understand how this stress affects the circadian system using the model plant *Arabidopsis thaliana* (*Arabidopsis*). As consistent maintenance of soil water potential during circadian imaging presented technical issues, we instead approximated the physiological consequences of water deficit stress through the addition of 200 mM mannitol to lower the water potential of the agar substrate (Figure 1, Verslues et al., 2006). One of the initial metabolic consequences of water-deficit stress is the accumulation of PAP due to the redox-induced inactivation of SAL1 (Estavillo et al., 2011; Chan et al., 2016), and so we measured PAP levels under our experimental conditions. As previously reported for soil-grown plants, the application of osmotic stress was sufficient to induce accumulation of PAP,

with comparable increases in PAP observed in plants transferred to mannitol and grown under either constant white (cW) light or constant red+blue light provided by LEDs (cR+B, Figure 1A, Estavillo et al., 2011). Interestingly, we observed a 1-hour increase in the circadian free running period (FRP) in wild-type plants subjected to osmotic stress under cR+B light (Fig. 1B and 1C, $P < 0.01$, Dunnett's test). A comparable FRP extension of approximately 1 hour was observed in plants transferred to Murashige and Skoog (MS) plates infused with an alternate osmoticum (PEG 8000, Fig. 1B and 1C).

Loss of SAL1 activity results in lengthening of circadian period in *sal1* mutants

We hypothesised that the accumulation of PAP during osmotic stress contributed to the observed extension of the circadian period. Intracellular PAP accumulation can be increased through disruption of *SAL1*, a gene that encodes a redox-sensitive phosphatase, and so we examined the FRP of *sal1* mutant alleles to test our hypothesis (Figure 2, Kim and von Arnim, 2009; Wilson et al., 2009; Rodríguez et al., 2010). Circadian rhythms can be routinely monitored by measuring changes in photosystem II (PSII) operating efficiency (F_q'/F_m' , Litthauer et al., 2015). Using this technique, we observed a significant increase in FRP under constant blue (cB) light in *alx8-1*, *fry1-6*, and *fou8* alleles of *SAL1* compared to wild type (Fig. 2A-D, Figure S1). This long-period phenotype was rescued by introducing a wild-type copy of the *SAL1* coding region along with a 1-kb region of upstream genomic sequence into the *alx8-1* background (Figure 2E), suggesting that a mutation in *SAL1* underlies the phenotype observed in the mutant lines.

To better document the *sal1* mutant circadian phenotype, we introduced a *CCA1::LUC2* luciferase reporter into the *fry1-6* background (Figure 2F). When measuring luciferase bioluminescence, we observed a one-hour extension of FRP under cB light (23.80 ± 0.17 hrs in wild type compared to 25.23 ± 0.14 hrs in *fry1-6*; $p < 0.025$, Student's t test). We also observed a modest yet significant increase in FRP under cR+B light (22.90 ± 0.05 hrs in wild type compared to 23.52 ± 0.07 hrs in *fry1-6*, $p < 0.05$, Student's t test, Figure 2F). Interestingly, we did not observe a difference in FRP between wild type and *sal1* plants under constant red (cR) light or constant darkness (Figure 2F). Such data suggest that FRP is delayed in a blue light-dependent manner in *sal1* mutants.

PAP accumulates *in vivo* in response to osmotic stress due to a change in the redox state within the chloroplast that inhibits the enzymatic activity of SAL1 (Chan et al., 2016). We therefore sought to assess the consequences of these oxidative stresses upon the nuclear

circadian system (Figure 2G). We first used 50 μ M methyl viologen (MV, which induces reactive oxygen species production at PSII) to induce oxidative stress within the chloroplast (Figure 2G, Lai et al., 2012; Chan et al., 2016). Application of MV induced a 1.5 hr lengthening of circadian period in wild-type seedlings ($\tau = 22.66 \pm 0.08$ and 24.25 ± 0.15 hrs on mock- and MV-treated plates, respectively), with a more modest lengthening observed in *fry1-6* seedlings ($\tau = 23.23 \pm 0.21$ and 24.57 ± 0.16 hrs in the absence or presence of MV, respectively). Intriguingly, FRP in wild type and *sal1* seedlings was indistinguishable after MV treatment. Therefore, the application of oxidative stress using MV lengthens circadian period.

Recent work has suggested that PAP acts as a secondary messenger during ABA signalling to promote stomatal closure; therefore, we next examined whether exogenous ABA was able to reconstitute the long-period phenotype of *sal1* plants (Fig. 2H, Pornsiriwong et al., 2017). In contrast to our hypothesis, the circadian FRP was reduced in both wild-type and *sal1* plants in the presence of exogenous ABA when compared to mock-treated controls (Figure 2H). In the presence of ABA, *sal1* lines retained their extended FRP phenotype (post-hoc Bonferroni adjusted t test), although the magnitude of the phenotype was less than in mock-treated plants (Figure 2H). Our data suggest that enhanced ABA signalling does not contribute to the delayed FRP of *sal1* plants, although additional work will be required to fully understand the interaction between SAL1 and ABA signalling in a circadian context.

It has been proposed that increased fluence rates enhance the accumulation of PAP *in vivo* (Estavillo et al., 2011); therefore, we completed fluence rate response curves under either cB or cR light to determine whether increased fluence rates would exacerbate the *sal1* circadian phenotype (Figure 3). Under dim blue light ($5 \mu\text{mol m}^{-2} \text{s}^{-1}$), we did not observe a difference in FRP between wild-type and *fry1-6* plants (Figure 3A). However, we did observe a significant difference in *sal1* plants' response to increasing blue light compared to wild type ($p < 0.001$), which resulted in a lengthening of FRP in *sal1* plants transferred to $\geq 20 \mu\text{mol m}^{-2} \text{s}^{-1}$ cB light (Figure 3A). By contrast, we did not observe a significant lengthening of FRP in *fry1-6* plants transferred to any tested fluence rate of cR light, as was suggested by our initial studies under cR light (Fig. 2F and 3B).

In order to better understand the extended circadian FRP phenotype, we examined the accumulation of clock-regulated transcripts under either $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light or $30 \mu\text{mol m}^{-2} \text{s}^{-1}$ red light (Figure 3C-F, Figure S2). In agreement with our luciferase data (Fig.

2F and 3A-B), we observed that *CCA1* and *TOC1* transcript accumulation was delayed by approximately 6-9 hrs under cB light (Figure 3C and 3E). This phase shift was less apparent in plants transferred to cR light (Fig. 3D and 3F). Since *sal1* plants present a blue-light dependent phenotype, we monitored the accumulation of *CRYPTOCHROME1* (*CRY1*), *CRY2*, and *ZEITLUPE* transcripts to confirm that accumulation of these blue photoreceptors was not repressed by the loss of *SAL1*. However, we found that neither *CRY1*, *CRY2*, nor *ZEITLUPE* transcript accumulation were significantly repressed in *sal1* plants (Figure S3).

We next examined PAP accumulation in plants grown under our experimental light conditions. As previously reported, we were unable to detect PAP in any of our wild-type samples (Fig. 3G-H, Chen et al., 2011; Estavillo et al., 2011; Lee et al., 2012). In *fry1-6* plants, the absence of a circadian phenotype under lower cB light intensities was correlated with a significant reduction in PAP accumulation (Figure 3G), with less than 2 nmol g⁻¹ PAP accumulating in *fry1-6* plants under dim blue light compared to in excess of 5 nmol g⁻¹ above 20 μmol m⁻² s⁻¹ blue light (Figure 3G). Interestingly, PAP accumulation remained greater in *fry1-6* plants transferred to different intensities of cR or cR+B despite the circadian phenotype being less pronounced under these conditions (Fig. 3H, 3I, S4). Such data suggest that PAP acts to delay the circadian system via a blue light-induced pathway, or that cR light stimulates an opposing or compensatory signalling cascade.

Reduced PAP accumulation rescues the circadian phenotype of *sal1* mutants

Loss of *SAL1* activity results in a lengthened circadian phenotype (Figure 2) and so we examined whether PAP levels correlated with FRP in *sal1* mutants (Figure 4). Given the correlation between PAP accumulation and the *sal1* circadian phenotype under cB light, we hypothesized that exogenous application of PAP would be sufficient to extend FRP. Application of PAP to intact wild-type seedlings did not induce gene expression, presumably because of the enzymatic activity of the endogenous *SAL1* protein (Estavillo et al., 2011; Pornsiriwong et al., 2017). Therefore, we examined whether the application of additional PAP was sufficient to extend FRP in *sal1* plants that have a compromised ability to degrade exogenous PAP (Figure 4A). Following entrainment, plants were transferred to cB light for imaging before PAP was applied. As expected, there was no significant difference in FRP in wild-type plants following PAP application (p=0.953, Figure 4A). However, we did observe a lengthening of circadian periodicity from 24.52 ± 0.14 hrs to 25.25 ± 0.12 hrs in *sal1* plants following treatment with PAP (Figure 4A, p<0.025).

We next tested whether reduced PAP accumulation in the *sal1* background was sufficient to rescue the circadian phenotype (Figure 4B-D). SAL1 is a bi-functional enzyme with PAP phosphatase and inositol polyphosphate 1-phosphatase activities *in vitro* (Quintero et al., 1996; Xiong et al., 2001). In order to specifically reduce PAP levels *in vivo*, we over-expressed *ARABIDOPSIS HAL2-LIKE (AHL)*, a paralogue of *SAL1* with only PAP phosphatase activity (Kim and von Arnim, 2009; Chen and Xiong, 2010; Hirsch et al., 2011). Transgenic lines over-expressing *AHL* in an *alx8-1* background had wild-type levels of PAP under cB light (Figure 4B). We also found that overexpression of the *AHL* paralogue was able to rescue the mutant circadian phenotype of *alx8-1* seedlings (Fig. 4C and 4D). Such data demonstrate that PAP phosphatase activity is sufficient to reduce PAP accumulation and complement the *sal1* circadian phenotype.

SAL1 is a constitutively expressed protein

As *sal1* mutants have not previously been characterized as having a circadian phenotype, we explored the regulation of *SAL1* transcripts and protein over diel and circadian timescales (Figure 5). Consistent with previous microarray studies (Figure S5, Mockler et al., 2007), we found that *SAL1* transcripts accumulate gradually over the course of the day under entraining conditions (Figure 5A), whereas under cW light, we did not observe a discernible rhythm in *SAL1* transcript (Figure 5C, Mockler et al., 2007). Interestingly, no changes in protein accumulation were apparent in our transgenic lines expressing *SAL1-GFP* under the control of its native promoter (Figure 5B). *SAL1* oscillations consequently appear to be primarily driven by the diel cycle rather than *SAL1* being a classical output of the core nuclear circadian system.

Lengthening of circadian period in *sal1* mutants is not induced by sulfur deprivation

The *sal1* mutation induces accumulation of its substrate, PAP, and to a lesser extent the PAP precursor, 3'-phosphoadenosine 5'-phosphosulfate (PAPS, Figure 6, Chen et al., 2011; Estavillo et al., 2011; Lee et al., 2012). As a consequence, *sal1* seedlings present a sulfur-deprived phenotype, presumably derived from disruption of sulfate assimilation pathways (Figure 6A, Mugford et al., 2009; Lee et al., 2012). Sulfate assimilation is vital for plant metabolism (Takahashi et al., 2011), and so we investigated whether the long-period circadian phenotype of the *sal1* mutant is induced via the reduced accumulation of sulfate. We first evaluated the consequences of gross sulfate starvation on nuclear rhythms using a collection of luciferase reporter lines in a wild-type background (Figure 6B and Figure S6).

Our sulfate-deprived growth conditions were sufficient to induce accumulation of the sulfate starvation marker *APS REDUCTASE1* and *SULTR4;2* (Figure S6). However, despite the induction of a sulfate-starvation response, there was no difference in the period of bioluminescence rhythms driven by the *CCA1*, *LHY*, or *TOC1* promoters (Figure 6B and Figure S6B-D). As sulfates are necessary for the maintenance of photosynthesis (Terry, 1976), we next examined rhythms of PSII operating efficiency in wild-type and *fry1-6* plants under sulfate-deprived conditions (Figure 6C and Figure S6E). As with our studies using luciferase reporters (Figure 6B), we did not observe any significant difference in FRP in wild-type plants in the presence or absence of sulfates ($\tau = 23.96 \pm 0.32$ and 23.74 ± 0.30 hrs on MS media and MS media lacking sulfates, respectively). Sulfate deprivation is therefore insufficient to extend the period of the circadian system.

In order to further evaluate the contribution of sulfur limitation to the *sal1* circadian phenotype, we examined additional mutant lines deficient in sulfate metabolism (Fig. 6A, 6D and 6E). Plants lacking *ARABIDOPSIS 5-PHOSPHOSULFATE KINASE1* (*APK1*) and *APK2* are less able to phosphorylate adenosine 5'-phosphosulfate and so accumulate fewer glucosinolates, with a commensurate increase in desulfo-glucosinolates - a phenotype that is also observed in *sal1* alleles (Mugford et al., 2009; Lee et al., 2012). As when we examined plants grown on sulfate-deficient media, we observed no difference in FRP in *apk1 apk2* mutants compared to wild type (Figure 6D, $\tau = 23.87 \pm 0.11$ hrs in *apk1 apk2* compared to 23.70 ± 0.09 hrs in wild type). Similarly, *cad2-1* plants, which accumulate less glutathione than wild-type plants (Cobbett et al., 1998), had a comparable phenotype to wild-type plants (Figure 6E). These data demonstrate that gross deficiencies in sulfate metabolism do not extend FRP and thus is unlikely to account for the mechanism by which SAL1/PAP signaling regulates the circadian rhythm.

Plants lacking exoribonucleases have a comparable long-period phenotype to the *sal1* mutant

The accumulation of PAP as a consequence of *SAL1* inactivation leads to the inhibition of XRN exoribonuclease activity (Chan et al., 2016). As mis-regulation of RNA processing frequently leads to an altered circadian free-running period (Jones et al., 2012; Wang et al., 2012; Macgregor et al., 2013; Perez-Santángelo et al., 2014), we examined whether a mutation of XRN exoribonucleases to genetically simulate SAL1-mediated XRN inhibition was sufficient to alter nuclear circadian rhythms (Figure 7).

The *Arabidopsis* genome expresses three *XRN* genes, with *XRN2* and *XRN3* acting within the nucleus whereas *XRN4* accumulates in the cytosol (Kastenmayer and Green, 2000). Transcripts from these genes did not accumulate with a daily rhythm (Figure S5). As *XRNs* display a degree of functional redundancy and are likely all inhibited by PAP accumulation (Gy et al., 2007; Nagarajan et al., 2013), we examined circadian rhythms in the *xrn2 xrn3 xrn4* (*xrn234*) triple mutant (Figure 7). As observed in *sal1* plants, these *xrn234* seedlings have a long FRP compared to the wild type (Figure 7A, $\tau = 24.71 \pm 0.20$ hrs in *xrn234* compared to 23.88 ± 0.19 hrs in wild type). This long-period phenotype corresponded to a delayed phase of *CCA1* transcript accumulation under cB light in both *fry1-6* and *xrn234* seedlings (Figure 7B). Interestingly, this phase delay in transcript accumulation in *xrn234* seedlings was less apparent in cW light, similar to the more subtle phenotype observed in *sal1* alleles under cW light or cR+B light (Fig. 2F and 7C). These data support the current model for PAP signalling that suggests that PAP accumulation in *sal1* lines represses *XRN* activity, rather than direct targeting of individual transcripts (Wilson et al., 2009; Rodríguez et al., 2010; Lee et al., 2012).

Loss of *SAL1* mimics the clock's response to osmotic stress via a blue light-induced pathway

In order to assess the contribution of *SAL1* and PAP to circadian timekeeping during osmotic stress, we revisited our experimental design outlined in Figure 1. The transfer to media containing mannitol did not induce the accumulation of additional PAP in *sal1* plants under either cR+B or cB light, although PAP accumulation increased in the wild type as previously observed (Fig. 1A, 8A, 8B). Under cR+B light, *sal1* seedlings had a longer FRP than wild-type controls in the presence or absence of mannitol, which correlated with the increased accumulation of PAP in these lines (Figure 8C). Both wild-type and *sal1* plants retained a modest (yet significant) circadian response to osmotic stress under cR+B and cR light (Fig. 8C and S7). Interestingly, *sal1* seedlings did not have an extended circadian FRP when transferred to mannitol under cB light, although wild-type plants retained this response (Figure 8D). Comparable mannitol-induced shifts in circadian phase were observed when we examined the accumulation of *CCA1*, *PRR5*, and *GI* transcripts under cB light (Fig. 8E and S8). Such data suggest that PAP accumulation is sufficient to extend circadian FRP under cB light, but that additional red light-induced factors also coordinate the circadian system's response to mannitol.

Discussion

PAP accumulation is sufficient to extend circadian period in the presence of blue light

Inactivation of SAL1 through either mutation or application of oxidative stress within the chloroplast induces the accumulation of PAP (Estavillo et al., 2011; Chan et al., 2016). We were able to detect PAP in *sal1* seedlings under all conditions tested, but only observed an extension of circadian FRP under cB or cR+B light (Fig. 1, 3, 4, and 8). PAP accumulated to a greater extent in *sal1* seedlings transferred to 20 or 40 $\mu\text{mol m}^{-2} \text{s}^{-1}$ cB light compared to those moved to 5 $\mu\text{mol m}^{-2} \text{s}^{-1}$ cB light (Figure 3G). This increase in PAP accumulation was correlated with the presentation of the mutant circadian phenotype, with *sal1* seedlings having an FRP indistinguishable from the wild type when transferred to very dim blue light (Figure 3A). Despite this correlation, PAP levels remained higher in *sal1* mutants than in the wild type under these low light conditions (Figure 3G). Such data suggest that either a threshold concentration of PAP is necessary *in vivo* to delay the molecular clock, or that the blue light-dependent signal perturbed by PAP is only significant at higher fluence rates of blue light.

Intriguingly, we also noted that *sal1* and *xrn234* seedlings had a less pronounced circadian defect when transferred to constant conditions that included red wavelengths of light (Fig. 2F, 7, and 8), although elevated PAP levels were also observed in *sal1* plants transferred to cR light (Figure 3H, Figure S4). Such data suggest that either a red light-mediated signal supersedes or acts in parallel with the PAP-derived pathway, or that the PAP-derived signal specifically affects a blue light-mediated response. The role of phytochrome-related factors in chloroplast retrograde signaling has previously been demonstrated (Salomé et al., 2013; Norén et al., 2016). Plants lacking iron have an extended FRP that is dependent upon phytochromes, suggesting an additional role for iron within the circadian system beyond the maintenance of photosynthesis (Chen et al., 2013; Hong et al., 2013; Salomé et al., 2013). As a consequence, it is likely that multiple signals relay information regarding the metabolic status of the chloroplast to the nucleus.

Exogenous applications of PAP alone to intact leaves of wild-type plants has previously been shown to be ineffective, presumably because endogenous SAL1 is sufficient to metabolize this exogenous PAP (Estavillo et al., 2011; Pornsiriwong et al., 2017). However, the exogenous application of PAP was sufficient to extend FRP in *fry1-6* seedlings (Figure 4A). The exogenous application of PAP is therefore sufficient to extend the long-

period circadian phenotype of *sal1* mutant plants that are unable to degrade this metabolite, suggesting that accumulation of PAP underlies the circadian phenotype of *sal1* plants.

Mutation of *SAL1* has a pleiotropic effect upon plant development, with auxin hyposensitivity and ABA hyper-sensitivity being reported in *sal1* alleles (Xiong et al., 2001; Chen and Xiong, 2010; Rodríguez et al., 2010). In particular, the accumulation of PAP in *sal1* plants up-regulates specific ABA signalling components to induce stomatal closure (Pornsiriwong et al., 2017). ABA induces a complex circadian response, with exogenous ABA having no effect or increasing the circadian period in plants grown in the presence of sucrose (Hanano et al., 2006; Liu et al., 2013). Conversely, ABA shortens the circadian period in the absence of exogenous sucrose (Lee et al., 2016). We were able to recapitulate this latter phenotype in wild-type plants under our conditions (grown in the absence of exogenous sucrose, Fig. 2H). Similarly, an accelerated FRP was observed in *sal1* seedlings in response to ABA (Fig. 2H), demonstrating that these lines retain a sensitivity to this hormone. Although we cannot completely discount a role for altered ABA signalling in the *sal1* phenotype, we do note that a reduction of PAP accumulation in *sal1* mutants over-expressing *AHL* was sufficient to rescue the *sal1* mutant phenotype (Fig. 4B-D). Therefore, we propose that the perturbations in plant hormone accumulation and sensitivity in *sal1* alleles are part of the global developmental consequences of increased PAP accumulation, rather than altered ABA sensitivity inducing the extended circadian FRP observed.

Regulation of the circadian system by SAL1 does not arise as a consequence of sulfate limitation in *sal1*

Sulfate assimilation occurs via a branching pathway, part of which culminates in the production of PAPS that acts as a donor of activated sulfate for many sulfation reactions (Figure 6A, Takahashi et al., 2011). *sal1* mutants also accumulate desulfo-glucosinolates, presumably because of a homeostatic disruption within this branch of the sulfate assimilation pathway as a consequence of the accumulation of PAPS and PAP (Lee et al., 2012). Consequently, *sal1* lines also accumulate jasmonic acid (Rodríguez et al., 2010). Previous work has demonstrated that jasmonic acid production is regulated by the circadian system (Goodspeed et al., 2012), and jasmonate signaling is gated by the clock, leading to improved resistance against herbivory and infection (Covington et al., 2008; Goodspeed et al., 2012; Shin et al., 2012; Ingle et al., 2015).

A recent study demonstrated that perturbed glucosinolate accumulation shortens FRP under cR+B light (Kerwin et al., 2011). In order to test the consequences of perturbed sulfate

metabolism upon FRP under our experimental conditions, we examined plants grown on sulfate-deficient media (Fig. 6B and 6C) and mutants with perturbed sulphate metabolism (*apk1 apk2* and *cad2-1*, Fig. 6D and 6E). In agreement with Kerwin *et al.*, we did not observe a significant extension of FRP in sulfate-deprived conditions (Fig. 6B and 6C). Instead, we observed a modest shortening of FRP in one of our luciferase lines (*TOC1::LUC*, Figure 6B). Our data subsequently suggest that sulfate limitation does not induce a long FRP and is not the mechanism by which PAP regulates circadian rhythm.

Loss of XRN activity replicates the circadian phenotypes of *sal1* mutants

The importance of post-transcriptional regulation of circadian gene expression is increasingly being recognized, particularly in response to environmental changes (Garbarino-Pico and Green, 2007; Kojima *et al.*, 2011; Sanchez *et al.*, 2011). Mis-regulation of RNA processing frequently leads to an altered circadian FRP (Jones *et al.*, 2012; Wang *et al.*, 2012; Macgregor *et al.*, 2013; Perez-Santángelo *et al.*, 2014) and alternate splicing contributes to modifications to the circadian system in response to temperature and drought (James *et al.*, 2012; Filichkin *et al.*, 2015). Similarly, circadian regulation of exosome activity has previously been implicated in the circadian system of the bread mold *Neurospora crassa* (Guo *et al.*, 2009; Zhang *et al.*, 2015).

Inactivation of SAL1 by oxidative stress induces the intracellular accumulation of PAP that inhibits the activity of XRN exoribonucleases (Dichtl *et al.*, 1997; Mechold *et al.*, 2006; Estavillo *et al.*, 2011). Redundancy between XRN family members has previously been reported and the accumulated PAP in mutant alleles of *SAL1* have been suggested to simultaneously inhibit all three XRNs (Gy *et al.*, 2007; Nagarajan *et al.*, 2013). A role for cytoplasmic XRN activity within the circadian system of the green algae *Chlamydomonas reinhardtii* has previously been reported (Matsuo *et al.*, 2008). In this case, loss of XRN activity led to a lengthened FRP, similar to that observed in *sal1* and *xrn234* plants (Fig. 2 and 7). Therefore, it is possible that the loss of XRN activity induces global changes in circadian transcript abundance and/or stability that likely account for the delayed FRP of the circadian system.

Osmotic stress delays the circadian system

Many stress responses are typically associated with different times of day, leading to speculation that such responses are modulated by the circadian system (Walley *et al.*, 2007; Mizuno and Yamashino, 2008; Sanchez *et al.*, 2011; Grundy *et al.*, 2015). While the role of

the circadian system in modulating plants' tolerance of water-deprived conditions is beginning to be elucidated (Fukushima et al., 2009; Legnaioli et al., 2009; Wilkins et al., 2010; Nakamichi et al., 2016), comparatively little is understood regarding how osmotic stress influences the circadian system (Grundy et al., 2015). Multiple clock transcripts accumulate to a greater extent in response to osmotic stress in barley (*Hordeum vulgare*), although a consistent change in the phase of gene expression was not observed under diel conditions in this previous work (Habte et al., 2014). One of the consequences of drought and osmotic stress is the increased generation of reactive oxygen species during photosynthesis, leading to changes in the redox state of the chloroplast (Apel, 2004; Chan et al., 2016). Recent work has revealed that the redox status of peroxiredoxins within the chloroplast varies with a circadian rhythm (Edgar et al., 2012), while a similar circadian pattern of H₂O₂ accumulation and catalase activity is also apparent (Lai et al., 2012).

Our work demonstrated that osmotic stress is sufficient to extend circadian period in *Arabidopsis* under either cR+B, cR, or cB light, although a much more pronounced effect was observed under monochromatic blue light conditions (Fig. 1A, 8C-8D, S7). In addition, the extension of the circadian period was correlated with the accumulation of PAP in wild-type plants (Fig. 1 and 8). While we do not consider plants' perception and response to osmotic stress to occur solely via the regulation of SAL1 activity, it was noteworthy that a significant response to osmotic stress was not observed in *sall* lines transferred to cB light (Figure 8D); such data are consistent with the pronounced circadian phenotype of *sall* alleles, specifically under these conditions (Fig. 2 and 3), and reveal an additional contribution of SAL1 to plants' responses to osmotic stress.

Recently, it has been suggested that a delay of the circadian system acts to slow metabolism, consequently improving survival during sub-optimal conditions (Syed et al., 2015). Such data are consistent with our observations that increased levels of PAP (due to the mutation of SAL1 or the application of osmotic stress) lengthens FRP and delays flowering (Fig. 1, 3, 4, and 8, Wilson et al., 2009). Therefore, we propose that the accumulation of PAP in response to environmental stress leads to the inhibition of XRN exonucleases, leading to enhanced stability of specific transcripts and a consequent delay in circadian timing. This mechanism enables environmental signals to be integrated with the circadian clock to adjust plants' response to stressful conditions.

Materials and Methods

Plant materials and growth conditions

Mutant alleles of *SAL1* have been reported previously (Rossel et al., 2006; Gy et al., 2007; Rodríguez et al., 2010). *alx8-1* and *fry1-6* alleles of *SAL1*, as well as *xrn2-1* and *xrn3-3*, were re-isolated from seed provided by the Nottingham Arabidopsis Stock Centre (Scholl et al., 2000). *xrn4* alleles have previously been reported (Roman et al., 1995; Gazzani et al., 2004; Olmedo et al., 2006; Potuschak et al., 2006), as has the *xrn2-1 xrn3-3 xrn4-6* triple mutant (Hirsch et al., 2011). *apk1 apk2* lines were a kind gift from the Farmer lab (University of Lausanne, Switzerland, Rodríguez et al., 2010). *cad2-1* seeds (Cobbett et al., 1998) were provided by Prof. Phil Mullineaux (University of Essex, UK). *fry1-6 CCA1::LUC2* lines were generated by crossing *fry1-6* to a previously reported Columbia *CCA1::LUC2* line (Jones et al., 2015). Transgenic plants were generated as follows. The *SAL1* coding sequence and a 900-bp region upstream of the transcriptional start site were transferred into pCR8/GW/TOPO (Invitrogen) via the TOPO cloning method using oligonucleotides described in Table S1. A binary vector containing this *SAL1* genomic fragment was created by LR recombination with pGWB4 (Nakagawa et al., 2007) to generate pGWB4 *SAL1*. The *AHL* coding sequence was similarly transferred into pCR8/GW/TOPO using oligonucleotides described in Table S1. A binary vector containing the *AHL* cDNA fragment was created by LR recombination with pGWB41 (Nakagawa et al., 2007) to generate pGWB41 *AHL*. Plasmids were moved into *Agrobacterium tumefaciens* strain GV3101 and transformed into *alx8-1* plants using standard protocols (Narusaka et al., 2010). Transformants were selected on Murashige and Skoog (MS) media supplemented with 50 µg/mL hygromycin (Fisher Scientific).

All wild-type and transgenic lines were in the *Arabidopsis thaliana* ecotype Columbia-0 (Col-0) background. Seeds were surface sterilized and sown on soil or 0.8% agar plates containing half-strength MS medium (Sigma Aldrich M5524). For sulfate deficiency experiments, sulfate salts in MS medium M5524 (<http://www.sigmaaldrich.com/technical-documents/protocols/biology/murashige-skoog.html>) were replaced with chloride salts as follows: 10.31 mM NH₄NO₃, 0.05 mM H₃BO₃, 1.50 mM CaCl₂, 0.05 µM CoCl₂, 0.05 µM CuCl₂, 0.05 mM EDTA, 0.05 mM FeCl₃, 0.75 mM MgCl₂, 0.05 mM MnCl₂, 0.52 µM Na₂MoO₄, 2.50 µM KI, 9.40 mM KNO₃, 0.63 mM KH₂PO₄, 15 µM ZnCl₂, 0.8% agar, pH 5.7. Plants were entrained under 12 h white light: 12 h dark cycles.

Application of osmotic stress

In order to apply osmotic stress (-0.5 MPa), 5-day-old seedlings were transferred from MS plates to those containing either osmotica 24 h before transfer to constant light for imaging. Treatment plates contained either 200 mM mannitol or were infused with PEG 8000 as described by Verslues et al., 2006. In brief, 1.5% agar plates containing half-strength MS medium and 6 mM MES buffer were solidified and then overlaid with a solution of 250 g/l PEG 8000. The solution was allowed to sit for 24 h, producing an osmotic potential of -0.5 MPa, before the excess solution was removed from the plates prior to transplant.

Luciferase imaging

To complete luciferase imaging, individual seedlings were entrained for 6 days in 12 h:12 h light:dark cycles under white light on half-strength MS media without supplemental sucrose (unless transferred to constant darkness, in which case 3% (w/v) sucrose was added to the media). Plants were sprayed with 3 mM D-luciferin in 0.01% Triton X-100 before being transferred to free-running conditions under the indicated fluence rate as previously described (Litthauer et al., 2015). Data was processed using ImageJ software (Schneider et al., 2012). Patterns of bioluminescence were fitted to cosine waves using Fourier Fast Transform-Non-Linear Least Squares (FFT-NLLS, Plautz et al., 1997) to estimate the length of the circadian period. RAE is a measure of rhythmic robustness, with a value of 0 indicating an exact fit to a cosine wave (Plautz et al., 1997). Sample size was chosen to achieve a power of 0.8 in a two-sample t test at $\alpha = 0.05$. Previously collected data was used to estimate $\sigma = 0.6$.

Abscisic acid treatment

Six-day-old seedlings were entrained for luciferase imaging (as described above) before being transplanted to half-strength MS media plates containing either 10 μ M abscisic acid (Acros Organics #133485000) or a mock treatment (0.1% DMSO) at dawn (ZT0). Plants were sprayed with 3 mM D-luciferin in 0.01% Triton X-100 before being transferred to free-running conditions under a combination of 30 μ mol m⁻² s⁻¹ red light and 20 μ mol m⁻² s⁻¹ blue light for circadian imaging.

Methyl viologen treatment

Six-day-old seedlings were entrained for luciferase imaging (as described above) before being sprayed with 50 μ M methyl viologen (Sigma Aldrich) and 3 mM D-luciferin in 0.01%

Triton X-100 at dawn (ZT0). Plants were transferred to free-running conditions under a combination of 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ red light and 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue light for circadian imaging.

Application of PAP to seedlings

Twelve-day-old seedlings were prepared for luciferase imaging (as described above) and transferred into constant blue light (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$). 1 mM PAP was applied to seedlings in 0.01% Triton X-100 at Zeitgeber (ZT) 29. Luciferase imaging and circadian analysis was completed as described above.

Chlorophyll fluorescence imaging

Chlorophyll fluorescence parameters were recorded with a Fluorimager imaging system (Technologica Ltd, UK) as previously described (Litthauer et al., 2015). Patterns of F_q'/F_m' were fitted to cosine waves using FFT-NLLS (Plautz et al., 1997) to estimate circadian period length and additional circadian parameters. Sample size was chosen to achieve a power of 0.8 in a two-sample t test at $\alpha = 0.05$. Previously collected data was used to estimate $\sigma = 0.6$.

RT-qPCR

Following entrainment, plants were transferred to constant light at the indicated fluence rate and quality. Tissue was harvested at the indicated time directly onto liquid nitrogen before RNA was isolated from 10 to 15 seedlings for each data point using Tri Reagent according to the manufacturer's protocol (Sigma Aldrich, Dorset, UK, <http://www.sigmaaldrich.com>). Reverse transcription was performed using RevertAid reverse transcriptase following DNase treatment (Fisher Scientific, Loughborough, UK). RT-qPCR was performed using a BioRad CFX96 Real-Time system following MIQE guidelines (Bustin et al., 2009). PCR was completed for 40 cycles using the following protocol: 95°C for 15 s, 55°C for 15 s, 72°C for 30 s. Each biological sample was run in triplicate, with starting quantity estimated from critical thresholds using the standard curve of amplification using BioRad CFX Manager 5.1. Calibration curves were run as an internal control within each RT-qPCR run, with data only accepted if experimental samples fell within the linear range of amplification, and if quality criteria were met ($r^2 > 0.97$, PCR efficiency $\pm 15\%$, as determined from the calibration curve). Data for each sample were normalized to *PP2A* as an internal control. Primer sets used are described in Table S1.

Protein extraction and immunoblot analysis

Twelve-day-old seedlings were frozen in liquid nitrogen, ground into powder, and extracted in homogenisation buffer (25 mM MOPS, 0.25 M sucrose, 0.1 mM MgCl₂, 8 mM L-Cys, pH 7.8). After quantifying the total protein concentrations with Bradford Reagent (Sigma Aldrich), equal amounts of proteins were separated on 12.5% SDS-PAGE gels and then semi-dry transferred onto a 0.45 µm nitrocellulose membrane (Amersham). SAL1-GFP and actin were immunodetected by anti-GFP (Ab290, 1:10 000 dilution, Abcam) and anti-actin (mAB1501, 1:2 000, Sigma Aldrich) antibodies, respectively. IgG (H+L) HRP conjugates (Promega) were used to detect the primary antibodies. Immunoreactive bands were quantified by scanning the membrane with a Fusion FX imaging system (Vilber Lourmat).

Extraction and HPLC analysis of PAP

PAP was extracted from whole seedlings as previously described (Bürstenbinder et al., 2007; Estavillo et al., 2011). Metabolites were extracted from 150-300 mg ground tissue using 1 mL 0.1 M HCl with incubation on ice for 15 min, and centrifuged twice at 16,000 x g at 4°C for 5 min. 150 µL of the supernatant was added to 770 µL CP buffer (620 mM citric acid and 760 mM Na₂HPO₄, pH 4) and derivatised using 80 µL 50% (w/v) chloroacetaldehyde solution with incubation at 80°C for 10 min, and centrifuged for 45 min at 16,000 x g at 4°C. Analysis of PAP was performed as previously described (Bürstenbinder et al., 2007; Estavillo et al., 2011). 20 µL of the supernatant was injected into an Agilent 1100 HPLC system connected to a FLD G1321A (Agilent) fluorescent detector. PAP was analysed by reverse-phase HPLC using a Luna 5 µm C18(2) 100 Å column (Phenomenex). The column was equilibrated for 0.2 min with 95% (v/v) of buffer A (5.7mM [CH₃(CH₂)₃]₄NHSO₄ and 30.5 mM KH₂PO₄, pH 5.8) and 5% (v/v) buffer B (67% [v/v] acetonitrile and 33% [v/v] buffer A), followed by a linear gradient for 53 min up to 50% (v/v) of buffer B. The column was re-equilibrated for 7 min with 5% (v/v) buffer B. PAP concentration was calculated relative to a commercially available standard (Santa Cruz Biotechnology, sc-210760).

Accession numbers

Genes examined in this article can be found in the Arabidopsis Genome Initiative database under the following accession numbers: *APK1*, At2g14750; *APK2*, At4g39940; *AHL*, At5g54390; *CAD2/GSH1*, At4g23100; *CCA1*, At2g46830; *CRY1*, At4g08920; *CRY2*, At1g04400; *ELF4*, At2g40080; *GIGANTEA*, At1g22770; *LHY*, At1g01060; *PRR5*,

At5g24470; *SAL1*, At5g63980; *TOC1*, At5g61380; *XRN2*, At5g42540; *XRN3*, At1g75660; *XRN4*, At1g54490; *ZTL*, At5g57360.

Supplemental Data

Figure S1. *sal1* alleles have an extended circadian period.

Figure S2. Abundance of clock transcripts in *sal1* seedlings.

Figure S3. Abundance of blue photoreceptors in *sal1* seedlings.

Figure S4. PAP accumulation under constant red light.

Figure S5. Transcript accumulation of *SAL1* and *XRN* ribonucleases in entraining and constantly lit conditions.

Figure S6. Sulfate deprivation induces the accumulation of genes associated with sulfur anabolism.

Figure S7. Circadian rhythms in response to osmotic stress under constant red light.

Figure S8. Abundance of clock transcripts in response to osmotic stress.

Table S1. Oligos used in this study.

Acknowledgements

We thank Prof. Barry Pogson (ANU, Australia) and Dr. Markus Wirtz (Universitat Heidelberg), as well as Drs. Uli Bechold and Stuart Fisk (University of Essex) for useful discussions. We are grateful to Prof. Edward Farmer (University of Lausanne, Switzerland), Drs. Pascal Genschik and Thomas Potuschak (CNRS, France), Prof. Antony Hall (Earlham Institute, UK), Prof. Phil Mullineaux (University of Essex, UK), Prof. Albrecht von Arnim (University of Tennessee, USA), and Prof. Alex Webb (University of Cambridge, UK) for the kind gifts of seed.

Figure legends

Figure 1. Osmotic stress induces the accumulation of PAP and extends the circadian period. (A) Accumulation of PAP in *Columbia* (Col-0) seedlings in the presence of 200 mM mannitol under constant light conditions. Plants were grown for 11 days under 12:12 h L/D cycles before being transferred to plates containing 200 mM mannitol 24 hours before transfer to constant light. Plates were transferred to either 24 $\mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue light supplemented with 36 $\mu\text{mol m}^{-2} \text{s}^{-1}$ red light (cR+B) or to 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ constant white light (cW) at dawn of day 12. Seedlings were harvested at ZT96. Data are the mean of three biological replicates and are representative of two independent experiments. Error bars indicate standard deviation. Asterisks indicate a significant difference compared with the respective mock control ($p < 0.025$, Bonferroni adjusted Student's t test). (B) Representative bioluminescence data of luciferase activity in *Columbia* (Col-0) plants carrying a *CCA1::LUC2* reporter construct in the presence of 200 mM mannitol or PEG 8000. Plants were grown on half-strength MS medium for 5 days under 12:12 h L/D cycles before being transferred to either a mock-treated control, 200 mM mannitol, or plates infused with PEG 8000 24 hrs before imaging under cR+B light (30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ red and 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue light). Data are representative of three independent experiments. Error bars represent standard error of the mean and are presented every 10 hrs for clarity. $n=10$. (C) Circadian period estimates of luciferase activity in the presence of 200 mM mannitol or PEG 8000. *Columbia* (Col-0) seedlings carrying either the *CCA1*, *LHY*, or *TOC1* promoters fused to a *LUCIFERASE* reporter were assessed. Data are representative of three independent experiments. Error bars express standard error of the mean, $n=10$. Asterisks indicate $p < 0.01$ compared with mock controls (Dunnett's test).

Figure 2. *sal1* alleles have an extended circadian period. Rhythms of PSII operating efficiency (F_q'/F_m') measured over circadian time in *alx8-1* (A) and *fry1-6* (C) mutant alleles of *SAL1*. Plants were grown for 12 days under 12:12 h L/D cycles before being transferred to constant blue light (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Data represent mean values of multiple seedlings ($n=8$) and are representative of at least three independent experiments. Standard error of the mean is presented every 5 hours for clarity. (B, D) Circadian period estimates of F_q'/F_m' circadian rhythms presented in (A and C) using FFT-NLLS with baseline de-trending (Plautz et al., 1997). Asterisks indicate a significant difference compared with the Col-0 control ($p < 0.05$, Student's t test). (E) Circadian period estimates of F_q'/F_m' circadian rhythms in *Columbia* (Col-0), *alx8-1*, and *alx8-1* seedlings transformed with a *SAL1::SAL1-GFP* construct. Period

estimates are plotted against Relative Amplitude Error (RAE), which is a measure of rhythmic robustness (a value of 0 indicates an exact fit to a cosine wave, Plautz et al., 1997), $n = 8$. Data from one of three independent experiments are shown. Asterisks indicate a significant difference compared with the *alx8-1* mutant ($p < 0.05$, Dunnett's test). **(F)** Circadian period estimates of luciferase activity in Columbia (Col-0) and *fry1-6* plants carrying a *CCA1::LUC2* reporter construct. Plants were grown on half-strength MS medium for 6 days before transfer to either $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light (cB), a combination of $30 \mu\text{mol m}^{-2} \text{s}^{-1}$ red light and $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light (cR+B), or $30 \mu\text{mol m}^{-2} \text{s}^{-1}$ red light (cR). Plants transferred to constant darkness (Darkness) were grown on half-strength MS medium supplemented with 3% (w/v) sucrose. Data are representative of at least three independent experiments. Error bars express standard error of the mean, $n=10$. Asterisks indicate a significant difference compared with the Columbia control ($p < 0.025$, Bonferroni adjusted Student's t test). **(G)** Circadian period estimates of luciferase activity in *Columbia* (Col-0) and *fry1-6* plants carrying a *CCA1::LUC2* reporter construct in the presence of methyl viologen. Plants were grown on half-strength MS medium for 6 days before application of $50 \mu\text{M}$ methyl viologen and transfer to $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light and $30 \mu\text{mol m}^{-2} \text{s}^{-1}$ red light for imaging. Data are representative of three independent experiments. Error bars express standard error of the mean, $n = 10$. Asterisks indicate $p < 0.025$ compared with the respective Columbia control (post-hoc Bonferroni adjusted Student's t test). **(H)** Circadian period estimates of luciferase activity in the presence of $10 \mu\text{M}$ ABA. Data are representative of three independent experiments. Error bars express standard error of the mean, $n=10$. Asterisks indicate a significant difference compared with indicated controls ($p < 0.025$, post-hoc Bonferroni adjusted Student's t test).

Figure 3. The sal1 circadian phenotype is exacerbated under blue light. **(A)** Fluence rate response curves to measure the free-running circadian period under constant blue (cB) light in Columbia (Col-0) and *fry1-6* seedlings carrying a *CCA1::LUC2* reporter. Seedlings were entrained in 12:12 h L/D cycles for 6 days before being transferred to the indicated fluence rate of cB light. Data are representative of three independent experiments, standard error of the mean is shown, $n = 10$. Asterisks highlight a significant difference between Col-0 and *fry1-6* at the indicated fluence rate (post-hoc Student's t-test, $p < 0.05$). **(B)** Fluence rate response curves to measure the free-running circadian period under constant red (cR) light. Seedlings were entrained as described in (A) before being transferred to the indicated fluence

rate of constant red light. Data are representative of three independent experiments, standard error of the mean is shown, $n = 10$. **(C-F)** Accumulation of circadian clock-regulated transcripts under constant blue (cB) (C, E) or constant red (cR) (D, F) light in *salI* seedlings using RT-qPCR. Levels of *CCA1* (C, D) and *TOC1* (E, F) mRNA were assessed. Plants were entrained to 12:12 h light:dark cycles for 12 days on MS medium before being moved to constant conditions with either $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue (cB) or $30 \mu\text{mol m}^{-2} \text{s}^{-1}$ red (cR) light. Data for each gene were normalized with an internal control (*PP2a*) and are the mean of at least two biological replicates. Error bars indicate standard error of the mean. **(G, H, I)** Accumulation of PAP in *fryI-6* seedlings under different fluence rates of blue (G), red (H), or red+blue (I) light. Seedlings were entrained in 12:12 h L/D cycles for 12 days before being transferred to the indicated fluence rate and quality of light for 4 days. Seedlings were harvested at ZT96. UD, PAP levels were below the detection threshold in Col-0 in each measurement. Asterisks highlight a significant difference for the selected comparison ($P < 0.025$, Bonferroni adjusted Student's T-test).

Figure 4. PAP accumulation is correlated with circadian defects in *salI* mutants. **(A)** Circadian period of *fryI-6* seedlings following application of PAP. Seedlings were entrained in 12:12 h L/D cycles for 6 days before being transferred to $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue (cB) light. PAP was applied at ZT29. Asterisks indicate a significant difference compared with the mock-treated control ($p < 0.025$, Bonferroni adjusted Student's t test). Standard error of the mean is shown, $n > 19$. Data are representative of three independent experiments. **(B)** Accumulation of PAP in Columbia (Col-0), *alx8-1*, and *alx8-1* seedlings transformed with a *35S::AHL* construct. Plants were grown for 12 days under 12:12 h light:dark cycles before being transferred to constant $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue (cB) light. Seedlings were harvested at ZT96. Data are the mean of three biological replicates and are representative of two independent experiments. UD, PAP levels were below the detection threshold. **(C, D)** F_q/F_m rhythms and circadian period estimates in *alx8-1* seedlings transformed with a *35S::AHL* construct. Plants were grown for 12 days under 12:12 h L/D cycles before being transferred to constant blue (cB) light ($20 \mu\text{mol m}^{-2} \text{s}^{-1}$). Data are representative of three independent experiments, standard error of the mean is shown, $n=8$. Asterisks indicate a significant difference compared with *alx8-1* plants ($p < 0.0125$, Bonferroni adjusted Student's t test).

Figure 5. *SAL1* transcript and protein accumulation are not altered by the circadian system. (A) *SAL1* transcript accumulation in *Columbia* (Col-0) plants under 16:8 h L/D cycles. Data were normalized to an internal control (*PP2a*). Data are the average of three biological replicates, error bars show standard error of the mean. (B) Immunoblot analysis of *SAL1* protein levels under 16:8 h LD cycles. Plants were grown as described in (A). Data are the average of three biological replicates, error bars show standard error of the mean. (C) *SAL1* transcript accumulation over circadian time. *Columbia* (Col-0) plants were entrained under 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ white light in 12:12 h L/D cycles for 12 days before being transferred to constant white (cW) light. Data are the mean of three biological replicates. Error bars express standard error of the mean.

Figure 6. Sulfate deprivation does not extend circadian period. (A) Schematic of sulfate metabolism in *Arabidopsis*, adapted from (Bohrer et al., 2014). Abbreviations of metabolites: APS, adenosine-5'-phosphosulfate; GLS, glucosinolate; GSH, glutathione; PAP, 5'-phosphoadenosine 3'-phosphate; PAPS, 3-phosphoadenosine 5-phosphosulfate. Abbreviations of enzymes and transporters: APK, APS KINASE; APR, APS REDUCTASE; GSH1, GLUTAMATE-CYSTEINE LIGASE. (B) Circadian period estimates of luciferase activity in the presence or absence of sulfate salts. *Columbia* (Col-0) seedlings carrying either the *CCA1*, *LHY*, or *TOC1* promoters fused to a *LUCIFERASE* reporter were assessed. Plants were entrained for 12 d before transfer to constant blue (cB) light (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$) for imaging. Standard error of the mean is shown, $n = 10$. Data from one of three independent experiments are shown. (C) Circadian rhythms of F_q'/F_m' in plants grown on agar lacking sulfate salts. Plants were grown as described in (B). Data represent mean values of multiple seedlings ($n=8$) and are representative of at least three independent experiments. (D, E) Circadian period estimates of F_q'/F_m' in *apk1 apk2* (D) and *cad2-1* (E) plants. Data are representative of at least two independent experiments. Standard error of the mean is shown, $n = 8$.

Figure 7. Circadian phenotypes of *xrn* mutants. (A) Rhythms of PSII operating efficiency (F_q'/F_m') were measured over circadian time in *Columbia* (Col-0) and *xrn2-1 xrn3-3 xrn4-6* (*xrn234*) mutant seedlings. Period estimates are plotted against Relative Amplitude Error (RAE). Plants were grown for 12 days under 12:12 h L/D cycles before being transferred to constant blue (cB) light (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Data represent mean values of multiple seedlings

(n = 7) and are representative of three independent experiments. Error bars indicate standard error, asterisks indicate a significant difference compared with Col-0 plants ($p < 0.05$, Student's t test). **(B)** Assessment of *CCA1* transcript accumulation under constant blue (cB) light in *Columbia* (Col-0), *fry1-6*, and *xrn234* seedlings using RT-qPCR. Plants were entrained in 12:12 h L/D cycles before being moved to constant conditions with $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light. Data for each gene were normalized with an internal control (*PP2a*) and are the mean of three biological replicates. **(C)** *CCA1* transcript accumulation under constant white (cW) light in *fry1-6* and *xrn234* seedlings. Transcript accumulation in *Columbia* (Col-0), *fry1-6*, and *xrn234* seedlings was compared using RT-qPCR. Plants were entrained as in (B) before being moved to constant conditions with $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ white light. Data for each gene were normalized with an internal control (*PP2a*) and are the mean of three biological replicates. Error bars indicate standard error.

Figure 8. PAP levels correlate with a lengthened circadian period under osmotic stress and broad-spectrum blue light. **(A, B)** Accumulation of PAP in *Columbia* (Col-0) and *fry1-6* seedlings in the presence of 200 mM mannitol. Plants were grown for 11 days under 12:12 h L/D cycles before being transferred to plates containing 200 mM mannitol 24 hours before transfer to constant light. Plates were subsequently transferred to either constant red and blue light ($30 \mu\text{mol m}^{-2} \text{s}^{-1}$ constant red light supplemented with $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light (cR+B, A) or to $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue light (cB, B) at dawn of day 12. Seedlings were harvested at ZT96. UD, PAP levels were below the detection threshold. Data are the mean of three biological replicates and are representative of two independent experiments. Standard deviation is shown. Asterisks indicate a significant difference compared to a mock-treated control ($p < 0.025$, Bonferroni adjusted Student's t test). **(C, D)** Circadian period estimates of luciferase activity in *Columbia* (Col-0) and *fry1-6* plants carrying a *CCA1::LUC2* reporter construct in the presence of 200 mM mannitol. Plants were entrained and transferred to growth substrate containing 200 mM mannitol as in (A). Seedlings were transferred to either constant red and blue light (cR+B, C) or constant blue light (cB, D) for imaging. Data are representative of three independent experiments. Error bars express standard error of the mean, n = 10. Asterisks indicate a significant difference compared with a mock-treated control ($p < 0.025$, Bonferroni adjusted Student's t test). **(E)** Accumulation of *CCA1* transcript following transfer to 200 mM mannitol. Plants were entrained to 12:12 h light:dark cycles for 11 days on MS medium before being transferred to either 200 mM mannitol or a mock

treatment at dawn. Seedlings were moved to constant conditions with 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue light at dawn of day 12. Data were normalized with an internal control (*PP2a*) and are the mean of at least two biological replicates. Error bars indicate standard error of the mean.

Literature cited

- Adams S, Manfield I, Stockley P, Carré IA (2015) Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. PLoS ONE 10: e0143943
- Alabadi D, Oyama T, Yanovsky M, Harmon F, Mas P, Kay S (2001) Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. Science 293: 880-883
- Apel K (2004) REACTIVE OXYGEN SPECIES: Metabolism, Oxidative Stress, and Signal Transduction. Annu Rev Plant Biol 55: 373-399
- Bohrer A-S, Kopriva S, Takahashi H (2014) Plastid-cytosol partitioning and integration of metabolic pathways for APS/PAPS biosynthesis in Arabidopsis thaliana. Frontiers in Plant Science 5: 751
- Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT (2009) The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. Clinical Chemistry 55: 611-622
- Bürstenbinder K, Rzewuski G, Wirtz M, Hell R, Sauter M (2007) The role of methionine recycling for ethylene synthesis in Arabidopsis. Plant J 49: 238-249
- Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroehler E, Grassl J, Arlt W, Estavillo GM, Jackson CJ, Pogson BJ (2016) Sensing and signaling of oxidative stress in chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase. PNAS 113: E4567-4576
- Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ (2016) Learning the Languages of the Chloroplast: Retrograde Signaling and Beyond. Ann Rev Plant Biol 67: 25-53
- Chen H, Xiong L (2010) The bifunctional abiotic stress signalling regulator and endogenous RNA silencing suppressor FIERY1 is required for lateral root formation. Plant, Cell & Env 33: 2180-2190
- Chen H, Zhang B, Hicks LM, Xiong L (2011) A nucleotide metabolite controls stress-responsive gene expression and plant development. PLoS ONE 6: e26661
- Chen Y-Y, Wang Y, Shin L-J, Wu J-F, Shanmugam V, Tsednee M, Lo J-C, Chen C-C, Wu S-H, Yeh K-C (2013) Iron is involved in the maintenance of circadian period length in Arabidopsis. Plant Physiol 161: 1409-1420
- Chow BY, Helfer A, Nusinow DA, Kay SA (2012) ELF3 recruitment to the PRR9 promoter requires other Evening Complex members in the Arabidopsis circadian clock. Plant Sig Beh 7: 170-173
- Christie JM, Blackwood L, Petersen J, Sullivan S (2015) Plant Flavoprotein Photoreceptors. Plant & Cell Phys 56: 401-413
- Cobbett CS, May MJ, Howden R, Rolls B (1998) The glutathione-deficient, cadmium-sensitive mutant, cad2-1, of Arabidopsis thaliana is deficient in gamma-glutamylcysteine synthetase. Plant J 16: 73-78
- Covington MF, Maloof JN, Straume M, Kay SA, Harmer SL (2008) Global transcriptome analysis reveals circadian regulation of key pathways in plant growth and development. Genome Biology 9: R130

819 Dichtl B, Stevens A, Tollervey D (1997) Lithium toxicity in yeast is due to the inhibition of
820 RNA processing enzymes. *EMBO J* 16: 7184-7195

821 Dong MA, Farré EM, Thomashow MF (2011) Circadian clock-associated 1 and late
822 elongated hypocotyl regulate expression of the C-repeat binding factor (CBF)
823 pathway in *Arabidopsis*. *PNAS* 108: 7241-7246

824 Edgar RS, Green EW, Zhao Y, Van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja
825 UK, Feeney KA, Maywood ES, Hastings MH, Baliga NS, Merrow M, Millar AJ,
826 Johnson CH, Kyriacou CP, O'Neill JS, Reddy AB (2012) Peroxiredoxins are
827 conserved markers of circadian rhythms. *Nature* 485: 459-464

828 Eriksson ME, Webb AAR (2011) Plant cell responses to cold are all about timing. *Current*
829 *Opin Plant Biol* 14: 731-737

830 Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E, Whelan
831 J, David P, Javot H, Brearley C, Hell R, Marin E, Pogson BJ (2011) Evidence for a
832 SAL1-PAP Chloroplast Retrograde Pathway That Functions in Drought and High
833 Light Signaling in *Arabidopsis*. *Plant Cell* 23: 3992-4012

834 Fankhauser C, Staiger D (2002) Photoreceptors in *Arabidopsis thaliana*: light perception,
835 signal transduction and entrainment of the endogenous clock. *Planta* 216: 1-16

836 Farré EM, Kay SA (2007) PRR7 protein levels are regulated by light and the circadian clock
837 in *Arabidopsis*. *Plant J* 52: 548-560

838 Filichkin SA, Cumbie JS, Dharmawardhana P, Jaiswal P, Chang JH, Palusa SG, Reddy ASN,
839 Megraw M, Mockler TC (2015) Environmental Stresses Modulate Abundance
840 and Timing of Alternatively Spliced Circadian Transcripts in *Arabidopsis*. *Molecular*
841 *Plant* 8: 207-227

842 Fukushima A, Kusano M, Nakamichi N, Kobayashi M, Hayashi N, Sakakibara H, Mizuno T,
843 Saito K (2009) Impact of clock-associated *Arabidopsis* pseudo-response regulators in
844 metabolic coordination. *PNAS* 106: 7251-7256

845 Garbarino-Pico E, Green CB (2007) Posttranscriptional regulation of mammalian circadian
846 clock output. *Cold Spring Harb Symp Quant Biol* 72: 145-156

847 Gazzani S, Lawrenson T, Woodward C, Headon D, Sablowski R (2004) A link between
848 mRNA turnover and RNA interference in *Arabidopsis*. *Science* 306: 1046-1048

849 Gendron JM, Pruneda-Paz JL, Doherty CJ, Gross AM, Kang SE, Kay SA (2012) *Arabidopsis*
850 circadian clock protein, TOC1, is a DNA-binding transcription factor. *PNAS* 109:
851 3167-3172

852 Goodspeed D, Chehab EW, Min-Venditti A, Braam J, Covington MF (2012) *Arabidopsis*
853 synchronizes jasmonate-mediated defense with insect circadian behavior. *PNAS* 109:
854 4674-4677

855 Grundy J, Stoker C, Carré IA (2015) Circadian regulation of abiotic stress tolerance in plants.
856 *Frontiers in Plant Science* 6: 201

857 Guo J, Cheng P, Yuan H, Liu Y (2009) The exosome regulates circadian gene expression in a
858 posttranscriptional negative feedback loop. *Cell* 138: 1236-1246

859 Gy I, Gascioli V, Laressergues D, Morel J-B, Gombert J, Proux F, Proux C, Vaucheret H,
860 Mallory AC (2007) *Arabidopsis* FIERY1, XRN2, and XRN3 are endogenous RNA
861 silencing suppressors. *Plant Cell* 19: 3451-3461

862 Habte E, Müller LM, Shtaya M, Davis SJ, von Korff M (2014) Osmotic stress at the barley
863 root affects expression of circadian clock genes in the shoot. *Plant, Cell & Env* 37:
864 1321-1327

865 Hanano S, Domagalska MA, Nagy F, Davis SJ (2006) Multiple phytohormones influence
866 distinct parameters of the plant circadian clock. *Genes Cells* 11: 1381-1392

867 Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland G,
868 Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY

- FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. *Plant Cell* 24: 428-443
- Hirsch J, Misson J, Crisp PA, David P, Bayle V, Estavillo GM, Javot H, Chiarenza S, Mallory AC, Maizel A, Declerck M, Pogson BJ, Vaucheret H, Crespi M, Desnos T, Thibaud M-C, Nussaume L, Marin E (2011) A novel *fry1* allele reveals the existence of a mutant phenotype unrelated to 5'→3' exoribonuclease (XRN) activities in *Arabidopsis thaliana* roots. *PLoS ONE* 6: e16724
- Hong S, Kim SA, Guerinot ML, McClung CR (2013) Reciprocal Interaction of the Circadian Clock with the Iron Homeostasis Network in *Arabidopsis*. *Plant Physiol* 161: 893-903
- Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. *Trends in Plant Science* 19: 240-249
- Huang W, Perez-Garcia P, Pokhilko A, Millar AJ, Antoshechkin I, Riechmann JL, Mas P (2012) Mapping the core of the *Arabidopsis* circadian clock defines the network structure of the oscillator. *Science* 336: 75-79
- Hut RA, Beersma DGM (2011) Evolution of time-keeping mechanisms: early emergence and adaptation to photoperiod. *Philos Trans R Soc Lond B Biol Sci* 366: 2141-2154
- Ingle RA, Stoker C, Stone W, Adams N, Smith R, Grant M, Carré I, Roden LC, Denby KJ (2015) Jasmonate signalling drives time-of-day differences in susceptibility of *Arabidopsis* to the fungal pathogen *Botrytis cinerea*. *Plant J* 84: 937-948
- James AB, Syed NH, Bordage S, Marshall J, Nimmo GA, Jenkins GI, Herzyk P, Brown JWS, Nimmo HG (2012) Alternative splicing mediates responses of the *Arabidopsis* circadian clock to temperature changes. *Plant Cell* 24: 961-981
- Jones M (2009) Entrainment of the *Arabidopsis* circadian clock. *J Plant Biol* 52: 202-209
- Jones M (2017) Interplay of Circadian Rhythms and Light in the Regulation of Photosynthesis-Derived Metabolism. *In*, Vol 79. https://link.springer.com/chapter/10.1007/124_2017_2, p In press
- Jones M, Hu W, Litthauer S, Lagarias JC, Harmer SL (2015) A Constitutively Active Allele of Phytochrome B Maintains Circadian Robustness in the Absence of Light. *Plant Physiology* 169: 814-825
- Jones M, Williams BA, McNicol J, Simpson CG, Brown JWS, Harmer SL (2012) Mutation of *Arabidopsis* spliceosomal timekeeper locus1 causes circadian clock defects. *Plant Cell* 24: 4066-4082
- Kastenmayer JP, Green PJ (2000) Novel features of the XRN-family in *Arabidopsis*: evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. *PNAS* 97: 13985-13990
- Kerwin RE, Jiménez-Gómez JM, Fulop D, Harmer SL, Maloof JN, Kliebenstein DJ (2011) Network quantitative trait loci mapping of circadian clock outputs identifies metabolic pathway-to-clock linkages in *Arabidopsis*. *Plant Cell* 23: 471-485
- Kim B-H, von Arnim AG (2009) FIERY1 regulates light-mediated repression of cell elongation and flowering time via its 3'(2'),5'-bisphosphate nucleotidase activity. *Plant J* 58: 208-219
- Kojima S, Shingle DL, Green CB (2011) Post-transcriptional control of circadian rhythms. *J Cell Sci* 124: 311-320
- Kurihara Y, Schmitz RJ, Nery JR, Schultz MD, Okubo-Kurihara E, Morosawa T, Tanaka M, Toyoda T, Seki M, Ecker JR (2012) Surveillance of 3' Noncoding Transcripts Requires FIERY1 and XRN3 in *Arabidopsis*. *G3* 2: 487-498
- Lai AG, Doherty CJ, Mueller-Roeber B, Kay SA, Schippers JHM, Dijkwel PP (2012) CIRCADIAN CLOCK-ASSOCIATED 1 regulates ROS homeostasis and oxidative stress responses. *PNAS* 109: 17129-17134

- Lee B-R, Huseby S, Koprivova A, Chételat A, Wirtz M, Mugford ST, Navid E, Brearley C, Saha S, Mithen R, Hell R, Farmer EE, Kopriva S (2012) Effects of *fou8/fry1* Mutation on Sulfur Metabolism: Is Decreased Internal Sulfate the Trigger of Sulfate Starvation Response? *PLoS ONE* 7: e39425
- Lee HG, Más P, Seo PJ (2016) MYB96 shapes the circadian gating of ABA signaling in *Arabidopsis*. *Scientific Reports* 6: 17754
- Legnaioli T, Cuevas J, Más P (2009) TOC1 functions as a molecular switch connecting the circadian clock with plant responses to drought. *EMBO J* 28: 3745-3757
- Li J, Li G, Wang H, Wang Deng X (2011) Phytochrome signaling mechanisms. *The Arabidopsis book / American Society of Plant Biologists* 9: e0148
- Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust circadian oscillation of PSII operating efficiency under blue light. *Plant J* 83: 1034-1045
- Liu T, Carlsson J, Takeuchi T, Newton L, Farré EM (2013) Direct regulation of abiotic responses by the *Arabidopsis* circadian clock component PRR7. *Plant J* 76: 101-114
- Lu SX, Webb CJ, Knowles SM, Kim SHJ, Wang Z, Tobin EM (2012) CCA1 and ELF3 Interact in the control of hypocotyl length and flowering time in *Arabidopsis*. *Plant Physiol* 158: 1079-1088
- Macgregor DR, Gould P, Foreman J, Griffiths J, Bird S, Page R, Stewart K, Steel G, Young J, Paszkiewicz K, Millar AJ, Halliday KJ, Hall AJ, Penfield S (2013) HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENES1 Is Required for Circadian Periodicity through the Promotion of Nucleo-Cytoplasmic mRNA Export in *Arabidopsis*. *Plant Cell* 25: 4391-4404
- Matsuo T, Okamoto K, Onai K, Niwa Y, Shimogawara K, Ishiura M (2008) A systematic forward genetic analysis identified components of the *Chlamydomonas* circadian system. *Genes & Development* 22: 918-930
- McWatters H, Kolmos E, Hall A, Doyle M, Amasino R, Gyula P, Nagy F, Millar A, Davis S (2007) ELF4 is required for oscillatory properties of the circadian clock. *Plant Physiol* 144: 391-401
- Mechold U, Ogryzko V, Ngo S, Danchin A (2006) Oligoribonuclease is a common downstream target of lithium-induced pAp accumulation in *Escherichia coli* and human cells. *Nucleic Acids Research* 34: 2364-2373
- Millar AJ (2016) The Intracellular Dynamics of Circadian Clocks Reach for the Light of Ecology and Evolution. *Ann Rev Plant Biol* 67: 595-618
- Mittler R, Vanderauwera S, Suzuki N, MILLER G, Tognetti VB, Vandepoele K, Gollery M, Shulaev V, Van Breusegem F (2011) ROS signaling: the new wave? *Trends in plant science* 16: 300-309
- Mizuno T, Yamashino T (2008) Comparative transcriptome of diurnally oscillating genes and hormone-responsive genes in *Arabidopsis thaliana*: insight into circadian clock-controlled daily responses to common ambient stresses in plants. *Plant Cell Physiol* 49: 481-487
- Mockler T, Michael T, Priest H, Shen R, Sullivan C, Givan S, McEntee C, Kay S, Chory J (2007) The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. *Cold Spring Harb Symp Quant Biol* 72: 353-363
- Mugford SG, Yoshimoto N, Reichelt M, Wirtz M, Hill L, Mugford ST, Nakazato Y, Noji M, Takahashi H, Kramell R, Gigolashvili T, Flügge U-I, Wasternack C, Gershenzon J, Hell R, Saito K, Kopriva S (2009) Disruption of adenosine-5'-phosphosulfate kinase in *Arabidopsis* reduces levels of sulfated secondary metabolites. *Plant Cell* 21: 910-927

967 Nagarajan VK, Jones CI, Newbury SF, Green PJ (2013) XRN 5'→3' exoribonucleases:
968 structure, mechanisms and functions. *Biochimica et Biophysica Acta* 1829: 590-603

969 Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka
970 K, Jinbo T, Kimura T (2007) Development of series of gateway binary vectors,
971 pGWBs, for realizing efficient construction of fusion genes for plant transformation. *J*
972 *Biosci Bioeng* 104: 34-41

973 Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-
974 RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in the
975 *Arabidopsis* circadian clock. *Plant Cell* 22: 594-605

976 Nakamichi N, Takao S, Kudo T, Kiba T, Wang Y, Kinoshita T, Sakakibara H (2016)
977 Improvement of *Arabidopsis* biomass and cold-, drought-, and salinity-stress
978 tolerance by modified circadian clock-associated PSEUDO-RESPONSE
979 REGULATORS. *Plant & Cell Physiology* 57: 1085-1097

980 Narusaka M, Shiraishi T, Iwabuchi M, Narusaka Y (2010) The floral inoculating protocol: a
981 simplified *Arabidopsis thaliana* transformation method modified from floral dipping.
982 *Plant Biotechnology* 27: 349-351

983 Norén, L., Kindgren, P., Stachula, P., Rühl, M., Eriksson, M. E., Hurry, V., & Strand, Å.
984 (2016). Circadian and Plastid Signaling Pathways Are Integrated to Ensure Correct
985 Expression of the CBF and COR Genes during Photoperiodic Growth. *Plant*
986 *Physiology*, 171: 1392–1406

987 Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farré EM, Kay SA
988 (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of
989 hypocotyl growth. *Nature* 475: 398-402

990 Olmedo G, Guo H, Gregory BD, Nourizadeh SD, Aguilar-Henonin L, Li H, An F, Guzman P,
991 Ecker JR (2006) ETHYLENE-INSENSITIVE5 encodes a 5'→3' exoribonuclease
992 required for regulation of the EIN3-targeting F-box proteins EBF1/2. *PNAS* 103:
993 13286-13293

994 Perez-Santángelo S, Mancini E, Francey LJ, Schlaen RG, Chernomoretz A, Hogenesch JB,
995 Yanovsky MJ (2014) Role for LSM genes in the regulation of circadian rhythms.
996 *PNAS* 111: 15166-15171

997 Plautz JD, Straume M, Stanewsky R, Jamison CF, Brandes C, Dowse HB, Hall JC, Kay SA
998 (1997) Quantitative analysis of *Drosophila* period gene transcription in living animals.
999 *J Biol Rhythms* 12: 204-217

1000 Pornsiriwong W, Estavillo GM, Chan KX, Tee EE, Ganguly D, Crisp PA, Phua SY, Zhao C,
1001 Qiu J, Park J, Yong MT, Nisar N, Yadav AK, Schwessinger B, Rathjen J, Cazzonelli
1002 CI, Wilson PB, Gilliam M, Chen Z-H, Pogson BJ (2017) A chloroplast retrograde
1003 signal, 3'-phosphoadenosine 5'-phosphate, acts as a secondary messenger in abscisic
1004 acid signaling in stomatal closure and germination. *eLife* 6: e23361

1005 Potuschak T, Vansiri A, Binder BM, Lechner E, Vierstra RD, Genschik P (2006) The
1006 exoribonuclease XRN4 is a component of the ethylene response pathway in
1007 *Arabidopsis*. *The Plant cell* 18: 3047-3057

1008 Quintero FJ, Garcíadeblás B, Rodríguez-Navarro A (1996) The SAL1 gene of *Arabidopsis*,
1009 encoding an enzyme with 3'(2'),5'-bisphosphate nucleotidase and inositol
1010 polyphosphate 1-phosphatase activities, increases salt tolerance in yeast. *Plant Cell* 8:
1011 529-537

1012 Robles P, Fleury D, Candela H, Cnops G, Alonso-Peral MM, Anami S, Falcone A, Caldana
1013 C, Willmitzer L, Ponce MR, Van Lijsebettens M, Micol JL (2010) The
1014 RON1/FRY1/SAL1 gene is required for leaf morphogenesis and venation patterning
1015 in *Arabidopsis*. *Plant Physiol* 152: 1357-1372

- Rodríguez VM, Chételat A, Majcherczyk P, Farmer EE (2010) Chloroplastic phosphoadenosine phosphosulfate metabolism regulates basal levels of the prohormone jasmonic acid in Arabidopsis leaves. *Plant Physiol* 152: 1335-1345
- Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene signal transduction in Arabidopsis thaliana: five novel mutant loci integrated into a stress response pathway. *Genetics* 139: 1393-1409
- Rossel JB, Walter PB, Hendrickson L, Chow WS, Poole A, Mullineaux PM, Pogson BJ (2006) A mutation affecting ASCORBATE PEROXIDASE 2 gene expression reveals a link between responses to high light and drought tolerance. *Plant, Cell Env* 29: 269-281
- Salomé PA, Oliva M, Weigel D, Krämer U (2013) Circadian clock adjustment to plant iron status depends on chloroplast and phytochrome function. *EMBO J* 32: 511-523
- Sanchez A, Shin J, Davis SJ (2011) Abiotic stress and the plant circadian clock. *Plant Sig Beh* 6: 223-231
- Sanchez SE, Petrillo E, Kornblihtt AR, Yanovsky MJ (2011) Alternative splicing at the right time. *RNA Biology* 8: 954-959
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671-675
- Scholl RL, May ST, Ware DH (2000) Seed and molecular resources for Arabidopsis. *Plant Physiol* 124: 1477-1480
- Shin J, Heidrich K, Sanchez-Villarreal A, Parker JE, Davis SJ (2012) TIME FOR COFFEE represses accumulation of the MYC2 transcription factor to provide time-of-day regulation of jasmonate signaling in Arabidopsis. *Plant Cell* 24: 2470-2482
- Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T (2015) Photoperiodic flowering: time measurement mechanisms in leaves. *Ann Rev Plant Biol* 66: 441-464
- Steduto P, Faurès JM, Hoogeveen J (2012) Coping with water scarcity: An action framework for agriculture and food security. Food and Agriculture Organization.
- Syed NH, Prince SJ, Mutava RN, Patil G, Li S, Chen W, Babu V, Joshi T, Khan S, Nguyen HT (2015) Core clock, SUB1, and ABAR genes mediate flooding and drought responses via alternative splicing in soybean. *J Exp Bot* 66: 7129-7149
- Takahashi H, Kopriva S, Giordano M, Saito K, Hell R (2011) Sulfur assimilation in photosynthetic organisms: molecular functions and regulations of transporters and assimilatory enzymes. *Ann Rev Plant Biol* 62: 157-184
- Terry N (1976) Effects of sulfur on the photosynthesis of intact leaves and isolated chloroplasts of sugar beets. *Plant Physiol* 57: 477-479
- Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu J-K (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 45: 523-539
- Walley JW, Coughlan S, Hudson ME, Covington MF, Kaspi R, Banu G, Harmer SL, Dehesh K (2007) Mechanical stress induces biotic and abiotic stress responses via a novel cis-element. *PLoS Genet* 3: 1800-1812
- Wang X, Wu F, Xie Q, Wang H, Wang Y, Yue Y, Gahura O, Ma S, Liu L, Cao Y, Jiao Y, Puta F, McClung CR, Xu X, Ma L (2012) SKIP is a component of the spliceosome linking alternative splicing and the circadian clock in Arabidopsis. *Plant Cell* 24: 3278-3295
- Wilkins O, Bräutigam K, Campbell MM (2010) Time of day shapes Arabidopsis drought transcriptomes. *Plant J* 63: 715-727
- Wilson PB, Estavillo GM, Field KJ, Pornsiriwong W, Carroll AJ, Howell KA, Woo NS, Lake JA, Smith SM, Harvey Millar A, von Caemmerer S, Pogson BJ (2009) The

1065 nucleotidase/phosphatase SAL1 is a negative regulator of drought tolerance in
 1066 Arabidopsis. *Plant J* 58: 299-317
 1067 Xiong L, Bh L, Ishitani M, Lee H, Zhang C, Zhu J-K (2001) FIERY1 encoding an inositol
 1068 polyphosphate 1-phosphatase is a negative regulator of abscisic acid and stress
 1069 signaling in Arabidopsis. *Genes & Development* 15: 1971-1984
 1070 Yu X, Liu H, Klejnot J, Lin C (2010) The Cryptochrome Blue Light Receptors. *The*
 1071 *Arabidopsis book* / American Society of Plant Biologists 8: e0135
 1072 Zhang L, Wan Y, Huang G, Wang D, Yu X, Huang G, Guo J (2015) The exosome controls
 1073 alternative splicing by mediating the gene expression and assembly of the
 1074 spliceosome complex. *Scientific Reports* 5: 13403
 1075
 1076
 1077





Figure 1. Osmotic stress induces the accumulation of PAP and extends circadian period.

(A) Accumulation of PAP in Columbia (Col-0) seedlings in the presence of 200 mM mannitol under constant light conditions. Plants were grown for 11 days under 12:12 L/D cycles before being transferred to plates containing 200 mM mannitol 24 hours before transfer to constant light. Plates were transferred to either 24 $\mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue light supplemented with 36 $\mu\text{mol m}^{-2} \text{s}^{-1}$ red light (cR+B) or to 60 $\mu\text{mol m}^{-2} \text{s}^{-1}$ constant white light (cW) at dawn of day 12. Seedlings were harvested at ZT96. Data are the mean of three biological replicates and are representative of two independent experiments. Error bars indicate standard deviation. Asterisks indicate a significant difference compared with respective mock control ($p < 0.025$, Bonferroni adjusted Student's *t*-test). **(B)** Representative bioluminescence data of luciferase activity in Columbia (Col-0) plants carrying a *CCA1::LUC2* reporter construct in the presence of 200 mM mannitol or PEG 8000. Plants were grown on half-strength MS medium for 5 days under 12:12 L/D cycles before being transferred to either a mock-treated control, 200 mM mannitol, or plates infused with PEG 8000, 24 hrs before imaging under cR+B light (30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ red and 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue light). Data are representative of three independent experiments. Error bars represent standard error of the mean and are shown every 10 hrs for clarity. $n=10$. **(C)** Circadian period estimates of luciferase activity in the presence of 200 mM mannitol or PEG 8000. Columbia (Col-0) seedlings carrying either the *CCA1*, *LHY*, or *TOC1* promoters fused to a *LUCIFERASE* reporter were assessed. Data are representative of three independent experiments. Error bars express standard error of the mean, $n=10$. Asterisks indicate $p < 0.01$ compared with mock controls (Dunnett's test).






Figure 3. The *sal1* circadian phenotype is exacerbated under blue light. (A) Fluence rate response curves to measure free-running circadian period under constant blue (cB) light in Columbia (Col-0) and *fry1-6* seedlings carrying a *CCA1::LUC2* reporter. Seedlings were entrained in 12:12 L/D cycles for 6 days before being transferred to the indicated fluence rate of cB light. Data are representative of three independent experiments, standard error of the mean is shown, $n = 10$. Asterisks highlight a significant difference between Col-0 and *fry1-6* at the indicated fluence rate (post-hoc Student's T-test, $p < 0.05$). (B) Fluence rate response curves to measure free-running circadian period under constant red (cR) light. Seedlings were entrained as described in (A) before being transferred to the indicated fluence rate of cR light. Data are representative of three independent experiments, standard error of the mean is shown, $n = 10$. (C-F) Accumulation of circadian clock-regulated transcripts under constant blue (cB) (C, E) or constant red (cR) (D, F) light in *sal1* seedlings using RT-qPCR. Levels of *CCA1* (C, D), and *TOC1* (E, F) mRNA were assessed. Plants were entrained to 12:12 h light:dark cycles for 12 days on MS medium before being moved to constant conditions with either 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue (cB) or 30 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (cR) red light. Data for each gene were normalized with an internal control (*PP2a*) and are the mean of at least two biological replicates. Error bars indicate standard error of the mean. (G, H, I) Accumulation of PAP in *fry1-6* seedlings under different fluence rates of blue (G), red (H), or red+blue (I) light. Seedlings were entrained to 12:12 h light:dark cycles for 12 days on MS medium before being transferred to the indicated fluence rate and quality of light for 4 days. Seedlings were harvested at ZT96. UD, PAP levels were below the detection threshold in Col-0 in each measurement. Asterisks highlight a significant difference for the selected comparison ($p < 0.025$, Bonferroni adjusted Student's t-test).




Figure 4. PAP accumulation is correlated with circadian defects in *sal1* mutants. (A) Circadian period of *fry1-6* seedlings following application of PAP. Seedlings were entrained in 12:12 h L/D cycles for 6 days before being transferred to 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue light. PAP was applied at ZT29. Asterisks indicate a significant difference compared with mock treated control ($p < 0.025$, Bonferroni adjusted Student's t-test). Standard error of the mean is shown, $n > 19$. Data are representative of three independent experiments. (B) Accumulation of PAP in Columbia (Col-0), *alx8-1*, and *alx8-1* seedlings transformed with a 35S::AHL construct. Plants were grown for 12 days under 12:12 light:dark cycles before being transferred to constant 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$ blue (cB) light. Seedlings were harvested at ZT96. Data are the mean of three biological replicates and are representative of two independent experiments. UD, PAP levels were below the detection threshold. (C, D) F_q'/F_m' rhythms and circadian period estimates in *alx8-1* seedlings transformed with a 35S::AHL construct. Plants were grown for 12 days under 12:12 h L/D cycles before being transferred to constant blue (cB) light (20 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Data are representative of three independent experiments, standard error of the mean is shown, $n = 8$. Asterisks indicate a significant difference compared with *alx8-1* plants ($P < 0.0125$, Bonferroni adjusted Student's t-test).




Figure 5. *SAL1* transcript and protein accumulation are not altered by the circadian system. (A) *SAL1* transcript accumulation in wild type (Col-0) plants under 16:8 h LD cycles. Data were normalized to an internal control (*PP2a*). Data are the average of three biological replicates, error bars show standard error of the mean. **(B)** Immunoblot analysis of *SAL1* protein levels under 16:8 h LD cycles. Plants were grown as described in (A). Data are the average of three biological replicates, error bars show standard error of the mean. **(C)** *SAL1* transcript accumulation over circadian time. Wild type (Col-0) plants were entrained under $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ white light in 12:12 h L/D cycles for 12 days before being transferred to constant white (cW) light. Data are the mean of three biological replicates. Error bars express standard error of the mean. See also Figure S4.






Figure 6. Sulfate deprivation does not extend circadian period. (A) Schematic of sulfate metabolism in Arabidopsis, adapted from (Bohrer et al., 2014). Abbreviations of metabolites: APS, adenosine-5'-phosphosulfate; GLS, glucosinolate; GSH, glutathione; PAP, 5'-phosphoadenosine 3'-phosphate; PAPS, 3-phosphoadenosine 5-phosphosulfate. Abbreviations of enzymes and transporters: APK, APS KINASE; APR, APS REDUCTASE; GSH1, GLUTAMATE-CYSTEINE LIGASE. (B) Circadian period estimates of luciferase activity in the presence or absence of sulfate salts. Columbia (*Col-0*) seedlings carrying either the *CCA1*, *LHY*, or *TOC1* promoters fused to a *LUCIFERASE* reporter were assessed. Plants were entrained for 12 d before transfer to constant blue (cB) light ($20 \mu\text{mol m}^{-2} \text{s}^{-1}$) for imaging. Standard error of the mean is shown, $n = 10$. Data from one of three independent experiments are shown. (C) Circadian rhythms of F_q'/F_m' in plants grown on agar lacking sulfate salts. Plants were grown as described in (B). Data represent mean values of multiple seedlings ($n=8$) and are representative of at least three independent experiments. (D, E) Circadian period estimates of F_q'/F_m' in *apk1 apk2* (D), and *cad2-1* (E) plants. Data are representative of at least two independent experiments. Standard error of the mean is shown, $n = 8$. See also Figure S6.




Figure 7. Circadian phenotypes of *xrn234* mutants. (A) Rhythms of PSII operating efficiency (F_q'/F_m') were measured over circadian time in Columbia (Col-0) and *xrn2-1 xrn3-3 xrn4-6* (*xrn234*) mutant seedlings. Period estimates are plotted against Relative Amplitude Error (RAE). Plants were grown for 12 days under 12:12 h L/D cycles before being transferred to constant blue (cB) light ($20 \mu\text{mol m}^{-2} \text{s}^{-1}$). Data represent mean values of multiple seedlings ($n=7$) and are representative of three independent experiments. Error bars indicate standard error, asterisks indicate a significant difference compared with Col-0 plants ($p<0.05$, Student's t test). (B) Assessment of *CCA1* transcript accumulation under constant blue light in *fry1-6* and *xrn234* seedlings using RT-qPCR. Plants were entrained in 12:12 h L/D cycles before being moved to constant conditions with $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light. Data for each gene were normalized with an internal control (*PP2A*) and are the mean of three biological replicates. (C) *CCA1* transcript accumulation under constant white light in *fry1-6* and *xrn234* seedlings. Transcript accumulation in Columbia (Col-0), *fry1-6* and *xrn234* seedlings was compared using RT-qPCR. Plants were entrained as in (B) before being moved to constant conditions with $60 \mu\text{mol m}^{-2} \text{s}^{-1}$ white (cW) light. Data for each gene were normalized with an internal control (*PP2A*) and are the mean of three biological replicates. Error bars indicate standard error.




Figure 8. PAP levels correlate with a lengthened circadian period under osmotic stress and broad spectrum blue light. (A,B) Accumulation of PAP in Columbia (Col-0) and *fry1-6* seedlings in the presence of 200 mM mannitol. Plants were grown for 11 days under 12:12 L/D cycles before being transferred to plates containing 200 mM mannitol 24 hours before transfer to constant light. Plates were subsequently transferred to either constant red and blue light ($30 \mu\text{mol m}^{-2} \text{s}^{-1}$ constant red light supplemented with $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light (cR+B, A) or to $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ constant blue light (cB, B) at dawn of day 12. Seedlings were harvested at ZT96. UD, PAP levels were below the detection threshold. Data are the mean of three biological replicates and are representative of two independent experiments. Standard deviation is shown. Asterisks indicate a significant difference compared with a mock treated control ($p < 0.025$, Bonferroni adjusted Student's t-test). (C, D) Circadian period estimates of luciferase activity in Columbia (Col-0) and *fry1-6* plants carrying a *CCA1::LUC2* reporter construct in the presence of 200mM mannitol. Plants were entrained and transferred to growth substrate containing 200mM mannitol as in (A). Seedlings were transferred to either constant red and blue light (cR+B, C) or constant blue light (cB, D) for imaging. Data are representative of three independent experiments. Error bars express standard error of the mean, $n=10$. Asterisks indicate a significant difference compared with a mock treated control ($P < 0.025$, Bonferroni adjusted Student's t test). (E) Accumulation of *CCA1* transcript following transfer to 200 mM mannitol. Plants were entrained to 12:12 h light:dark cycles for 11 days on MS medium before being transferred to either 200 mM mannitol or a mock treatment at dawn. Seedlings were moved to constant conditions with $20 \mu\text{mol m}^{-2} \text{s}^{-1}$ blue light at dawn of day 12. Data were normalized with an internal control (*PP2a*) and are the mean of at least two biological replicates. Error bars indicate standard error of the mean.

Parsed Citations

Adams S, Manfield I, Stockley P, Carré IA (2015) Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. PLoS ONE 10: e0143943

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Alabadi D, Oyama T, Yanovsky M, Harmon F, Mas P, Kay S (2001) Reciprocal regulation between TOC1 and LHY/CCA1 within the Arabidopsis circadian clock. Science 293: 880-883

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Apel K (2004) REACTIVE OXYGEN SPECIES: Metabolism, Oxidative Stress, and Signal Transduction. Annu Rev Plant Biol 55: 373-399

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bohrer A-S, Kopriva S, Takahashi H (2014) Plastid-cytosol partitioning and integration of metabolic pathways for APS/PAPS biosynthesis in Arabidopsis thaliana. Frontiers in Plant Science 5: 751

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bustin SA, Benes V, Garson JA, Hellems J, Huggett J, Kubista M, Mueller R, Nolan T, Pfaffl MW, Shipley GL, Vandesompele J, Wittwer CT (2009) The MIQE guidelines: Minimum information for publication of quantitative real-time PCR experiments. Clinical Chemistry 55: 611-622

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Bürstenbinder K, Rzewuski G, Wirtz M, Hell R, Sauter M (2007) The role of methionine recycling for ethylene synthesis in Arabidopsis. Plant J 49: 238-249

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chan KX, Mabbitt PD, Phua SY, Mueller JW, Nisar N, Gigolashvili T, Stroehrer E, Grassl J, Arlt W, Estavillo GM, Jackson CJ, Pogson BJ (2016) Sensing and signaling of oxidative stress in chloroplasts by inactivation of the SAL1 phosphoadenosine phosphatase. PNAS 113: E4567-4576

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ (2016) Learning the Languages of the Chloroplast: Retrograde Signaling and Beyond. Ann Rev Plant Biol 67: 25-53

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen H, Xiong L (2010) The bifunctional abiotic stress signalling regulator and endogenous RNA silencing suppressor FIERY1 is required for lateral root formation. Plant, Cell & Env 33: 2180-2190

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen H, Zhang B, Hicks LM, Xiong L (2011) A nucleotide metabolite controls stress-responsive gene expression and plant development. PLoS ONE 6: e26661

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chen Y-Y, Wang Y, Shin L-J, Wu J-F, Shanmugam V, Tsednee M, Lo J-C, Chen C-C, Wu S-H, Yeh K-C (2013) Iron is involved in the maintenance of circadian period length in Arabidopsis. Plant Physiol 161: 1409-1420

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Chow BY, Helfer A, Nusinow DA, Kay SA (2012) ELF3 recruitment to the PRR9 promoter requires other Evening Complex members in the Arabidopsis circadian clock. Plant Sig Beh 7: 170-173

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Christie JM, Blackwood L, Petersen J, Sullivan S (2015) Plant Flavoprotein Photoreceptors. Plant & Cell Phys 56: 401-413

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Cobbett CS, May MJ, Howden R, Rolls B (1998) The glutathione-deficient, cadmium-sensitive mutant, cad2-1, of Arabidopsis thaliana is deficient in gamma-glutamylcysteine synthetase. Plant J 16: 73-78

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Covington MF, Maloof JN, Straume M, Kay SA, Harmer SL (2008) Global transcriptome analysis reveals circadian regulation of key pathways in plant growth and development. Genome Biology 9: R130

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dichtl B, Stevens A, Tollervey D (1997) Lithium toxicity in yeast is due to the inhibition of RNA processing enzymes. EMBO J 16: 7184-7195

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Dong MA, Farré EM, Thomashow MF (2011) Circadian clock-associated 1 and late elongated hypocotyl regulate expression of the C-repeat binding factor (CBF) pathway in Arabidopsis. PNAS 108: 7241-7246

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Edgar RS, Green EW, Zhao Y, Van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja UK, Feeney KA, Maywood ES, Hastings MH, Baliga NS, Merrow M, Millar AJ, Johnson CH, Kyriacou CP, O'Neill JS, Reddy AB (2012) Peroxiredoxins are conserved markers of circadian rhythms. Nature 485: 459-464

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Eriksson ME, Webb AAR (2011) Plant cell responses to cold are all about timing. Current Opin Plant Biol 14: 731-737

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud E, Whelan J, David P, Javot H, Brearley C, Hell R, Marin E, Pogson BJ (2011) Evidence for a SAL1-PAP Chloroplast Retrograde Pathway That Functions in Drought and High Light Signaling in Arabidopsis. Plant Cell 23: 3992-4012

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fankhauser C, Staiger D (2002) Photoreceptors in Arabidopsis thaliana: light perception, signal transduction and entrainment of the endogenous clock. Planta 216: 1-16

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Farré EM, Kay SA (2007) PRR7 protein levels are regulated by light and the circadian clock in Arabidopsis. Plant J 52: 548-560

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Filichkin SA, Cumbie JS, Dharmawardhana P, Jaiswal P, Chang JH, Palusa SG, Reddy ASN, Megraw M, Mockler TC (2015) Environmental Stresses Modulate Abundance and Timing of Alternatively Spliced Circadian Transcripts in Arabidopsis. Molecular Plant 8: 207-227

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Fukushima A, Kusano M, Nakamichi N, Kobayashi M, Hayashi N, Sakakibara H, Mizuno T, Saito K (2009) Impact of clock-associated Arabidopsis pseudo-response regulators in metabolic coordination. PNAS 106: 7251-7256

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Garbarino-Pico E, Green CB (2007) Posttranscriptional regulation of mammalian circadian clock output. Cold Spring Harb Symp Quant Biol 72: 145-156

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gazzani S, Lawrenson T, Woodward C, Headon D, Sablowski R (2004) A link between mRNA turnover and RNA interference in Arabidopsis. Science 306: 1046-1048

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gendron JM, Pruneda-Paz JL, Doherty CJ, Gross AM, Kang SE, Kay SA (2012) Arabidopsis circadian clock protein, TOC1, is a DNA-binding transcription factor. PNAS 109: 3167-3172

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Goodspeed D, Chehab EW, Min-Venditti A, Braam J, Covington MF (2012) Arabidopsis synchronizes jasmonate-mediated defense with insect circadian behavior. PNAS 109: 4674-4677

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Grundy J, Stoker C, Carré IA (2015) Circadian regulation of abiotic stress tolerance in plants. Frontiers in Plant Science 6: 201

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Guo J, Cheng P, Yuan H, Liu Y (2009) The exosome regulates circadian gene expression in a posttranscriptional negative feedback loop. Cell 138: 1236-1246

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Gyl, Gasciolli V, Lauressergues D, Morel J-B, Gombert J, Proux F, Proux C, Vaucheret H, Mallory AC (2007) Arabidopsis FIERY1, XRN2, and XRN3 are endogenous RNA silencing suppressors. Plant Cell 19: 3451-3461

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Habte E, Müller LM, Shtaya M, Davis SJ, von Korff M (2014) Osmotic stress at the barley root affects expression of circadian clock genes in the shoot. Plant, Cell & Env 37: 1321-1327

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hanano S, Domagalska MA, Nagy F, Davis SJ (2006) Multiple phytohormones influence distinct parameters of the plant circadian clock. Genes Cells 11: 1381-1392

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Herrero E, Kolmos E, Bujdoso N, Yuan Y, Wang M, Berns MC, Uhlworm H, Coupland G, Saini R, Jaskolski M, Webb A, Gonçalves J, Davis SJ (2012) EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. Plant Cell 24: 428-443

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hirsch J, Misson J, Crisp PA, David P, Bayle V, Estavillo GM, Javot H, Chiarenza S, Mallory AC, Maizel A, Declerck M, Pogson BJ, Vaucheret H, Crespi M, Desnos T, Thibaud M-C, Nussaume L, Marin E (2011) A novel fry1 allele reveals the existence of a mutant phenotype unrelated to 5'→3' exoribonuclease (XRN) activities in Arabidopsis thaliana roots. PLoS ONE 6: e16724

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hong S, Kim SA, Guerinet ML, McClung CR (2013) Reciprocal Interaction of the Circadian Clock with the Iron Homeostasis Network in Arabidopsis. Plant Physiol 161: 893-903

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Hsu PY, Harmer SL (2014) Wheels within wheels: the plant circadian system. Trends in Plant Science 19: 240-249

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Huang W, Perez-Garcia P, Pokhilko A, Wilmarth A, Antoshchik I, Richman JL, Mas P (2012) Mapping the core of the Arabidopsis

circadian clock defines the network structure of the oscillator. Science 336: 75-79

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Hut RA, Beersma DGM (2011) Evolution of time-keeping mechanisms: early emergence and adaptation to photoperiod. Philos Trans R Soc Lond B Biol Sci 366: 2141-2154

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Ingle RA, Stoker C, Stone W, Adams N, Smith R, Grant M, Carré I, Roden LC, Denby KJ (2015) Jasmonate signalling drives time-of-day differences in susceptibility of Arabidopsis to the fungal pathogen Botrytis cinerea. Plant J 84: 937-948

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

James AB, Syed NH, Bordage S, Marshall J, Nimmo GA, Jenkins GI, Herzyk P, Brown JWS, Nimmo HG (2012) Alternative splicing mediates responses of the Arabidopsis circadian clock to temperature changes. Plant Cell 24: 961-981

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jones M (2009) Entrainment of the Arabidopsis circadian clock. J Plant Biol 52: 202-209

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jones M (2017) Interplay of Circadian Rhythms and Light in the Regulation of Photosynthesis-Derived Metabolism. In, Vol 79. https://link.springer.com/chapter/10.1007/124_2017_2, p In press

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jones M, Hu W, Litthauer S, Lagarias JC, Harmer SL (2015) A Constitutively Active Allele of Phytochrome B Maintains Circadian Robustness in the Absence of Light. Plant Physiology 169: 814-825

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Jones M, Williams BA, McNicol J, Simpson CG, Brown JWS, Harmer SL (2012) Mutation of Arabidopsis spliceosomal timekeeper locus1 causes circadian clock defects. Plant Cell 24: 4066-4082

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kastenmayer JP, Green PJ (2000) Novel features of the XRN-family in Arabidopsis: evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. PNAS 97: 13985-13990

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kerwin RE, Jiménez-Gómez JM, Fulop D, Harmer SL, Maloof JN, Kliebenstein DJ (2011) Network quantitative trait loci mapping of circadian clock outputs identifies metabolic pathway-to-clock linkages in Arabidopsis. Plant Cell 23: 471-485

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kim B-H, von Arnim AG (2009) FIERY1 regulates light-mediated repression of cell elongation and flowering time via its 3'(2'),5'-bisphosphate nucleotidase activity. Plant J 58: 208-219

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kojima S, Shingle DL, Green CB (2011) Post-transcriptional control of circadian rhythms. J Cell Sci 124: 311-320

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Kurihara Y, Schmitz RJ, Nery JR, Schultz MD, Okubo-Kurihara E, Morosawa T, Tanaka M, Toyoda T, Seki M, Ecker JR (2012) Surveillance of 3' Noncoding Transcripts Requires FIERY1 and XRN3 in Arabidopsis. G3 2: 487-498

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only Title Only Author and Title](#)

Lai AG, Doherty CJ, Mueller-Roeber B, Kales A, Schippers JHM, Dijkwel PP (2012) CIRCADIAN CLOCK-ASSOCIATED 1 regulates ROS

Downloaded from [http://www.jstor.org/](#) on April 19, 2018
Copyright © 2018 American Society of Plant Biologists. All rights reserved.

homeostasis and oxidative stress responses. PNAS 109: 17129-17134

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lee B-R, Huseby S, Koprivova A, Chételat A, Wirtz M, Mugford ST, Navid E, Brearley C, Saha S, Mithen R, Hell R, Farmer EE, Kopriva S (2012) Effects of *fou8/fry1* Mutation on Sulfur Metabolism: Is Decreased Internal Sulfate the Trigger of Sulfate Starvation Response? PLoS ONE 7: e39425

Lee HG, Más P, Seo PJ (2016) MYB96 shapes the circadian gating of ABA signaling in Arabidopsis. Scientific Reports 6: 17754

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Legnaioli T, Cuevas J, Más P (2009) TOC1 functions as a molecular switch connecting the circadian clock with plant responses to drought. EMBO J 28: 3745-3757

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Li J, Li G, Wang H, Wang Deng X (2011) Phytochrome signaling mechanisms. The Arabidopsis book / American Society of Plant Biologists 9: e0148

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Litthauer S, Battle MW, Lawson T, Jones MA (2015) Phototropins maintain robust circadian oscillation of PSII operating efficiency under blue light. Plant J 83: 1034-1045

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Liu T, Carlsson J, Takeuchi T, Newton L, Farré EM (2013) Direct regulation of abiotic responses by the Arabidopsis circadian clock component PRR7. Plant J 76: 101-114

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Lu SX, Webb CJ, Knowles SM, Kim SHJ, Wang Z, Tobin EM (2012) CCA1 and ELF3 Interact in the control of hypocotyl length and flowering time in Arabidopsis. Plant Physiol 158: 1079-1088

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Macgregor DR, Gould P, Foreman J, Griffiths J, Bird S, Page R, Stewart K, Steel G, Young J, Paszkiewicz K, Millar AJ, Halliday KJ, Hall AJ, Penfield S (2013) HIGH EXPRESSION OF OSMOTICALLY RESPONSIVE GENES1 Is Required for Circadian Periodicity through the Promotion of Nucleo-Cytoplasmic mRNA Export in Arabidopsis. Plant Cell 25: 4391-4404

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Matsuo T, Okamoto K, Onai K, Niwa Y, Shimogawara K, Ishiura M (2008) A systematic forward genetic analysis identified components of the Chlamydomonas circadian system. Genes & Development 22: 918-930

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

McWatters H, Kolmos E, Hall A, Doyle M, Amasino R, Gyula P, Nagy F, Millar A, Davis S (2007) ELF4 is required for oscillatory properties of the circadian clock. Plant Physiol 144: 391-401

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mechold U, Ogryzko V, Ngo S, Danchin A (2006) Oligoribonuclease is a common downstream target of lithium-induced pAp accumulation in Escherichia coli and human cells. Nucleic Acids Research 34: 2364-2373

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Millar AJ (2016) The Intracellular Dynamics of Circadian Clocks Reach for the Light of Ecology and Evolution. Ann Rev Plant Biol 67: 595-618

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mittler R, Vanderauwera S, Suzuki N, MILLER G, Tognetti VB, Vandepoele K, Gollery M, Shulaev V, Van Breusegem F (2011) ROS signaling: the new wave? Trends in plant science 16: 300-309

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mizuno T, Yamashino T (2008) Comparative transcriptome of diurnally oscillating genes and hormone-responsive genes in *Arabidopsis thaliana*: insight into circadian clock-controlled daily responses to common ambient stresses in plants. Plant Cell Physiol 49: 481-487

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mockler T, Michael T, Priest H, Shen R, Sullivan C, Givan S, McEntee C, Kay S, Chory J (2007) The DIURNAL project: DIURNAL and circadian expression profiling, model-based pattern matching, and promoter analysis. Cold Spring Harb Symp Quant Biol 72: 353-363

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Mugford SG, Yoshimoto N, Reichelt M, Wirtz M, Hill L, Mugford ST, Nakazato Y, Noji M, Takahashi H, Kramell R, Gigolashvili T, Flügge U-I, Wasternack C, Gershenzon J, Hell R, Saito K, Kopriva S (2009) Disruption of adenosine-5'-phosphosulfate kinase in *Arabidopsis* reduces levels of sulfated secondary metabolites. Plant Cell 21: 910-927

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nagarajan VK, Jones CI, Newbury SF, Green PJ (2013) XRN 5'→3' exoribonucleases: structure, mechanisms and functions. Biochimica et Biophysica Acta 1829: 590-603

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakagawa T, Kurose T, Hino T, Tanaka K, Kawamukai M, Niwa Y, Toyooka K, Matsuoka K, Jinbo T, Kimura T (2007) Development of series of gateway binary vectors, pGWBs, for realizing efficient construction of fusion genes for plant transformation. J Biosci Bioeng 104: 34-41

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakamichi N, Kiba T, Henriques R, Mizuno T, Chua N-H, Sakakibara H (2010) PSEUDO-RESPONSE REGULATORS 9, 7, and 5 are transcriptional repressors in the *Arabidopsis* circadian clock. Plant Cell 22: 594-605

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nakamichi N, Takao S, Kudo T, Kiba T, Wang Y, Kinoshita T, Sakakibara H (2016) Improvement of *Arabidopsis* biomass and cold-, drought-, and salinity-stress tolerance by modified circadian clock-associated PSEUDO-RESPONSE REGULATORS. Plant & Cell Physiology 57: 1085-1097

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Narusaka M, Shiraishi T, Iwabuchi M, Narusaka Y (2010) The floral inoculating protocol: a simplified *Arabidopsis thaliana* transformation method modified from floral dipping. Plant Biotechnology 27: 349-351

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Norén, L., Kindgren, P., Stachula, P., Rühl, M., Eriksson, M. E., Hurry, V., & Strand, Å. (2016). Circadian and Plastid Signaling Pathways Are Integrated to Ensure Correct Expression of the CBF and COR Genes during Photoperiodic Growth. Plant Physiology, 171: 1392–1406

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Nusinow DA, Helfer A, Hamilton EE, King JJ, Imaizumi T, Schultz TF, Farré EM, Kay SA (2011) The ELF4-ELF3-LUX complex links the circadian clock to diurnal control of hypocotyl growth. Nature 475: 398-402

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Olmedo G, Guo H, Gregory BD, Nourizadeh SD, Aguilar-Henonin L, Li H, An F, Guzman P, Ecker JR (2006) ETHYLENE-INSENSITIVE5 encodes a 5'→3' exoribonuclease required for regulation of the EIN3-targeting F-box proteins EBF1/2. PNAS 103: 13286-13293

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Perez-Santángelo S, Mancini E, Francey LJ, Schlaen RG, Chernomoretz A, Hogenesch JB, Yanovsky MJ (2014) Role for LSM genes in the regulation of circadian rhythms. PNAS 111: 15166-15171

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Plautz JD, Straume M, Stanewsky R, Jamison CF, Brandes C, Dowse HB, Hall JC, Kay SA (1997) Quantitative analysis of Drosophila period gene transcription in living animals. J Biol Rhythms 12: 204-217

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Pornsiriwong W, Estavillo GM, Chan KX, Tee EE, Ganguly D, Crisp PA, Phua SY, Zhao C, Qiu J, Park J, Yong MT, Nisar N, Yadav AK, Schwessinger B, Rathjen J, Cazzonelli CI, Wilson PB, Gilliam M, Chen Z-H, Pogson BJ (2017) A chloroplast retrograde signal, 3'-phosphoadenosine 5'-phosphate, acts as a secondary messenger in abscisic acid signaling in stomatal closure and germination. eLife 6: e23361

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Potuschak T, Vansiri A, Binder BM, Lechner E, Vierstra RD, Genschik P (2006) The exoribonuclease XRN4 is a component of the ethylene response pathway in Arabidopsis. The Plant cell 18: 3047-3057

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Quintero FJ, Garcíadeblás B, Rodríguez-Navarro A (1996) The SAL1 gene of Arabidopsis, encoding an enzyme with 3'(2'),5'-bisphosphate nucleotidase and inositol polyphosphate 1-phosphatase activities, increases salt tolerance in yeast. Plant Cell 8: 529-537

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Robles P, Fleury D, Candela H, Cnops G, Alonso-Peral MM, Anami S, Falcone A, Caldana C, Willmitzer L, Ponce MR, Van Lijsebettens M, Micol JL (2010) The RON1/FRY1/SAL1 gene is required for leaf morphogenesis and venation patterning in Arabidopsis. Plant Physiol 152: 1357-1372

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rodríguez VM, Chételat A, Majcherczyk P, Farmer EE (2010) Chloroplastic phosphoadenosine phosphosulfate metabolism regulates basal levels of the prohormone jasmonic acid in Arabidopsis leaves. Plant Physiol 152: 1335-1345

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Roman G, Lubarsky B, Kieber JJ, Rothenberg M, Ecker JR (1995) Genetic analysis of ethylene signal transduction in Arabidopsis thaliana: five novel mutant loci integrated into a stress response pathway. Genetics 139: 1393-1409

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Rossel JB, Walter PB, Hendrickson L, Chow WS, Poole A, Mullineaux PM, Pogson BJ (2006) A mutation affecting ASCORBATE PEROXIDASE 2 gene expression reveals a link between responses to high light and drought tolerance. Plant, Cell Env 29: 269-281

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Salomé PA, Oliva M, Weigel D, Krämer U (2013) Circadian clock adjustment to plant iron status depends on chloroplast and phytochrome function. EMBO J 32: 511-523

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sanchez A, Shin J, Davis SJ (2011) Abiotic stress and the plant circadian clock. Plant Sig Beh 6: 223-231

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Sanchez SE, Petrillo E, Kornblihtt AR, Yanovsky MJ (2011) Alternative splicing at the right time. RNA Biology 8: 954-959

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)

Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Schneider CA, Rasband WS, Elieciari KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9: 671-675

Pubmed: [Author and Title](#)

CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Scholl RL, May ST, Ware DH (2000) Seed and molecular resources for Arabidopsis. Plant Physiol 124: 1477-1480

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Shin J, Heidrich K, Sanchez-Villarreal A, Parker JE, Davis SJ (2012) TIME FOR COFFEE represses accumulation of the MYC2 transcription factor to provide time-of-day regulation of jasmonate signaling in Arabidopsis. Plant Cell 24: 2470-2482

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T (2015) Photoperiodic flowering: time measurement mechanisms in leaves. Ann Rev Plant Biol 66: 441-464

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Steduto P, Faurès JM, Hoogeveen J (2012) Coping with water scarcity: An action framework for agriculture and food security. Food and Agriculture Organization.

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Syed NH, Prince SJ, Mutava RN, Patil G, Li S, Chen W, Babu V, Joshi T, Khan S, Nguyen HT (2015) Core clock, SUB1, and ABAR genes mediate flooding and drought responses via alternative splicing in soybean. J Exp Bot 66: 7129-7149

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Takahashi H, Kopriva S, Giordano M, Saito K, Hell R (2011) Sulfur assimilation in photosynthetic organisms: molecular functions and regulations of transporters and assimilatory enzymes. Ann Rev Plant Biol 62: 157-184

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Terry N (1976) Effects of sulfur on the photosynthesis of intact leaves and isolated chloroplasts of sugar beets. Plant Physiol 57: 477-479

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu J-K (2006) Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. Plant J 45: 523-539

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Walley JW, Coughlan S, Hudson ME, Covington MF, Kaspi R, Banu G, Harmer SL, Dehesh K (2007) Mechanical stress induces biotic and abiotic stress responses via a novel cis-element. PLoS Genet 3: 1800-1812

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wang X, Wu F, Xie Q, Wang H, Wang Y, Yue Y, Gahura O, Ma S, Liu L, Cao Y, Jiao Y, Puta F, McClung CR, Xu X, Ma L (2012) SKIP is a component of the spliceosome linking alternative splicing and the circadian clock in Arabidopsis. Plant Cell 24: 3278-3295

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wilkins O, Bräutigam K, Campbell MM (2010) Time of day shapes Arabidopsis drought transcriptomes. Plant J 63: 715-727

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Wilson PB, Estavillo GM, Field KJ, Pornsiriwong W, Carroll AJ, Howell KA, Woo NS, Lake JA, Smith SM, Harvey Millar A, von Caemmerer S, Pogson BJ (2009) The nucleotidase/phosphatase SAL1 is a negative regulator of drought tolerance in Arabidopsis. Plant J 58: 299-317

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Xiong L, Bh L, Ishitani M, Lee H, Zhang C, Zhu J-K (2001) FIERY1 encoding an inositol polyphosphate 1-phosphatase is a negative regulator of abscisic acid and stress signaling in Arabidopsis. Genes & Development 15: 1971-1984

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Yu X, Liu H, Klejnot J, Lin C (2010) The Cryptochrome Blue Light Receptors. The Arabidopsis book / American Society of Plant Biologists 8: e0135

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)

Zhang L, Wan Y, Huang G, Wang D, Yu X, Huang G, Guo J (2015) The exosome controls alternative splicing by mediating the gene expression and assembly of the spliceosome complex. Scientific Reports 5: 13403

Pubmed: [Author and Title](#)
CrossRef: [Author and Title](#)
Google Scholar: [Author Only](#) [Title Only](#) [Author and Title](#)