

AID-expressing epithelium is protected from oncogenic transformation by an NKG2D surveillance pathway

Arantxa Pérez-García¹, Pablo Pérez-Durán^{1,†}, Thomas Wossning^{1,††}, Isora V Hernandez^{1,†††}, Sonia M Mur¹, Marta Cañamero², Francisco X Real³ & Almudena R Ramiro^{1,*}

Abstract

Activation-induced deaminase (AID) initiates secondary antibody diversification in germinal center B cells, giving rise to higher affinity antibodies through somatic hypermutation (SHM) or to isotype-switched antibodies through class switch recombination (CSR). SHM and CSR are triggered by AID-mediated deamination of cytosines in immunoglobulin genes. Importantly, AID activity in B cells is not restricted to Ig loci and can promote mutations and pro-lymphomagenic translocations, establishing a direct oncogenic mechanism for germinal center-derived neoplasias. AID is also expressed in response to inflammatory cues in epithelial cells, raising the possibility that AID mutagenic activity might drive carcinoma development. We directly tested this hypothesis by generating conditional knock-in mouse models for AID overexpression in colon and pancreas epithelium. AID overexpression alone was not sufficient to promote epithelial cell neoplasia in these tissues, in spite of displaying mutagenic and genotoxic activity. Instead, we found that heterologous AID expression in pancreas promotes the expression of NKG2D ligands, the recruitment of CD8⁺ T cells, and the induction of epithelial cell death. Our results indicate that AID oncogenic potential in epithelial cells can be neutralized by immunosurveillance protective mechanisms.

Keywords activation-induced deaminase; cancer; epithelium; NKG2D; pancreas

Subject Categories Cancer; Immunology

DOI 10.15252/emmm.201505348 | Received 16 April 2015 | Revised 16 July 2015 | Accepted 21 July 2015 | Published online 17 August 2015

EMBO Mol Med (2015) 7: 1327–1336

Introduction

Activation-induced deaminase (AID) is the enzyme that initiates the reactions of secondary antibody diversification: somatic hypermutation (SHM) and class switch recombination (CSR) (Muramatsu *et al*, 2000). These reactions enable the generation of antibodies with increased affinity for antigen (SHM) and with diversified, specialized functions for antigen removal (CSR), and are therefore critical for a competent immune response (Di Noia & Neuberger, 2007; Stavnezer *et al*, 2008; Alt *et al*, 2013; Robbiani & Nussenzweig, 2013). Accordingly, defective AID activity promotes a Hyper-IgM immunodeficiency syndrome in humans (Revy *et al*, 2000). AID triggers SHM and CSR by direct deamination of cytosine nucleosides in the DNA of immunoglobulin genes, resulting in the generation of U:G mismatches (Petersen-Mahrt *et al*, 2002; Alt *et al*, 2013; Robbiani & Nussenzweig, 2013). These U:G mismatches are in turn processed by alternative repair pathways that ultimately lead in SHM to the fixation of a mutation, and in CSR to a DNA double-strand break (DSB) and a recombination reaction (Di Noia & Neuberger, 2007; Stavnezer *et al*, 2008; Alt *et al*, 2013; Robbiani & Nussenzweig, 2013).

Activation-induced deaminase activity is not confined to immunoglobulin genes and can promote mutations and DSB followed by illegitimate chromosomal translocations in other regions of the genome (Ramiro *et al*, 2004, 2006; Liu *et al*, 2008; Robbiani *et al*, 2008, 2009). Importantly, chromosomal translocations are the hallmark of mature B-cell lymphomas, the most frequent of all human lymphomas. Indeed, AID deficiency delays the onset of lymphomagenesis in the mouse (Ramiro *et al*, 2004; Kovalchuk *et al*, 2007; Pasqualucci *et al*, 2008), establishing a direct link between AID collateral genotoxic activity and neoplastic transformation in B lymphocytes.

In the last few years, it has become clear that AID expression is not, as originally thought, exclusively restricted to activated B cells. AID expression has been reported in several tissues, including

1 B Cell Biology Lab, Centro Nacional de Investigaciones Cardiovasculares Carlos III (CNIC), Madrid, Spain

2 Roche Diagnostics GmbH, Penzberg, Germany

3 Epithelial Carcinogenesis Group, Spanish National Cancer Research Centre (CNIO), Madrid, Spain

*Corresponding author. Tel: +34 91 4531200; Fax: +34 91 4531245; E-mail: aramiro@cnic.es

†Present address: Department of Pathology, NYU Cancer Institute, New York University School of Medicine, New York, NY, USA

††Present address: Klinik für Tumorbologie, Freiburg, Germany

†††Present address: Otsuka Pharmaceutical, Barcelona, Spain

gastric, hepatic, and gut epithelia (Endo *et al*, 2007, 2008; Matsumoto *et al*, 2007; reviewed in Marusawa *et al* (2011)). AID expression in these tissues is most frequently associated with inflammatory conditions and the activation of the NF- κ B pathway (Endo *et al*, 2007; Matsumoto *et al*, 2007) and has been claimed to promote the accumulation of mutations in epithelial cells (Matsumoto *et al*, 2007, 2010; Takai *et al*, 2009; reviewed in Marusawa *et al* (2011)). Given that chronic inflammation in epithelial tissues predisposes to cancer development (Mantovani *et al*, 2008), the finding that the mutagenic activity of AID can be induced in an inflammatory context has fostered the idea that AID might contribute to or even constitute the link between inflammation and cancer (Takai *et al*, 2012; reviewed in Marusawa *et al* (2011)).

Several gain-of-function mouse models have been generated to address the contribution of AID to neoplastic transformation. Ubiquitous AID overexpression led mostly to early T cell neoplasia (Okazaki *et al*, 2003), hampering a thorough analysis of other malignancies. In contrast, B-cell-specific AID overexpression did not result in lymphomagenesis (Muto *et al*, 2006; Robbiani *et al*, 2009) unless the tumor suppressor p53 was removed (Robbiani *et al*, 2009). However, to date, the impact of specific AID expression in epithelial tissues, classically subject to inflammation-induced neoplastic transformation, has not been addressed. Here, we aimed to test this possibility directly by generating conditional knock-in models of AID overexpression. AID expressed in colon and pancreas epithelia was not sufficient to promote carcinogenesis, in spite of being expressed at high levels and displaying genotoxic activity. Instead, AID triggered the expression of NKGD2 ligands and the recruitment of immune cells and promoted a cytotoxic response and cell death. Our data indicate that the oncogenic potential of AID in epithelial cells is neutralized by an immunosurveillance pathway that prevents the expansion of pretumoral cells.

Results

Inflammation-induced AID does not contribute to carcinogenesis

Inflammation is known to play a critical role in the etiology of colorectal and pancreatic ductal adenocarcinoma (reviewed in Feagins *et al* (2009); Vonderheide & Bayne (2013)). To investigate whether inflammatory conditions promote AID expression in these tissues, we treated human epithelial cell lines derived from colorectal adenocarcinoma (LoVo and SW480) and pancreatic adenocarcinoma (AsPC and PaTU) with the pro-inflammatory cytokine TNF- α and measured AID expression by qRT-PCR. TNF- α stimulation increased AID mRNA expression in all cell lines analyzed (Fig 1A and B). To assess whether primary, non-transformed cells were also able to express AID in response to inflammatory stimuli, we generated explants from mouse pancreatic acinar cells and treated them with TNF- α . As with the human tumor cells, mouse primary epithelial cells expressed AID upon exposure to TNF- α (Fig 1C). TNF- α treatment typically induced 4–30-fold increases in AID mRNA levels in the different cell types tested, consistent with previous findings in liver, gastric and colorectal cell lines (Endo *et al*, 2007, 2008; Matsumoto *et al*, 2007). Together, these data confirm previous results showing that inflammatory stimuli can trigger AID expression in cell lines originated from human colorectal adenocarcinoma (Endo




Figure 1. Inflammation-induced AID expression does not contribute to carcinogenesis.

A–C AID expression was analyzed in colon and pancreatic human cell lines and in pancreas explants from C57BL/6 mice. Samples were treated as indicated with 50 ng/ml TNF- α . (A) qRT-PCR analysis of AID expression in LoVo and SW480 colon cell lines. $n = 5$ (LoVo); 3 (SW480). ** P -value: LoVo: 0.0017; SW480: 0.0079. (B) qRT-PCR analysis of AID expression in AsPC and PaTU-8988S pancreatic cell lines ($n = 2$). * P -value: AsPC: 0.0369; PaTU-8988S: 0.0119. (C) qRT-PCR analysis of AID expression in pancreatic explants from wild-type mice ($n = 3$). * $P = 0.0242$. D AID^{-/-} or AID^{+/-} mice were treated with 3% DSS for 10 cycles, and colonic sections were analyzed by histologic inspection after H/E staining. Graphs represent mean frequency values of adenoma and adenocarcinoma lesions of five independent experiments. $n = 28$ (AID^{-/-} males); 35 (AID^{+/-} males); 23 (AID^{-/-} females); 25 (AID^{+/-} females). P -value: male: 0.8; female: 0.246.

Data information: All data are mean values \pm SEM. Statistical differences were analyzed by two-tailed unpaired Student's t -test.

et al, 2008), and show that pancreatic adenocarcinoma cells and primary pancreatic cells are also responsive to TNF- α treatment.

Inflammation-induced AID expression has been proposed to contribute to or even be the leading cause of some epithelium-derived tumors, such as colorectal adenocarcinoma (Marusawa *et al*, 2011). To address whether endogenous AID expressed in epithelium under inflammatory conditions could contribute to carcinogenesis, we made use of the well-established model of dextran sulfate sodium (DSS)-induced colitis-associated cancer (CAC) (Cooper *et al*, 2000). AID^{-/-} mice or AID^{+/-} littermates were treated for 10 cycles with 3% DSS and evaluated by pathological criteria. We found that the frequency of oncogenic lesions was not significantly different in AID^{-/-} versus AID^{+/-} mice (Fig 1D). We conclude that endogenous AID does not significantly contribute to colorectal adenocarcinoma in the DSS-induced CAC model.

Conditional AID expression in epithelial cells does not promote adenocarcinoma development

The absence of a significant contribution of endogenous AID to carcinogenesis in DSS-treated mice could be explained by an insufficient amount of AID in this model. Indeed, AID expression is known to be limiting for its activity in B cells (Sernandez *et al*, 2008), and AID levels in B cells are typically 100–1,000 fold higher than those detected in epithelial cells under inflammatory conditions (unpublished observations). To directly evaluate whether AID expression

can contribute to carcinogenesis, we generated two mouse models for conditional AID expression in epithelial cells of colonic and pancreatic origin (Fig 2A). We introduced an AID-GFP-encoding




Figure 2. Heterologous AID expression does not promote carcinoma development.

- A** Schematic of the constructs used for conditional expression of AID in epithelial cells. An AID-IRES-GFP cassette preceded by a transcriptional STOP flanked by LoxP sites was introduced by homologous recombination within the endogenous Rosa26 locus (R26AID^{+/KI} mice, top). R26AID^{+/KI} mice were bred with Villin-CRE and p48-CRE mice to achieve specific AID expression in colon and pancreas, respectively (bottom).
- B** GFP immunofluorescence in colonic and pancreatic tissue from R26AID^{+/KI} Villin-CRE^{+/TG} and R26AID^{+/KI} p48-CRE^{+/KI} mice. Scale bar: 50 μm.
- C** qRT-PCR analysis of AID expression in colonic and pancreatic tissue from R26AID^{+/KI} Villin-CRE^{+/TG} and R26AID^{+/KI} p48-CRE^{+/KI} mice. $n = 5$ (R26AID^{+/+} Villin-CRE^{+/TG}); 4 (R26AID^{+/KI} Villin-CRE^{+/TG}); 2 (R26AID^{+/+} p48-CRE^{+/KI}); 2 (R26AID^{+/KI} p48-CRE^{+/KI}). LPS+IL4-stimulated B cells are shown as a positive control ($n = 2$). Bars show mean values \pm SEM normalized to LPS+IL4-treated B cells.
- D** Kaplan-Meier survival curves for R26AID^{+/KI} Villin-CRE^{+/TG} (left) ($n = 47$ (R26AID^{+/+} Villin-CRE^{+/TG}); 38 (R26AID^{+/KI} Villin-CRE^{+/TG})) and R26AID^{+/KI} p48-CRE^{+/KI} mice (right) ($n = 39$ (R26AID^{+/+} p48-CRE^{+/KI}); 23 (R26AID^{+/KI} p48-CRE^{+/KI})).

cassette in the endogenous Rosa26 locus preceded by a transcriptional stop flanked by two loxP sites (R26AID^{+/KI} mice). To achieve specific expression of AID in epithelial cells, we bred R26AID^{+/KI} mice with mice expressing the Cre-recombinase under a villin promoter, which specifically drives expression in colon (el Marjou *et al*, 2004) (R26AID^{+/KI} Villin-CRE^{+/TG} mice), or the pancreas-specific Ptf1 (p48) gene (Kawaguchi *et al*, 2002) (R26AID^{+/KI} p48-CRE^{+/KI} mice). R26AID^{+/+} Villin-CRE^{+/TG} and R26AID^{+/+} p48-CRE^{+/KI} mice were used as controls. To confirm that the Rosa26 AID-GFP cassette was functional, we first evaluated the expression of the reporter protein GFP by immunofluorescence in colon of R26AID^{+/KI} Villin-CRE^{+/TG} mice and pancreas of R26AID^{+/KI} p48-CRE^{+/KI} mice (Fig 2B). GFP was expressed in R26AID^{+/KI} Villin-CRE^{+/TG} colon and R26AID^{+/KI} p48-CRE^{+/KI} pancreas but not in control mice (Fig 2B) or in other tissues (not shown). We next measured AID transcript levels by qRT-PCR. In R26AID^{+/KI} Villin-CRE^{+/TG} and R26AID^{+/KI} p48-CRE^{+/KI} mice, the amount of AID in the targeted epithelial tissues was similar to that found in B cells activated *in vitro* with LPS + IL4, whereas AID expression in control mice remained at background level (Fig 2C). AID is thus expressed in the epithelium of R26AID^{+/KI} Villin-CRE^{+/TG} colon and R26AID^{+/KI} p48-CRE^{+/KI} pancreas at levels known to be functional in B cells.

To assess the contribution of AID to adenocarcinoma development, we monitored tumor incidence in R26AID^{+/KI} Villin-CRE^{+/TG} and R26AID^{+/KI} p48-CRE^{+/KI} mice. The onset of pancreatic and colorectal adenocarcinoma in a variety of mouse models ranges from 5–6 months to 1–1.5 years (Fodde & Smits, 2001; Aguilar *et al*, 2004; Martinelli *et al*, 2015). Therefore, to avoid confounding results arising from spontaneous tumorigenesis in very old mice, we set analysis end points at 75–100 weeks. Survival of R26AID^{+/KI} Villin-CRE^{+/TG} mice was indistinguishable from that of R26AID^{+/+} Villin-CRE^{+/TG} littermate controls (Fig 2D, left). Likewise, survival of R26AID^{+/KI} p48-CRE^{+/KI} did not differ from that of R26AID^{+/+} p48-CRE^{+/KI} controls (Fig 2D, right). To rule out the presence of early malignancies in aged animals, we performed thorough pathological analysis of colon and pancreas sections of all animals, but could not detect any tumor development in R26AID^{+/KI} Villin-CRE^{+/TG} and R26AID^{+/KI} p48-CRE^{+/KI} animals at 75–100 weeks (Fig EV1). Expression of AID in colon or pancreatic epithelial cells is thus not sufficient to promote tumor development.

AID generates mutations and DNA double-strand breaks in pancreatic epithelium

The failure of AID expression to trigger tumorigenesis prompted us to evaluate its activity in epithelial cells. We first analyzed the *in vivo* mutagenic activity of ectopically expressed AID. The primary target sequences for AID mutagenic activity are immunoglobulin genes; although other genes are known to be susceptible to AID-induced mutagenesis, this occurs at much lower rates ($\sim 10^{-4}$ mutations/bp) and the mechanisms responsible for this susceptibility are poorly understood. One of the best-characterized requirements for AID activity is that the target sequence be transcriptionally active (Chaudhuri *et al*, 2003; Ramiro *et al*, 2003; Pavri & Nussenzweig, 2011). To simplify the mutagenesis analysis, we made use of the p48 pancreatic AID expression model to take advantage of the known low complexity transcriptome of acinar cells (MacDonald




Figure 3. AID expression in pancreas promotes DNA lesions.

A, B Analysis of AID mutagenic activity in Elastase1 (A) and Elastase2a (B) by next-generation sequencing. Pancreatic DNA was isolated from pools of R26AID^{+/KI}p48CRE^{+/KI} and control R26AID^{+/+}p48CRE^{+/KI} 20-week-old mice, and then PCR-amplified with specific primers and sequenced as previously described (Perez-Duran *et al*, 2012). Graphs show cytosine mutation frequency overall (total) or at AGCT hotspots. *n* = 2. **P*-value: Elastase1: 0.0382; Elastase2a: 0.009.

C HTM-mediated quantification of γ H2AX intensities per nuclei in pancreas explants cultured *in vitro* for 6 days. Red lines show mean values. Results of two independent experiments are shown. *****P* < 0.0001.

D Representative images of γ H2AX staining in pancreas explants from R26AID^{+/+}p48CRE^{+/KI} (top) or R26AID^{+/KI}p48CRE^{+/KI} mice (bottom) (40 \times magnification). White arrow points a representative positive cell.

Data information: Statistical differences were analyzed by two-tailed unpaired Student's *t*-test.

et al, 2010). AID preferentially targets the consensus hotspots WRCY/RGYW and particularly AGCT motifs (Rogozin & Kolchanov, 1992; Pham *et al*, 2003; Perez-Duran *et al*, 2012). Based on this, we analyzed the presence of mutations in 800–900 bp downstream of the transcriptional start site of two highly transcribed genes in pancreas, Elastase1 (Ela1) and Elastase2 (Ela2a), by next-generation sequencing, which allows large number of mutations to be analyzed at a very high depth (Perez-Duran *et al*, 2012). This analysis revealed that mutations are specifically accumulated at the Ela1 and Ela2 genes in R26AID^{+/KI}p48CRE^{+/KI} animals (Fig 3A and B) at frequencies similar to those of other non-Ig genes in B cells (Liu *et al*, 2008). AID activity was verified for Ela1 by conventional Sanger sequencing (Table 1). In contrast to previous reports (Matsumoto *et al*, 2007), we did not detect AID-induced mutations

at the tumor suppressor gene Trp53, where mutation frequency was identical in R26AID^{+/KI}p48CRE^{+/KI} mice and R26AID^{+/+}p48CRE^{+/KI} controls (Table 1).

To assess whether AID activity in R26AID^{+/KI}p48CRE^{+/KI} mice leads not only to mutations but also to more aggressive lesions, such as DSBs, we quantified γ -H2AX, a histone phosphorylation produced in response to this type of DNA damage. For this analysis, we generated acinar-cell explants from R26AID^{+/KI}p48CRE^{+/KI} and control mice, stained them with anti- γ H2AX, and quantified the intensity of staining per nucleus by high-throughput microscopy (HTM). We found that AID expression in R26AID^{+/KI}p48CRE^{+/KI} mice promoted a significant increase in the levels of γ -H2AX (Fig 3C and D), indicating that AID generates DSBs in this cellular context.

AID induces NKG2D ligands, T cell recruitment and apoptotic cell death in pancreas

Activation of the DNA damage response (DDR) pathway induces the expression of NKG2D ligands in epithelial cells, which are in turn recognized by NKG2D receptors expressed by NK cells and subsets of T cells (Dieffenbach *et al*, 2001; Gasser *et al*, 2005; Champsaur & Lanier, 2010; Raulet *et al*, 2013). This cross talk promotes the elimination of precancerous cells and is therefore a mechanism to prevent tumor development (Guerra *et al*, 2008). Given that AID expression in pancreas promotes mutations and DNA damage without leading to tumor development, we sought for the evidence of precancerous cells and found that pancreas from aged R26AID^{+/KI}p48CRE^{+/KI} mice contained more proliferating cells, as assessed by Ki67 staining, than control pancreas (Fig 4A), indicating that pancreatic AID expression leads to an abnormal rate of cell division. The epithelial identity of Ki67⁺ cells was confirmed both by morphology (Fig 4A, magnified micrograph on the right) and by staining with the epithelium-specific anti-cytokeratin 8 antibody (Figs 4B and EV2). We next asked whether the NKG2D immune surveillance pathway could be in play in R26AID^{+/KI}p48CRE^{+/KI} mice. To test this hypothesis, we first analyzed the expression of the Rae1 NKG2D ligand in epithelial cells from pancreatic explants of R26AID^{+/KI}p48CRE^{+/KI} and control mice by flow cytometry. Acinar cells from R26AID^{+/KI}p48CRE^{+/KI} mice expressed higher levels of Rae1 than their control littermates, although the difference was not statistically significant (Fig 4C). To assess whether RAE ligands were expressed by pancreatic cells *in vivo*, we prepared pancreas extracts from aged (75-week-old) R26AID^{+/KI}p48CRE^{+/KI} mice and controls and measured the amount of five RAE isoforms by droplet digital PCR (ddPCR). With this technique, each sample is fractionated into thousands of droplets, in which PCR amplification reactions occur independently, thereby increasing the sensitivity and quantitative potential of the amplification. Amplification of RAE isoforms was detected in more drops from R26AID^{+/KI}p48CRE^{+/KI}

Table 1. Analysis of AID mutagenic activity by Sanger sequencing.

	Genotype	Total clones analyzed	Mutations	Total bp sequenced	Frequency ($\times 10^4$)
Elastase1	R26AID ^{+/+} p48CRE ^{+/KI}	84	4	70,018	0.571
	R26AID ^{+/KI} p48CRE ^{+/KI}	82	13	69,355	1.87
Trp53	R26AID ^{+/+} p48CRE ^{+/KI}	66	0	59,472	0
	R26AID ^{+/KI} p48CRE ^{+/KI}	59	1	53,936	0.185




Figure 4. AID expression in pancreas promotes proliferation and NKG2D ligand expression.

- A Representative images of Ki67 staining in pancreas from aged (75-week-old) R26AID^{+/+}p48CRE^{+/KI} (top) or R26AID^{+/KI}p48CRE^{+/KI} mice (bottom). Detail is shown on the right. Scale bar: 100 μm. Graph shows quantification of Ki67-positive epithelial cells per mm² of tissue (n = 8). *P = 0.0266.
- B Representative immunofluorescence staining of 20-week-old R26AID^{+/KI}p48CRE^{+/KI} mice: blue, DAPI; red, Ki67; green, CK8. Scale bar: 50 μm. Detail is shown on the bottom.
- C Quantitative FACS analysis of RAE expression in pancreatic explants from R26AID^{+/+}p48CRE^{+/KI} and R26AID^{+/KI}p48CRE^{+/KI} mice. Left: Graph shows mean fluorescence intensity. Each dot represents an individual mouse. n = 9 (R26AID^{+/+}p48CRE^{+/KI}); 8 (R26AID^{+/KI}p48CRE^{+/KI}). P = 0.077. Right: Representative FACS staining for RAE in explants from R26AID^{+/+}p48CRE^{+/KI} (red) and R26AID^{+/KI}p48CRE^{+/KI} mice (black).
- D Analysis of RAE expression by ddPCR in aged (75-week-old) mice. Data are presented as the percentage of positive drops normalized to the mean control value. Each point represents an individual mouse and shows the mean amplification from two independent experiments n = 7 (R26AID^{+/+}p48CRE^{+/KI}); 9 (R26AID^{+/KI}p48CRE^{+/KI}). *P = 0.012.
- E Analysis of killing activity. Primary explants of pancreatic cells from R26AID^{+/+}p48CRE^{+/KI} or R26AID^{+/KI}p48CRE^{+/KI} mice were cultured with primary NK cells, and killing activity was assessed as described in Materials and Methods (n = 4). P = 0.19.

Data information: Statistical differences were analyzed by two-tailed unpaired Student's *t*-test.

samples than from controls (Fig 4D), indicating that AID promotes the expression of NKG2D ligands in pancreas, most likely as a result of DSB and DDR. We found that primary explants from R26AID^{+/KI}p48-CRE^{+/KI} tended to be more sensitive to NK-mediated killing than R26AID^{+/+}p48-CRE^{+/KI} littermate controls (Fig 4E), indicating that NKG2D ligand expression in AID-expressing pancreas is functional.

We next asked whether the expression of RAE ligands promoted the recruitment of immune cells to R26AID^{+/KI}p48-CRE^{+/KI} pancreas *in vivo*. Hematoxylin–eosin staining of pancreas sections from aged mice clearly revealed the presence of immune infiltrates in AID-expressing pancreas of R26AID^{+/KI}p48-CRE^{+/KI} mice (Fig 5A). The composition of these immune infiltrates was analyzed by antibody staining to detect macrophages (F4/80), B cells (Pax5) and T cells (CD3). The vast majority of cells in the immune infiltrates of R26AID^{+/KI}p48-CRE^{+/KI} mice were CD3⁺ T cells (Fig 5B), with only a negligible contribution from B cells and macrophages (not shown). To discount age-related effects, we analyzed 20-week-old mice, finding that the accumulation of T cell infiltrates is detectable in these young animals (Fig 5C). The main NKG2D-expressing T cell subset is the CD8⁺ population (Raulet *et al*, 2013), and immunofluorescence analysis of immune infiltrates revealed that a high proportion of the CD3⁺ infiltrate is composed of CD8⁺ T cells (Fig 5C), a finding consistent with the reported recruitment of CTL

cells to pancreatic islets transgenically expressing Rae ϵ (Markiewicz *et al*, 2012). Finally, we found that aged R26AID^{+/KI}p48-CRE^{+/KI} mice had significantly higher levels of pancreatic TNF- α mRNA than control littermates (Fig 5D), indicating that AID promotes the expression of effector cytotoxicity. Consistently, R26AID^{+/KI}p48-CRE^{+/KI} pancreas contained cells undergoing apoptotic cell death, detected by caspase-3 immunohistochemistry (Fig 5E, P = 0.054). Together, these results indicate that heterologous AID expression in pancreas promotes a cytotoxic response, most likely arising from the generation of genotoxic activity and NKG2D ligand expression and the recruitment of NKG2D-expressing CTL cells.

Discussion

In recent years, the finding that inflammatory cues induce AID expression in epithelial cells has boosted interest in the notion that AID might promote carcinogenesis and even be the causative link between inflammation and neoplastic transformation (Marusawa *et al*, 2011; Takai *et al*, 2012). In this regard, our analyses of AID expression in response to TNF- α have confirmed previous data in colonic cell lines and have expanded these observations to pancreatic cell lines. In addition, we found that primary pancreatic epithelium is also sensitive to TNF- α , suggesting that AID




Figure 5. AID expression in pancreas promotes immune infiltration and cell death.

A Hematoxylin–eosin (HE) staining of pancreas from aged (75-week-old) R26AID^{+/+}p48CRE^{+/Kl} and R26AID^{+Kl}p48CRE^{+/Kl} mice. Left: Representative HE staining showing an immune infiltrate in a R26AID^{+Kl}p48CRE^{+/Kl} mouse. Scale bar: 200 μm. Right: Quantification of number of foci per mm² of tissue. *n* = 13 (R26AID^{+/+}p48CRE^{+/Kl}); 21 (R26AID^{+Kl}p48CRE^{+/Kl}). ***P* = 0.0098.

B CD3 immunohistochemistry of pancreas from 75-week-old R26AID^{+/+}p48CRE^{+/Kl} and R26AID^{+Kl}p48CRE^{+/Kl} mice. Left: Representative image of a CD3 infiltrate in a R26AID^{+Kl}p48CRE^{+/Kl} mouse. Scale bar: 100 μm. Right: Quantification of the number of CD3-positive cells per mm² of tissue. *n* = 12 (R26AID^{+/+}p48CRE^{+/Kl}); 13 (R26AID^{+Kl}p48CRE^{+/Kl}). **P* = 0.0149.

C Representative immunofluorescence staining of CD3 and CD8 in pancreas from 20-week-old R26AID^{+/+}p48CRE^{+/Kl} and R26AID^{+Kl}p48CRE^{+/Kl} mice. Scale bar: 20 μm.

D TNF-α expression. Total RNA was isolated from pancreas of aged (75-week-old) R26AID^{+/+}p48CRE^{+/Kl} and control mice, and TNF-α expression was quantified by qRT-PCR. Each dot represents an individual mouse. *n* = 8 (R26AID^{+/+}p48CRE^{+/Kl}); 15 (R26AID^{+Kl}p48CRE^{+/Kl}). **P* = 0.0392.

E Cell death detection. Pancreas from 75-week-old R26AID^{+/+}p48CRE^{+/Kl} and R26AID^{+Kl}p48CRE^{+/Kl} mice were stained with anti-caspase-3. Left: Representative staining from an R26AID^{+/+}p48CRE^{+/Kl} mouse (top) and an R26AID^{+Kl}p48CRE^{+/Kl} mouse (bottom). Scale bar: 50 μm. Right: Quantification of number of cells per mm² of tissue. Each dot represents an individual mouse. *n* = 9 (R26AID^{+/+}p48CRE^{+/Kl}); 11 (R26AID^{+Kl}p48CRE^{+/Kl}). *P* = 0.0545.

Data information: Statistical differences were analyzed by two-tailed unpaired Student's *t*-test.

expression can indeed take place *in vivo* in pro-inflammatory contexts.

We wanted to further explore the physiological relevance of AID expression in promoting epithelium malignant transformation. We found that AID deficiency does not reduce the incidence of oncogenic lesions in an inflammation-induced carcinoma model. Our results contrast with the finding that AID deficiency reduces colon carcinogenesis in IL10^{-/-} mice (Takai *et al*, 2012). It may be that the effect reported by Takai *et al* is not specifically driven by epithelial cells, but rather by B cells in IL10^{-/-} mice, a possibility that could be tested using conditionally rather than constitutively AID-deficient animals.

Previous reports have claimed that AID heterologous expression leads to epithelial cell neoplasia in various tissues (Endo *et al*, 2008; Takai *et al*, 2009, 2012; Marusawa *et al*, 2011). In those studies, overexpression was always achieved with transgenes, none of which was epithelium specific, and the incidence of neoplasias was extremely low, and varied with the insertion sites and with ongoing mouse generations (Okazaki *et al*, 2003). In contrast, here we have

developed epithelium-specific conditional knock-in models, thus avoiding both widespread expression and transgene-derived artifacts. Remarkably, our models allowed B-cell-like AID expression levels, a basic prerequisite for AID to be competent, given that its levels are rate limiting in its native context (Sernandez *et al*, 2008; Takizawa *et al*, 2008). We found that AID expressed ectopically in pancreatic cells is able to mutate non-immunoglobulin genes and to generate genotoxic DSBs, suggesting that in this context, there are no obvious mechanisms for negative regulation of AID activity, as they were previously proposed for transgenic AID expression in B cells (Muto *et al*, 2006). AID mutagenic activity was detected in two highly expressed pancreatic genes (Ela1 and Ela2), which is expected from the well-established link between AID activity and transcription of its target genes (Chaudhuri *et al*, 2003; Ramiro *et al*, 2003; MacDonald *et al*, 2010; Pavri & Nussenzweig, 2011).

Our data indicate that AID activity in pancreas does not promote pancreatic carcinogenesis; instead, it triggers an NKG2D-mediated cytotoxic response that would eliminate pretumoral cells and prevent carcinoma development, in line with the published finding

that AID promotes an NKG2D immune response in B cells infected with the Abelson murine leukemia virus (Gourzi *et al*, 2006). The lymphomagenic potential of AID was previously shown to be dampened in B cells (Muto *et al*, 2006; Robbiani *et al*, 2009), where p53 exerts a cell-intrinsic tumor suppressor function (Robbiani *et al*, 2009). Here, we provide evidence of a further protective mechanism triggered by AID activity and carried out through an extrinsic immunosurveillance pathway in epithelial tissues. These data highlight the diversity of safeguarding events in AID-expressing cells and encourage a refined view of the previously acknowledged contribution of endogenous AID to epithelial-derived tumors.

Materials and Methods

Mice

R26AID mice were generated by insertion of a construct encoding mouse AID cDNA into the Rosa26 IRES-GFP targeting vector (Nyabi *et al*, 2009). AID cDNA was PCR-amplified from C57BL/6 mice (primers: forward 5'-TTCTGTGAAGACCGCAAGGCT-3'; reverse 5'-CCCTTCCCAGGCTTTGAAA-3'), cloned into the pENTR/D-TOPO Gateway vector (Invitrogen), and subsequently recombined into the Rosa 26 targeting vector, in which the cloned construct is preceded by a loxP-flanked transcriptional stop cassette and followed by an internal ribosomal entry site and GFP. The construct was linearized with PvuI before electroporation into hybrid 129/C57BL/6 ES cells. Clones positive for homologous recombination in the Rosa26 locus were identified by Southern blot of EcoRV-digested genomic DNA hybridized with a 5'-arm Rosa26 probe. R26AID mice were backcrossed to C57BL/6 background for 5 generations. R26AID mice were crossed with Villin-CRE^{+/K1} mice (el Marjou *et al*, 2004) and p48-CRE^{+/K1} mice (Kawaguchi *et al*, 2002) to promote expression of AID in colonic and pancreatic epithelial cells, respectively. Both Villin-CRE and p48-CRE mice were backcrossed to C57BL/6 for 5 generations. Balb/c AID^{-/-} mice were generated by backcrossing AID^{-/-} mice (Muramatsu *et al*, 2000) for 6 generations by speed congenics (Ramiro *et al*, 2004). Mice of both genders were used unless specified otherwise. All animals were housed in the Centro Nacional de Investigaciones Cardiovasculares animal facility under a 12-h light/dark cycle with food *ad libitum*.

Number of animals per group to detect biologically significant effect sizes was calculated using appropriate statistical sample size formula and indicated in the biometrical planning section of the animal license submitted to the governing authority. Blinding and randomization was not applicable to the animal studies. All animal procedures conformed to EU Directive 2010/63EU and Recommendation 2007/526/EC regarding the protection of animals used for experimental and other scientific purposes, enforced in Spanish law under RD 53/2013.

Cell lines and primary acinar cell culture

Primary pancreatic acinar cells were isolated and cultured as described in Gout *et al* (2013). Briefly, complete pancreas from 8-week-old mice was mechanically and enzymatically digested with collagenase to obtain isolated acinar structures. Acini were grown in Waymouth's medium supplemented with 2.5% FBS, 10 mM HEPES,

0.25 mg/ml trypsin inhibitor (Sigma) and 25 ng/ml of recombinant human epidermal growth factor (Sigma). PaTU-8988S, AsPC-1, LoVo and SW480 cells were grown in DMEM supplemented with 10% FCS and 10 mM HEPES. PaTU-8988S cell line was kindly provided by Dr Thomas Gress (University of Marburg). AsPC-1, LoVo and SW480 cell lines were obtained from the ATCC. All of them were mycoplasma negative. TNF- α (50 ng/ml) was added when indicated.

DSS-induced colitis-associated cancer (CAC) experiments

8–10-week-old Balb/c AID^{+/-} and AID^{-/-} (Ramiro *et al*, 2004) mice were given 3% dextran sulfate sodium salt (DSS, Sigma) in their drinking water for 5 days followed by regular drinking water for 10 days. Colon samples were obtained from these mice after 10 cycles of DSS treatment, and Swiss roll preparations were fixed, embedded in paraffin for section, and stained with hematoxylin/eosin (H/E).

Immunofluorescence

Pancreas and colon specimens were fixed with 4% paraformaldehyde, incubated with 30% sucrose, embedded in OCT compound (Olympus), and frozen in dry ice. 10- μ m sections were permeabilized and blocked with Image-It FX signal enhancer (Invitrogen, Molecular Probes). The following antibodies were used: rabbit anti-GFP (Abcam, 1/100), rat anti-mouse CD8a (BD Pharmingen, 1/100), mouse anti-Cytokeratin 8 (TROMA I, 1/50), goat anti-rabbit Alexa Fluor 488 (Molecular Probes, 1/500), goat anti-mouse Alexa Fluor 488 (Molecular Probes, 1/500), and goat anti-rat Cy3 (Jackson ImmunoResearch, 1/500). Slides were mounted with Vectashield mount medium containing DAPI (Vector Laboratories).

Immunohistochemistry

Pancreases were fixed in neutral-buffered 10% formalin solution (Sigma), embedded in paraffin blocks, and cut in 5- μ m sections. H/E staining was performed using standard protocols. For immunohistochemistry, sodium citrate buffer was used for antigen retrieval. The following antibodies were used: polyclonal rabbit anti-human CD3 (Dako, 1/200), rabbit anti-Ki67 (Abcam, 1/100), and biotinylated goat anti-rabbit (Abcam, 1/200). Biotinylated antibody was detected with the ABC system using diaminobenzidine as substrate (Vector Laboratories). Images were acquired with a Leica DM2500 microscope fitted with a 20 \times magnification lens. Reactive cells from 10 microscope fields per pancreas were counted using ImageJ. Results are shown as the number of reactive cells per mm² of tissue.

qRT-PCR

RNA was extracted from colon samples with TRIzol (Sigma). RNA from pancreas samples was extracted with GTC solution, following a standard phenol-chloroform protocol (reagents from Sigma), followed by DNase treatment (Qiagen). cDNA was synthesized using Random Hexamers (Roche) and SuperScript II reverse transcriptase. mRNA was quantified by SYBR green assay (Applied Biosystems), with normalization to GAPDH. The following primers were used: human-AID (forward) 5'-AAA TGT CCG CTG GGC TAA GG-3', (reverse) 5'-GGA GGA AGA GCA ATT CCA CGT-3';

human-GAPDH (forward) 5'-GAA GGT GAA GGT CGG AGT C-3', (reverse) 5'-GAA GAT GGT GAT GGG ATT TC-3'; mouse AID (forward) 5'-ACC TTC GCA ACA AGT CTG GCT-3', (reverse) 5'-AGC CTT GCG GTC TTC ACA GAA-3'; mouse-GAPDH (forward) 5'-TGA AGC AGG CAT CTG AGG G-3', (reverse) 5'-CGA AGG TGG AAG AGT GGG AG-3'; mouse-TNF- α (forward) 5'-AGC CCA CGT CGT AGC AAA CCA-3', (reverse) 5'-ACA ACC CA CGG CTG GCA CC-3'.

Next-generation sequencing for detection of mutations

DNA from three R26AID^{+/+}p48-CRE^{+/KI} and three R26AID^{+/KI}p48-CRE^{+/KI} mice was extracted and amplified using the following oligonucleotides: Elastase1 (forward) 5'-GCA CAG CAT CTT TTG TTT GGG TAA-3', (reverse) 5'-GGG GAC AGT GGT CTA CTC TCT-3'; Elastase2a (forward) 5'-ACG AGT CCA GGA CAA TCA GAG A-3', (reverse) 5'-TGA TAA GGC CAC TCA TAA AAA GGA-3'. Amplification reactions were carried out with 2.5 U of Pfu Ultra (Stratagene) in a 50- μ l reaction with the following profile: 94°C for 5 min followed by 25 cycles at 94°C for 10 s, 60°C for 30 s, and 72°C for 1 min. For NGS sequencing, the PCR product from two reactions per mouse were pooled, and an equimolar amount of DNA from each of the two mouse genotypes was pooled and mixed.

γ -H2AX staining and Opera acquisition

Primary pancreatic acinar cells were grown on 96-well plates (Perkin Elmer), and γ -H2AX (Millipore, 1/500) immunofluorescence was performed using standard procedures. Images were automatically acquired from each well with the Opera High-Content Screening System (Perkin Elmer). A 40 \times magnification lens was used, and pictures were taken at nonsaturating conditions. Images were segmented using DAPI staining to generate masks matching cell nuclei, and the mean per-cell γ -H2AX signal was calculated.

Droplet digital PCR

Each 20 μ l ddPCR contained 10 μ l of 2 \times ddPCR Supermix (Bio-Rad), 1 μ l of cDNA generated from 1.4 μ g of RNA, and 500 nM of each primer: pan-RAE (forward) 5'-TGG ACA CTC ACA AGA CCA ATG-3', (reverse) 5'-CCC AGG TGG CAC TAG GAG T-3'; and 250 nM Taqman probe (panRAE-FAM 5'-CCA TGA TTT ATC CGC AAA GCC AGG GCC-3'). After droplet generation, a mean of 12,000 droplets was obtained per sample.

Samples were transferred into a 96-well plate (Eppendorf) and cycled in a thermal cycler (Bio-Rad) under the following conditions: 95°C for 10 min followed by 38 cycles of 94°C for 30 s, 58°C for 1 min, and a final step at 98°C for 10 min. After amplification, samples were transferred to a droplet reader (QX100 Droplet Digital PCR, Bio-Rad) from which positive-drop data were extracted with QuantaSoft software. Results are represented as the proportion of positive drops in duplicates of each sample.

NK killing assay *in vitro*

Primary pancreatic acinar cells from R26AID^{+/+}p48-CRE^{+/KI} and R26AID^{+/KI}p48-CRE^{+/KI} mice were isolated as described above. After 6 days of culture, acinar cells were trypsinized and stained with CFSE. NK cells were isolated by cell sorting from wild-type

The paper explained

Problem

AID is a DNA-modifying enzyme expressed in germinal center B cells and required for the adaptive immune response, and its activity is linked to B cell neoplasias. A number of studies have reported AID expression in epithelial cells, usually associated with inflammatory cues. However, the contribution of AID expression to carcinogenesis remains controversial.

Results

We have developed conditional mouse models to promote epithelial-specific AID expression in pancreas and colon and found that despite high levels of AID expression and activity, these mice do not develop carcinomas. Instead, AID genotoxicity in pancreas promotes a response that involves NKG2D ligand expression, CD8⁺ T cell recruitment, and epithelial cell death. Therefore, our data provide evidence that AID-expressing epithelia are protected from neoplastic transformation by an NKG2D surveillance pathway.

Impact

Our results uncover a new protective mechanism counter-acting AID oncogenic potential through an extrinsic immunosurveillance pathway in epithelial cells. These findings challenge the view that AID could contribute significantly to inflammation-related carcinogenesis.

C57BL/6 mice. Both male and female aged 6–8 weeks were used as NK cells donors. NK cells and acinar pancreatic cells were co-cultured for 4 h in the presence of IL2 (2,000 U/ml, Peprotech) at a 1:10 (target cell:NK effector cell) ratio. Killing was analyzed by staining with DAPI by flow cytometry. Data are presented as the proportion of CFSE⁺DAPI⁺ cells normalized to the same population in cultures lacking NK cells.

Statistics

Statistical analyses were performed with GraphPad Prism (version 6.01 for Windows, GraphPad Software, San Diego, CA, USA) using two-tailed Student's *t*-test. $P \leq 0.05$ was considered statistically significant. Error bars in figures represent standard error of the mean (SEM). Normal distribution of data was assessed by applying a D'Agostino & Pearson omnibus normality test. F-test was used to compare variances between groups. For the survival analyses, GraphPad Prism was used and the Mantel–Cox test was applied. Differences were considered statistically significant at $P \leq 0.05$.

Expanded View for this article is available online:

<http://embomolmed.embopress.org>

Acknowledgements

We thank all members of the B Cell Biology Laboratory for helpful discussions; L Belder and A Alvarez-Prado for advice on next-generation sequencing; VG de Yébenes for critical reading of the manuscript; D Pisano, A Lopez-Contreras, N del Pozo, D Megias, A de Molina, and JM Ligos for technical advice; R Cuellar and Bio-Rad for kind support on ddPCR; and S Bartlett for English editorial support. AP-G is a fellow of the research training program (FPU- AP2009-1732) funded by the Ministerio de Educación, Cultura y Deporte, PP-D was an FPI fellow from the Ministerio de Ciencia e Innovación. ARR is supported by Centro Nacional de Investigaciones Cardiovasculares (CNIC). This work was funded by

grants from the Ministerio de Economía y Competitividad (SAF2010-21394, SAF2013-42767-R) and the European Research Council Starting Grant program (BCLYM-207844) to ARR. The CNIC is supported by the Ministerio de Economía y Competitividad and the Pro-CNIC Foundation. FXR is supported by SAF2011-29530 and ONCOBIO Consolider grants from Ministerio de Economía y Competitividad (Madrid, Spain), RTICC from Instituto de Salud Carlos III, and grant 256974 from European Union Seventh Framework Programme to FXR.

Author contributions

AP-G and ARR designed the research, analyzed data, and wrote the manuscript. AP-G, PP-D, TW, IVS, and SMM performed experiments. MC performed histopathological analyses. FXR designed the research and provided expertise and reagents on pancreatic adenocarcinoma.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Aguilar S, Corominas JM, Malats N, Pereira JA, Dufresne M, Real FX, Navarro P (2004) Tissue plasminogen activator in murine exocrine pancreas cancer: selective expression in ductal tumors and contribution to cancer progression. *Am J Pathol* 165: 1129–1139
- Alt FW, Zhang Y, Meng FL, Guo C, Schwer B (2013) Mechanisms of programmed DNA lesions and genomic instability in the immune system. *Cell* 152: 417–429
- Champsaur M, Lanier LL (2010) Effect of NKG2D ligand expression on host immune responses. *Immunol Rev* 235: 267–285
- Chaudhuri J, Tian M, Khuong C, Chua K, Pinaud E, Alt FW (2003) Transcription-targeted DNA deamination by the AID antibody diversification enzyme. *Nature* 422: 726–730
- Cooper HS, Murthy S, Kido K, Yoshitake H, Flanigan A (2000) Dysplasia and cancer in the dextran sulfate sodium mouse colitis model. Relevance to colitis-associated neoplasia in the human: a study of histopathology, B-catenin and p53 expression and the role of inflammation. *Carcinogenesis* 21: 757–768
- Di Noia JM, Neuberger MS (2007) Molecular mechanisms of antibody somatic hypermutation. *Annu Rev Biochem* 76: 1–22
- Diefenbach A, Jensen ER, Jamieson AM, Raulet DH (2001) Rae1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. *Nature* 413: 165–171
- Endo Y, Marusawa H, Kinoshita K, Morisawa T, Sakurai T, Okazaki IM, Watashi K, Shimotohno K, Honjo T, Chiba T (2007) Expression of activation-induced cytidine deaminase in human hepatocytes via NF- κ B signaling. *Oncogene* 26: 5587–5595
- Endo Y, Marusawa H, Kou T, Nakase H, Fujii S, Fujimori T, Kinoshita K, Honjo T, Chiba T (2008) Activation-induced cytidine deaminase links between inflammation and the development of colitis-associated colorectal cancers. *Gastroenterology* 135: 889–898
- Feagins LA, Souza RF, Spechler SJ (2009) Carcinogenesis in IBD: potential targets for the prevention of colorectal cancer. *Nat Rev Gastroenterol Hepatol* 6: 297–305
- Fodde R, Smits R (2001) Disease model: familial adenomatous polyposis. *Trends Mol Med* 7: 369–373
- Gasser S, Orsulic S, Brown EJ, Raulet DH (2005) The DNA damage pathway regulates innate immune system ligands of the NKG2D receptor. *Nature* 436: 1186–1190
- Gourzi P, Leonova T, Papavasiliou FN (2006) A role for activation-induced cytidine deaminase in the host response against a transforming retrovirus. *Immunity* 24: 779–786
- Gout J, Pommier RM, Vincent DF, Kaniewski B, Martel S, Valcourt U, Bartholin L (2013) Isolation and culture of mouse primary pancreatic acinar cells. *J Vis Exp* doi: 10.3791/50514
- Guerra N, Tan YX, Joncker NT, Choy A, Gallardo F, Xiong N, Knoblauch S, Cado D, Greenberg NM, Raulet DH (2008) NKG2D-deficient mice are defective in tumor surveillance in models of spontaneous malignancy. *Immunity* 28: 571–580
- Kawaguchi Y, Cooper B, Gannon M, Ray M, MacDonald RJ, Wright CV (2002) The role of the transcriptional regulator Ptf1a in converting intestinal to pancreatic progenitors. *Nat Genet* 32: 128–134
- Kovalchuk AL, duBois W, Mushinski E, McNeil NE, Hirt C, Qi CF, Li Z, Janz S, Honjo T, Muramatsu M et al (2007) AID-deficient Bcl-xL transgenic mice develop delayed atypical plasma cell tumors with unusual Ig/Myc chromosomal rearrangements. *J Exp Med* 204: 2989–3001
- Liu M, Duke JL, Richter DJ, Vinuesa CG, Goodnow CC, Kleinstein SH, Schatz DG (2008) Two levels of protection for the B cell genome during somatic hypermutation. *Nature* 451: 841–845
- MacDonald RJ, Swift GH, Real FX (2010) Transcriptional control of acinar development and homeostasis. *Prog Mol Biol Transl Sci* 97: 1–40
- Mantovani A, Allavena P, Sica A, Balkwill F (2008) Cancer-related inflammation. *Nature* 454: 436–444
- el Marjou F, Janssen KP, Chang BH, Li M, Hindie V, Chan L, Louvard D, Chambon P, Metzger D, Robine S (2004) Tissue-specific and inducible Cre-mediated recombination in the gut epithelium. *Genesis* 39: 186–193
- Markiewicz MA, Wise EL, Buchwald ZS, Pinto AK, Zafirova B, Polic B, Shaw AS (2012) RAE1epsilon ligand expressed on pancreatic islets recruits NKG2D receptor-expressing cytotoxic T cells independent of T cell receptor recognition. *Immunity* 36: 132–141
- Martinelli P, Madriles F, Canamero M, Pau EC, Pozo ND, Guerra C, Real FX (2015) The acinar regulator Gata6 suppresses KrasG12V-driven pancreatic tumorigenesis in mice. *Gut* doi: 10.1136/gutjnl-2014-308042
- Marusawa H, Takai A, Chiba T (2011) Role of activation-induced cytidine deaminase in inflammation-associated cancer development. *Adv Immunol* 111: 109–141
- Matsumoto Y, Marusawa H, Kinoshita K, Endo Y, Kou T, Morisawa T, Azuma T, Okazaki IM, Honjo T, Chiba T (2007) Helicobacter pylori infection triggers aberrant expression of activation-induced cytidine deaminase in gastric epithelium. *Nat Med* 13: 470–476
- Matsumoto Y, Marusawa H, Kinoshita K, Niwa Y, Sakai Y, Chiba T (2010) Up-regulation of activation-induced cytidine deaminase causes genetic aberrations at the CDKN2b-CDKN2a in gastric cancer. *Gastroenterology* 139: 1984–1994
- Muramatsu M, Kinoshita K, Fagarasan S, Yamada S, Shinkai Y, Honjo T (2000) Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* 102: 553–563
- Muto T, Okazaki IM, Yamada S, Tanaka Y, Kinoshita K, Muramatsu M, Nagaoka H, Honjo T (2006) Negative regulation of activation-induced cytidine deaminase in B cells. *Proc Natl Acad Sci USA* 103: 2752–2757
- Nyabi O, Naessens M, Haigh K, Gembarska A, Goossens S, Maetens M, De Clercq S, Drogat B, Haenebalcke L, Bartunkova S et al (2009) Efficient mouse transgenesis using Gateway-compatible ROSA26 locus targeting vectors and F1 hybrid ES cells. *Nucleic Acids Res* 37: e55
- Okazaki IM, Hiai H, Kakazu N, Yamada S, Muramatsu M, Kinoshita K, Honjo T (2003) Constitutive expression of AID leads to tumorigenesis. *J Exp Med* 197: 1173–1181

- Pasqualucci L, Bhagat G, Jankovic M, Compagno M, Smith P, Muramatsu M, Honjo T, Morse HC 3rd, Nussenzweig MC, Dalla-Favera R (2008) AID is required for germinal center-derived lymphomagenesis. *Nat Genet* 40: 108–112
- Pavri R, Nussenzweig MC (2011) AID targeting in antibody diversity. *Adv Immunol* 110: 1–26
- Perez-Duran P, Belver L, de Yébenes VG, Delgado P, Pisano DG, Ramiro AR (2012) UNG shapes the specificity of AID-induced somatic hypermutation. *J Exp Med* 209: 1379–1389
- Petersen-Mahrt SK, Harris RS, Neuberger MS (2002) AID mutates *E. coli* suggesting a DNA deamination mechanism for antibody diversification. *Nature* 418: 99–103
- Pham P, Bransteitter R, Petruska J, Goodman MF (2003) Processive AID-catalysed cytosine deamination on single-stranded DNA simulates somatic hypermutation. *Nature* 424: 103–107
- Ramiro AR, Jankovic M, Callen E, Difilippantonio S, Chen HT, McBride KM, Eisenreich TR, Chen J, Dickins RA, Lowe SW et al (2006) Role of genomic instability and p53 in AID-induced c-myc-IgH translocations. *Nature* 440: 105–109
- Ramiro AR, Jankovic M, Eisenreich T, Difilippantonio S, Chen-Kiang S, Muramatsu M, Honjo T, Nussenzweig A, Nussenzweig MC (2004) AID is required for c-myc/IgH chromosome translocations in vivo. *Cell* 118: 431–438
- Ramiro AR, Stavropoulos P, Jankovic M, Nussenzweig MC (2003) Transcription enhances AID-mediated cytosine deamination by exposing single-stranded DNA on the nontemplate strand. *Nat Immunol* 4: 452–456
- Raulet DH, Gasser S, Gowen BG, Deng W, Jung H (2013) Regulation of ligands for the NKG2D activating receptor. *Annu Rev Immunol* 31: 413–441
- Revy P, Muto T, Levy Y, Geissmann F, Plebani A, Sanal O, Catalan N, Forveille M, Dufourcq-Labeau R, Gennery A et al (2000) Activation-induced cytosine deaminase (AID) deficiency causes the autosomal recessive form of the Hyper-IgM syndrome (HIGM2). *Cell* 102: 565–575
- Robbiani DF, Bothmer A, Callen E, Reina-San-Martin B, Dorsett Y, Difilippantonio S, Bolland DJ, Chen HT, Corcoran AE, Nussenzweig A et al (2008) AID is required for the chromosomal breaks in c-myc that lead to c-myc/IgH translocations. *Cell* 135: 1028–1038
- Robbiani DF, Bunting S, Feldhahn N, Bothmer A, Camps J, Deroubaix S, McBride KM, Klein IA, Stone G, Eisenreich TR et al (2009) AID produces DNA double-strand breaks in non-Ig genes and mature B cell lymphomas with reciprocal chromosome translocations. *Mol Cell* 36: 631–641
- Robbiani DF, Nussenzweig MC (2013) Chromosome translocation, B cell lymphoma, and activation-induced cytosine deaminase. *Annu Rev Pathol* 8: 79–103
- Rogozin IB, Kolchanov NA (1992) Somatic hypermutagenesis in immunoglobulin genes. II. Influence of neighbouring base sequences on mutagenesis. *Biochim Biophys Acta* 1171: 11–18
- Sernandez IV, de Yébenes VG, Dorsett Y, Ramiro AR (2008) Haploinsufficiency of activation-induced deaminase for antibody diversification and chromosome translocations both in vitro and in vivo. *PLoS ONE* 3: e3927
- Stavnezer J, Guikema JE, Schrader CE (2008) Mechanism and regulation of class switch recombination. *Annu Rev Immunol* 26: 261–292
- Takai A, Marusawa H, Minaki Y, Watanabe T, Nakase H, Kinoshita K, Tsujimoto G, Chiba T (2012) Targeting activation-induced cytosine deaminase prevents colon cancer development despite persistent colonic inflammation. *Oncogene* 31: 1733–1742
- Takai A, Toyoshima T, Uemura M, Kitawaki Y, Marusawa H, Hiai H, Yamada S, Okazaki IM, Honjo T, Chiba T et al (2009) A novel mouse model of hepatocarcinogenesis triggered by AID causing deleterious p53 mutations. *Oncogene* 28: 469–478
- Takizawa M, Tolarova H, Li Z, Dubois W, Lim S, Callen E, Franco S, Mosaico M, Feigenbaum L, Alt FW et al (2008) AID expression levels determine the extent of cMyc oncogenic translocations and the incidence of B cell tumor development. *J Exp Med* 205: 1949–1957
- Vonderheide RH, Bayne LJ (2013) Inflammatory networks and immune surveillance of pancreatic carcinoma. *Curr Opin Immunol* 25: 200–205



License: This is an open access article under the terms of the Creative Commons Attribution 4.0 License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.