

Marshall, P., Bayliss, A., Farid, S., Tyers, C., Bronk Ramsey, C., Cook, G., Doğan, T., Freeman, S.P.H.T., İlkmen, E. and Knowles, T. (2019) 14C wiggle-matching of short tree-ring sequences from post-medieval buildings in England. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 438, pp. 218-226. (doi:10.1016/j.nimb.2018.03.018)

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

http://eprints.gla.ac.uk/207513/

Deposited on 10 January 2020

Enlighten – Research publications by members of the University of Glasgow <u>http://eprints.gla.ac.uk</u>

¹⁴C wiggle-matching of short tree-ring sequences from post-medieval buildings in England

Marshall, P^{a*}, Bayliss, A^b, Farid, S^a, Tyers, C^a, Bronk Ramsey, C^c, Cook, G^d, Doğan, T^e, Freeman, S P H T^d, İlkmen, E^e, and Knowles, T^f

^a Historic England, Cannon Bridge House, London, EC4R 2YA, UK

^b Historic England, Cannon Bridge House, London, EC4R 2YA and Biological and

Environmental Sciences, University of Stirling, Stirling FK9 4LA, UK

 $^\circ$ Oxford Radiocarbon Accelerator Unit, Dyson Perrins Building, South Parks Road, Oxford, OX1 3QY, UK

^d SUERC Radiocarbon Dating Laboratory, Scottish Enterprise Technology Park, Rankine Avenue, East Kilbride G75 0QF, UK

^e Tübitak National 1 MV AMS Laboratory, Marmara Research Center, Earth and Marine Science Institute, Barış Mahallesi, 1 Dr Zeki Acar Caddesi, PK 21 41470, Gebze Kocaeli, Turkey

^f School of Chemistry, University of Bristol, Cantock's Close, Clifton, Bristol, BS8 1T5, UK ^{*} Corresponding author. Address: Historic England, Cannon Bridge House, 25 Dowgate Hill, London, EC4R 2YA, UK. Email: peter.Marshall@HistoricEngland.org.uk

<u>Abstract</u>

This study tests whether accurate dating by AMS radiocarbon wigglematching short tree-ring series (c. 30 annual rings) in the period after AD 1510 can be achieved routinely. Such an approach has proved problematic for some intervals in the period AD 1160–1541 (Bayliss *et al.* 2017), which are before single-year calibration data are available (Stuiver 1993). We suggest that such calibration data are essential if this approach is to be employed for the informed conservation of standing buildings.

<u>Keywords</u>

¹⁴C; AMS; radiocarbon wiggle match; radiocarbon calibration

1. Introduction

Over the past 25 years scientific dating has become an integral part of the processes for conservation and repair of historic buildings in England. Precise dating informs decisions about the preservation of buildings, allows us to identify significant fabric, and aids in the specification of appropriate repair

strategies. Small differences in date can lead to great differences in the significance of the extant building, and thus to great differences in the costs of the agreed solution for a particular structure.

In providing the required precise dating for historic buildings, the scientific dating method of choice is dendrochronology. In some cases, however, treering analysis does not provide calendar dating, usually either because there are insufficient growth rings in the timbers or because they are of a species that is unsuitable for dendrochronology. In these cases, radiocarbon wigglematching is needed to provide an equivalent level of precision and reliability.

2. Material and Methods

Radiocarbon wiggle-matching has previously been undertaken on part of a 303-ring pine series dated by dendrochronology to AD 1367–1670 from Jermyn Street, London (Tyers *et al.* 2009). In this study, measurements from three participating AMS laboratories all provided accurate wiggle-matches from ring series covering a century or more. Insufficient data are available from this core, however, to undertake wiggle-matching on shorter sequences.

For this reason, new measurements were obtained on a 149-ring core, LED-A22, from the east principal rafter in truss 3 of one of the roofs of Ledston Hall, West Yorkshire (1.34104 W, 53.75494 N). The growth-rings in this timber span AD 1520–1668, with 20 rings of sapwood and bark edge surviving. It is part of a site sequence, LEDASQ01, which is dated by dendrochronology to AD 1424–1668 (Table 1). The ring-width data of the series are provided in Arnold *et al.* (2015).

Radiocarbon measurements were made on a total of 60 single-year tree-ring samples from this core in 2016–17. The 17 dated at the Scottish Universities Environmental Research Centre were prepared to α -cellulose, combusted, graphitised, and dated by AMS as described by Dunbar *et al.* (2016). The 18 dated at the Oxford Radiocarbon Accelerator Unit were processed using an acid-alkali-acid pretreatment followed by bleaching with sodium chlorite as described by Brock *et al.* (2010, table 1 (UW)), graphitised (Dee and Bronk

Ramsey 2000), and measured by AMS (Bronk Ramsey *et al.* 2004). The 14 dated at the Bristol Radiocarbon Accelerator Mass Spectrometry Facility were processed using the base-acid-base-acid-bleach method as described by Němec *et al.* (2010), graphitised using an IonPlus AGE3 graphitisation system (Wacker *et al.* 2010), and measured using a MICADAS AMS (Synal *et al.* 2007). The 15 dated at the TUBITAK were pretreated using the acid-alkali-acid method modified from Hadjas *et al.* (2004), graphitised (Wacker *et al.* 2010) and measured by AMS on a 1 MV NEC Pelletron accelerator.

At Oxford and SUERC, δ^{13} C values, relative to VPDB, were obtained by IRMS from the gas combusted for graphitisation; at BRAMS and TUBITAK δ^{13} C values were measured by AMS.

The conventional radiocarbon ages reported for these samples, along with the rings dated, are listed in Table 2. The quoted errors are each laboratory's estimates of the total error in their dating systems. Five pairs of replicate radiocarbon measurements are available on rings dated to the same calendar year, all of which are statistically consistent at 95% confidence (Ward and Wilson 1978; Table 2). This scatter is in line with statistical expectation. Only two of the replicate δ^{13} C values are statistically consistent at 95% confidence, one is inconsistent at 95% confidence, but consistent at 95% confidence, and two are significantly different at more than 99% confidence (Ward and Wilson 1978; Table 2). These results are more scattered than would be expected on statistical grounds. This suggests that the different pre-treatment protocols used for wood samples by the Oxford and SUERC laboratories may be affecting the δ^{13} C values, especially since replicate δ^{13} C values on bone samples reported by the two laboratories in the same period are in much better agreement (Bayliss *et al.* 2016, fig. 14).

Forty-two pairs of replicate measurements, 17 groups of triplicate measurements, and one quadruple group of measurements are also available on rings dated by AMS (this study) and gas proportional counting Stuiver (1993) to the same calendar year (Figure 1). Of these 60 sets of radiocarbon ages, 52 groups are consistent at 95% confidence, 6 groups are consistent at

99% confidence, and two inconsistent at more than 99%. This scatter is rather more than would be expected simply on the basis of statistics. Stuiver (1993) reported counting errors only, however, and, when the errors quoted for this dataset are inflated using the laboratory error multiplier suggested by Stuiver *et al.* (1998, 1045), 56 sets of measurements are consistent at 95% confidence and the remaining four at 99% confidence, which is within statistical expectation.

3. Wiggle-matching the entire sequence

Wiggle-matching has been undertaken using the Bayesian approach first described by Christen and Litton (1995), implemented using OxCal v4.2 (Bronk Ramsey 2009) and the IntCal113 atmospheric calibration data for the northern hemisphere (Reimer *et al.* 2013).

Figure 2 shows the model for the entire sequence of data from Ledston Hall. This has good overall agreement (Acomb: 70.2, An:9.1, n:60; Bronk Ramsey *et al.* 2001), and estimates the final ring of the sequence to have been formed in *cal AD 1664–1671 (95% probability; SUERC-68040*; Fig 2). This is compatible with the date of AD 1668 produced for this ring by dendrochronology (Table 1).

Wiggle-matching was then undertaken separately of the radiocarbon results quoted by each laboratory (Figs. 3a–b and 4a–b). The model composed of measurements made in Bristol has good overall agreement (Acomb: 102.8, An: 18.9, n: 14; Fig. 3a) and estimates that the last ring of the timber formed in *cal AD 1658–1672 (95% probability; Ring 149*), probably in *cal AD 1662–1670 (68% probability)*. The model of measurements made at Oxford also has good overall agreement (Acomb: 30.9, An: 16.7, n: 18; Fig. 3b) and estimates the last ring of the timber to have formed in *cal AD 1655–1670 (95% probability; Ring 149*), probably in *cal AD 1655–1670 (95% probability; Ring 149*), probably in *cal AD 1657–1664 (68% probability)*. Results from SUERC also have good overall agreement (Acomb: 109.9, An: 17.1, n: 17; Fig. 4a) and suggest that the last ring of the timber formed in *cal AD 1656–1674 (95% probability; SUERC-68040*), probably in *cal AD 1662–1671 (68% probability)*. The wiggle match for TUBITAK measurements also

have good overall agreement (Acomb: 75.0, An: 18.3, n: 15; Fig. 4b) and suggest that the last ring of the timber formed in *cal AD 1661–1679 (95% probability; ring_149*), probably in *cal AD 1665–1674 (68% probability)*. In all cases the estimated date of the final ring *(at 95% probability)* includes the felling date provided by dendrochronology of AD 1668.

4. Wiggle-matching partial sequences

Given that the length of the available tree-ring sequence is the most common limitation on successful dendrochronology in historic buildings from England, we ran a series of 25 short wiggle-matches on blocks consisting of between 29 and 30 rings. The results on the seven dated rings in each block were incorporated into a wiggle-match model that estimated the date of the final ring of the complete core. These estimates could then be compared with the known date for the final ring as derived from dendrochronology (AD 1668) to determine the accuracy of the short wiggle-matches.

The results are given in Table 3 and summarised in Figure 5. The Highest Posterior Density interval at 95% probability includes the tree-ring date for the final ring of LED-A22 (AD 1668) in all cases, and the interval at 68% probability includes the tree-ring date in all but four cases. This correspondence between the results of the wiggle-matching and the dendrochronology is greater than statistical expectation.

5. The long wiggle-match (AD 1160–1668)

As illustrated in Figure 2, a wiggle-match comprising the radiocarbon measurements on the 45 dated rings from Ledston Hall has good overall agreement (Acomb: 70.2; An: 9.1; n: 60) and produces posterior distributions that are compatible with the dendrochronology (AD 1668). Similarly, the wiggle-match including the results on the 18 dated rings from Jermyn Street (Tyers *et al.* 2009, fig. 4) has good overall agreement when recalculated using IntCal13 (Acomb: 35.3; An: 16.7; n: 18) and also produces posterior distributions that are compatible with the dendrochronology (AD 1670).

In contrast, the wiggle-match including the radiocarbon measurements on the 79 dated rings from the sites considered by Bayliss *et al.* (2017, fig 9) has poor overall agreement (Acomb: 1.6; An: 8.0; n: 79), and the Highest Posterior Density interval for the final ring does not include the date obtained for this ring by dendrochronology (AD 1544) even at 99% probability.

When the entire dataset is combined to form a wiggle-match sequence covering AD 1160–1668, the model has poor overall agreement (Acomb: 1.7; An: 6.4; n: 121), and the Highest Posterior Density interval for the final ring does not include the date obtained for this ring by dendrochronology (AD 1668) even at 99% probability.

6. Discussion

These studies in combination suggest that wiggle-matching of either short (*c.* 30-ring) or long (more 100-ring) tree-ring sequences produces results that are compatible with dendrochronology in the period after AD 1510. In the centuries before this, there appears to be time periods when wiggle-matching does not produce such accuracy (AD 1240–1306 and AD 1396–1532; Bayliss *et al.* 2017, table 6).

This pattern is observed when considering the datasets measured at Oxford and SUERC separately (Bayliss *et al.* 2017, table 5; Tyers *et al.* 2009, table 3; Figs. 3b and 4a), and so must derive from the calibration curve used, IntCal13 (Reimer *et al.* 2013).

Figures 6–7 shows the radiocarbon ages obtained on single known-age treerings as part of this study and those reported by Tyers *et al.* (2009) and Bayliss *et al.* (2017) in comparison to the radiocarbon ages covering this period included in IntCal13 (Reimer *et al.* 2013). The latter are on decadal samples (Wk; Hogg *et al.* 2002), single-year and decadal samples (QL; Stuiver 1993, corrected as described by Stuiver and Becker 1993; Stuiver *et al.* 1998), decadal and bi-decadal samples (UB; Hogg *et al.* 2002; Pearson *et al.* 1986), and decadal and 23-year and 24-year samples (GrN; van der Plicht *et al.* 1995). Single-year data clearly dominate the period after AD 1510 (Fig 7), which is the period when wiggle-matching appears to be accurate within the precision quoted and the test data scatter within statistical expectation around the IntCal envelope. In the earlier period, there is more deviation between the test data and the IntCal envelope. As the statistical method of curve construction is the same in both periods (Niu *et al.* 2013), this is unlikely to be the cause of this difference, but rather there appears to be detailed structure in the atmospheric concentration of radiocarbon in the problematic periods which is not apparent from the calibration data currently available.

It should be noted that, if accurate wiggle-matching is possible in the post AD 1510 period because of the availability of single-year calibration data, then it is possible to accurately match measurements on samples of European wood against calibration data measured largely on Douglas fir trees that grew on the west coast of America. This would suggest neither intra-hemispheric locational offsets in the ¹⁴C concentration of wood (McCormac *et al.* 1995) nor the translocation of ¹⁴C between annual growth rings in trees (Grootes *et al.* 1989) are significant factors preventing accurate wiggle-matching.

A calibration curve based on ¹⁴C measurements on single tree-rings appears to be required for wiggle-matching to provide estimates of calendar date that are accurate to within the quoted uncertainty. Such accuracy is essential if the results are to inform the long-term preservation and conservation of historic buildings.

Acknowledgements

We would like to thank Alison Arnold and Robert Howard for their work on the tree-ring dating of Ledston Hall.

References

A. Arnold, R. Howard, Auckland Castle, Bishop Auckland, County Durham: Tree-Ring Analysis of Timbers English Heritage Research Report Series, English Heritage, Portsmouth, 2013.

A. Arnold, R. Howard, C. Tyers, Ledston Hall, Hall Lane, Ledston, Leeds, West Yorkshire.Tree-Ring Analysis of Timbers, Research Report Series, English Heritage, Portsmouth, 2015.

A.J. Arnold, R.E. Howard, C.D. Litton, Tree-ring Analysis of Timbers from the Main Guard, Pontefract Castle, Pontefract, West Yorkshire, Centre for Archaeology Report, English Heritage, Portsmouth, 2005b.

A.J. Arnold, R.E. Howard, C.D. Litton, Nottingham Tree-ring Dating Laboratory: additional dendrochronology dates, no 32 Vernacular Architecture, 39 (2008) 107–111.

A.J. Arnold, R.E. Howard, C.D. Litton, Tree-ring Analysis of Timbers from the Roof of the Keep, or Little Castle, Bolsover Castle, Derbyshire, Centre for Archaeology Report, English Heritage, Portsmouth, 2003.

A.J. Arnold, R.E. Howard, C.D. Litton, Tree-ring Analysis of Timbers from the Riding School, Bolsover Castle, Bolsover, Derbyshire, Centre for Archaeology Report, English Heritage, Portsmouth, 2005a.

A. Bayliss, N. Beavan, D. Hamilton, K. Köhler, K., É.Á.Nyerges, C. Bronk Ramsey, E. Dunbar, M. Fecher, T. Goslar, B. Kromer, P. Reimer, E. Bánffy, T. Marton, K. Oross, A. Osztás, I. Zalai-Gaál, A. Whittle, Peopling the past: creating a site biography in the Hungarian Neolithic, Bericht der Römisch-Germanischen Kommission 94 (2016) 23–91.

A. Bayliss, P. Marshall, C. Tyers, C. Bronk Ramsey, G. Cook, S.P.H.T. Freeman, S. Griffiths, Informing Conservation: Towards ¹⁴C Wiggle-Matching of Short Tree-Ring Sequences from Medieval Buildings in England,

Radiocarbon, 59 (2017), 985–1007. https://doi.org/10.1017/RDC.2016.61 F. Brock, T. Higham, P. Ditchfield, C. Bronk Ramsey, Current pretreatment methods for AMS radiocarbon dating at the Oxford Radiocarbon Accelerator Unit (ORAU), Radiocarbon, 52 (2010) 103–112.

https://doi.org/10.1017/S0033822200045069

C. Bronk Ramsey, Bayesian analysis of radiocarbon dates, Radiocarbon, 51 (2009) 337–360. https://doi.org/10.1017/S0033822200033865

C. Bronk Ramsey, J. van der Plicht, B. Weninger, 'Wiggle matching' radiocarbon dates, Radiocarbon 43(2A) (2001) 381–389.

radiocarbon dates, Radiocarbon 43(2A) (2001) 381-38

https://doi.org/10.1017/S0033822200038248

C. Bronk Ramsey, T. Higham, P. Leach, Towards high precision AMS: progress and limitations, Radiocarbon 46 (2004), 17–24.

https://doi.org/10.1017/S0033822200039308

A. Christen, C. Litton, A Bayesian approach to wiggle-matching, Journal of Archaeological Science 22(6) (1995) 719–725. https://doi.org/10.1016/0305-4403(95)90002-0

M. Dee, C. Bronk Ramsey, Refinement of graphite target production at ORAU, Nuclear Instruments and Methods in Physics Research B, 172 (2000) 449–453. https://doi.org/10.1016/S0168-583X(00)00337-2

E. Dunbar, G.T. Cook, P. Naysmith, B.G. Tipney, S. Xu, AMS ¹⁴C dating at the Scottish Universities Environmental Research Centre (SUERC) Radiocarbon Dating Laboratory, Radiocarbon, 58 (2016) 9–23. https://doi.org/10.1017/RDC.2016.14

P.M. Grootes, G.W. Farwell, F.H. Schmidt, D.D. Leach, Rapid response of tree cellulose radiocarbon content to changes in atmospheric $^{14}CO_2$, concentration, Tellus, 41 (1989) 134-148. I. Hajdas, G. Bonani, J. Thut, G. Leone, R. Pfenninger, C. Maden A report on sample preparation at the ETH/PSI AMS facility in Zurich, Nuclear Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 223-224 (2004) 267-271. https://doi.org/10.1016/j.nimb.2004.04.054 A.G. Hogg, F.G. McCormac, T.F.G. Higham, P.J. Reimer, M.G.L. Baillie, J.G. Palmer, High-precision radiocarbon measurements of contemporaneous treering dated wood from the British Isles and New Zealand: AD 1850–950, Radiocarbon 44 (2002) 633-640. https://doi.org/10.1017/S0033822200032082 R.E. Howard, R.R. Laxton, C.D. Litton, Tree-ring analysis of timbers from Clumpcliff Farm, Methley Lane, Rothwell, nr Oulton, West Yorkshire, Ancient Monuments Labratory Report, English Heritage, Portsmouth, 2000. F.G. McCormac, M.G.L. Baillie, J.R. Pilcher, Location-dependent differences in the ¹⁴C content of wood, Radiocarbon, 37 (1995) 395–407. M. Němec, L. Wacker, I. Hajdas, H. Gäggeler, Alternative methods for cellulose preparation for AMS measurement, Radiocarbon 52 (2010) 1358-1370. https://doi.org/10.1017/S0033822200046440 M. Niu, T.G. Heaton, P.G. Blackwell, C.E. Buck, The Bayesian approach to radiocarbon calibration curve estimation: the IntCal13, Marine13, and ShCal13 methodologies, Radiocarbon, 55 (2013) 1905–1922. https://doi.org/10.2458/azu js rc.55.17222 G.W. Pearson, J.R. Pilcher, M.G.L. Baillie, D.M. Corbett, F. Qua, Highprecision ¹⁴C measurements of Irish oaks to show the natural ¹⁴C variations from AD 1840 to 5210 BC, Radiocarbon, 28 (1986) 911-934. https://doi.org/10.1017/S0033822200060197 P.J. Reimer, E. Bard, A. Bayliss, J.W. Beck, P. Blackwell, C. Bronk Ramsey, C.E. Buck, H. Cheng, R.L. Edwards, M. Friedrich, P.M. Grootes, T.P. Guilderson, H. Haflidason, I. Hajdas, C. Hatté, T.J. Heaton, D.L. Hoffmann, A.G. Hogg, K.A. Hughen, K.F. Kaiser, B. Kromer, S.W. Manning, M. Niu, R.W. Reimer, D.A. Richards, E.M. Scott, J.R. Southon, R.A. Staff, C.S.M. Turney, J. van der Plicht, IntCal13 and Marine13 radiocarbon age calibration curves 0-50,000 years cal BP, Radiocarbon, 55 (2013) 1869-1887. https://doi.org/10.2458/azu js rc.55.16947 M. Stuiver, A note on single-year calibration of the radiocarbon timescale AD 1510-1954, Radiocarbon, 35 (1993) 67-72. https://doi.org/10.1017/S0033822200013813 M. Stuiver, B. Becker, High-precision decadal calibration of the radiocarbon time scale, AD 1950-6000 BC, Radiocarbon, 35 (1993) 35-66. https://doi.org/10.1017/S0033822200013801 M. Stuiver, P.J. Reimer, T.F. Braziunas, High-precision radiocarbon age calibration for terrestrial and marine samples, Radiocarbon 40(3) (1998) 1127-1151. https://doi.org/10.1017/S0033822200019172 M. Stuiver, P.J. Reimer, E. Bard, W. Beck, G. S. Burr, K.A. Hughen, B. Kromer, G. McCormac, J. van der Plicht, and M. Spurk, IntCal98 radiocarbon age calibration, 24,000-0 cal BP, Radiocarbon 40 (3) (1998) 1041-1083. https://doi.org/10.1017/S0033822200019123

65

H.-A. Synal, M. Stocker, M. Suter, MICADAS: A new compact radiocarbon AMS system, Nucl. Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 259 (2007) 7–13.

https://doi.org/10.1016/j.nimb.2007.01.138

C. Tyers, J. Sidell, J. van der Plicht, P. Marshall, G. Cook, C. Bronk Ramsey, A. Bayliss, Wiggle-matching using known-age pine from Jermyn Street, London, Radiocarbon, 51 (2009) 385–396.

https://doi.org/10.1017/S0033822200055788

J. van der Plicht, E. Jansma, H. Kars, The 'Amsterdam Castle': a case study of wiggle matching and the proper calibration curve, Radiocarbon 37(3) (1995) 965–968. https://doi.org/10.1017/S0033822200015058

L. Wacker, M. Němec, J. Bourquin, A revolutionary graphitisation system: Fully automated, compact and simple, Nuclear Instruments Methods Phys. Res. Sect. B Beam Interact. with Mater. Atoms, 268 (2010) 931–934. https://doi.org/10.1016/j.nimb.2009.10.067

G.K. Ward, S.R. Wilson, Procedures for comparing and combining radiocarbon age determinations: a critique, Archaeometry, 20 (1978) 19–32. DOI: 10.1111/j.1475-4754.1978.tb00208.x

Table 1: Results of cross-matching site chronology LEDASQ01 with relevant independent site reference chronologies

Reference chronology	<i>t</i> -value	Span of chronology	Reference
Riding House, Bolsover	12.0	AD 1494–1744	Arnold <i>et al.</i> 2005a
Castle, Derbyshire			
Pontefract Castle,	11.0	AD 1507–1656	Arnold <i>et al.</i> 2005b
Pontefract, West			
Yorkshire			
Little Castle, Bolsover	10.4	AD 1532–1749	Arnold <i>et al.</i> 2003
Castle, Derbyshire			
Clumpcliff Farm,	10.2	AD 1452–1613	Howard <i>et al.</i> 2000
Rothwell, West Yorkshire			
Auckland Castle, Bishop	10.1	AD 1425–1698	Arnold and Howard 2013
Auckland, County			
Durham			
Nun Appleton, Tadcaster,	9.6	AD 1478–1657	Arnold <i>et al.</i> 2008
West Yorkshire			

Table 2:	Details	of sampled	l tree-rings	and radio	ocarbon	results fro	m LED-A22
----------	---------	------------	--------------	-----------	---------	-------------	-----------

Laboratory code	Material	^{1₄} C age (BP)	δ ¹³ C _{IRMS} (‰)	Tree-ring date (AD)
SUERC-68046	<i>Quercus</i> sp. heartwood, ring 1	306±29	-23.5±0.2	1520
TUBITAK-127	<i>Quercus</i> sp. heartwood, ring 3	321±29	-24.1±1.0	1522
OxA-34316	Quercus sp. heartwood, ring 4	311±33	-23.4±0.2	4500
OxA-34317	<i>Quercus</i> sp. heartwood, ring 4	357±35	-23.8±0.2	1523
Ring 4	333±25BP, T′=0.9; −23.6±0.14 ‰, T′	=2.0; T′(5%)=3.8, v=1	
BRAMS-1230	<i>Quercus</i> sp. heartwood, ring 7	315±26	-21.9±0.2	1526
SUERC-68056	<i>Quercus</i> sp. heartwood, ring 9	324±29	-24.0±0.2	4500
OxA-34321	<i>Quercus</i> sp. heartwood, ring 9	359±33	-24.9±0.2	1528
Ring 9	339±22BP; T'=0.6; −24.5±0.14 ‰, T'	=10.1; T'(59	%)=3.8, ν=1	
TUBITAK-128	<i>Quercus</i> sp. heartwood, ring 13	276±27	- 23.5±1.0	1532
OxA-34275	<i>Quercus</i> sp. heartwood, ring 14	262±25	-24.9±0.2	1533
BRAMS-1231	<i>Quercus</i> sp. heartwood, ring 15	319±26	-22.9±0.2	1534
SUERC-68044	<i>Quercus</i> sp. heartwood, ring 19	282±29	-24.1±0.2	1538
TUBITAK-129	<i>Quercus</i> sp. heartwood, ring 21	352±50	- 24.5±1.7	1540
BRAMS-1232	<i>Quercus</i> sp. heartwood, ring 22	294±26	- 25.0± 0.2	1541
SUERC-68054	<i>Quercus</i> sp. heartwood, ring 24	315±29	- 24.2±0.2	1510
OxA-34323	<i>Quercus</i> sp. heartwood, ring 24	380±33	-24.9±0.2	1545
Ring 24	344±22BP; T′=2.2; −24.6±0.14 ‰, T′	=6.1; T′(5%)=3.8, v=1	
SUERC 68053	<i>Quercus</i> sp. heartwood, ring 29	322±29	-25.2±0.2	4540
OxA-34281	<i>Quercus</i> sp. heartwood, ring 29	255±25	-26.2±0.2	1548
Ring 29	284±19BP; T'=3.1; −25.7±0.14 ‰, T'	=12.5; T′(59	‰)=3.8, v=1	
BRAMS-1233	Quercus sp. heartwood, ring 31	302±26	-26.1±0.2	1550

OxA-34277	Quercus sp. heartwood, ring 34	328±24	-24.3± 0.2	1553
TUBITAK-130	<i>Quercus</i> sp. heartwood, ring 36	307±27	-25.5±0.8	1555
SUERC-68051	<i>Quercus</i> sp. heartwood, ring 39	335±29	-24.6±0.2	1558
OxA-34319	Quercus sp. heartwood, ring 44	334±32	-25.4±0.2	1563
TUBITAK-131	Quercus sp. heartwood, ring 45	310±27	-25.3±0.8	1564
BRAMS-1234	Quercus sp. heartwood, ring 47	353±26	-22.4±0.2	1566
SUERC-68034	<i>Quercus</i> sp. heartwood, ring 49	306±29	-25.1±0.2	1568
OxA-34457	Quercus sp. heartwood, ring 54	364±27	-25.9±0.2	1573
BRAMS-1235	Quercus sp. heartwood, ring 56	382±26	-28.4±0.2	1575
TUBITAK-132	<i>Quercus</i> sp. heartwood, ring 58	290±48	-25.1±1.5	1577
SUERC-68035	Quercus sp. heartwood, ring 59	346±29	-25.8±0.2	1578
TUBITAK-133	Quercus sp. heartwood, ring 62	286±27	-26.7±0.8	1581
OxA-34279	Quercus sp. heartwood, ring 64	331±26	-25.9±0.2	1583
BRAMS-1236	Quercus sp. heartwood, ring 67	336±26	-25.8±0.2	1586
SUERC-68052	<i>Quercus</i> sp. heartwood, ring 69	310±29	-25.9±0.2	1588
OxA-34282	Quercus sp. heartwood, ring 74	297±26	-27.1±0.2	1593
TUBITAK-134	Quercus sp. heartwood, ring 77	396±29	-26.3±0.9	1596
SUERC-68036	Quercus sp. heartwood, ring 79	334±29	-25.6±0.2	1598
OxA-34283	Quercus sp. heartwood, ring 84	336±26	-26.4±0.2	1603
TUBITAK-135	<i>Quercus</i> sp. heartwood, ring 85	336±28	-24.9±0.9	1604
BRAMS-1238	Quercus sp. heartwood, ring 86	365±26	-24.4± 0.2	1605
SUERC-68050	<i>Quercus</i> sp. heartwood, ring 89	347±29	-25.2± 0.2	1608
OxA-34278	Quercus sp. heartwood, ring 94	341±25	-24.8±0.2	1613
TUBITAK-136	<i>Quercus</i> sp. heartwood, ring 96	416±42	-26.3±0.9	1615
BRAMS-1239	<i>Quercus</i> sp. heartwood, ring 98	370±26	-23.1±0.2	1617
SUERC-68042	Quercus sp. heartwood, ring 99	364±29	-23.7±0.2	1618
OxA-34318	Quercus sp. heartwood, ring 104	374±31	-24.1± 0.2	1623
TUBITAK-137	Quercus sp. heartwood, ring 107	351±27	-23.0±0.7	1626
SUERC-68055	Quercus sp. heartwood, ring 109	313±29	-23.8±0.2	4000
OxA-34322	Quercus sp. heartwood, ring 109	351±34	-24.1±0.2	1628
Ring 109	284±19BP; T'=3.1; -24.0±0.14 ‰, T'	=1.1; T'(5%)=3.8, v=1	
BRAMS-1240	Quercus sp. heartwood, ring 110	348±26	-23.3±0.2	1629
OxA-34280	Quercus sp. heartwood, ring 114	321±26	-25.0±0.2	1633
TUBITAK-138	Quercus sp. heartwood, ring 115	312±26	-22.7±0.7	1634
SUERC-68043	Quercus sp. heartwood, ring 119	320±29	-22.9±0.2	1638
BRAMS-1241	Quercus sp. heartwood, ring 120	291±26	-22.7±0.2	1639
TUBITAK-139	Quercus sp. heartwood, ring 122	271±28	-24.0±0.9	1641
OxA-34284	Quercus sp. heartwood, ring 124	275±25	-25.7±0.2	1643
BRAMS-1242	Quercus sp. heartwood, ring 125	263±26	-25.7±0.2	1644
SUERC-68041	Quercus sp. heartwood, ring 129	214±29	-23.7±0.2	1648
L		1	1	

BRAMS-1243	<i>Quercus</i> sp. heartwood, ring 131	301±26	-21.7±0.2	1650
OxA-34320	Quercus sp. heartwood, ring 134	292±32	-24.3±0.2	1653
TUBITAK-140	<i>Quercus</i> sp. heartwood, ring 135	212±29	-23.8±1.1	1654
SUERC-68045	<i>Quercus</i> sp. heartwood, ring 139	267±29	-23.4±0.2	1658
OxA-34276	Quercus sp. heartwood, ring 144	214±24	-24.2±0.2	1663
TUBITAK-141	Quercus sp. heartwood, ring 146	248±55	-26.3±2.3	1665
BRAMS-1244	<i>Quercus</i> sp. heartwood, ring 148	214±26	-21.9±0.2	1667
SUERC-68040	<i>Quercus</i> sp. heartwood, ring 149	222±29	-24.0±0.2	1668

Table 3: Summary of the estimated dates for the final ring of LED-A22 (dated by dendrochronology to AD 1668) from wiggle-matching 29–30-year blocks (An: 26.7, n: 7 for all).

Dinge	Acomb	Highest Posterior De	nsity interval (cal AD)
Killys	ACOILD	fighest Fosterior De	05% probability
1 00	40.0	66% probability	95% probability
1–29	48.0	1000-1070 (59%) Of	1055-1709 (89%) Of
4 0 4		1083-1088 (9%)	1760-1770 (6%)
4-34	51.5	1659-1675	1655–1693 (93%) OF
0.00	50.0	1660 1675 (609/) or	1700-1706 (2%)
9–39	50.8	1000-1075 (00%) OF	1655—1706
4 4 4 4	00.0	1063-1067 (6%)	4057 4004
14-44	03.Z	1004-1070	1657-1694
19-49	83.3	1655-1689	1645-1699
24-54	12.3	1650—1657 (10%) Or	1643–1706
00 50	00.0	1005-1094 (58%)	
29-59	90.6	1653—1678 (58%) Or	1647—1700
24 04	400.7	1080-1092 (10%)	4500 4000 (449()
34-64	102.7	1655-1704	1596–1620 (11%) or
	05.4	4500 4045 (000())	1634—1723 (84%)
39-69	95.1	1599–1615 (20%) or	4500 4000 (070()
		1651—1682 (38%) OF	1593—1636 (27%) OF
		1697—1704 (8%) or	1642-1725 (68%)
44 74	05.7	1/19—1/21 (2%) 1509 1617 (10%) or	1505 1621 (119()
44—74	95.7	1598—1617 (42%) Or	1595—1631 (44%) OF
		7002-7070 (9%) OF	1640—1677 (26%) OF
40.70	04 5	1709–1722 (17%) 4500-4647 (200() -=	1695-1724 (25%)
49-79	91.5	1599–1617 (39%) or	1592–1675 (80%) or
54.04	00.4	1649-1670 (29%)	1694-1715 (15%)
54-84	98.4	1597—1618 (43%) Or	1576–1675 (81%) OF
		1051-1008 (10%) OF	1692-1711 (14%)
50.00	102.6	1090-1707 (9%)	1576 1596 (29/) or
59-69	103.0	1605-1613 (6%) 01	1570 - 1500 (2%) 01
		1027-1008 (0078)	1696 - 1702 (1%)
64 04	110 /	1621 1650	1690-1702 (178)
60.00	110.4	1021-1009	1607 1672
74 104	110.0	1623-1654	1562 1560 (1%) or
74-104	114.1	1037-1004	$1611_{1672} (94\%)$
70 100	117.0	1555 1567 (18%) or	1511 - 1072 (3478)
79-109	117.0	1637_1667 (50%)	1617_1675 (20%) 01
8/ 11/	115 5	1551_1568 (31%) or	$1538_{1573}(37\%)$ or
04-114	115.5	1640_1668 (37%)	1616_1677 (58%)
80_110	117 5	1548_1568 (35%) or	1529 - 1574 (46%) or
03-113	117.5	1650-1670 (33%)	1609 - 1635 (11%) or
		1000-1010 (0070)	1646_1675 (38%)
01_121	12/ 8	1553-1564 (36%) or	1546 - 1574 (50%) or
34-124	124.0	1658_1670 (32%)	1640-1677 (45%)
00_120	027	1664_1674	$1555_{1557}(1\%)$ or
33-123	92.1	1004-1074	1657_1670 (0/%)
10/_13/	78.3	1662_1674	1552_1563 (6%) or
104-134	10.5	1002-10/4	1656_1679 (80%)
100_120	07.7	1664_1673	1656_1678
11/ 1//	10/ 1	1665_1673	1650_1675
110 140	104.1	1667 1671	1650 1675
119-149	90.0	1004-1071	1009-1070

Figure 1: Offsets between radiocarbon ages on single tree-ring from this study and measurements on single tree-rings of the same calendar date reported by Stuiver (1993, corrected as described by Stuiver and Becker 1993). Colour figure



Figure 2: Probability distributions of dates from LED-A22. Each distribution represents the relative probability that an event occurs at a particular time. For each of the dates two distributions have been plotted: one in outline, which is the result of simple radiocarbon calibration, and a solid one, based on the wiggle-match sequence. Distributions other than those relating to particular samples, correspond to aspects of the model. For example, the distribution '*SUERC-68040*' is the estimated date of the final ring of this core. The large square brackets down the left-hand side of the diagram along with the CQL2 keywords (Bronk Ramsey 2009) define the model exactly.

R_Date SUERC-68040 [A:118]						-
R_Date BRAMS-1244 [A:145]	3 <u></u>		~		-	
R Date TUBITAK-141 [A:170]						-
Gap 2 R Date OxA-34276 [A:135]	1				-	<u> </u>
Gap 5 R Date SUERC 68045 (A:128)						
Gap 4 TUDITAK 140 (A:01)						
Gap1						
R_Date OxA-34320 [A:60] Gap 3			10000			
R_Date BRAMS-1243 [A:37]				- 		
R_Date SUERC-68041 [A:41]					-	-
R Date BRAMS-1242 [A:152]	S			-		-
R Date OxA-34284 [A:181]	2 <u>-</u>					-
R Date TUBITAK-1241 IA:1601						
Gap 2 R Date BRAMS-1241 (A 153)						L
Gap 1 P Date SUEPC 68042 [A:107]						
Gap 4 TUDITAK 428 (A.427)			-			
Gap 1						
R_Date OxA-34280 [A:122] Gap 5						
R_Date BRAMS-1240 [A:106]						
R Combine Ring_109 [A:125]						
R_Date TUBITAK-137 [A:103]						
Gap 3 R Date OxA-34318 [A:64]				-		
Gap 5 R Date SUERC-68042 [A:121]						
Gap 1 Gap 1 D D D D A A S 1220 (A 119)						
Gap 2 TUDITAK 400 (4 70)			-			
R_Date TUBITAK-136 [A:70]						
R_Date OxA-34278 [A:99]						
R_Date SUERC-68050 [A:100]			·			
R Date BRAMS-1238 [A:136]						
R_Date TUBITAK-135 [A:95]						-
Gap 1 R Date OxA-34283 [A:103]						
R Date SUFRC-68036 (A.116)			~~~			
Gap 2 Date TUDITAK 124 (A:20)			~			
Gap 3 C A GAODO GA GOT						
R_Date OxA-34282 [A:60] Gap 5	2 DA 0					
R [*] Date SUERC-68052 [A:101]						
R Date BRAMS-1236 [A:122]			~ ~			
R_Date OxA-34279 [A:122]			~~			-
R_Date TUBITAK-133 [A:41]						-
Gap 3 R Date SUERC-68035 [A:113]			~~~			-
Gap 1 R Date TUBITAK-132 (A 104)						
Gap 2 P Date BRAMS 1235 (A:23)			~			
Gap 2						
Gap 5						
R_Date SUERC-68034 [A:113]						
R Date BRAMS-1234 [A:77]						
R Date TUBITAK-131 [A:110]						-
Gap R_Date OxA-34319 [A:117]			~~~			<u> </u>
R Date SUERC-68051 [A 104]			~~~			
Gap 3 R Date TUBITAK-130 (A.132)			\sim			
Gap 2			~			
Gap 3 DDAMO 1000			~			
Gap 2						
R_Combine Ring_29 [A:86]			~			<u> </u>
R_Combine Ring_24 [A:39]			~			
R_Date BRAMS-1232 [A:144]			\sim			
Gap 1 R Date TUBITAK-129 (A-77)						<u> </u>
Gap 2 Data SUERC 68044 [A:143]						
Gap 4						
Gap]			~			
Gap 1			~			
R_Date TUBITAK-128 [A:115]						
R_Combine Ring_9 [A:81]			~~			
R_Date BRAMS-1230 [A:121]		-	~~~			-
R_Combine Ring 4 [A:122]			\sim			
R Date TUBITAK-127 IA 1081			\sim			
Gap 2 R Date SUERC_68046 (A-70)		-				
Sequence Lodoton Hell (Acomb=70.0: A:	-0.1					
_ocquence Leusion Hall [Acomb-70.2; Al	-5.1, 11-00]	1	1	1		1

Posterior Density Estimate (cal AD)

Figure 3: Probability distributions of dates from LED-A22 (a) BRAMS-, and (b) OxA-. The format is identical to that of Figure 2. The large square brackets down the left-hand side of the diagram along with the CQL2 keywords define the model exactly



Posterior Density Estimate (cal AD)

Figure 4: Probability distributions of dates from LED-A22 (a) SUERC, and (b) TUBITAK. The format is identical to that of Figure 2. The large square brackets down the left-hand side of the diagram along with the CQL2 keywords define the model exactly







Posterior Density Estimate (cal AD)



1643.5 1653.5 1663.5

Posterior Density Estimate (cal AD)

1683.5 1693.5 1703.5 1713.5 1753.5 -1763.5 -

1723.5 1733.5 1743.5 1783.5

1773.5

Figure 5: Posterior density estimates for the final ring of LED-A22, derived from the short wiggle-matches based on sequences of 29–30 tree-rings.

0.04

0.02

1503.5 1513.5 1523.5 1533.5

1553.5 -1563.5 - 1573.5 1583.5 1593.5

1543.5

1603.5

1613.5 1623.5 1633.5 **Figure 6:** Radiocarbon ages of known-age tree-ring rings AD 1150–1668: single years (OxA-, SUERC-, GrA-, BRAMS-, TUBITAK-; this study, Bayliss *et al.* 2017, Tyers *et al.* 2009), decadal samples (Wk; Hogg *et al.* 2002), single-year and decadal samples (QL; Stuiver 1993 as corrected by Stuiver and Becker 1993; Stuiver *et al.* 1998), decadal and bi-decadal samples (UB; Hogg *et al.* 2002; Pearson *et al.* 1986), decadal and 23-year and 24-year samples (GrN: van der Plicht *et al.* 1995). Colour figure



Figure 7: Radiocarbon ages of known-age tree-ring rings AD 1510–1670: single years (OxA-, SUERC-, GrA-, BRAMS-, TUBITAK-; this study, Bayliss *et al.* 2017, Tyers *et al.* 2009), decadal samples (Wk; Hogg *et al.* 2002), single-year samples (QL; Stuiver 1993 as corrected by Stuiver and Becker 1993; Stuiver *et al.* 1998), decadal and bi-decadal samples (UB; Hogg *et al.* 2002; Pearson *et al.* 1986), decadal and 23-year and 24-year samples (GrN: van der Plicht *et al.* 1995). Colour figure

