

Original research

Dysregulation of prostaglandins, leukotrienes and lipoxin A₄ in bronchiectasis

Pallavi Bedi,¹ Kerstin Ziegler,² Phil D Whitfield,³ Donald Davidson,¹ Adriano Giorgio Rossi,¹ Adam T Hill⁴

excessive neutrophilic inflammation. Lipid mediators such

as prostaglandins and leukotrienes have crucial roles in

the inflammatory response. Further characterisation of

inflammatory and proinflammatory lipid mediators could

lipids obtained from serum and airways in patients with

these lipids and understanding the interplay of anti-

lead to the development of novel anti-inflammatory

Aim The aim of our study was to characterise the

Methods Six healthy volunteers, 10 patients with

mild bronchiectasis, 15 with moderate bronchiectasis

participants had 60 mL of blood taken and underwent a

bronchoscopy while in the stable state. Lipidomics was

done on serum and bronchoalveolar lavage fluid (BALF).

and 9 with severe bronchiectasis were recruited. All

Results In the stable state, in serum there were

15-hydroxyeicosatetranoic acid (15-HETE) and

significantly higher levels of prostaglandin E₂ (PGE₂),

leukotriene B₄ (LTB₄) in patients with moderate-severe

disease compared with healthy volunteers. There was

a significantly lower level of lipoxin A, (LXA,) in severe

ABSTRACT Introduction Bronchiectasis is characterised by

therapies for bronchiectasis.

bronchiectasis in the stable state.

► Additional supplemental material is published online only. To view, please visit the journal online (http://dx.doi. org/10.1136/thoraxjnl-2020-216475).

¹MRC Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK ²Department of Lipidomics, University of the Highlands and Islands, Inverness, UK ³Department of Lipidomics, University of Glasgow, Glasgow, UK ⁴Respiratory Medicine, MRC

Respiratory Medicine, MRC Centre for Inflammation Research, University of Edinburgh, Edinburgh, UK

Correspondence to

Dr Pallavi Bedi, University of Edinburgh MRC Centre for Inflammation Research, Edinburgh EH16 4TJ, UK; drpallavibedi@gmail.com

Received 28 October 2020 Accepted 21 September 2021 Published Online First 17 November 2021



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY. Published by BMJ.

To cite: Bedi P, Ziegler K, Whitfield PD, *et al. Thorax* 2022;**77**:960–967.



In BALF, there were significantly higher levels of PGE_2 , 5-HETE, 15-HETE, 9-hydroxyoctadecadienoic acid and LTB_4 in moderate–severe patients compared with healthy volunteers.

bronchiectasis.

In the stable state, there was a negative correlation of PGE_2 and LTB_4 with % predicted forced expiratory volume in 1 s and a positive correlation with antibiotic courses. LXA_4 improved blood and airway neutrophil phagocytosis and bacterial killing in patients with bronchiectasis. Additionally LXA_4 reduced neutrophil activation and degranulation.

Conclusion There is a dysregulation of lipid mediators in bronchiectasis with excess proinflammatory lipids. LXA_4 improves the function of reprogrammed neutrophils. The therapeutic efficacy of LXA_4 in bronchiectasis warrants further studies.

INTRODUCTION

Excessive inflammation is widely accepted to be a unifying component in many chronic diseases, including bronchiectasis, vascular diseases, metabolic syndrome and neurological diseases, and thus is a public health concern. Understanding endogenous control points within the inflammatory response could potentially provide us with new perspectives on disease pathogenesis and treatment

KEY MESSAGES

WHAT IS THE KEY QUESTION?

⇒ Is there a dysregulation of lipids in bronchiectasis that leads to unremitting and chronic inflammation?

WHAT IS THE BOTTOM LINE?

 \Rightarrow Ex vivo, lipoxin A₄ can improve the function of reprogrammed blood and airway neutrophils in bronchiectasis.

WHY READ ON?

⇒ This is the first study assessing the role of lipids in bronchiectasis in depth. Here we have demonstrated that dysregulation of the proinflammatory and anti-inflammatory lipids in bronchiectasis contribute to failure of resolution of inflammation. Targeting the lipid pathways to initiate the resolution process in bronchiectasis may lead to development of novel nonantibiotic therapy in the stable state.

approaches. Break in the barrier, trauma and microbial invasion encourage the host to clear microbial pathogens, remodel and regenerate tissue. The acute inflammatory response is protective and usually self-limiting. Oxidative lipid products including, notably, the products of unsaturated lipids are increasingly being recognised as important contributors to chronic inflammatory diseases.^{1 2} Eicosanoids^{3 4} produced from the n-6 polyunsaturated fatty acid, arachidonic acid (20:4), as well as many cytokines and chemokines,⁵⁶ have crucial roles in the initial response. Interactions among prostaglandins, leukotrienes and proinflammatory cytokines amplify inflammation, the signs and effects of which can be reduced by pharmacological inhibition and receptor antagonists.³⁻⁵ However, given that excessive inflammation contributes to several widely occurring diseases, improvements are required in treatment and in our understanding of the mechanisms involved. Eicosanoids also include lipoxins that possess potent anti-inflammatory properties.⁷ Further, n-3 fatty acids (eicosapentaenoic acid 20:5, and docosahexaenoic acid, 22:6) are precursors for proresolving mediators such as resolvins and maresins that limit the duration and magnitude of inflammatory responses.⁸ Recent advances have facilitated more detailed profiling of lipid mediators in serum and at the sites of inflammation.9-11

Characterisation of these lipids could thereby lead to the development of novel anti-inflammatory therapies for chronic inflammatory conditions.

Bronchiectasis is characterised by recurrent cough, daily sputum production and recurrent chest infections. There is excessive neutrophilic inflammation, but the driver for this unremitting inflammation is not known. We hypothesise that there is a failure of resolution of inflammation in bronchiectasis. Recently, we have been investigating the lipid pathway to establish if there is a dysregulation of the lipids in bronchiectasis contributing to the chronic inflammation. While this could be contributing to the persistent chronic inflammatory state in bronchiectasis, there is currently no data in the literature to indicate the interplay of lipid mediators in bronchiectasis.

The aim of our study was to characterise the lipids obtained from blood and airway samples in patients with bronchiectasis in the stable state and to assess the efficacy of lipoxin A_4 (LXA₄) on neutrophil function.

METHODS

Six healthy volunteers (partners of the patients recruited in the study with no background medical conditions and not currently on medication), 10 patients with mild bronchiectasis, 15 with moderate and 9 with severe bronchiectasis were recruited. All participants had 60 mL of blood taken and underwent a bronchoscopy. Two segments of the lungs were washed out in patients with bronchiectasis, an area affected by bronchiectasis and an area unaffected by bronchiectasis, predetermined by CT scan of chest. This led to patients acting as their own internal control. Lipidomics was done on blood and bronchoalveolar samples were obtained.

Bronchiectasis severity

The severity of bronchiectasis was calculated using the Bronchiectasis Severity Index (BSI).¹² The BSI is a risk stratification tool for morbidity and mortality in bronchiectasis. The minimum score is 0 and the maximum score is 26. A score between 0 and 4 indicates mild disease; 5–8 indicates moderate disease; and a score of ≥ 9 indicate severe disease. The BSI was calculated in all patients with bronchiectasis taking part in the study.

Inclusion criteria

Inclusion criteria include idiopathic or postinfective bronchiectasis, age >18 years and no infective exacerbation of bronchiectasis for at least 4 weeks prior to giving serum/bronchoscopy.

Exclusion criteria

Patients on statin, aspirin, inhaled corticosteroids and long-term macrolides were excluded.

Bronchoscopy

All participants underwent a bronchoscopy. Participants were sedated with midazolam±fentanyl. Bronchoalveolar lavage (BAL) and brushings were obtained. For patients with bronchiectasis, BAL was done in an area *affected* by bronchiectasis and in an area *unaffected* by bronchiectasis as identified on CT scans done prior to bronchoscopy.

Liquid chromatography-tandem mass spectrometry (LC-MS/ MS) analysis of eicosanoids

Quantification of eicosanoids and related lipid mediators in patient bronchoalveolar lavage fluid (BALF) and serum samples

was performed using LC-MS/MS¹³ (see online supplemental file, online supplemental figure 1 and online supplemental table T1).

LXA₄ detection by ELISA

Serum LXA₄ was measured as per manufacturer's instruction (Neogen).

 LXA_4 was the lipid of choice for all the experiments outlined in this article as pilot data (data not shown in this paper) using sputum and blood from bronchiectasis exacerbations showed that LXA_4 was the only lipid that improved after treatment with antibiotics.

Isolation of blood and airway neutrophils

Freshly drawn blood was collected into 3.8% sodium citrate, and granulocytes were subsequently isolated by dextran sedimentation and discontinuous Percoll gradient.¹⁴ Sputum and BAL were washed and treated with sputolysin, and airway neutrophils were isolated. Anti-CD16 antibodies (Abcam) were used to identify neutrophils by flow cytometry.

LXA₄ function on reprogrammed bronchiectasis neutrophils

We assessed phagocytosis and killing of green fluorescent protein (GFP)–*Pseudomonas* O1 (PAO1), spontaneous neutrophil apoptosis, neutrophil activation (CD62L/CD11b expression) and neutrophil degranulation (myeloperoxidase release). Further information on the specific experiments is provided in the online supplemental file.

Table 1 Baseline demographics of the study population				
	Patients with I N=34	Healthy		
Parameters	Mild n=10	Moderate n=15	Severe n=9	volunteers N=6
Age (years)	55 (4.1)	65 (2.2)	64 (2.2)	52 (6.8)
Biological sex (% female)	40	60	22	80
Aetiology				
Idiopathic	10 (100%)	12 (80%)	6 (67%)	
Postinfective		3 (20%)	3 (33%)	
Total WCC (×10 ⁹ /L)	6 (0.5)	6.3 (0.4)	9.3 (1.1)	5.9 (0.5)
Neutrophils	3.3 (0.3)	4.1 (0.3)	6.6 (1.1)	3.5 (0.3)
Eosinophils	0.2 (0.04)	0.3 (0.07)	0.2 (0.06)	0.2 (0.06)
Monocytes	0.5 (0.03)	0.6 (0.05)	0.7 (1)	0.5 (0.05)
ESR (mm/hour)	6.7 (1.8)	13.2 (2.9)	19.6 (6.8)	4.8 (1)
CRP (mg/L)	2.8 (0.5)	4 (1)	16 (7.4)	3.2 (1.1)
FEV ₁ % predicted (L)	95 (5.5)	82 (4)	55 (6.5)	_
FVC % predicted (L)	111 (6)	97 (4)	84 (6)	-
TLCO % predicted (SI)	94% (4.9)	82% (4.2.)	74% (7.8)	-
KCO % predicted (SI)	106% (4.5)	97% (3.7)	100% (7.2)	_
Chronic colonisation	5 (50%)	12 (86%)	8 (89%)	-
Exacerbations in the last year	0.4 (0.3)	2.4 (0.5)	4.2 (0.9)	-
Hospital admissions in the last year	0	0.05 (0.05)	0.7 (0.2)	-

Data presented as mean (±SE of mean).

CRP, C reactive protein; ESR, erythrocyte sedimentation rate; FEV,, forced expiratory volume in 1 s; FVC, forced vital capacity; KCO, transfer coefficient corrected for alveolar volume; TLCO, transfer factor for the lung for carbon monoxide; WCC, white cell count.



Figure 1 Significantly higher levels of $PGE_{2^{\prime}}$ 15-HETE and LTB_4 detected in patients with moderate–severe disease compared with healthy volunteers; p=0.03, p=0.03 and p=0.02, respectively. higher levels of 5-HETE and 9-HODE detected in patients with more moderate–severe disease compared with healthy volunteers, but not statistically significant. Lipidomics were obtained by mass spectrometry and liquid chromatography. Pooled data presented as mean±SEM. One-way analysis of variance used for comparisons. Healthy=6 volunteers, mild=9 patients, moderate–severe=15 patients. 5-HETE, 5-hydroxyeicosatetranoic acid; 15-HETE, 15-hydroxyeicosatetranoic acid; 9-HODE, 9-hydroxyoctadecadienoic acid; $LTB_{4^{\prime}}$ leukotriene $B_{4^{\prime}}$; PGE₂, prostaglandin E₂.



Figure 2 Significantly lower level of LXA₄ is severe bronchiectasis. Healthy, n=6; mild, n=10; moderate, n=15; severe, n=9. Pooled data represented as mean \pm SEM. LXA₄, lipoxin A₄.

Statistical analysis

Flow cytometry analysis was performed using FlowJo V.10.0.4 (Tree Star, Ashland, Oregon, USA). Results are presented as mean \pm SEM. Paired and unpaired t-tests were used to compare the two groups, where applicable. Data were analysed by one-way analysis of variance (ANOVA) with Bonferonni's multiple comparison post hoc test (GraphPad Prism V.6; GraphPad Software, La Jolla, California, USA), when three groups were involved. A repeated measures ANOVA was used where samples from the same participant receive multiple treatments; significance was accepted with p values: *p<0.05.

RESULTS

Baseline demographics of the participants are shown in table 1.

Serum lipidomics

LC-MS/MS was done on all serum samples obtained, where available. Only data that passed quality control were used, and thus we have a smaller sample size. The demographics are shown in online supplemental table 1. Samples were divided into mild⁹ and moderate–severe¹⁵ bronchiectasis groups (moderate and severe groups were combined as lipids were not detected in all



Figure 3 Significantly higher levels of PGE_2 , 5-HETE, 15-HETE, 9-HODE and LTB_4 detected in patients with moderate–severe disease compared with healthy volunteers. Lipidomics obtained by mass spectrometry and liquid chromatography. pooled data presented as mean±SEM. One-way analysis of variance used for comparisons. Healthy=6 volunteers, mild= 9 patients, moderate–severe=15 patients. 5-HETE, 5-hydroxyeicosatetranoic acid; 15-HETE, 15-hydroxyeicosatetranoic acid; 9-HODE, 9-hydroxyoctadecadienoic acid; LTB_4 , leukotriene B_4 ; PGE₂, prostaglandin E_2 .

samples by lipidomics, and hence data were combined to obtain meaningful interpretation. Baseline demographics of mild⁹ and moderate–severe¹⁵ groups are provided in online supplemental table T2.

The main lipids assessed were LXA₄, resolvins, maresins, prostaglandin E₂ (PGE₂), 5-hydroxyeicosatetranoic acid (5-HETE), 15-hydroxyeicosatetranoic acid (15-HETE), leukotriene B₄ (LTB₄) and 9-hydroxyoctadecadienoic acid (9-HODE). 15-HETE is a precursor of LXA₄ and is a proinflammatory mediator. LTB₄ is a proinflammatory cytokine and 9-HODE is a proinflammatory metabolite produced from arachidonic acid.

There were significantly higher levels of PGE₂, 15-HETE and LTB₄ in patients with moderate–severe bronchiectasis compared with healthy controls (p=0.03, p=0.03 and p=0.02, respectively). Although there was a trend towards higher 5-HETE and 9-HODE levels in moderate–severe bronchiectasis compared with healthy volunteers, this failed to reach statistical significance (p=0.3 and 0.2, respectively). Although there was a trend towards higher PGE₂, 15-HETE, LTB₄, 5-HETE and 9-HODE levels between the mild bronchiectasis and healthy volunteers, this was not statistically significant (figure 1). LXA₄, resolvins and maresins were not detected in any of the samples by LC-MS/MS.

Blood LXA₄ measured by ELISA

 LXA_4 was detected by ELISA. Using one-way ANOVA, we found that there was a significantly lower level of LXA_4 in severe bronchiectasis, p=0.04 (figure 2), as measured by ELISA.

BALF lipidomics

There were significantly higher levels of PGE₂, 5-HETE, 15-HETE, 9-HODE and LTB₄ in patients with moderate–severe bronchiectasis compared with healthy volunteers (p < 0.0001, p=0.004, p=0.005, p=0.04 and p < 0.0001, respectively) (figure 3). Although there was a trend, there was no statistically significant difference in the PGE₂, 15-HETE, LTB₄, 5-HETE and 9-HODE levels between mild bronchiectasis and healthy volunteers. LXA₄, resolvins and maresins were not detected in any of the samples by LC-MS/MS. PGE₂ was not detectable in BALF from healthy individuals.

Correlation of blood $\mathrm{PGE}_{\mathrm{2}}$ and $\mathrm{LTB}_{\mathrm{4}}$ to markers of disease severity

There was a positive correlation of PGE₂ with antibiotic courses (r=0.78, 95% CI 0.61 to 0.88, p<0.0001) and a negative correlation with % predicted forced expiratory volume in 1 s (FEV₁) (r=-0.46, 95% CI -0.15 to -0.69, p=0.004). Similarly, there was a positive correlation of LTB4 with antibiotic



Figure 4 Using Pearson two-tailed correlation coefficient, we found that there was a correlation between serum PGE₂ and antibiotic courses received in the preceding year and % predicted FEV₁ (A,B). Also, there was a correlation between serum LTB₄ and antibiotic courses received in the preceding year and % predicted FEV₁ (C,D). FEV₁, forced expiratory volume in 1 s; LTB₄, leukotriene B₄; PGE₂, prostaglandin E₂.

courses (r=0.76, 95% CI 0.56 to 0.87, p<0.0001) and a negative correlation with % predicted FEV₁ (r=-0.58, 95% CI -0.3 to -0.76, p=0.0003; figure 4). There was no correlation of PGE₂ or LTB₄ to other parameters of the disease severity on the BSI. There was no correlation of 15-HETE, 5-HETE and 9-HODE to markers of disease severity.

Effect of biological sex on PGE,

There was biological sex imbalance in our cohort, and to investigate this further, there was a subanalysis of blood PGE₂. The blood PGE₂ was 1.1 (±0.3) for male healthy volunteers and 1.3 (±0.5) for female healthy volunteers (p=0.2), 4.2 (±1.4) for male mild bronchiectasis and 3.9 (±0.1) for female mild bronchiectasis (p=0.8), 2.9 (±0.6) for male moderate–severe bronchiectasis, and 4.2 (±0.7) for female moderate–severe bronchiectasis (p=0.2). This is a small study and so detailed further subanalysis was not carried out.

Effect of LXA₄ on blood neutrophil function

As we detected significantly lower levels of LXA₄ in blood of patients with severe bronchiectasis, we investigated the effect of LXA₄ on reprogrammed neutrophil function.¹⁵ With the complexity of the experiments, the authors studied healthy controls and patients with mild and severe bronchiectasis only. Blood neutrophils from healthy volunteers (n=6) and patients with mild (n=10) and severe bronchiectasis (n=9) were pretreated ex vivo with varying concentrations of LXA₄ for 30 min and then coincubated with GFP–PAO1. Phagocytosis of bacteria (GFP–PAO1) was assessed after 15 min and killing after 24 hours. Total phagocytosis was calculated by assessing neutrophils positive for GFP. Data were analysed by gating the overall phagocytosis in healthy controls and in patients with mild and severe bronchiectasis. Next, we gated the neutrophils that had taken

up much higher number of bacteria, as indicated by the mean fluorescence (fluorescence intensity, FLI). This was done at 50% of the total phagocytosis and we called this high MFLI phagocytosis. In healthy controls, mild and severe bronchiectasis, LXA₄ was able to significantly improve phagocytosis (comparing only the high MFLI and not total phagocytosis) (figure 5A) and killing of GFP–PAO1 in a concentration-dependent manner with statistical significance achieved at 100 nM LXA₄ (figure 5B).

Effect of LXA₄ on airway neutrophil function

Airway neutrophils were isolated from BALF from patients with bronchiectasis (no airway neutrophils could be isolated from healthy volunteers), and phagocytosis and killing assays were performed using GFP–PAO1.

Neutrophils collected from affected lung segments demonstrated statistically significant improvement in phagocytosis (neutrophil gating done as in previous experiment) and killing after treatment with 100 nM LXA₄ in mild, moderate and severe bronchiectases (figure 6A,B). The neutrophils collected from unaffected lung segments displayed significant improvement in bacterial killing after treatment with 100 nM LXA₄ in patients with severe bronchiectasis only (p=0.02).

Effect of LXA₄ on stable disease-state peripheral blood neutrophil spontaneous apoptosis, surface expression of CD11b and CD62L, and myeloperoxidase release

For this experiment, blood neutrophils from healthy volunteers and patients with stable-state bronchiectasis with either mild and moderate–severe disease were pretreated with LXA_4 1, 10 and 100 nM and assessed for the onset of spontaneous apoptosis. LXA_4 did not significantly modulate spontaneous apoptosis or viability of neutrophils (p=0.4, p=0.5 and p=0.4, respectively),



Figure 5 (A) There was a concentration-dependent increase in phagocytosis induced by LXA_4 , in healthy volunteers, mild and severe bronchiectasis. One-way ANOVA with a Bonferroni correction for multiple comparisons, with p values representing the comparison of control to 1, 10 and 100 nM of LXA_4 . Pooled % neutrophil phagocytosis data, showing means±SEM. (B). There was dose-dependent increase in killing with LXA_4 in healthy volunteers, mild and severe bronchiectasis. One-way ANOVA with Bonferonni's correction for multiple comparisons used, with p values representing the comparison of control to 1, 10 and 100 nM of LXA_4 . Pooled % neutrophil killing data showing means±SEM. ANOVA, analysis of variance; LXA_4 , lipoxin A_4 .

in contrast to roscovitine, used as a positive control for apoptosis induction (online supplemental figure S2).

related to the % predicted FEV_1 and directly related to the number of antibiotic courses for bronchiectasis exacerbations.

To evaluate the effect of LXA_4 on fMLF-induced neutrophil activation, surface expression of CD11b and CD62L was measured. In this experiment, LXA_4 treatment induced a small but statistically significant reduction in fMLF-induced upregulation of CD11b and shedding of CD62L in a concentrationdependent manner, in healthy volunteers and in patients with mild and moderate–severe bronchiectases (online supplemental figure S3 and 4).

Additionally, LXA_4 reduced neutrophil degranulation and release of myeloperoxidase in a concentration-dependent manner from neutrophils isolated from healthy volunteers and in patients with mild and moderate–severe bronchiectasis (online supplemental figure S5).

DISCUSSION

This is the first study characterising the lipid profile in bronchiectasis blood and airways in the stable state. We established that there is a dysregulation of the lipids in serum and airways in the stable state. In serum, there was significantly higher levels of the proinflammatory metabolites PGE₂, 15-HETE and LTB₄ and significantly lower levels of the anti-inflammatory mediator LXA₄ in severe disease compared with mild disease and healthy volunteers. Although LXA₄ was not detectable in serum using LC-MS/MS, it was detected by ELISA. This can be explained by level of sensitivity of the assays used. Detection limits were in nanogram per millilitre by LC-MS and in picogram per millilitre by ELISA. In the airways, there were significantly higher levels of PGE₂, 5-HETE, 15-HETE, 9-HODE and LTB₄. There was a correlation of PGE₂ and LTB₄ with antibiotic courses and % predicted FEV₁. Higher levels of PGE₂ and LTB₄ were inversely In the stable state, LXA_4 was able to significantly improve phagocytosis and killing of GFP *Pseudomonas aeruginosa* by blood and airway neutrophils, in a concentration-dependent manner. In addition, the authors demonstrated that LXA_4 reduced fMLF-induced neutrophil activation. However, there was no effect of LXA_4 on spontaneous neutrophil apoptosis.

To the authors' best knowledge, prostaglandins have not been studied in bronchiectasis. In COPD, PGE, levels are increased in the exhaled breath and are known to correlate with airflow obstruction.¹⁶¹⁷ Additionally, studies have demonstrated that PGE₂ is a critical component in amplifying and perpetuating senescence and inflammation in COPD fibroblasts.¹⁸ PGE, enhances LTB₄-mediated polymorphonuclear leucocyte extravasation and tissue injury that is blocked by topical administration of synthetic LXA_4^{19} However, prostaglandins are also key in the temporal switch of LTB₄ to LXA₄—a term coined as 'lipid mediator class switching'.²⁰ In the stable state, the authors demonstrated that PGE, and LTB, were significantly higher and LXA₄ was significantly lower in blood in severe bronchiectasis. Serum PGE₂ and LTB₄ levels were correlated to airflow obstruction as measured by FEV₁. Higher serum PGE₂ and LTB₄ levels were also correlated to more exacerbations requiring antibiotic courses in bronchiectasis, both markers of disease severity in bronchiectasis.¹² Studies have established that disease severity in bronchiectasis predicts mortality, hospital admissions, exacerbations, quality of life, respiratory symptoms, exercise capacity and lung function decline in bronchiectasis.²¹ This dysregulation between lipid mediators that the authors have shown here would thereby lead to more inflammation even in the stable state. Although PGE, levels are known to initiate the class switching



Figure 6 (A) Bacterial phagocytosis and (B) killing. LXA₄ significantly increased phagocytosis and killing of PAO1 by airway neutrophils isolated from the lung segments affected by Bx in mild, moderate and severe diseases. In unaffected segments, LXA₄ only had an improvement in bacterial killing in patients with severe Bx only. Pooled data presented as mean±SEM. Paired t-tests used for all comparisons. Bx, bronchiectasis; LXA₄, lipoxin A₄; PAO1, *Pseudomonas* O1.

during resolution of inflammation, this was not demonstrated in our study. Almost certainly, the levels of LXA_4 detected in bronchiectasis serum are unable to counter-regulate the production of the proinflammatory LTB_4 .

In addition to PGE₂ and LTB₄, the other metabolites (5-HETE, 15-HETE and 9-HODE) detected in our study are also proinflammatory mediators, and they remain elevated even in the stable state in moderate and severe bronchiectases compared with mild disease. LTB₄ has been studied in bronchiectasis and is known to be one of the major chemotactic factors in the bronchial airways in bronchiectasis.²² Additionally, there is evidence to suggest that LTB₄ is raised during an exacerbation and reduces with antibiotic therapy.²³ Persistently elevated lipid mediators in bronchiectasis, in part, may explain why more patients with moderate–severe bronchiectasis have a higher mortality rate and more hospital admissions compared with patients with mild disease.¹²

LXA₄ was able to improve blood and airway neutrophil bacterial phagocytosis and killing, thereby enhancing bacterial clearance and potentially having long-term consequences on infection, inflammation and resolution. The impact of LXA₄ on bronchiectasis airway neutrophil function was more pronounced when neutrophils were isolated from a disease-affected lung region than an unaffected region. The mechanism underpinning this observation remains to be determined, but this raises the intriguing possibility that regional lung patterning of LXA₄ deficiency might exist, with cells collected having had divergent in vivo exposure to LXA₄ before experimental use.

 $\rm LXA_4$ was able to reduce fMLF-induced CD11b upregulation, CD62L shedding and myeloperoxidase release. These

anti-inflammatory functions of LXA₄ have previously been demonstrated in the literature.¹⁷ However, this is the first time that the effects of lipoxins have been demonstrated on a subset of reprogrammed neutrophils from patients with bronchiectasis.

Biological sex may have an effect on the dynamics of the lipids in bronchiectatic airways. We did not identify differences with blood PGE_2 , but as this was a small study, further detailed subanalysis was not carried out. Further studies would be needed to explore this.

The authors have shown that serum neutrophils are reprogrammed in bronchiectasis, leading to persistent and unresolving inflammation.¹⁵ This study now demonstrates that there is a dysregulation of the lipid mediators and failure of class switching during inflammation in bronchiectasis, despite adequate levels of PGE₂. Whether there is a role of cyclo-oxygenase inhibitors in bronchiectasis to block the production of PGE₂ needs to be explored further. Certainly, with the emergence of antibiotic resistance, the role of novel specialised proresolving lipid mediators is promising, especially in bronchiectasis, where recurrent exacerbations requiring antibiotic therapy is one of the cardinal features of the disease.

Limitations of the study

There are a couple of limitations in this study: first, that this is a small study, and second, this study did not assess the role LXA_4 on bronchiectasis neutrophils during exacerbations.

CONCLUSION

There is a dysregulation of lipid mediators in bronchiectasis in the stable state with excess proinflammatory lipids. LXA_4 improves

the function of reprogrammed neutrophils. The therapeutic efficacy of LXA, in bronchiectasis warrants further studies.

Acknowledgements The authors thank T Tolker Nielsen, University of Copenhagen, for providing the GFP Pseudomonas aeruginosa, and the QMRI Flow Cytometry and Cell Sorting Facility for assistance with flow cytometric analysis.

Contributors PB performed the experiments, collected and interpreted the data and wrote the manuscript. KZ performed lipidomics and contributed to the writing of the manuscript. PDW contributed to interpretation of data and writing of the manuscript. DJD, AGR and ATH contributed to experimental design, interpretation of data and writing of the manuscript.

Funding PB was funded by Chief Scientist Office for this study (CAF/13/02). KZ and PDW gratefully acknowledge the financial support of the European Regional. DJD was supported by a Medical Research Council Senior Non-clinical Fellowship (G1002046). AGR was supported by Medical Research Council (grant MR/ K013386/1) Development Fund, Scottish Funding Council and Highlands and Islands Enterprise.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval Lothian Research Ethics Committee gave approval for the stud (10/S1402 /S1402/33).

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. Relevant data is included in the manuscript.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: https://creativecommons.org/ licenses/by/4.0/.

REFERENCES

- 1 Bonnans C, Levy BD. Lipid mediators as agonists for the resolution of acute lung inflammation and injury. *Am J Respir Cell Mol Biol* 2007;36:201–5.
- 2 Stables MJ, Gilroy DW. Old and new generation lipid mediators in acute inflammation and resolution. *Prog Lipid Res* 2011;50:35–51.
- 3 Flower RJ. Prostaglandins, bioassay and inflammation. Br J Pharmacol 2006;147 Suppl 1:S182–92.
- 4 Samuelsson B. Role of basic science in the development of new medicines: examples from the eicosanoid field. J Biol Chem 2012;287:10070–80.

- 5 Dinarello CA, Simon A, van der Meer JWM. Treating inflammation by blocking interleukin-1 in a broad spectrum of diseases. *Nat Rev Drug Discov* 2012;11:633–52.
- 6 Samuelsson B, Dahlén SE, Lindgren JA, et al. Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. Science 1987;237:1171–6.
- 7 Klawitter J, Zafar I, Klawitter J, et al. Effects of lovastatin treatment on the metabolic distributions in the Han:SPRD rat model of polycystic kidney disease. *BMC Nephrol* 2013;14:165.
- 8 Schwab JM, Chiang N, Arita M, et al. Resolvin E1 and protectin D1 activate inflammation-resolution programmes. *Nature* 2007;447:869–74.
- 9 Ollero M, Astarita G, Guerrera IC, et al. Plasma lipidomics reveals potential prognostic signatures within a cohort of cystic fibrosis patients. J Lipid Res 2011;52:1011–22.
- 10 Massey KA, Nicolaou A. Lipidomics of polyunsaturated-fatty-acid-derived oxygenated metabolites. *Biochem Soc Trans* 2011;39:1240–6.
- 11 Serhan CN. Novel lipid mediators and resolution mechanisms in acute inflammation: to resolve or not? *Am J Pathol* 2010;177:1576–91.
- 12 Chalmers JD, Goeminne P, Aliberti S, *et al.* The bronchiectasis severity index. An international derivation and validation study. *Am J Respir Crit Care Med* 2014;189:576–85.
- 13 Maskrey BH, Rushworth GF, Law MH, et al. 12-Hydroxyeicosatetraenoic acid is associated with variability in aspirin-induced platelet inhibition. J Inflamm 2014;11:33.
- 14 Haslett C, Guthrie LA, Kopaniak MM, et al. Modulation of multiple neutrophil functions by preparative methods or trace concentrations of bacterial lipopolysaccharide. Am J Pathol 1985;119:101–10.
- 15 Bedi P, Davidson DJ, McHugh BJ, et al. Blood neutrophils are reprogrammed in bronchiectasis. Am J Respir Crit Care Med 2018;198:880–90.
- 16 Montuschi P, Kharitonov SA, Ciabattoni G, *et al.* Exhaled leukotrienes and prostaglandins in COPD. *Thorax* 2003;58:585–8.
- 17 Chen Y, Chen P, Hanaoka M, et al. Enhanced levels of prostaglandin E2 and matrix metalloproteinase-2 correlate with the severity of airflow limitation in stable COPD. *Respirology* 2008;13:1014–21.
- 18 Dagouassat M, Gagliolo J-M, Chrusciel S, et al. The cyclooxygenase-2-prostaglandin E2 pathway maintains senescence of chronic obstructive pulmonary disease fibroblasts. Am J Respir Crit Care Med 2013;187:703–14.
- 19 Takano T, Clish CB, Gronert K, *et al*. Neutrophil-mediated changes in vascular permeability are inhibited by topical application of aspirin-triggered 15-epi-lipoxin A₄ and novel lipoxin B₄ stable analogues. *J Clin Invest* 1998;101:819–26.
- 20 Serhan CN. Pro-Resolving lipid mediators are leads for resolution physiology. *Nature* 2014;510:92–101.
- 21 McDonnell MJ, Aliberti S, Goeminne PC, *et al*. Multidimensional severity assessment in bronchiectasis: an analysis of seven European cohorts. *Thorax* 2016;71:1110–8.
- 22 Mikami M, Llewellyn-Jones CG, Bayley D, *et al*. The chemotactic activity of sputum from patients with bronchiectasis. *Am J Respir Crit Care Med* 1998;157:723–8.
- 23 Chalmers JD, Smith MP, McHugh BJ, et al. Short- and long-term antibiotic treatment reduces airway and systemic inflammation in non-cystic fibrosis bronchiectasis. Am J Respir Crit Care Med 2012;186:657–65.

Thorax: first published as 10.1136/thoraxjnl-2020-216475 on 17 November 2021. Downloaded from http://thorax.bmj.com/ on November 11, 2022 at University of Glasgow. Protected by

copyright

On line supplement

Solid Phase Extraction (SPE) Chromatography

ISOLUTE C18 500 mg/6 mL cartridges were conditioned with 3x 6 mL methanol and 2x 6 mL HPLC grade water. Samples were applied and allowed to run through by force of gravity. Afterwards cartridges were first washed with half the sample volume of HPLC grade water and then with 2x 6 mL hexane. Compounds were eluted with 6 mL ethylacetate and dried under vacuum. Samples were re-dissolved in 100 μ L methanol/water 50/50 and stored at -80°C until analysis.

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) Analysis

Samples were analysed on a Thermo Scientific TSQ Quantum Ultra triple quadrupole mass spectrometer system coupled to a Thermo Scientific Accela 1250 pump and autosampler. Separation was achieved on a Phenomenex Kinetex XB-C18 LC column of 100 x 2.1 mm with 1.7 μ m particle size, maintained at 32°C. The injection volume was 10 μ L. A solvent gradient was run with solvent A consisting of 90% water, 10% methanol and 0.1% acetic acid and solvent B consisting of 100% methanol and 0.1% acetic acid and solvent B consisting of 100% methanol and 0.1% acetic acid and solvent B consisting of 100% solvent B between 0 and 10 minutes, linear increase to 70% solvent B between 10 and 11 minutes, linear increase to 100% solvent B between 11 and 18 minutes, holding at 100% solvent B between 18 and 20 minutes, decrease back to 45% solvent B between 20 and 21 minutes and column equilibration at 45 % solvent B between 21 and 25 minutes.



Figure: S1. Graph of a representative sample.

The mass spectrometer was coupled to an electrospray ionisation (ESI) source operated in negative mode and compounds were detected by multiple reaction monitoring (MRM). 56 compounds were monitored, including six deuterium labelled internal standards (see table 6 below, internal standards shown in italics). Compounds were quantified through comparison to original standards. The LC-MS/MS system was controlled and data analysis was performed with Thermo Xcalibur 2.2 software. Results were normalised to the sample volume and recovery of the associated internal standard.

Compound	Precursor	Product	Compound	Precursor	Product
	ion	ion		ion	ion
6 keto PGF1a	369.2	163.2	9 HETE	319.2	151.3
20 OH LTB ₄	351.2	195.2	11,12 dHET	337.2	167.2
20 COOH LTB ₄	365.2	195.1	14,15 dHET d11	348.3	207.2
11 dehydro TxB ₂	367.2	161.2	15 HEPE	317.2	219.2
8IsoPGF2a	353.2	193.2	8,9 dHET	337.2	127.2
TxB2	369.2	169.1	12 HEPE	317.2	179.2
iPF2a VI	353.2	115.2	5 HEPE	317.2	115.2
11beta PGF2a	353.2	193.1	5,6 dHET	337.2	145.1
PGE2	351.2	271.2	13 HODE	295.3	195.2
dihydro PGE2	351.2	235.0	20 HETE	319.2	245.3
PGE2 d4	355.2	193.2	7 HDHA	343.2	141.2
PGD2	351.2	233.1	9 HODE	295.3	171.2
dihydro PGD2	351.2	175.2	15 OxoETE	317.2	113.3
PGF2a	353.3	193.2	15 HETE	319.2	219.2
RvD2	375.2	175.1	8 HETE	319.2	155.2
LXA ₄	351.2	115.3	11 HETE	319.2	167.2
RvD1	375.2	215.2	12 HETE	319.3	179.2
LTD ₄	495.3	177.0	15 HETE d8	327.3	226.1
5,6 EET	319.2	191.3	17 HDHA	343.3	281.3
Maresin 1	359.2	177.2	14 HDHA	343.3	205.2
LTB ₄	335.2	195.2	9,10 EpOME	295.2	171.2
LTB ₄ d4	339.2	197.2	5 OxoETE	317.2	203.3
10,17 diHDHA	359.2	153.2	5 HETE	319.3	115.2
LTE4	438.2	333.3	14,15 EET d11	330.3	175.2
LTE4 d5	443.3	338.2	11,12 EET	319.2	167.2
15deoxy D 12,14 PGJ2	315.2	271.2	8,9 EET	319.3	155.2
14,15 dHET	337.2	207.2	14,15 EET	319.3	175.2
9,10 diHOME	313.2	201.2			

Table T1: Compounds monitored by liquid chromatography mass spectrometry.

Effect of LXA₄ on stable disease state peripheral blood neutrophil spontaneous apoptosis, surface expression of CD11b and CD62L and myeloperoxidase release

For this experiment, blood neutrophils from healthy volunteers, and stable state bronchiectasis patients with either mild or severe disease were pre-treated with LXA₄ 1nM, 10nM, and 100nM and assessed for the onset of spontaneous apoptosis. LXA₄ did not significantly modulate spontaneous apoptosis or viability of neutrophils (p=0.4, p=0.5 and p=0.4 respectively), in contrast to Roscovitine, used a positive control for apoptosis induction (figure S1).



Figure S2. Blood neutrophils from (a) healthy volunteer, (b) mild and (c) severe bronchiectasis patients were cultured for 20 hours and cell viability, apoptosis and necrosis were assessed by flow cytometry. There was no effect of LXA_4 , even at a maximum dose of 100nM, on spontaneous apoptosis at 20 hours. Pooled data presented as mean +/- SEM; n=6 per group. Analysis by two way ANOVA. Roscovitine is a CDK-inhibitor that promotes apoptosis and was used as a positive control (5).

LX= Lipoxin A₄; NC= negative control; rosco= roscovitine.





Figure S3. There was a concentration-dependent reduction of fMLF-induced activation (30 minutes) and CD11b expression by LXA₄; p=0.005 healthy volunteers; p=0.008 mild bronchiectasis; p=0.01 severe bronchiectasis. Pooled data presented as mean \pm SEM; n=6 per group. Analysis by one-way ANOVA. BX= bronchiectasis; NC= negative control; fMLF= n- formyl-methyl- leucyl- phenylalanine; LX= Lipoxin A₄.

Effect of LXA₄ on fMLF induced shedding of CD62L



Figure S4. There was a concentration-dependent reduction of fMLF-induced activation and CD62L shedding by LXA_4 ; p=0.01 healthy volunteers; p=0.03 mild bronchiectasis; p=0.04 severe bronchiectasis. Pooled data presented as mean \pm SEM; n=6 per group. Analysis was done by one-way ANOVA. BX= bronchiectasis; NC= negative control; fMLF= n- formyl-methyl- leucyl- phenylalanine; LX= Lipoxin A₄.



Effect of LXA₄ of neutrophil degranulation and myeloperoxidase release

Figure S5. There was a concentration-dependent reduction of fMLF (30 minutes) and cytochalasin B (10 minutes) induced activation and release of MPO by LXA₄, (a) p=0.02 healthy volunteers; (b) p=0.04 mild bronchiectasis; (c) p=0.02 severe bronchiectasis (In severe bronchiectasis, LX 100nM on its own seems to increase MPO generation - this is due to one experimental data set, which skewed the overall result). Pooled data presented as mean \pm SEM. Analysis done by one way ANOVA. Cyt= Cytochalasin B; F=fMLF= n-formyl- methyl-leucyl-phenylalanine; NC= negative control; LX= LXA₄.

Parameters	Mild	Moderate-Severe
	(N=9)	(N=15)
Age	55	68
_	(5)	(2.2)
Gender (% female)	40%	53%
Aetiology		
Idiopathic	9 (100%)	11 (74%)
Post infective	0	4 (26%)
Total WCC (x10 ⁹ /L)	6.3	7
	(0.8)	(0.5)
Neutrophils	3.3	4.4
	(1)	(0.4)
Eosinophils	0.2	0.2
	(0.09)	(0.03)
Monocytes	0.5	0.6
	(0.06)	(0.07)
ESR (mm/hr)	6	17
	(2.1)	(4.4)
CRP (mg/L)	2.2	10
	(0.5)	(4)
FEV ₁ % predicted	101	73
	(6)	(5.6)
FVC % predicted	117	95
	(1.5)	(4.6)
TLCO % predicted	100	80
	(7)	(5)
KCO % predicted	104	97
	(6)	(4)
Chronic colonisation	2	12
	(22%)	(80%)
Exacerbations in the last	0.2	3.6
year	(0.2)	(0.6)
Hospital admissions in	0	0.3
the last year		(0.1)

Table T2. Baseline demographics of the sub study population. Data presented as mean (\pm standard error of mean).

CRP= c reactive protein; ESR= erythrocyte sedimentation rate; $FEV_1=$ forced expiratory volume in 1sec; FVC= forced vital capacity; KCO= transfer coefficient corrected for alveolar volume; TLCO= transfer factor for the lung for carbon monoxide; WCC= white cell count.

On line supplement

Solid Phase Extraction (SPE) Chromatography

ISOLUTE C18 500 mg/6 mL cartridges were conditioned with 3x 6 mL methanol and 2x 6 mL HPLC grade water. Samples were applied and allowed to run through by force of gravity. Afterwards cartridges were first washed with half the sample volume of HPLC grade water and then with 2x 6 mL hexane. Compounds were eluted with 6 mL ethylacetate and dried under vacuum. Samples were re-dissolved in 100 μ L methanol/water 50/50 and stored at -80°C until analysis.

Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) Analysis

Samples were analysed on a Thermo Scientific TSQ Quantum Ultra triple quadrupole mass spectrometer system coupled to a Thermo Scientific Accela 1250 pump and autosampler. Separation was achieved on a Phenomenex Kinetex XB-C18 LC column of 100 x 2.1 mm with 1.7 μ m particle size, maintained at 32°C. The injection volume was 10 μ L. A solvent gradient was run with solvent A consisting of 90% water, 10% methanol and 0.1% acetic acid and solvent B consisting of 100% methanol and 0.1% acetic acid and solvent B consisting of 100% methanol and 0.1% acetic acid and solvent B consisting of 100% solvent B between 0 and 10 minutes, linear increase to 70% solvent B between 10 and 11 minutes, linear increase to 100% solvent B between 11 and 18 minutes, holding at 100% solvent B between 18 and 20 minutes, decrease back to 45% solvent B between 20 and 21 minutes and column equilibration at 45 % solvent B between 21 and 25 minutes.



Figure: S1. Graph of a representative sample.

The mass spectrometer was coupled to an electrospray ionisation (ESI) source operated in negative mode and compounds were detected by multiple reaction monitoring (MRM). 56 compounds were monitored, including six deuterium labelled internal standards (see table 6 below, internal standards shown in italics). Compounds were quantified through comparison to original standards. The LC-MS/MS system was controlled and data analysis was performed with Thermo Xcalibur 2.2 software. Results were normalised to the sample volume and recovery of the associated internal standard.

Compound	Precursor	Product	Compound	Precursor	Product
	ion	ion		ion	ion
6 keto PGF1a	369.2	163.2	9 HETE	319.2	151.3
20 OH LTB ₄	351.2	195.2	11,12 dHET	337.2	167.2
20 COOH LTB ₄	365.2	195.1	14,15 dHET d11	348.3	207.2
11 dehydro TxB ₂	367.2	161.2	15 HEPE	317.2	219.2
8IsoPGF2a	353.2	193.2	8,9 dHET	337.2	127.2
TxB2	369.2	169.1	12 HEPE	317.2	179.2
iPF2a VI	353.2	115.2	5 HEPE	317.2	115.2
11beta PGF2a	353.2	193.1	5,6 dHET	337.2	145.1
PGE2	351.2	271.2	13 HODE	295.3	195.2
dihydro PGE2	351.2	235.0	20 HETE	319.2	245.3
PGE2 d4	355.2	193.2	7 HDHA	343.2	141.2
PGD2	351.2	233.1	9 HODE	295.3	171.2
dihydro PGD2	351.2	175.2	15 OxoETE	317.2	113.3
PGF2a	353.3	193.2	15 HETE	319.2	219.2
RvD2	375.2	175.1	8 HETE	319.2	155.2
LXA ₄	351.2	115.3	11 HETE	319.2	167.2
RvD1	375.2	215.2	12 HETE	319.3	179.2
LTD ₄	495.3	177.0	15 HETE d8	327.3	226.1
5,6 EET	319.2	191.3	17 HDHA	343.3	281.3
Maresin 1	359.2	177.2	14 HDHA	343.3	205.2
LTB ₄	335.2	195.2	9,10 EpOME	295.2	171.2
LTB ₄ d4	339.2	197.2	5 OxoETE	317.2	203.3
10,17 diHDHA	359.2	153.2	5 HETE	319.3	115.2
LTE4	438.2	333.3	14,15 EET d11	330.3	175.2
LTE4 d5	443.3	338.2	11,12 EET	319.2	167.2
15deoxy D 12,14 PGJ2	315.2	271.2	8,9 EET	319.3	155.2
14,15 dHET	337.2	207.2	14,15 EET	319.3	175.2
9,10 diHOME	313.2	201.2			

Table T1: Compounds monitored by liquid chromatography mass spectrometry.

Effect of LXA₄ on stable disease state peripheral blood neutrophil spontaneous apoptosis, surface expression of CD11b and CD62L and myeloperoxidase release

For this experiment, blood neutrophils from healthy volunteers, and stable state bronchiectasis patients with either mild or severe disease were pre-treated with LXA₄ 1nM, 10nM, and 100nM and assessed for the onset of spontaneous apoptosis. LXA₄ did not significantly modulate spontaneous apoptosis or viability of neutrophils (p=0.4, p=0.5 and p=0.4 respectively), in contrast to Roscovitine, used a positive control for apoptosis induction (figure S1).



Figure S2. Blood neutrophils from (a) healthy volunteer, (b) mild and (c) severe bronchiectasis patients were cultured for 20 hours and cell viability, apoptosis and necrosis were assessed by flow cytometry. There was no effect of LXA_4 , even at a maximum dose of 100nM, on spontaneous apoptosis at 20 hours. Pooled data presented as mean +/- SEM; n=6 per group. Analysis by two way ANOVA. Roscovitine is a CDK-inhibitor that promotes apoptosis and was used as a positive control (5).

LX= Lipoxin A₄; NC= negative control; rosco= roscovitine.





Figure S3. There was a concentration-dependent reduction of fMLF-induced activation (30 minutes) and CD11b expression by LXA₄; p=0.005 healthy volunteers; p=0.008 mild bronchiectasis; p=0.01 severe bronchiectasis. Pooled data presented as mean \pm SEM; n=6 per group. Analysis by one-way ANOVA. BX= bronchiectasis; NC= negative control; fMLF= n- formyl-methyl- leucyl- phenylalanine; LX= Lipoxin A₄.

Effect of LXA₄ on fMLF induced shedding of CD62L



Figure S4. There was a concentration-dependent reduction of fMLF-induced activation and CD62L shedding by LXA_4 ; p=0.01 healthy volunteers; p=0.03 mild bronchiectasis; p=0.04 severe bronchiectasis. Pooled data presented as mean \pm SEM; n=6 per group. Analysis was done by one-way ANOVA. BX= bronchiectasis; NC= negative control; fMLF= n- formyl-methyl- leucyl- phenylalanine; LX= Lipoxin A₄.



Effect of LXA₄ of neutrophil degranulation and myeloperoxidase release

Figure S5. There was a concentration-dependent reduction of fMLF (30 minutes) and cytochalasin B (10 minutes) induced activation and release of MPO by LXA₄, (a) p=0.02 healthy volunteers; (b) p=0.04 mild bronchiectasis; (c) p=0.02 severe bronchiectasis (In severe bronchiectasis, LX 100nM on its own seems to increase MPO generation - this is due to one experimental data set, which skewed the overall result). Pooled data presented as mean \pm SEM. Analysis done by one way ANOVA. Cyt= Cytochalasin B; F=fMLF= n-formyl- methyl-leucyl-phenylalanine; NC= negative control; LX= LXA₄.

Parameters	Mild	Moderate-Severe
	(N=9)	(N=15)
Age	55	68
_	(5)	(2.2)
Gender (% female)	40%	53%
Aetiology		
Idiopathic	9 (100%)	11 (74%)
Post infective	0	4 (26%)
Total WCC (x10 ⁹ /L)	6.3	7
	(0.8)	(0.5)
Neutrophils	3.3	4.4
	(1)	(0.4)
Eosinophils	0.2	0.2
	(0.09)	(0.03)
Monocytes	0.5	0.6
	(0.06)	(0.07)
ESR (mm/hr)	6	17
	(2.1)	(4.4)
CRP (mg/L)	2.2	10
	(0.5)	(4)
FEV ₁ % predicted	101	73
	(6)	(5.6)
FVC % predicted	117	95
	(1.5)	(4.6)
TLCO % predicted	100	80
	(7)	(5)
KCO % predicted	104	97
	(6)	(4)
Chronic colonisation	2	12
	(22%)	(80%)
Exacerbations in the last	0.2	3.6
year	(0.2)	(0.6)
Hospital admissions in	0	0.3
the last year		(0.1)

Table T2. Baseline demographics of the sub study population. Data presented as mean (\pm standard error of mean).

CRP= c reactive protein; ESR= erythrocyte sedimentation rate; $FEV_1=$ forced expiratory volume in 1sec; FVC= forced vital capacity; KCO= transfer coefficient corrected for alveolar volume; TLCO= transfer factor for the lung for carbon monoxide; WCC= white cell count.