

Mendelian randomization analyses implicate biogenesis of translation machinery in human aging

Sara Javidnia,¹ Stephen Cranwell,¹ Stefanie H. Mueller,² Colin Selman,³ Jennifer M.A. Tullet,⁴ Karoline Kuchenbaecker,^{5,6} and Nazif Alic¹

¹Institute of Healthy Ageing, Research Department of Genetics Evolution and Environment, University College London, London WC1E 6BT, United Kingdom; ²Institute of Health Informatics, University College London, London NW1 2DA, United Kingdom; ³Institute of Biodiversity, Animal Health and Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow G12 8QQ, United Kingdom; ⁴School of Biosciences, University of Kent, Canterbury CT2 7NZ, United Kingdom; ⁵UCL Genetics Institute, Research Department of Genetics Evolution and Environment, University College London, London WC1E 6BT, United Kingdom; ⁶Division of Psychiatry, University College London, London W1T 7NF, United Kingdom

Reduced provision of protein translation machinery promotes healthy aging in a number of animal models. In humans, however, inborn impairments in translation machinery are a known cause of several developmental disorders, collectively termed ribosomopathies. Here, we use causal inference approaches in genetic epidemiology to investigate whether adult, tissue-specific biogenesis of translation machinery drives human aging. We assess naturally occurring variation in the expression of genes encoding subunits specific to the two RNA polymerases (Pols) that transcribe ribosomal and transfer RNAs, namely Pol I and III, and the variation in expression of ribosomal protein (RP) genes, using Mendelian randomization. We find each causally associated with human longevity ($\beta = -0.15 \pm 0.047$, $P = 9.6 \times 10^{-4}$, $q = 0.015$; $\beta = -0.13 \pm 0.040$, $P = 1.4 \times 10^{-3}$, $q = 0.023$; $\beta = -0.048 \pm 0.016$, $P = 3.5 \times 10^{-3}$, $q = 0.056$, respectively), and this does not appear to be mediated by altered susceptibility to a single disease. We find that reduced expression of Pol III, RPs, or Pol I promotes longevity from different organs, namely visceral adipose, liver, and skeletal muscle, echoing the tissue specificity of ribosomopathies. Our study shows the utility of leveraging genetic variation in expression to elucidate how essential cellular processes impact human aging. The findings extend the evolutionary conservation of protein synthesis as a critical process that drives animal aging to include humans.

[Supplemental material is available for this article.]

The changing demographic of human societies, where the proportion of older people is rapidly increasing (Christensen et al. 2009), makes it imperative to understand the basic biology of aging as an avenue for improvement of the health in older people (Partridge et al. 2018). Work in model organisms has identified a number of genes and processes that drive animal aging (López-Otín et al. 2013; Partridge et al. 2020). These often show striking evolutionary conservation, strongly suggestive of their relevance to aging in humans. However, direct evidence for their involvement in human aging is not often sought owing to the inherent methodological limitations, long timescales, and excessive costs associated with studying aging in human populations. The general challenges of translating findings from model organisms to humans are only starting to be overcome, for example, with human population genetic studies or drug trials (Mannick et al. 2018; Deelen et al. 2019; Torres et al. 2021). Validation of candidate processes as drivers of human aging may prove essential to show their potential as targets for treatment of age-related diseases and dysfunctions.

Multiple lines of evidence obtained in model organisms indicate that the biogenesis of protein synthesis machinery is one such conserved processes that drives organismal aging. For example,

loss of function in the genes encoding ribosomal proteins (RPs) can extend life span in yeast and worms (Kaeberlein et al. 2005; Chen et al. 2007; Chiocchetti et al. 2007; Curran and Ruvkun 2007; Hansen et al. 2007). Our own work has shown that reducing the activity of Pol I, which supplies most ribosomal RNAs (rRNAs), and Pol III, which generates the 5S rRNA and transfer RNAs (tRNAs), can extend life span in fruit flies, with the effect of Pol III also conserved in worms and yeast (Filer et al. 2017; Martínez et al. 2020). However, the relevance of the biogenesis of translation machinery to human aging has not been examined.

Protein synthesis is essential for cellular and organismal function. Indeed, loss of function in genes required to make ribosomes, the macromolecular machines that synthesize cellular proteins, is a cause of disease in humans (Mills and Green 2017). A range of human developmental disorders results from mutations in genes required for ribosome biogenesis, including RP genes (Mills and Green 2017) and subunits of Pol I and III (Bernard et al. 2011; Dauwerse et al. 2011; Weaver et al. 2015; Paolacci et al. 2018). These disorders are collectively called ribosomopathies, and their occurrence appears to contradict evidence from aging studies in model organisms, where loss of function in genes required for protein synthesis promotes longevity. However, it is possible that human disease-causing mutations cause a substantial loss of function present during development, whereas the prolongevity

Corresponding authors: k.kuchenbaecker@ucl.ac.uk, n.alic@ucl.ac.uk

Article published online before print. Article, supplemental material, and publication date are at <https://www.genome.org/cgi/doi/10.1101/gr.275636.121>. Freely available online through the *Genome Research* Open Access option.

© 2022 Javidnia et al. This article, published in *Genome Research*, is available under a Creative Commons License (Attribution 4.0 International), as described at <http://creativecommons.org/licenses/by/4.0/>.

interventions in model organisms are induced by a more moderate, possibly tissue-specific, reduction in activity in the adult. This idea prompted us to seek ways to assess the impact of moderate reduction in expression of components of translation machinery in adult tissues on human aging.

In this study, we establish a new paradigm for *in homine* validation of candidate aging processes from model organisms. Using Mendelian randomization (MR), we examine if naturally occurring variation in the expression of the subunits of Pol III or Pol I complexes or of RPs in adult humans is causally associated with human longevity. We provide evidence that biogenesis of protein synthesis machinery in specific adult tissues drives human aging.

Results

Expression of Pol III-specific subunits in the adipose tissue is linked to human longevity

We initially focused on Pol III because limiting expression of its subunits has been shown to affect life span in multiple animal models (Filer et al. 2017). These effects are tissue specific; for example, reducing Pol III subunit expression specifically in the intestine of the fly and the worm is sufficient to extend their life span (Filer et al. 2017). For this reason, we focused on organs and tissues known to modulate life span in model organisms: the brain, pituitary gland, muscle, liver, pancreas, small and large intestine (colon), and adipose tissue (Libina et al. 2003; Bartke and Brown-Borg 2004; Giannakou et al. 2004; Hwangbo et al. 2004; Broughton et al. 2005; Taguchi et al. 2007; Demontis and Perrimon 2010; Templeman et al. 2017; Woodling et al. 2020). Where gene expression has been shown to be highly correlated between similar tissues (The GTEx Consortium 2020), we used only one representative organ (e.g., visceral adipose for adipose). Because of technical constraints of transcriptomic analyses, no information is available on the variation in expression of genes transcribed by Pol III (or Pol I for that matter), such as rRNAs or tRNAs, requiring us to focus on the expression of the genes that code for the polymerase. Pol III is a protein complex composed of 17 subunits, 11 of which are specific to the enzyme (Vannini and Cramer 2012). The expression of these 11 genes was ubiquitous across these tissues (Fig. 1A) as expected from Pol III's fundamental cellular role. Still, the expression levels differed between tissues, with, for example, the pancreas showing the overall lowest expression (Fig. 1A).

To assess how variation in the expression of Pol III subunits may impact human aging, we gathered expression quantitative trait loci (eQTLs) for all the Pol III-specific subunits in the noted tissues from the Genotype-Tissue Expression (GTEx) project (The




Figure 1. MR analysis of the association between the expression of Pol III-specific subunits and longevity. (A) Levels of mRNA coding for Pol III-specific subunits in the indicated tissues are shown. (TPM) Transcripts per million. (B) MR analysis was performed using Pol III-specific subunit expression in indicated tissues as exposures and longevity (survival beyond the 90th or 99th percentile) as outcomes. The values of β are given per tissue and the significant associations indicated as those that pass the 10% FDR threshold (accounting for eight tissues and two outcomes, i.e., 16 tests). (C) MR scatter plot of association between Pol III expression in visceral adipose and longevity (90th percentile), showing results of two MR methods. Each black point denotes a *cis*-eQTL; green bars show SEM. Longevity β is the natural log of the odds ratio, where lower values indicate a lower chance of surviving beyond the 90th percentile. Expression β is reported as normalized effect sizes (NESS), with larger values indicating higher expression, and refers to expression of multiple genes.

GTEx Consortium 2020). Because all the subunits participate in a common function, we pooled all the available *cis*-eQTLs associated with the expression of any of the subunits of the Pol III complex for further analysis. Note that this includes a set of paralogs, *POLR3G* and *POLR3GL*, for which recent work has indicated that previously described differences in their cellular roles most likely arise from expression-level differences (Wang et al. 2020). We used these *cis*-eQTLs as instruments in a two-sample MR analysis to assess their causal effects on human longevity. The outcome was based on a recent meta-analysis of genome-wide association studies of human longevity. It included 11,262 individuals (cases) that had survived beyond the 90th percentile in a given human cohort and 25,483 controls (Deelen et al. 2019). The study also assessed associations in 3484 cases that had survived beyond the 99th percentile.

MR revealed Pol III subunit expression in the visceral adipose as causally associated with the likelihood of surviving above the 90th percentile ($\beta = -0.13 \pm 0.040$, $P = 1.4 \times 10^{-3}$, $q = 0.023$) (Fig. 1B,C; for per-*cis*-eQTL effect sizes and gene annotation, see Supplemental Fig. S1A; for details of MR results for all tissues, see Supplemental Table S1). This was statistically significant at a

10% false-discovery rate (FDR), accounting for the number of tissues tested and the two case definitions considered as outcomes (i.e., 16 tests). Note that the interpretation of the absolute value of the causal estimate β is complicated by the fact that *cis*-eQTLs for multiple genes were combined in the MR; however, the direction of the association is not, and the negative β revealed that increased Pol III subunit expression within adipose tissue is detrimental for longevity (Fig. 1C). No evidence for horizontal pleiotropy (HP) was found (Egger intercept, $P > 0.05$; MR-PRESSO, $P > 0.05$). No significant association was detected between longevity and Pol III subunit expression in any other tissue, although a strong negative association could also be observed in the small intestine, where the number of *cis*-eQTLs available as instruments was low (Fig. 1B; Supplemental Table S1). Overall, MR analysis indicated that reduced expression of Pol III subunits in the adult adipose tissue is causally associated with human longevity.

RP gene expression in the liver is associated with human longevity

We next examined the relationship between expression of over 80 human genes coding for RPs and longevity, as loss of function in RP genes can delay aging in yeast and worms (Kaerberlein et al.

2005; Chen et al. 2007; Chiochetti et al. 2007; Curran and Ruvkun 2007; Hansen et al. 2007). RP gene expression was ubiquitous, and their overall expression levels varied across tissues (Fig. 2A). MR analysis, using RP gene *cis*-eQTLs as instruments and longevity as the outcome, revealed a negative, causal association with the likelihood of survival beyond the 90th percentile for RP expression in the liver, which was statistically significant at 10% FDR ($\beta = -0.048 \pm 0.016$, $P = 3.5 \times 10^{-3}$, $q = 0.056$) (Fig. 2B,C; for per *cis*-eQTL effect sizes and gene annotation, see Supplemental Fig. S1B; for details of MR results for all tissues, see Supplemental Table S1). We found no evidence of HP (Egger intercept, $P > 0.05$; Mendelian randomization pleiotropy residual sum and outlier [MR-PRESSO] $P > 0.05$). Hence, in addition to reduced Pol III subunit expression in visceral adipose, we found evidence for a causal association where reduced expression of RP-coding genes in the liver promotes human longevity.

Some RPs have secondary, nonribosomal roles, and indeed, not all RP gene mutants are long-lived in model organisms (Kaerberlein et al. 2005; Chen et al. 2007; Chiochetti et al. 2007; Curran and Ruvkun 2007; Hansen et al. 2007). For this reason, we additionally examined the relationship between the expression of a subset of RP genes in the liver and survival beyond the 90th percentile. Specifically, we looked at the RPs whose orthologs are




Figure 2. MR analysis of association between expression of RP genes and longevity. (A) Levels of mRNA coding for RPs in the indicated tissues are shown. (B) MR analysis was performed using RP gene expression in indicated tissues as exposures and longevity (survival beyond the 90th or 99th percentile) as outcomes. The values of β are given per tissue and the significant associations indicated as those that pass the 10% FDR threshold (accounting for eight tissues and two outcomes, i.e., 16 tests). (C) MR scatter plot of association between RP expression in liver and longevity (90th percentile), showing results of two MR methods. Each black point denotes a *cis*-eQTL; green bars show SEM. Longevity β is the natural log of the odds ratio, where lower values indicate a lower chance of surviving beyond the 90th percentile. Expression β is reported as NESs, with larger values indicating higher expression, and refers to expression of multiple genes.

implicated in longevity in model organisms and found that the variation in their expression in the liver was also negatively associated with longevity ($\beta = -0.046 \pm 0.018$, $P = 0.010$) (Supplemental Fig. S2), confirming our initial findings.

Human longevity is affected by Pol I subunit expression in skeletal muscle

To date, Pol I has only been shown to affect aging in the fruit fly (Martínez et al. 2020). Similarly to Pol III and RP, the genes encoding the subunits unique to the Pol I complex were expressed ubiquitously, with levels of expression differing between tissues (Fig. 3A). MR analysis revealed a negative, causal association, with the likelihood of survival beyond the 99th percentile in skeletal muscle that was statistically significant at 10% FDR ($\beta = -0.15 \pm 0.047$, $P = 9.6 \times 10^{-4}$, $q = 0.015$) (Fig. 3B,C; for per *cis*-eQTL effect sizes and gene annotation, see Supplemental Fig. S1C; for details of MR results for all tissues, see Supplemental Table S1). Again, this organ was distinct from those where Pol III subunit or RP gene expression was relevant for longevity. In this case, we found some evidence of HP (Egger intercept, $P > 0.05$; MR-PRESSO $P = 3 \times 10^{-3}$), indicating that the genetic variants used in

MR may not affect longevity solely through their effect on Pol I subunit expression in muscle or, alternatively, that the relationship between expression and longevity may not be linear, as indeed is the case in the fruit fly (Martínez et al. 2020). Using MR-PRESSO to correct for outliers, the negative association between muscle expression and longevity remained significant (corrected for one outlier, $\beta = -0.18$, $P = 1.4 \times 10^{-4}$, no significant distortion). Overall, we found evidence that the expression of Pol I subunits in skeletal muscle is causally associated with human longevity.

Effects on longevity are unlikely to be explained by susceptibility to a specific age-related disease and are not observed for Pol II-specific subunit expression

The longevity phenotype captures the relative time of death but not the cause of death. For this reason, the link between expression of genes required for protein synthesis and longevity could be explained by the genes' effects on one or few age-related diseases rather than on organismal aging more broadly. For example, if RP expression solely affected cancer susceptibility and progression, but not aging more broadly, it may still have a noticeable effect on longevity per se. To examine this, we performed MR analysis using

either the Pol III or Pol I subunit or RP gene expression as exposures and a range of age-related diseases as outcomes. We limited the analysis to tissues in which a significant effect on longevity was observed for a specific gene set, namely, Pol III in visceral adipose, RP in liver, and Pol I in skeletal muscle. The associations that were significant and passed the 10% FDR threshold (accounting for 10 tests per exposure) are shown in Table 1 (for full analysis, see Supplemental Table S2). The only strong associations identified were for Pol I subunit expression in skeletal muscle, where increased expression was associated with a reduced risk of coronary heart disease and a decreased risk of amyotrophic lateral sclerosis (ALS) but an increased risk of Alzheimer's disease. Note that in models of Alzheimer's disease and ALS, protein synthetic machinery has been implicated in diseases pathophysiology, albeit cell autonomously (Meier et al. 2016; Moens et al. 2019; Yamada et al. 2019). The complex modulation of disease susceptibility observed for Pol I skeletal muscle expression makes it unlikely, albeit possible, that the effects on longevity were owing to a differential susceptibility to a single disease, indicating they may be caused by a broad effect on aging itself.

We also examined several anthropometric and developmental traits, as well as smoking as a common




Figure 3. MR analysis of association between expression of Pol I-specific subunits and longevity. (A) Levels of mRNA coding for Pol I-specific subunits in the indicated tissues are shown. (B) MR analysis was performed using Pol I-specific subunit expression in the indicated tissues as exposures and longevity (survival beyond the 90th or 99th percentile) as outcomes. The values of β are given per tissue and the significant associations indicated as those that pass the 10% FDR threshold (accounting for eight tissues and two outcomes, i.e., 16 tests). (C) MR scatter plot of association between Pol I expression in skeletal muscle and longevity (99th percentile), showing results of two MR methods. Each black point denotes a *cis*-eQTL; green bars show SEM. Longevity β is the natural log of the odds ratio, where lower values indicate a lower chance of surviving beyond the 99th percentile. Expression β is reported as NESs, with larger values indicating higher expression, and refers to expression of multiple genes.

Table 1. Associations between Pol I expression and disease

Exposure		Outcome					
Gene set	Tissue	Disease	n	β	SE	P	q
Pol I	Muscle skeletal	Alzheimer's disease (ID: 297)	38	0.058	0.023	0.011	0.037
		Amyotrophic lateral sclerosis (ID: 1086)	47	-0.013	0.0041	0.0011	0.011
		Coronary heart disease (ID: 7)	50	-0.030	0.011	0.0048	0.024

Only associations that pass 10% FDR threshold are reported (adjusted for 10 tests performed per exposure). For full analysis, see Supplemental Table S2. (n) Number of eQTLs (instruments) in the analysis.

confounder, finding no associations that passed the 10% FDR threshold (Supplemental Table S2).

Pols I and III are more specifically involved in the production of translation machinery, whereas Pol II is required for expression of all coding, messenger RNAs (Vannini and Cramer 2012). To further examine if longevity was associated specifically with provision of translation machinery rather than general transcription, we performed MR analysis for Pol II-specific subunits. We found no significant associations with longevity for Pol II subunit expression in any tissue at 10% FDR (Supplemental Table S1), indicating that longevity is specifically associated with RNA Pols that are dedicated to the production of the translation machinery.

Discussion

The biogenesis of the translation machinery is an essential cellular process required for cell and organismal growth (Grewal et al. 2007; Marygold et al. 2007; Marshall et al. 2012). Indeed, mutations in human genes that are key to this process can be linked to a number of developmental diseases (Mills and Green 2017). In contrast, in this study we provide evidence that a reduced expression of these genes in particular tissues in adult humans is causally linked to human longevity. The magnitude of the reduction in expression that favors longevity corresponds in scale to the natural, genetically encoded variation observed in human populations. Hence, it is likely that the relative reduction in biogenesis of the translation machinery that promotes longevity has substantially less effect on overall cellular capacity for translation than do disease-causing mutations. The contrast between the essential role of the translation machinery in development and its proaging, detrimental effect on the adult is consistent with evolutionary theories of aging, specifically antagonistic pleiotropy (Williams 1957): The rate of production of translation machinery is selected for optimal growth and reproduction even if it comes at a cost of accelerated aging later in life.

Our analyses indicate that Pol I, Pol III, or RPs limit longevity from different adult tissues, namely, muscle, adipose, and liver, respectively. This is akin to tissue specificity of ribosomopathies that may be explained by different protein synthesis requirements for different cells or cellular functions (Mills and Green 2017; Farley-Barnes et al. 2019), and would imply that protein synthesis drives aging from (at least) each of these three tissues. Indeed, previous work has shown that protein translation capacity plays an important role in the biology of each tissue; for example, Pol III plays an important role in adipocyte differentiation (Chen et al. 2018); tight control of ribosome biogenesis is likely to underlie the liver's daily cycles of protein synthesis (Sinturel et al. 2017); and reduced hepatic protein synthesis is observed in long-lived mouse models (Thompson et al. 2016),

whereas rDNA transcription by Pol I appears pivotal in the aging of skeletal muscle (Kirby et al. 2015). Each of the three tissues plays an important role in whole-animal metabolism. Although the link between gene expression and longevity is observed in specific tissues, the underlying mechanisms may involve tissue-nonautonomous effects such as those mediated by metabolic changes or endocrine signaling. Indeed, altering translation in specific tissues during development can have nonautonomous, endocrine effects on organismal growth in *Drosophila* (Lin et al. 2011; Marshall et al. 2012). Additionally, a recent report indicates that improved fidelity of translation can also promote longevity in the fruit fly (Martinez-Miguel et al. 2021).

Ethical considerations and costs of clinical trials severely limit the possibilities to investigate fundamental mechanisms of aging in humans. Numerous studies have shown that biogenesis of translational machinery drives aging in a number of experimental models, including yeast, nematode worms, and fruit flies (Kaeberlein et al. 2005; Chen et al. 2007; Chiocchetti et al. 2007; Curran and Ruvkun 2007; Hansen et al. 2007; Filer et al. 2017; Tiku et al. 2017; Martínez et al. 2020). We believe our study to be the first to show that this process is likely to be similarly relevant in the context of human aging. The paradigm we established leverages natural human variation in gene expression and can assess subtle, tissue-specific effects on longevity. This approach is especially useful for essential genes, in which complete knockouts are very rare or not observed owing to a severe impact on development, and overcomes the issues of assigning functional consequences to coding sequence variants. One drawback of our method is that the causal estimate β is essentially unit-less, as we are using the variation in mRNA expression of multiple genes as a proxy for the activity of the complex formed by the genes' protein products. However, the method does allow us to identify significant associations and, importantly, define their direction. The application of MR and the utilization of public data resources make it possible for future studies to apply this paradigm to investigate the relevance of other pathways that have been implicated in aging in animal models to humans. Finally, our work indicates that further studies of the link between translation machinery and aging are warranted and may result in interventions and treatments for human aging and age-related conditions.

Methods

Instrument selection

The human genes encoding the specific subunits of Pol I, II, and III or RPs were identified based on published literature (Vannini and Cramer 2012; Khatter et al. 2015). We identified *cis*-eQTLs for each gene using GTEx version 8 (The GTEx Consortium 2020) and combined them together based on their belonging to Pol III, ribosome,

Pol I, or Pol II. The human orthologs of the genes implicated in yeast or worm aging were obtained from the *Saccharomyces* Genome Database and from WormBase. The lists of genes per complex are provided in Supplemental Table S3.

In short, *cis*-eQTLs analysis was performed by GTEx consortium using FastQTL (Ongen et al. 2016). Based on 838 donors with whole-genome sequencing available, genes were included in tissue-specific eQTL analysis if expression thresholds of >0.1 TPM and six or more reads were detected in at least 20% of samples. For each gene, variants with a MAF > 0.01 in the 838 donors and located in ± 1 -Mb window around the transcription start site (TSS) were considered for *cis*-eQTL analysis. GTEx contains up to 17% of individuals with non-European or admixed ancestry (Gay et al. 2020). The association analyses between SNPs and gene expression were adjusted for ancestry by including the first five principal components from a principal component analysis based on genetic similarity between the samples.

The GTEx tissue dashboard was used to query for single tissue *cis*-eQTLs in eight tissues of interest: (1) adipose (visceral), (2) colon (transverse), (3) small intestine, (4) pancreas, (5) liver, (6) muscle (skeletal), (7) pituitary, and (8) brain (cortex). Query results for all significantly associated *cis*-eQTLs (FDR < 0.05) containing GENCODE ID, gene symbol, variant ID, dbSNP ID, *P*-value, and effect size were downloaded for subsequent analyses. The *cis*-eQTL associations were assessed by the GTEx investigators using linear regression, implemented in FastQTL (Ongen et al. 2016) after removal of low expressing genes. Effect sizes for *cis*-eQTLs are reported as normalized effect sizes (NESs) calculated as effect of the alternative allele (ALT) relative to the reference allele (REF). NESs are computed in a normalized space as the slope in the linear regression; any noise from potential residual variation in the relative NES between genes is unlikely to result in false-positive associations as it is unlikely to be linked to the outcome.

Effect allele frequencies were obtained from gnomAD v2.1.1 for European (Non-Finnish) ancestry samples. To ensure independence of instruments, linkage disequilibrium (LD) pruning was performed for each gene using the 1000 genomes reference panel for European ancestry samples and excluding weaker instrument (determined as larger *P*-value for eQTL) in pairings exceeding $r^2 = 0.1$. The lists of all *cis*-eQTLs that were fed into the MR analysis are provided in Supplemental Table S4, including details of their genomic location and the affected gene.

Tissue-specific two-sample MR

We used R version 3.6.1 (R Core Team 2017) and the TwoSampleMR package version 0.5.3 (Hemani et al. 2018) to perform two-sample MR separately for the eight tissues of interest.

To assess causal effects of sets of genes coding for either subunits specific to Pol I, II, or III or genes encoding RPs on human longevity, we obtained the summary statistics of a meta-analysis of genome-wide association studies, where cases were defined as persons living to an age above the 90th or 99th percentile of their cohort, based on cohort life tables (Deelen et al. 2019). The meta-analysis was based on 11,262 case and 25,483 control samples of European ancestry from 18 studies. Longevity meta-analysis summary statistics were kindly provided by J. Deelen.

After data harmonization between tissue-specific pooled *cis*-eQTL instruments per gene set (Pol I, II, or III or RPs) and outcome meta-analysis of life span, MR estimates were obtained using MR methods inverse variance weighted (IVW) (Burgess et al. 2013), MR Egger regression, and weighted median. Harmonized data are provided in Supplemental Table S5.

We also investigated whether associations of Pol I or III and RP with longevity were mediated by effects on a single disease or

risk factor. We included common age-related diseases and smoking. GWAS summary statistics were extracted from the largest publicly available study. The antagonistic pleiotropy theory states aging is related to mutations that are beneficial for early fitness but harmful for the organism in later life. To assess whether we see such pleiotropic effects, we also included some developmental and anthropometric traits. MR on these other outcomes was performed with MR-Base online (Hemani et al. 2018). Please find the detailed description of studies included for age-related disease outcomes, as well as developmental and anthropometric trait outcomes, in Supplemental Table S6.

HP analysis

The presence of HP was investigated using MR Egger estimates (Bowden et al. 2015), as well as MR-PRESSO (Verbanck et al. 2018). In short, the intercept of a MR Egger regression gives an estimate of the average pleiotropic effect across all instruments. MR-PRESSO will also test for significant global HP and will additionally test individual instruments. This allows the detection of outliers, their removal, and the estimation of distortion in causal effect estimates owing to inclusion of instruments with significant pleiotropic effects.

Multiple testing correction and correlation analysis

The *P*-value threshold for 10% FDR was calculated using the Benjamini and Hochberg procedure (Benjamini and Hochberg 1995) to account for the number of tissues (eight) and the two case definitions (for longevity) or the number of outcomes (for disease or developmental and anthropometric traits). This means that for the longevity analysis, we accounted for 16 tests, whereas for disease or developmental and the anthropometric traits, we accounted for 10 tests. We calculated the *q*-values using the *qvalue* package version 2.16.0 in R with conservative parameters that approximate the Benjamini and Hochberg procedure ($\lambda = 0$) accounting for the same numbers of tests as indicated above (Storey et al. 2004).

Expression data

GTEx transcript abundance data were obtained from the Human Protein Atlas version 19.3 (Uhlén et al. 2005) for expression in adipose tissue, cerebral cortex, colon, liver, pancreas, pituitary gland, skeletal muscle, and small intestine.

Competing interest statement

The authors declare no competing interests.

Acknowledgments

We thank J. Deelen for summary statistics with rsID mapping. This work was funded in part by Biotechnology and Biological Sciences Research Council grants BB/R014507/1 to N.A. and BB/S014357/1 to N.A., J.M.A.T., and C.S. and Wellcome grant 212360/Z/18/Z to K.K. S.J. was a recipient of Persia Mirzakhani Scholarship for Women.

Author contributions: S.J. and S.C. performed the analysis. S.H.M., K.K., and N.A. supervised the study. K.K. and N.A. conceived the study. C.S., J.M.A.T., K.K., and N.A. obtained funding. N.A. wrote the manuscript with contributions from all authors.

References

- Bartke A, Brown-Borg H. 2004. Life extension in the dwarf mouse. *Curr Top Dev Biol* **63**: 189–225. doi:10.1016/S0070-2153(04)63006-7
- Benjamini Y, Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Ser B Stat Methodol* **57**: 289–300. doi:10.1111/j.2517-6161.1995.tb02031.x
- Bernard G, Chouery E, Putorti ML, Tétrault M, Takanohashi A, Carosso G, Clément I, Boespflug-Tanguy O, Rodriguez D, Delague V, et al. 2011. Mutations of *POLR3A* encoding a catalytic subunit of RNA polymerase Pol III cause a recessive hypomyelinating leukodystrophy. *Am J Hum Genet* **89**: 415–423. doi:10.1016/j.ajhg.2011.07.014
- Bowden J, Davey Smith G, Burgess S. 2015. Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* **44**: 512–525. doi:10.1093/ije/dyv080
- Broughton SJ, Piper MD, Ikeya T, Bass TM, Jacobson J, Drieger Y, Martinez P, Hafen E, Withers DJ, Leever SJ, et al. 2005. Longer lifespan, altered metabolism, and stress resistance in *Drosophila* from ablation of cells making insulin-like ligands. *Proc Natl Acad Sci* **102**: 3105–3110. doi:10.1073/pnas.0405775102
- Burgess S, Butterworth A, Thompson SG. 2013. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet Epidemiol* **37**: 658–665. doi:10.1002/gepi.121758
- Chen D, Pan KZ, Palter JE, Kapahi P. 2007. Longevity determined by developmental arrest genes in *Caenorhabditis elegans*. *Aging Cell* **6**: 525–533. doi:10.1111/j.1474-9726.2007.00305.x
- Chen CY, Lanz RB, Walkey CJ, Chang WH, Lu W, Johnson DL. 2018. Maf1 and repression of RNA polymerase III-mediated transcription drive adipocyte differentiation. *Cell Rep* **24**: 1852–1864. doi:10.1016/j.celrep.2018.07.046
- Chiocchetti A, Zhou J, Zhu H, Karl T, Haubenreisser O, Rinnerthaler M, Heeren G, Oender K, Bauer J, Hintner H, et al. 2007. Ribosomal proteins Rpl10 and Rps6 are potent regulators of yeast replicative life span. *Exp Gerontol* **42**: 275–286. doi:10.1016/j.exger.2006.11.002
- Christensen K, Doblhammer G, Rau R, Vaupel JW. 2009. Ageing populations: the challenges ahead. *Lancet* **374**: 1196–1208. doi:10.1016/S0140-6736(09)61460-4
- Curran SP, Ruvkun G. 2007. Lifespan regulation by evolutionarily conserved genes essential for viability. *PLoS Genet* **3**: e56. doi:10.1371/journal.pgen.0030056
- Dauwerse JG, Dixon J, Seland S, Ruivenkamp CA, van Haeringen A, Hoefsloot LH, Peters DJ, Boers AC, Daumer-Haas C, Maiwald R, et al. 2011. Mutations in genes encoding subunits of RNA polymerases I and III cause Treacher Collins syndrome. *Nat Genet* **43**: 20–22. doi:10.1038/ng.724
- Deelen J, Evans DS, Arking DE, Tesi N, Nygaard M, Liu X, Wojczynski MK, Biggs ML, van der Spek A, Atzmon G, et al. 2019. A meta-analysis of genome-wide association studies identifies multiple longevity genes. *Nat Commun* **10**: 3669. doi:10.1038/s41467-019-11558-2
- Demontis F, Perrimon N. 2010. FOXO/4E-BP signaling in *Drosophila* muscles regulates organism-wide proteostasis during aging. *Cell* **143**: 813–825. doi:10.1016/j.cell.2010.10.007
- Farley-Barnes KI, Ogawa LM, Baserga SJ. 2019. Ribosomopathies: old concepts, new controversies. *Trends Genet* **35**: 754–767. doi:10.1016/j.tig.2019.07.004
- Filer D, Thompson MA, Takhaviev V, Dobson AJ, Kotronaki I, Green JW, Heinemann M, Tullet JM, Alic N. 2017. RNA polymerase III limits longevity downstream of TORC1. *Nature* **552**: 263–267. doi:10.1038/nature25007
- Gay NR, Gloudemans M, Antonio ML, Abell NS, Balliu B, Park Y, Martin AR, Musharoff S, Rao AS, Aguet F, et al. 2020. Impact of admixture and ancestry on eQTL analysis and GWAS colocalization in GTEx. *Genome Biol* **21**: 233. doi:10.1186/s13059-020-02113-0
- Giannakou ME, Goss M, Jünger MA, Hafen E, Leever SJ, Partridge L. 2004. Long-lived *Drosophila* with overexpressed dFOXO in adult fat body. *Science* **305**: 361. doi:10.1126/science.1098219
- Grewal SS, Evans JR, Edgar BA. 2007. *Drosophila* TIF-IA is required for ribosome synthesis and cell growth and is regulated by the TOR pathway. *J Cell Biol* **179**: 1105–1113. doi:10.1083/jcb.200709044
- The GTEx Consortium. 2020. The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science* **369**: 1318–1330. doi:10.1126/science.aaz1776
- Hansen M, Taubert S, Crawford D, Libina N, Lee SJ, Kenyon C. 2007. Lifespan extension by conditions that inhibit translation in *Caenorhabditis elegans*. *Aging Cell* **6**: 95–110. doi:10.1111/j.1474-9726.2006.00267.x
- Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, Laurin C, Burgess S, Bowden J, Langdon R, et al. 2018. The MR-base platform supports systematic causal inference across the human genome. *eLife* **7**: e34408. doi:10.7554/eLife.34408
- Hwangbo DS, Gershman B, Tu MP, Palmer M, Tatar M. 2004. *Drosophila* dFOXO controls lifespan and regulates insulin signalling in brain and fat body. *Nature* **429**: 562–566. doi:10.1038/nature02549
- Kaerberlein M, Powers RW III, Steffen KK, Westman EA, Hu D, Dang N, Kerr EO, Kirkland KT, Fields S, Kennedy BK. 2005. Regulation of yeast replicative life span by TOR and Sch9 in response to nutrients. *Science* **310**: 1193–1196. doi:10.1126/science.1115535
- Khatter H, Myasnikov AG, Natchiar SK, Klaholz BP. 2015. Structure of the human 80S ribosome. *Nature* **520**: 640–645. doi:10.1038/nature14427
- Kirby TJ, Lee JD, England JH, Chaillou T, Esser KA, McCarthy JJ. 2015. Blunted hypertrophic response in aged skeletal muscle is associated with decreased ribosome biogenesis. *J Appl Physiol (1985)* **119**: 321–327. doi:10.1152/jappphysiol.00296.2015
- Libina N, Berman JR, Kenyon C. 2003. Tissue-specific activities of *C. elegans* DAF-16 in the regulation of lifespan. *Cell* **115**: 489–502. doi:10.1016/S0092-8674(03)00889-4
- Lin JI, Mitchell NC, Kalcina M, Tchoubrieva E, Stewart MJ, Marygold SJ, Walker CD, Thomas G, Leever SJ, Pearson RB, et al. 2011. *Drosophila* ribosomal protein mutants control tissue growth non-autonomously via effects on the prothoracic gland and ecdysone. *PLoS Genet* **7**: e1002408. doi:10.1371/journal.pgen.1002408
- López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. 2013. The hallmarks of aging. *Cell* **153**: 1194–1217. doi:10.1016/j.cell.2013.05.039
- Mannick JB, Morris M, Hockey HP, Roma G, Beibel M, Kulmatycki K, Watkins M, Shavlakadze T, Zhou W, Quinn D, et al. 2018. TORC1 inhibition enhances immune function and reduces infections in the elderly. *Sci Transl Med* **10**: aaq1564. doi:10.1126/scitranslmed.aaq1564
- Marshall L, Rideout EJ, Grewal SS. 2012. Nutrient/TOR-dependent regulation of RNA polymerase III controls tissue and organismal growth in *Drosophila*. *EMBO J* **31**: 1916–1930. doi:10.1038/emboj.2012.33
- Martínez CG, Filer D, Wenz KC, Rogan A, Phillips G, Li M, Fesha Y, Broughton SJ, Alic N. 2020. Partial inhibition of RNA polymerase I promotes animal health and longevity. *Cell Rep* **30**: 1661–1669.e4. doi:10.1016/j.celrep.2020.01.017
- Martínez-Miguel VE, Lujan C, Espie-Caullet T, Martínez-Martínez D, Moore S, Backes C, Gonzalez S, Galimov ER, Brown AEX, Halic M, et al. 2021. Increased fidelity of protein synthesis extends lifespan. *Cell Metab* **33**: 2288–2300.e12. doi:10.1016/j.cmet.2021.08.017
- Marygold SJ, Roote J, Reuter G, Lambertsson A, Ashburner M, Millburn GH, Harrison PM, Yu Z, Kenmochi N, Kaufman TC, et al. 2007. The ribosomal protein genes and minute loci of *Drosophila melanogaster*. *Genome Biol* **8**: R216. doi:10.1186/gb-2007-8-10-r216
- Meier S, Bell M, Lyons DN, Rodríguez-Rivera J, Ingram A, Fontaine SN, Mechas E, Chen J, Wolozin B, LeVine H III, et al. 2016. Pathological Tau promotes neuronal damage by impairing ribosomal function and decreasing protein synthesis. *J Neurosci* **36**: 1001–1007. doi:10.1523/JNEUROSCI.3029-15.2016
- Mills EW, Green R. 2017. Ribosomopathies: There's strength in numbers. *Science* **358**: eaan2755. doi:10.1126/science.aan2755
- Moens TG, Niccoli T, Wilson KM, Atlano ML, Birsá N, Gittings LM, Holbling BV, Dyson MC, Thoeng A, Neeves J, et al. 2019. *C9orf72* arginine-rich dipeptide proteins interact with ribosomal proteins in vivo to induce a toxic translational arrest that is rescued by eIF1A. *Acta Neuropathol* **137**: 487–500. doi:10.1007/s00401-018-1946-4
- Ongen H, Buil A, Brown AA, Dermitzakis ET, Delaneau O. 2016. Fast and efficient QTL mapper for thousands of molecular phenotypes. *Bioinformatics* **32**: 1479–1485. doi:10.1093/bioinformatics/btv722
- Paolacci S, Li Y, Agolini E, Bellacchio E, Arboleda-Bustos CE, Carrero D, Bertola D, Al-Gazali L, Alders M, Altmüller J, et al. 2018. Specific combinations of biallelic *POLR3A* variants cause Wiedemann–Rautenstrauch syndrome. *J Med Genet* **55**: 837–846. doi:10.1136/jmedgenet-2018-105528
- Partridge L, Deelen J, Slagboom PE. 2018. Facing up to the global challenges of ageing. *Nature* **561**: 45–56. doi:10.1038/s41586-018-0457-8
- Partridge L, Fuentealba M, Kennedy BK. 2020. The quest to slow ageing through drug discovery. *Nat Rev Drug Discov* **19**: 513–532. doi:10.1038/s41573-020-0067-7
- R Core Team. 2017. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org/>.
- Sinturel F, Gerber A, Mauvoisin D, Wang J, Gatfield D, Stubblefield JJ, Green CB, Gachon F, Schibler U. 2017. Diurnal oscillations in liver mass and cell size accompany ribosome assembly cycles. *Cell* **169**: 651–663.e14. doi:10.1016/j.cell.2017.04.015
- Storey JD, Taylor JE, Siegmund D. 2004. Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: a unified approach. *J R Stat Soc Series B Stat Methodol* **66**: 187–205. doi:10.1111/j.1467-9868.2004.00439.x

- Taguchi A, Wartschow LM, White MF. 2007. Brain IRS2 signaling coordinates life span and nutrient homeostasis. *Science* **317**: 369–372. doi:10.1126/science.1142179
- Templeman NM, Flibotte S, Chik JHL, Sinha S, Lim GE, Foster LJ, Nislow C, Johnson JD. 2017. Reduced circulating insulin enhances insulin sensitivity in old mice and extends lifespan. *Cell Rep* **20**: 451–463. doi:10.1016/j.celrep.2017.06.048
- Thompson AC, Bruss MD, Price JC, Khambatta CF, Holmes WE, Colangelo M, Dalidd M, Roberts LS, Astle CM, Harrison DE, et al. 2016. Reduced *in vivo* hepatic proteome replacement rates but not cell proliferation rates predict maximum lifespan extension in mice. *Aging Cell* **15**: 118–127. doi:10.1111/accel.12414
- Tiku V, Jain C, Raz Y, Nakamura S, Heestand B, Liu W, Späth M, Suchiman HED, Müller RU, Slagboom PE, et al. 2017. Small nucleoli are a cellular hallmark of longevity. *Nat Commun* **8**: 16083. doi:10.1038/ncomms16083
- Torres GG, Nygaard M, Caliebe A, Blanché H, Chantalat S, Galan P, Lieb W, Christiansen L, Deleuze JF, Christensen K, et al. 2021. Exome-wide association study identifies *FN3KRP* and *PGP* as new candidate longevity genes. *J Gerontol A Biol Sci Med Sci* **76**: 786–795. doi:10.1093/geron/ glab023
- Uhlén M, Björling E, Agaton C, Szgyarto CA, Amini B, Andersen E, Andersson AC, Angelidou P, Asplund A, Asplund C, et al. 2005. A human protein atlas for normal and cancer tissues based on antibody proteomics. *Mol Cell Proteomics* **4**: 1920–1932. doi:10.1074/mcp.M500279-MCP200
- Vannini A, Cramer P. 2012. Conservation between the RNA polymerase I, II, and III transcription initiation machineries. *Mol Cell* **45**: 439–446. doi:10.1016/j.molcel.2012.01.023
- Verbanck M, Chen CY, Neale B, Do R. 2018. Detection of widespread horizontal pleiotropy in causal relationships inferred from Mendelian randomization between complex traits and diseases. *Nat Genet* **50**: 693–698. doi:10.1038/s41588-018-0099-7
- Wang X, Gerber A, Chen WY, Roeder RG. 2020. Functions of paralogous RNA polymerase III subunits POLR3G and POLR3GL in mouse development. *Proc Natl Acad Sci* **117**: 15702–15711. doi:10.1073/pnas.1922821117
- Weaver KN, Watt KE, Hufnagel RB, Navajas Acedo J, Linscott LL, Sund KL, Bender PL, König R, Lourenco CM, Hehr U, et al. 2015. Acrofacial dysostosis, Cincinnati type, a mandibulofacial dysostosis syndrome with limb anomalies, is caused by *POLR1A* dysfunction. *Am J Hum Genet* **96**: 765–774. doi:10.1016/j.ajhg.2015.03.011
- Williams GC. 1957. Pleiotropy, natural selection, and the evolution of senescence. *Evolution (N Y)* **11**: 398–411. doi:10.1111/j.1558-5646.1957.tb02911.x
- Woodling NS, Aleyakpo B, Dyson MC, Minkley LJ, Rajasingam A, Dobson AJ, Leung KHC, Pomposova S, Fuentealba M, Alic N, et al. 2020. The neuronal receptor tyrosine kinase Alk is a target for longevity. *Aging Cell* **19**: e13137. doi:10.1111/accel.13137
- Yamada SB, Gendron TF, Niccoli T, Genuth NR, Grosely R, Shi Y, Glaria I, Kramer NJ, Nakayama L, Fang S, et al. 2019. RPS25 is required for efficient RAN translation of *C9orf72* and other neurodegenerative disease-associated nucleotide repeats. *Nat Neurosci* **22**: 1383–1388. doi:10.1038/s41593-019-0455-7

Received April 13, 2021; accepted in revised form December 15, 2021.



Mendelian randomization analyses implicate biogenesis of translation machinery in human aging

Sara Javidnia, Stephen Cranwell, Stefanie H. Mueller, et al.

Genome Res. 2022 32: 258-265 originally published online January 25, 2022

Access the most recent version at doi:[10.1101/gr.275636.121](https://doi.org/10.1101/gr.275636.121)

Supplemental Material <http://genome.cshlp.org/content/suppl/2022/01/10/gr.275636.121.DC1>

References This article cites 56 articles, 12 of which can be accessed free at:
<http://genome.cshlp.org/content/32/2/258.full.html#ref-list-1>

Open Access Freely available online through the *Genome Research* Open Access option.

Creative Commons License This article, published in *Genome Research*, is available under a Creative Commons License (Attribution 4.0 International), as described at <http://creativecommons.org/licenses/by/4.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Affordable, Accurate
Sequencing.



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>
