


# Anti-Müllerian hormone as a marker of ovarian reserve and premature ovarian insufficiency in children and women with cancer: a systematic review

Richard A. Anderson <sup>1,\*</sup>, David Cameron<sup>2</sup>, Florian Clatot<sup>3</sup>,  
Isabelle Demeestere<sup>4</sup>, Matteo Lambertini<sup>5,6</sup>, Scott M. Nelson <sup>7,8,9</sup>,  
and Fedro Peccatori<sup>10</sup>

<sup>1</sup>MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, UK <sup>2</sup>Edinburgh University Cancer Centre, IGMM, Edinburgh, UK <sup>3</sup>Centre Henri Becquerel, Rouen, France <sup>4</sup>Fertility clinic, CUB-Erasme Hospital, Université Libre de Bruxelles, Brussels, Belgium <sup>5</sup>Department of Medical Oncology, U.O. Clinica di Oncologia Medica, IRCCS Ospedale Policlinico San Martino, Genova, Italy <sup>6</sup>Department of Internal Medicine and Medical Sciences (DiMI), School of Medicine, University of Genova, Genova, Italy <sup>7</sup>School of Medicine, University of Glasgow, Glasgow, UK <sup>8</sup>NIHR Bristol Biomedical Research Centre, Bristol, UK <sup>9</sup>The Fertility Partnership, Oxford, UK <sup>10</sup>European Institute of Oncology IRCCS, Milan, Italy

\*Correspondence address. MRC Centre for Reproductive Health, The Queen's Medical Research Institute, Edinburgh BioQuarter, 47 Little France Crescent, Edinburgh EH16 4TJ, UK. Tel: +44-(0)-131-242-6386; E-mail:richard.anderson@ed.ac.uk  <https://orcid.org/0000-0002-7495-518X>

Submitted on August 17, 2021; resubmitted on November 18, 2021; editorial decision on January 11, 2022

## TABLE OF CONTENTS

- Introduction
- Methods
- Results
  - Study characteristics
  - Changes in AMH as a result of anticancer treatments
  - Recovery of AMH in longitudinal studies
  - Overall relationship of AMH with measures of POI and menstrual function after anticancer treatment
  - Post-treatment AMH and pregnancy
  - Age at treatment, ovarian reserve and extent of recovery
  - Studies investigating gonadotoxicity of anticancer treatments
  - AMH in patients with breast cancer
  - AMH in patients with lymphoma
  - Paediatric patients
  - Impact of duration of follow-up
- Discussion
- Conclusion

**BACKGROUND:** Female patients undergoing anticancer treatment are at elevated risk of adverse ovarian outcomes including infertility and premature ovarian insufficiency (POI), which is associated with short- and long-term health risks. Anti-Müllerian hormone (AMH) is a

key biomarker of ovarian reserve, but its role prior to and after cancer treatment is less well understood.

**OBJECTIVE AND RATIONALE:** To conduct a systematic review evaluating AMH as a biomarker of ovarian reserve and POI before and after anticancer treatment, which has become a pressing clinical issue in reproductive medicine. There are a large number of observational studies, but differences in patient groups, cancer diagnoses and study design make this a confusing field that will benefit from a thorough and robust review.

**SEARCH METHODS:** A systematic literature search for AMH in women with cancer was conducted in PubMed, Embase and Cochrane Central Register of Controlled Trials up to 1 April 2021. Bias review was conducted using the Risk of Bias In Non-randomized Studies of Interventions (ROBINS-I) protocol along with qualitative assessment of quality. Exploratory subgroups were established based on age, cancer type and length of follow-up.

**OUTCOMES:** Ninety-two publications (N=9183 patients) were included in this analysis after quality and bias review. Reduced/undetectable AMH was consistently identified in 69/75 studies (92%) following chemotherapy or radiotherapy, with reductions ranging from 42% to concentrations below the limit of detection, and many reporting mean or median declines of  $\geq 90\%$ . Where longitudinal data were analysed (42 studies), a majority (33/42 (79%)) of studies reported at least partial recovery of AMH at follow-up, however, effect estimates were highly variable, reflecting that AMH levels were strongly impacted by anticancer treatment (i.e. the chemotherapy regimen used and the number of treatment cycles need), with recovery and its degree determined by treatment regimen, age and pre-treatment AMH level. In 16/31 (52%) publications, oligo/amenorrhoea was associated with lower post-treatment AMH consistent with impending POI, although menstruation and/or pregnancy were reported in patients with low or undetectable AMH. Long-term (>5 years) follow-up of paediatric patients following cancer treatment also found significantly lower AMH compared with control groups in 14/20 (70%) of studies, with very variable effect sizes from complete loss of AMH to full recovery depending on treatment exposure, as in adult patients.

**WIDER IMPLICATIONS:** AMH can be used to identify the damaging effect of cancer treatments on ovarian function. This can be applied to individual women, including pre-pubertal and adolescent girls, as well as comparing different treatment regimens, ages and pre-treatment AMH levels in populations of women. While there was evidence for its value in the diagnosis of POI after cancer treatment, further studies across a range of diagnoses/treatment regimens and patient ages are required to clarify this, and to quantify its predictive value. A major limitation for the use of AMH clinically is the very limited data relating post-treatment AMH levels to fertility, duration of reproductive lifespan or time to POI; analysis of these clinically relevant outcomes will be important in further research.

**Key words:** anti-Müllerian hormone / AMH / fertility / cancer / gonadotoxicity / ovarian reserve / ovarian insufficiency / chemotherapy

## Introduction

Anticancer treatments are known to have substantial impact on ovarian reserve via mechanisms leading to depletion of growing and primordial follicles. This reduction in ovarian reserve may reduce reproductive lifespan and be sufficient in some women to induce premature ovarian insufficiency (POI; Fig. 1; [Spears et al., 2019](#)). This has profound implications for women's health and is associated with infertility, cardiovascular disease (CVD), neurological disease, osteoporosis and mood disorders, culminating in an increased risk of premature mortality ([Webber et al., 2016](#); [van Dorp et al., 2018](#); [Panay et al., 2020](#)). Consequently, knowing a patient's treatment-related risk of POI aids shared decision-making around the choice of anticancer treatment and the need for additional fertility preservation interventions.


Measurement of circulating anti-Müllerian hormone (AMH), produced by granulosa cells of growing follicles, is widely used as a surrogate for ovarian reserve and has potential as a diagnostic and predictive biomarker of POI ([Anderson and Su 2020](#)). AMH levels are often heavily impacted during anticancer treatment, followed by a highly variable recovery ([Anderson and Su 2020](#)). The diagnosis of POI is informed by monitoring a combination of biomarkers, including menstrual function and FSH ([Webber et al., 2016](#); [Panay et al., 2020](#)). However, confirming POI is more challenging in patients with cancer, as there is potential for recovery of ovarian/menstrual function, which can occur from several months to years after treatment completion ([Silva et al., 2016](#)).

Questions remain about the reliability of AMH for evaluating ovarian reserve in the context of different anticancer treatments, and at what point post-treatment it can provide useful predictive information which may guide later treatment decisions, such as choice of endocrine therapy in women with breast cancer. Our objective was to conduct a systematic review evaluating the value of AMH in assessing ovarian reserve in women with cancer undergoing anticancer treatments, and specifically in the prediction and diagnosis of POI.

## Methods

This study was conducted according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and registered with PROSPERO (CRD42020161717). PubMed, Embase and Cochrane Central Register of Controlled Trials (CENTRAL) were searched for articles published up to 1 April 2021 ([Supplementary Data](#)) and duplicate articles removed.

Search results were screened in stages by title, abstract and full text based on the following inclusion criteria: female, premenopausal patients who have prospectively or retrospectively undergone treatment for any cancer. Exclusion criteria included pathology other than cancer (with the exception of granulosa cell tumours), AMH/ovarian reserve not measured, cohort size <10, duplicate datasets/cohorts, poor quality (e.g. inconsistent reporting of results), review articles and congress abstracts.



Publications requiring evaluation of full text to determine inclusion underwent bias review via the Risk of Bias In Non-randomized Studies of Interventions (ROBINS-I) protocol (Sterne *et al.*, 2016). Bias conclusions were verified by a second author, with any disagreements mediated by a third author. All authors approved the final selection.

Data were extracted using the Synthesis Without Meta-analysis (SWiM) in systematic reviews reporting guideline (Campbell *et al.*, 2020). The main metrics assessed were differences in AMH following anticancer treatment (change versus pre-treatment or control groups), recovery of AMH (in longitudinal cohorts) and correlation of AMH with menstrual function (regular cycles, oligomenorrhoea or amenorrhoea as reported by patient at follow-up). Certainty of evidence was based on vote counting.

Extracted data also included description of study types (prospective/retrospective, cross-sectional, longitudinal), year of publication, cancer type, treatment, cohort mean/median age, timing of AMH measurement and assay type. Exploratory subgroups were established according to age (paediatric, adult,  $</\geq 35$  years), diagnosis (breast cancer or lymphoma) and length of follow-up.

The data underlying this article are available in the article and in its online [supplementary material](#), or as referenced publications.

## Results


### Study characteristics

Figure 2 shows the number of citations retrieved, the number after screening and the final number included in the analysis. The searches initially identified over 3700 citations. After removal of duplicates and assessment for quality and relevance to the study question at the title,

abstract and full-text level, 110 were included for bias review. Of these, a further 20 were assessed to be at severe or critical risk of bias and excluded (Supplementary Table S1), leaving 90 publications including 8916 patients (N) for systematic review, published between 2003 and 2021. During the publication review process, a further two relevant articles that were published after 1 April 2021 were identified (total 112 for bias review), resulting in a total of 92 publications and 9183 patients. An overview of publications included in this review is summarized in Table I, and details of the individual publications are summarized in Supplementary Table SII.

### Changes in AMH as a result of anticancer treatments

Papers reporting AMH before and after treatment (N=46; N=4117), and a clear majority of cross-sectional papers comparing post-treatment survivors with control groups (23/26, 88%; n/N=2283/3088) reported a large reduction in AMH following treatment (Table I; Supplementary Table SII). Effect sizes were variable depending on the treatment and diagnosis, but in the 26 papers that reported AMH values at both baseline and  $\leq 3$  months from end of treatment (Lutchman Singh *et al.*, 2007; Decanter *et al.*, 2010, 2018, 2021; Rosendahl *et al.*, 2010; Yu *et al.*, 2010; Brougham *et al.*, 2012; Henry *et al.*, 2014; Ben-Aharon *et al.*, 2015; Gharwan *et al.*, 2016; Gupta *et al.*, 2016; Anderson *et al.*, 2017, 2018; Bi *et al.*, 2017; D'Avila *et al.*, 2017; Leonard *et al.*, 2017; Trapp *et al.*, 2017; Cameron *et al.*, 2018; Evranos *et al.*, 2018; Yaish *et al.*, 2018; Passildas *et al.*, 2019; Silva *et al.*, 2019; Loubersac *et al.*, 2020; Berjeb *et al.*, 2021; Demeestere *et al.*, 2021; Martin *et al.*, 2021), reductions ranged from



42% to below the limit of detection (LOD) and 18 reported mean or median declines of  $\geq 90\%$  (Rosendahl et al., 2010; Brougham et al., 2012; Dillon et al., 2013a, b; Henry et al., 2014; Gharwan et al., 2016; Ben-Aharon et al., 2015; D'Avila et al., 2017; Dezellus et al., 2017; Trapp et al., 2017; Cameron et al., 2018; Decanter et al., 2018; Lambertini et al., 2019; Palinska-Rudzka et al., 2019; Passildas et al., 2019; Silva et al., 2019; Oktay et al., 2020; Ruddy et al., 2021; Yu et al., 2010). The six papers that did not detect a significant difference

in AMH versus controls after treatment were all in paediatric populations (i.e. under 18 years of age at cancer diagnosis) and represented long-term follow-up periods of 10–30 years in heterogeneous populations of childhood cancer survivors (Lie Fong et al., 2009; Nielsen et al., 2013; van den Berg et al., 2018; Nystrom et al., 2019; Nies et al., 2020; Elitzur et al., 2021). A shorter-term longitudinal study (follow-up of up to 43 months) in this population did, however, show clear reductions in AMH following treatment (Brougham et al., 2012),

**Table 1** Summary of publications included in review.

Subgroups	Number of publications	Number of patients	AMH reduction reported during/after anticancer treatment	AMH correlated with menstrual function*
All publications	92	9183	69/75 (92%)	15/30 (50%)
Longitudinal (pre-/post-treatment)	56	4825	46/49 (94%)	11/18 (61%)
Cross-sectional (post-treatment data only)	36	4358	23/26 (88%)	4/12 (33%)
Cancer type <sup>†</sup>				
Breast	38	3600	31/31 (100%)	11/18 (61%)
Lymphoma	11	1199	9/9 (100%)	3/43 (75%)
Multiple	33	3877	21/25 (84%)	2/8 (25%)
Other <sup>‡</sup>	11	573	9/11 (82%)	0/1 (0%)
Follow-up period (mean/median)				
< 1 year	11	796	9/9 (100%)	4/6 (67%)
1–2 years	36	2943	33/33 (100%)	7/15 (47%)
>2–5.5 years	19	2010	15/15 (100%)	3/6 (50%)
>5.5 years	26	3434	12/18 (67%)	1/4 (25%)
Cohort age				
Paediatric	24	2392	14/19 (74%)	1/6 (17%)
Adult	61	5991	48/49 (98%)	13/22 (59%)
Mixed	7	800	7/7 (100%)	1/3 (33%)
<35 years old at follow-up <sup>§</sup>	57	5917	43/47 (91%)	6/16 (38%)
≥35 years old at follow-up <sup>§</sup>	33	2999	24/26 (92%)	8/13 (62%)

\*Regular menstruation, oligomenorrhoea or amenorrhoea at follow-up.

<sup>†</sup>N = 91 as Palinska-Rudzka *et al.* (2019) included separate analysis for breast cancer and lymphoma.

<sup>‡</sup>Acute lymphoblastic leukaemia, differentiated thyroid cancer and gestational trophoblastic neoplasia.

<sup>§</sup>Mean or median as reported.

AMH, anti-Müllerian hormone.

with large differences in the degree of recovery of AMH by treatment gonadotoxicity: thus in those exposed to high-risk treatment, AMH remained undetectable, while recovering to levels similar to pre-treatment in those exposed to lower risk treatment.

### Recovery of AMH in longitudinal studies

In publications examining longitudinal data, post-treatment recovery of AMH was described in 33/42 (79%). With several notable exceptions described below, recovery was typically partial (i.e. to lower than pre-treatment level) or only occurred in a subset of patients. Twelve publications specifically evaluated the association of pre- and post-treatment AMH levels, with all reporting a significant association (Rosendahl *et al.*, 2010; Dillon *et al.*, 2013b; D'Avila *et al.*, 2015; Dezellus *et al.*, 2017; Anderson *et al.*, 2018; Lee *et al.*, 2018, 2020; Palinska-Rudzka *et al.*, 2019; Silva *et al.*, 2019; Celebi *et al.*, 2020; Loubersac *et al.*, 2020; Decanter *et al.*, 2021).

### Overall relationship of AMH with measures of POI and menstrual function after anticancer treatment

Ten publications specifically evaluated either the value of post-treatment AMH in the diagnosis of POI, or pre-treatment AMH in predicting the likelihood of POI. These studies were either in breast cancer or across multiple diagnoses, representing 762 patients

(Nielsen *et al.*, 2013; Lunsford *et al.*, 2014; Elchuri *et al.*, 2016; Anderson *et al.*, 2017; Nystrom *et al.*, 2019; Passildas *et al.*, 2019; Silva *et al.*, 2019; Zhong *et al.*, 2019; Demeestere *et al.*, 2021; Parissonne *et al.*, 2021). Six studies (two of which were in paediatric cohorts) reported post-treatment AMH levels <1.0 ng/ml or undetectable in patients with POI (Nielsen *et al.*, 2013; Lunsford *et al.*, 2014; Anderson *et al.*, 2017; Silva *et al.*, 2019; Zhong *et al.*, 2019; Parissonne *et al.*, 2021) but the diagnostic accuracy of AMH for post-treatment POI has only been reported in one study (Anderson *et al.*, 2017). Two papers found that lower pre-treatment AMH levels were associated with higher risk of POI (as assessed) post-treatment (Passildas *et al.*, 2019; Zhong *et al.*, 2019), but no studies have assessed post-treatment AMH as a predictor of time to POI.

Of 37 papers that specifically investigated menstrual function after treatment, all used prospective evaluation, and 14 (38%) found that reduced post-treatment AMH was associated with oligomenorrhoea or amenorrhoea (Rosendahl *et al.*, 2008; Su *et al.*, 2010; Lunsford *et al.*, 2014; Ruddy *et al.*, 2014; D'Avila *et al.*, 2015; Gharwan *et al.*, 2016; Palinska-Morarji *et al.*, 2017; Weners *et al.*, 2017; Decanter *et al.*, 2018, 2021; Kim *et al.*, 2018; Palinska-Rudzka *et al.*, 2019; Silva *et al.*, 2019; Li *et al.*, 2020). In some of these studies, the observations were nuanced, for example Su *et al.* (2010) found that post-treatment AMH was associated with amenorrhoea but was not associated with recovery of menses (at 5.2 years post-treatment follow-up). Palinska-Rudzka *et al.* (2019) found that AMH was lower in patients with amenorrhoea,

but that some (7/17) patients with AMH <LOD had menses. Gharwan et al. (2016) found AMH was associated with menstrual status only in woman aged 21–25 years old. Dezellus et al. (2017) found an association between AMH and menstrual status at 6 months of post-treatment follow-up, but not at 12 months of follow-up. Further studies are required to assess with rigour whether measurement of AMH adds to or can replace current diagnostic criteria for POI.

### Post-treatment AMH and pregnancy

A small number of studies have reported pregnancies in some patients despite low or undetectable AMH levels after cancer treatment (Hamy et al., 2016; Dezellus et al., 2017; Anderson et al., 2018; Loubersac et al., 2020; Demeestere et al., 2021). A recent analysis of AMH in women treated for advanced Hodgkin lymphoma in a randomized controlled trial showed similar pregnancy rates in the two treatment arms after 5 years of follow-up, despite much lower AMH levels and a higher POI rate in the arm receiving higher doses of alkylating agents (Demeestere et al., 2021). This is consistent with the lack of predictive value of AMH for pregnancy in healthy women (Steiner et al., 2017), but these limited data on a clinically highly relevant topic highlight the need for further prospective studies to inform on this subject.

### Age at treatment, ovarian reserve and extent of recovery

Of 39 publications that evaluated whether patient age at treatment was associated with post-treatment AMH levels, 30 (77%) found higher post-treatment AMH in younger individuals (Lutchman Singh et al., 2007; Lie Fong et al., 2009; Charpentier et al., 2014; Henry et al., 2014; Ben-Aharon et al., 2015; D'Avila et al., 2015; Acibucu et al., 2016; Elchuri et al., 2016; Anderson et al., 2017, 2018; Dezellus et al., 2017; Morarji et al., 2017; Shandley et al., 2017; Wengers et al., 2017; Al-Janabi et al., 2018; Evranos et al., 2018; Lee et al., 2018; Malisic et al., 2018; Yaish et al., 2018; Cameron et al., 2019; Silva et al., 2019; Celebi et al., 2020; Li et al., 2020; Loubersac et al., 2020; Su et al., 2020; van Velsen et al., 2020; Berjeb et al., 2021; Decanter et al., 2021; Elitzur et al., 2021; Martin 2021). Thirteen of these studies also stratified patient groups by <35 versus ≥35 years old, with 10 (83%) reporting lower AMH and poorer recovery in the older age group (Yu et al., 2010; Ben-Aharon et al., 2015; Acibucu et al., 2016; Dezellus et al., 2017; Trapp et al., 2017; Al-Janabi et al., 2018; Anderson et al., 2018; van den Berg et al., 2018; Yaish et al., 2018; Li et al., 2020; Mittica et al., 2020; van Velsen et al., 2020; Decanter et al., 2021). However, only five publications both stratified these two age groups and evaluated menstrual function, with three also showing a reduced chance of recovery of menstruation in older women (Yu et al., 2010; Dezellus et al., 2017; Decanter et al., 2021).

### Studies investigating gonadotoxicity of anticancer treatments

Fifty-two papers specifically investigated gonadotoxicity of treatments either by comparison of groups receiving specific regimens or by cumulative toxicity scores (e.g. cyclophosphamide equivalent dose [CED]; summarized in Supplementary Table SIII). In order to specifically evaluate treatment effect, papers that evaluated patient risk of gonadotoxicity based on other factors, such as diagnosis and disease

stage, were not counted. Forty papers (77%) reported a treatment effect on AMH, with higher gonadotoxicity correlating with lower post-treatment AMH. In general, chemotherapy regimens containing alkylating agents, such as cyclophosphamide (e.g. ACVBP, BEACOPP, CHOP, FEC), resulted in lower levels of AMH and poorer recovery of AMH after treatment than non-alkylating treatment regimens (Rosendahl et al., 2008; Gracia et al., 2012; Thomas-Teinturier et al., 2015; Morarji et al., 2017; Anderson et al., 2018; Leiper et al., 2020; Decanter et al., 2021).

Twenty-one publications reported gonadotoxicity scores or ranked treatments as higher/lower toxicity or higher/lower risk, based on treatment type and/or cumulative chemotherapy. In every study where higher versus lower toxicity was assessed, higher toxicity therapies and more treatment cycles resulted in lower post-treatment AMH compared with lower overall toxicity exposure (Rosendahl et al., 2010; Brougham et al., 2012; Gracia et al., 2012; Hamre et al., 2012; Di Paola et al., 2013; Nielsen et al., 2013; Krawczuk-Rybak et al., 2013a, 2013b, 2019; Charpentier et al., 2014; Lunsford et al., 2014; Thomas-Teinturier et al., 2015; Elchuri et al., 2016; van der Kooi et al., 2017, 2019; van den Berg et al., 2018; Cameron et al., 2019; George et al., 2019; Su et al., 2020; van Velsen et al., 2020; Parissonne et al., 2021).

The evidence regarding radiotherapy was less conclusive as its use was only cited in 10 publications (van Beek et al., 2007; Lie Fong et al., 2009; Brougham et al., 2012; Gracia et al., 2012; Dillon et al., 2013b; Miyoshi et al., 2013; Elchuri et al., 2016; van den Berg et al., 2018; George et al., 2019; Nies et al., 2020). However, in studies where radiotherapy did result in significantly lower AMH, this was typically when targeted to pelvic/abdominal regions or total body irradiation (Lie Fong et al., 2009; Gracia et al., 2012; Miyoshi et al., 2013; Elchuri et al., 2016; van den Berg et al., 2018; George et al., 2019).

### AMH in patients with breast cancer

Thirty-eight papers evaluated AMH levels in women following treatment for breast cancer, representing a total of 3600 patients (Table II). All reported a negative treatment effect on AMH, and in 12 papers with data available, reductions in AMH from pre-treatment to <3 months post-treatment ranged from ~80% to 99% (Lutchman Singh et al., 2007; Yu et al., 2010; Henry et al., 2014; Ben-Aharon et al., 2015; Anderson et al., 2017; D'Avila et al., 2017; Leonard et al., 2017; Trapp et al., 2017; Decanter et al., 2018; Passildas et al., 2019; Silva et al., 2019; Loubersac et al., 2020). Of those publications with available data, 13/16 found that post-treatment AMH levels were associated with age at baseline (Lutchman Singh et al., 2007; Henry et al., 2014; Ben-Aharon et al., 2015; D'Avila et al., 2015; Anderson et al., 2017; Dezellus et al., 2017; Wengers et al., 2017; Al-Janabi et al., 2018; Lee et al., 2018; Silva et al., 2019; Celebi et al., 2020; Li et al., 2020; Loubersac et al., 2020). Some degree of recovery in average AMH levels during follow-up was reported in 14/20 publications; however, the extent of this recovery was marginal (<15% average recovery from nadir levels as a percentage of baseline values) in 12 publications, and only partial in the remainder (range of follow-up 0–5 years), with none demonstrating a return of AMH to pre-treatment levels (Yu et al., 2010; Henry et al., 2014; Ben-Aharon et al., 2015; Anderson et al., 2017; Dezellus et al., 2017; Leonard et al., 2017; Decanter et al., 2018; Lambertini et al., 2019; Palinska

**Table II Characteristics of studies in breast cancer cohorts included in the systematic review (N = 38).**

Publication	Study design	Adult/ paediatric	Cohort size	Timing of AMH assessment	AMH assay	LOD (ng/ml)	Reduced/low AMH following anticancer treatment*	Correlation of post-treatment AMH with menstrual function†	Correlation of post-treatment AMH recovery with age
<b>Al-Janabi et al. (2018)</b>	Prospective longitudinal	Adult	60	BL, after first menses after 1st dose, after 4th dose	Gen II ELISA (Beckman Coulter)	Not reported	Yes	N/A	Yes
<b>Anderson et al. (2011)</b>	Prospective longitudinal	Adult	42	BL, PTFU 2, 3, 4, 5 years	ELISA (Beckman Coulter)	0.05	Yes	Yes	N/A
<b>Anderson et al. (2013)</b>	Prospective longitudinal	Adult	55	BL, after 1st and 2nd cycle, PTFU 1 year	Gen II ELISA (Beckman Coulter)	0.16	Yes	Yes	N/A
<b>Anderson et al. (2017)</b>	Prospective longitudinal	Adult	101	BL, PTFU 0, 12, 24 months	Elecsys® AMH (Roche Diagnostics)	0.01	Yes	N/A	Yes
<b>Ben-Aharon et al. (2015)</b>	Prospective longitudinal	Adult	20	BL, PTFU 3–4 days, 6, 12 months	ELISA (Beckman Coulter)	0.01	Yes	N/A	Yes
<b>Çelebi et al. (2020)</b>	Prospective cross-sectional	Adult	31	BL, during/after treatment every 3 months up to 12 months	ELISA	Not reported	Yes	N/A	Yes
<b>D'Avila et al. (2015)</b>	Prospective longitudinal	Adult	52	BL, PTFU 6 months	ELISA (Beckman Coulter)	Not reported	Yes	Yes	Yes
<b>D'Avila et al. (2017)</b>	Prospective longitudinal	Adult	52	BL, PTFU 2, 6 months	ELISA (Beckman Coulter—Immunotech)	Not reported	Yes	Yes	N/A
<b>Decanter et al. (2018)</b>	Prospective longitudinal	Adult	32	BL, 15 days before last cycle, PTFU 3, 6, 9 and 12 months	picoAMH ELISA (Ansh)	0.0224 (0.16 pmol/l)	Yes	Yes	N/A
<b>Dezellus et al. (2017)</b>	Prospective cross-sectional	Adult	249	BL, before each of 6 cycles, PTFU 6, 12 and 24 months	ELISA (Beckman Coulter—Immunotech)	0.14	Yes	No	Yes
<b>Eigindy et al. (2013)</b>	Prospective cross-sectional	Adult	100	BL, PTFU 12 months	Not reported	Not reported	Yes	N/A	N/A
<b>Goldfarb et al. (2020)</b>	Prospective cross-sectional	Adult	142	BL, PTFU 12, 18 and 24 months	picoAMH ELISA (Ansh Labs)	Not reported	Yes	N/A	N/A
<b>Hamy et al. (2016)</b>	Prospective cross-sectional	Adult	134	BL, during treatment, PTFU 4–66 months	ELISA (Beckman Coulter—Immunotech)	0.14	N/A	N/A	N/A
<b>Henry et al. (2014)</b>	Prospective cross-sectional	Adult	27	BL, mean PTFU 4.9 weeks, 13.6 months	Gen II ELISA (Beckman Coulter)	0.16	Yes	No	Yes
<b>Kim et al. (2018)</b>	Prospective cross-sectional	Adult	82	PTFU <2 months	ELISA (USCN Life Science)	0.003	N/A	Yes	N/A
<b>Lambertini et al. (2019)</b>	Prospective longitudinal	Adult	148	BL, PTFU 1, 3 years	Elecsys AMH (Roche Diagnostics)	0.01	Yes	N/A	N/A

Continued

Table II Continued

Publication	Study design	Adult/ paediatric	Cohort size	Timing of AMH assessment	AMH assay	LOD (ng/ml)	Reduced/low AMH following anticancer treatment*	Correlation of post-treatment AMH with menstrual function†	Correlation of post-treatment AMH recovery with age
<b>Lee et al. (2018)</b>	Prospective cross-sectional	Adult	105	PTFU 508 days	Gen II ELISA (Beckman Coulter)	0.16	N/A	No	Yes
<b>Lee et al. (2020)</b>	Prospective cross-sectional	Adult	67	BL, PTFU 1 year	Gen II ELISA (Beckman Coulter)	0.16	Yes	No	No
<b>Leonard et al. (2017)</b>	Prospective cross-sectional	Adult	202	BL, during treatment (C3), PTFU 0, 9, 12, 24 months	Elecsys AMH (Roche Diagnostics)	Not reported	Yes	N/A	N/A
<b>Li et al. (2020)</b>	Prospective cross-sectional	Adult	144	BL, PTFU 6 months	Human AMH ELISA (Cusabio Biotech)	Not reported	Yes	Yes	Yes
<b>Lutchman Singh et al. (2007)</b>	Prospective cross-sectional	Adult	35	BL, PTFU after re- sumption of menses	ELISA (Beckman Coulter— Immunotech)	Not reported	Yes	N/A	Yes
<b>Malisic et al. (2018)</b>	Prospective cross-sectional	Adult	40	PTFU 1–66 months	ELISA (Ansh Labs)	0.023	N/A	N/A	Yes
<b>Morarji et al. (2017)</b>	Prospective longitudinal	Adult	100	PTFU 2 years	Gen II ELISA (Beckman Coulter)	0.08	Yes	Yes	Yes
<b>Oktaç et al. (2020)</b>	Prospective cross-sectional	Adult	108	BL, PTFU <24 months	picoAMH ELISA (Ansh Labs)	0.003	Yes	N/A	N/A
<b>Palinska-Rudzka et al. (2019)</b>	Prospective longitudinal	Adult	54	BL, PTFU 1, 5 years	Gen II ELISA (Beckman Coulter)	0.02	Yes	Yes	N/A
<b>Partridge et al. (2010)</b>	Prospective cross-sectional	Adult	20	PTFU 5 years	ELISA (Diagnostic Systems Laboratories)	0.03	Yes	N/A	N/A
<b>Passidas et al. (2019)</b>	Prospective longitudinal	Adult	58	BL, PTFU 1 month	Elecsys AMH (Roche Diagnostics)	0.03	Yes	No	N/A
<b>Ruddy et al. (2014)</b>	Retrospective cross-sectional	Adult	124	BL, PTFU 12, 18 months	ELISA (Diagnostic Systems Laboratories)	Not reported	N/A	Yes	N/A
<b>Ruddy et al. (2021)</b>	Prospective longitudinal	Adult	277	BL, PTFU 1 year	Ultrasensitive ELISA (Ansh Labs)	Not reported	Yes	N/A	N/A
<b>Shandley et al. (2017)</b>	Prospective longitudinal	Adult	108	PTFU 7 years	Ultrasensitive ELISA and pico AMH ELISA (Ansh Labs)	0.006	N/A	N/A	Yes
<b>Silva et al. (2019)</b>	Prospective longitudinal	Adult	38	BL, during treatment, PTFU 1, 6 and >9 months up to 18 months	Ultrasensitive ELISA (Ansh Labs)	0.06	Yes	Yes	Yes
<b>Su et al. (2014)</b>	Prospective longitudinal	Adult	109	BL, PTFU 1–1009 days	Gen II ELISA (Beckman Coulter)	0.17	N/A	N/A	N/A

Continued



**Table II Continued**

Publication	Study design	Adult/ paediatric	Cohort size	Timing of AMH assessment	AMH assay	LOD (ng/ml)	Reduced/low AMH following anticancer treatment*	Correlation of post-treatment AMH with menstrual function†	Correlation of post-treatment AMH recovery with age
<b>Su et al. (2010)</b>	Prospective longitudinal	Adult	127	PTFU 1–4 years	ELISA (Diagnostic Systems Laboratories)	0.025	Yes	Yes	N/A
<b>Trapp et al. (2017)</b>	Prospective cross-sectional	Adult	170	BL, PTFU 2 years	ELISA (Beckman Coulter)	0.08	Yes	No	No
<b>Wenners et al. (2017)</b>	Prospective longitudinal	Adult	51	BL, during treatment, PTFU 6, 12, 24 months	Elecsys AMH (Roche Diagnostics)	Not reported	Yes	Yes	Yes
<b>Yu et al. (2010)</b>	Prospective longitudinal	Adult	26	BL, PTFU 6, 12, 24, 36, 52 weeks	ELISA (Diagnostic Systems Laboratories)	Not reported	Yes	No	No
<b>Zhong et al. (2019)</b>	Prospective cross-sectional	Adult	96	BL, PTFU 6, 12 months	Immunoassay (Beckman Coulter Unicel DX1800)	Not reported	Yes	N/A	N/A

\*Between baseline and follow-up in longitudinal studies, and versus controls in cross-sectional studies.

†Presence of regular cycles, oligomenorrhoea or amenorrhoea.

AMH: anti-Müllerian hormone; BL, baseline; LOD, limit of detection; N/A, not applicable; PTFU, post-treatment follow-up.

Rudzka et al., 2019; Silva et al., 2019; Zhong et al., 2019; Loubersac et al., 2020; Oktay et al., 2020; Goldfarb et al., 2021). In all studies reporting no recovery of AMH levels, patients received treatment with alkylating agents, such as cyclophosphamide-based chemotherapy regimens; however, higher pre-treatment AMH and lower age were identified as factors associated with greater recovery after these regimens (Yu et al., 2010; Anderson and Cameron 2011; Anderson et al., 2013; Elgindy et al., 2013; D’Avila et al., 2017; Wenness et al., 2017; Palinska-Rudzka et al., 2019; Zhong et al., 2019).

In patients with breast cancer, post-treatment AMH was also linked to menstrual function at follow-up in 11/18 publications (Su et al., 2010; Ruddy et al., 2014; D’Avila et al., 2015, 2017; Morarji et al., 2017; Wenness et al., 2017; Decanter et al., 2018; Kim et al., 2018; Palinska-Rudzka et al., 2019; Silva et al., 2019; Li et al., 2020). Patients with undetectable AMH generally had a higher risk of amenorrhoea, although several studies reported patients with low or undetectable AMH who did recover menstrual function.

The impact of BRCA status was investigated in two studies (Lambertini et al., 2019; Oktay et al., 2020), with one identifying increased loss of ovarian reserve and greater reduction in AMH post-treatment recovery in BRCA+ individuals (Oktay et al., 2020).

### AMH in patients with lymphoma

Of 11 studies in lymphoma (Table III), all nine publications evaluating treatment effect on AMH in patients with lymphoma (including Hodgkin lymphoma), representing a total of 521 patients, found a significant impact (van Beek et al., 2007; Decanter et al., 2010, 2021; Nitzschke et al., 2010; Demeestere et al., 2016, 2021; Gharwan et al., 2016; Anderson et al., 2018; Palinska-Rudzka et al., 2019). Eight papers reported significant decline in AMH values from pre-treatment to ≤3 months’ post-treatment (Decanter et al., 2010, 2021; Demeestere et al., 2016, 2021; Gharwan et al., 2016; Anderson et al., 2018; Palinska-Rudzka et al., 2019). An additional study also reported significant decline in post-treatment AMH compared with controls (Nitzschke et al., 2010). All eight papers evaluating gonadotoxicity found a regimen- or dose-dependent effect on AMH following treatment and during recovery (van Beek et al., 2007; Decanter et al., 2010, 2021; Hamre et al., 2012; Behringer et al., 2013; Anderson et al., 2018; Palinska-Rudzka et al., 2019; Demeestere et al., 2021). Five of these studies identified that patients receiving doxorubicin, bleomycin, vinblastine, dacarbazine (ABVD)-based regimens had at least partial post-treatment recovery of AMH at follow-up compared with those receiving cyclophosphamide-containing regimens (Decanter et al., 2010, 2021; Anderson et al., 2018; Palinska-Rudzka et al., 2019; Demeestere et al., 2021). One paper described limited recovery of AMH in older ABVD-treated patients (Anderson et al., 2018) whereas another found that the impact of age on recovery was only found in women treated with alkylating agents (Decanter et al., 2021).

### Paediatric patients

The majority (20/29; 69%) of papers including paediatric cancer patients were cross-sectional studies reporting on adult childhood cancer survivors (Bath et al., 2003; van Beek et al., 2007; Lie Fong et al., 2009; Gracia et al., 2012; Hamre et al., 2012; El-Shalakany et al., 2013; Miyoshi et al., 2013; Nielsen et al., 2013; Krawczuk-Rybak et al., 2013b, 2019; Charpentier et al., 2014; Lunsford et al., 2014; Thomas-

Table III Characteristics of studies in lymphoma cohorts included in the systematic review (N = 11).

Publication	Study design	Adult/ paediatric	Cohort size	Timing of AMH assessment	AMH assay	LOD (ng/ml)	Reduced/low AMH following anticancer treatment*	Correlation of post-treatment AMH with menstrual function†	Correlation of post-treatment AMH recovery with age
<b>Anderson et al. (2018)</b>	Prospective longitudinal	Adult	67	BL, after 2nd cycle, PTFU 0, 1, 2, 3 years	Elecsys AMH (Roche Diagnostics)	0.01	Yes	N/A	Yes
<b>Behringer et al. (2013)</b>	Prospective cross-sectional	Adult	562	PTFU 46 months	Gen II ELISA (Beckman Coulter)	Not reported	N/A	N/A	Yes
<b>Decanter et al. (2010)</b>	Prospective longitudinal	Adult	30	BL, 15 days after 1st cycle, 15 days before last cycle, PTFU 0, 3, 6, 9, 12 months	Gen II ELISA (Beckman Coulter)	0.098 (0.7 pmol/L)	Yes	N/A	N/A
<b>Decanter et al. (2021)</b>	Prospective longitudinal	Adolescent, Young adult, Adult	122	BL, 15 days after 1st cycle, 15 days before last cycle, PTFU 3, 6, 9, 12 months	Gen II ELISA (Beckman Coulter)	Not reported	Yes	Yes	Yes
<b>Demeestere et al. (2016)</b>	Prospective longitudinal	Adult	67	BL, PTFU 2, 3, 4, 5–7 years	Elecsys AMH (Roche Diagnostics)	Not reported	Yes	N/A	N/A
<b>Demeestere et al. (2021)</b>	Prospective longitudinal	Adult	145	BL, PTFU 0, 1, 2, 3, 4, 5 years	Elecsys AMH (Roche Diagnostics)	0.01 ng/ml	Yes	N/A	Yes
<b>Gharwan et al. (2016)</b>	Prospective longitudinal	Adult	28	BL, PTFU 4–18 months	Gen II ELISA (Beckman Coulter)	Not reported	Yes	Yes (in 21- to 25-year-old group)	N/A
<b>Hamre et al. (2012)</b>	Prospective cross-sectional	Paediatric	62	PTFU 18.1 years	ELISA (Beckman Coulter)	0.15	N/A	No	N/A
<b>Nitzsche et al. (2010)</b>	Prospective longitudinal	Adult	20	PTFU 30.6–32.2 months	ELISA (Diagnostic Systems Laboratories)	0.025	Yes	N/A	N/A
<b>Palinska-Rudzka et al. (2019)</b>	Prospective longitudinal	Adult	12	BL, PTFU 1, 5 years	Gen II ELISA (Beckman Coulter)	0.02	Yes	Yes	N/A
<b>van Beek et al. (2007)</b>	Prospective longitudinal	Paediatric	30	PTFU 11.6 years	ELISA (Diagnostic Systems Laboratories)	Not reported	Yes (in patients receiving MOPP)	N/A	No

\*Between baseline and follow-up in longitudinal studies, and vs controls in cross-sectional studies.

†Presence of regular cycles, oligomenorrhoea or amenorrhoea.

AMH: anti-Müllerian hormone; BL, baseline; LOD, limit of detection; N/A, not applicable; PTFU, post-treatment follow-up.

Teinturier *et al.*, 2015; Elchuri *et al.*, 2016; van der Kooi *et al.*, 2017; van den Berg *et al.*, 2018; Cameron *et al.*, 2019; George *et al.*, 2019; Nystrom *et al.*, 2019; Nies *et al.*, 2020), of which 19 had follow-up periods of >5 years. Reductions in AMH compared with pre-treatment levels or versus controls were reported by 14/20 (70%) publications (Bath *et al.*, 2003; van Beek *et al.*, 2007; Brougham *et al.*, 2012; El-Shalakany *et al.*, 2013; Krawczuk-Rybak *et al.*, 2013a, 2013b, 2019; Elchuri *et al.*, 2016; Gupta *et al.*, 2016; Morse *et al.*, 2016; van der Kooi *et al.*, 2017, 2019; Cameron *et al.*, 2019; George *et al.*, 2019). As with adult cancer patients, the degree of reduction varied with treatment received, with complete loss of AMH in some patients. Low AMH levels were found in some cancer survivors despite normal FSH levels (e.g. in 20% of survivors (George *et al.*, 2019)), indicating the added value of AMH in detecting partial ovarian damage. Additional studies reported trends to lower AMH, or low AMH levels in patients following childhood cancer that were not compared with a control group or baseline data (Hamre *et al.*, 2012; Miyoshi *et al.*, 2013; Nielsen *et al.*, 2013; Charpentier *et al.*, 2014; Lunsford *et al.*, 2014; van den Berg *et al.*, 2018; George *et al.*, 2019). While these studies support that long-term AMH levels can be reduced in survivors of childhood cancer, important outcomes such as relationships with fertility and age at menopause have not been evaluated.

Most studies (10/12 (90.9%)) (van Beek *et al.*, 2007; Lie Fong *et al.*, 2009; Brougham *et al.*, 2012; Hamre *et al.*, 2012; Miyoshi *et al.*, 2013; Krawczuk-Rybak *et al.*, 2013a, 2019; Charpentier *et al.*, 2014; Thomas-Teinturier *et al.*, 2015; van der Kooi *et al.*, 2019) which had data for both pre- and post-menarchal patients did not detect significant differences in post-treatment AMH by this parameter, and one reported better AMH recovery in pre-menarchal patients (van der Kooi *et al.*, 2019). Two publications also reported delayed puberty in patients with low post-treatment AMH (El-Shalakany *et al.*, 2013; Lunsford *et al.*, 2014).

### Impact of duration of follow-up

To assess the impact of cancer treatment on AMH levels during long-term follow-up, studies with <1 year of follow-up were excluded owing to the potential for a residual effect of cancer treatment and incomplete recovery. Based on median reported follow-up post-treatment periods, there were 35 studies with 1- to 2-year follow-up (Bath *et al.*, 2003; Rosendahl *et al.*, 2010; Yu *et al.*, 2010; Brougham *et al.*, 2012; Anderson *et al.*, 2013, 2017; Elgindy *et al.*, 2013; Iwase *et al.*, 2013; Dillon *et al.*, 2013b; Henry *et al.*, 2014; Ruddy *et al.*, 2014, 2021; Ben-Aharon *et al.*, 2015; Gharwan *et al.*, 2016; Gupta *et al.*, 2016; Dezellus *et al.*, 2017; Leonard *et al.*, 2017; Morarji *et al.*, 2017; Trapp *et al.*, 2017; Wengers *et al.*, 2017; Decanter *et al.*, 2010, 2018; Evranos *et al.*, 2018; Lee *et al.*, 2018, 2020; Yaish *et al.*, 2018; Silva *et al.*, 2019; van der Kooi *et al.*, 2019; Zhong *et al.*, 2019; Celebi *et al.*, 2020; Loubersac *et al.*, 2020; Oktay *et al.*, 2020; Berjeb *et al.*, 2021; Goldfarb *et al.*, 2021; Martin, 2021). Most (33/35; 96%) found that AMH was reduced after treatment, 19/25 (76%) later then identified some degree of increase in AMH at follow-up. Although only a few studies found a (very modest) increase in AMH after 12 months, several studies with a maximum post-treatment follow-up period of 12 months showed the highest post-treatment AMH value at 12 months, raising the possibility that further recovery could have been observed (Brougham *et al.*, 2012; Ben-Aharon *et al.*, 2015; Gupta

*et al.*, 2016; Decanter *et al.*, 2018; Yaish *et al.*, 2018). It therefore appears that 12 months after completion of treatment is a minimum period to allow complete or near-complete recovery of ovarian function: little recovery is seen thereafter and becomes confounded with the normal decline in AMH with increasing age.

Of 18 studies with a median >2–5.5 years of follow-up, 15/15 with available data (100%) identified AMH reductions in cancer survivors (Nitzschke *et al.*, 2010; Partridge *et al.*, 2010; Su *et al.*, 2010; Anderson and Cameron 2011; El-Shalakany *et al.*, 2013; Acibucu *et al.*, 2016; Demeestere *et al.*, 2016, 2021; Morse *et al.*, 2016; Anderson *et al.*, 2018; Cameron *et al.*, 2019; Lambertini *et al.*, 2019; Palinska-Rudzka *et al.*, 2019; Chemerinski *et al.*, 2020; van Velsen *et al.*, 2020), 8/10 (80%) identified at least partial recovery at follow-up in  $\geq 1$  patient group (Demeestere *et al.*, 2016, 2021; Morse *et al.*, 2016; Anderson *et al.*, 2018; Malisic *et al.*, 2018; Cameron *et al.*, 2019; Lambertini *et al.*, 2019; Palinska-Rudzka *et al.*, 2019) and 3/5 (60%) reported correlation of post-treatment AMH with menstrual function in some patients (Rosendahl *et al.*, 2008; Su *et al.*, 2010; Palinska-Rudzka *et al.*, 2019). Of 24 cross-sectional studies with follow-up >5.5 years, 11/17 (65%) found AMH reductions with treatment when compared with controls (van Beek *et al.*, 2007; Gracia *et al.*, 2012; Di Paola *et al.*, 2013; Krawczuk-Rybak *et al.*, 2013a, 2013b, 2019; Thomas-Teinturier *et al.*, 2015; Elchuri *et al.*, 2016; Leiper *et al.*, 2020; Mittica *et al.*, 2020; Roshandel *et al.*, 2021) and 1/4 (25%) linked AMH to menstrual function in some patients (Lunsford *et al.*, 2014). There are very limited data regarding continuing changes in AMH after cancer treatment and whether cancer survivors show an accelerated decline in AMH in the later reproductive years. In a mixed longitudinal/cross-sectional analysis, AMH levels were shown to be reduced according to gonadotoxicity but maintained a plateau for a prolonged period after treatment (Su *et al.*, 2020). Supporting that, cancer survivors showed no evidence of an increased rate of decline in AMH compared to age-matched controls (Cameron *et al.*, 2019).

## Discussion

Anticancer treatment has a clear negative impact on women's ovarian reserve, with reduced AMH levels observed across studies and diagnoses, and reductions of  $\geq 90\%$  commonly identified. Some degree of post-treatment recovery in AMH levels was described in many patient groups, although full- or near-complete restoration to pre-treatment levels was rare and typically occurred in patients receiving milder treatment. Figure 3 gives a representation of the key findings of this review. It is notable that there are relatively few prospective studies, and among those with longitudinal data, several had a maximum 1 year of follow-up. While AMH levels measured 1–2 years after treatment are likely to be reliable at indicating longer-term menstrual outcomes, further research is required to confirm this, especially in older women where recovery is slower and more limited (van Beek *et al.*, 2007; Decanter *et al.*, 2010; Behringer *et al.*, 2013; Anderson *et al.*, 2018; Palinska-Rudzka *et al.*, 2019).

Treatments known to be more gonadotoxic, whether this be drug or cumulative exposure related, generally caused a greater reduction in AMH. The clearest evidence comes from lymphoma, where patients receiving ABVD-based regimens showed more complete AMH recovery and resumption of menstruation compared to more severe

treatment strategies (van Beek et al., 2007; Decanter et al., 2010, 2021; Behringer et al., 2013; Anderson et al., 2018; Palinska-Rudzka et al., 2019). Studies in breast cancer also identified taxanes as contributing to lower AMH (Hamy et al., 2016; Lambertini et al., 2019; Silva et al., 2019). However, studies evaluating cumulative dose of chemotherapy, regardless of regimen, show this to be a key factor in ovarian outcomes (Charpentier et al., 2014; Elchuri et al., 2016; van den Berg et al., 2018; Cameron et al., 2019; George et al., 2019).

Pre-treatment AMH values were associated with post-treatment AMH in all 11 studies that evaluated this, supporting the predictive value of AMH at baseline. All of these studies were in post-menarchal patients. Several studies also identified low pre-treatment AMH in some patients (Lutchman Singh et al., 2007; Cameron et al., 2018; Palinska-Rudzka et al., 2019; van der Kooi et al., 2019), which was not limited to a particular diagnosis.

An important question is whether the lower AMH levels after cancer treatment then decline at a different rate compared to other women. Three studies found no significant difference in rates between patients and age-matched controls (van der Kooi et al., 2017; Cameron et al., 2019; Elitzur et al., 2021). However, there is a higher rate of decline in older women, both cancer survivors and controls (Dezellus et al., 2017; Cameron et al., 2019). These studies highlight the need for research investigating the relationships between post-treatment AMH, age and age of POI/menopause, which would be of substantial clinical value.

While the picture regarding treatment impact on AMH is clear, there was no clear relationship between AMH and menstrual status at follow-up, with menstruation reported in patients with undetectable AMH across different diagnoses and treatments (Decanter et al., 2010; Rosendahl et al., 2010; Ben-Aharon et al., 2015). The reliability of AMH in predicting menstrual function may be better in older patients, consistent with AMH becoming undetectable in advance of natural menopause in healthy women. In contrast, in younger women, menstrual cycles may continue despite a very limited ovarian reserve and undetectable AMH. Although several studies have reported a clear relationship between low/undetectable AMH and diagnosis of POI, the diagnostic validity of AMH for post-cancer treatment POI has not been sufficiently evaluated, either alone or in combination with established criteria.

Interpretation of AMH in paediatric cohorts is potentially complicated by the natural arc of AMH, which increases during childhood and through the early adult years before a decline to the menopause (Kelsey et al., 2011). With the exception of one publication (van der Kooi et al., 2019), there were no differences in the effect of treatment on post-treatment AMH levels between pre- and post-menarchal patients, suggesting the reliability of AMH as a biomarker in younger populations. It should also be noted that all six studies that did not find a treatment influence on AMH were in adult survivors of childhood cancer (Lie Fong et al., 2009; Nielsen et al., 2013; van den Berg et al., 2018; Nystrom et al., 2019; Nies et al., 2020; Elitzur et al., 2021), whereas other studies did find an effect. Further longitudinal studies, from diagnosis through to adulthood, are required to further explore the value of AMH for predicting late ovarian function shortly after completion of therapy in paediatric and adolescent patients with cancer.

Across the different publications, spanning nearly two decades, multiple AMH assay types, most commonly ELISAs or


electrochemiluminescence immunoassays (ECLIA), from a variety of manufacturers have been used. There is currently no internationally agreed calibration standardization for AMH, and the improvements in sensitivity afforded by automated assays in comparison with historical methods, as well as inherent variation and non-linear conversion between assays, can complicate interpretation of the data (Li et al., 2021). Whether variations in assay calibration and performance influence the clinical value of AMH assessment was not formally assessed, but the overall consistency of the main findings indicates that this is not a major issue. It is possible though that the use of assays with greater sensitivity (Chai et al., 2014; Decanter et al., 2014) will impact on analysis of AMH as a diagnostic test for POI.

A variety of additional independent factors can influence levels of AMH (Dafopoulos et al., 2010; Dölleman et al., 2013). Although publications featuring potentially AMH-modifying conditions, such as systemic illness, endometriosis, polycystic ovary syndrome and granulosa cell tumours, were excluded, studies with patients who use cigarettes or oral contraceptives (Elitzur et al., 2021), as well as *BRCA* mutation carriers (Lambertini et al., 2019; Oktay et al., 2020), were included. The studies described here represent highly heterogeneous patient populations, incorporating a wide range of ages, diagnoses and treatment types. While this may limit the ability of this review to make specific recommendations, it reinforces the wider applicability of our observations.

This review confirms AMH as a marker of ovarian reserve but does not specifically address assessment of time to menopause, which may be clinically useful. This has previously been addressed in relation to natural menopause via probability (Finkelstein et al., 2020) and estimated rate of decline-based models (de Kat et al., 2019; Ramezani Tehrani et al., 2020). As AMH has been shown to correlate with patient age and treatment gonadotoxicity and declines at a similar rate to healthy individuals following any post-treatment recovery, it should be possible to develop a model for menopause risk over time based on pre- and post-treatment AMH measurements. Additional longitudinal studies would support such a model and help to further establish the time course of any AMH recovery, and in which patient groups it occurs, especially whether a time point <2 years' post-treatment can reliably provide a long-term predictive marker.

## Conclusion

In this systematic review, we confirm that anticancer treatment substantially impacts AMH, with large reductions during treatment followed by a period of partial recovery in some women, peaking within 1–2 years (Fig. 3). Low post-treatment AMH was predicted by lower pre-treatment AMH, as well as by older age and gonadotoxicity of treatment. Consequently, discussions around treatment and ovarian outcomes should consider the risk of POI and need for family planning and fertility preservation, informed by AMH. While low post-treatment AMH correlated with increased likelihood of POI, the evidence with respect to oligo- or amenorrhoea was less clear and there were very little data relating AMH to fertility or time to/age at POI or menopause. AMH is therefore a clinically useful biomarker of ovarian reserve before and after cancer treatment and it is likely to be of value in the diagnosis of POI after cancer treatment but relationships with post-treatment fertility and reproductive lifespan require further investigation.



**Figure 3. Summary of findings of systematic review represented graphically.** The three lines represent women with high (green), average (blue) and low (red) AMH levels before treatment, with treatment represented by the shaded area and a threshold indicating POI is also represented. (1) AMH concentrations are reduced by cancer treatments. (2) Recovery is variable, depending on patient age, treatment regimen and pre-treatment AMH levels. Recovery can be near complete or absent, with the latter resulting in permanent POI. The relationship of post-treatment AMH to POI needs to be explored further. (3) Prediction of permanent POI at the time of end of treatment may be possible in some situations, but further studies are required to determine the patient groups for which this may and may not be possible. (4) There are insufficient data to be able to predict the duration of post-recovery ovarian function using AMH levels before or after treatment, which will interact with the physiological decline in AMH with increasing age. Reproduced from Jayasinghe *et al.* (2018) with permission.

## Supplementary data

Supplementary data are available at *Human Reproduction Update* online.

## Data availability

There are no new data associated with this article.

## Acknowledgements

The authors thank Simon Lott of Elements Communications Limited for editorial assistance with the preparation of the manuscript.

## Authors' roles

All authors developed the strategy for the literature search, reviewed the outputs of the searches and reviewed and approved the manuscript. R.A.A. guided the data analysis and development of the manuscript.

## Funding

All authors attended an Advisory Board meeting in December 2019 to discuss the project, for which they or their institution received an honorarium sponsored by Roche Diagnostics. Medical writing support and article processing charges were funded by Roche Diagnostics. M.L. was supported by the Italian Ministry of Health (5 × 1000 funds 2017; no grant number) and the Italian Association for Cancer Research (AIRC; MFAG 2020 ID 24698).

## Conflict of interest

D.C.'s institution has received research grants from Roche; payment for participation in independent data monitoring committees from Roche. F.P. has received speaker's honoraria from Ipsen and Merck. F.C.'s institution has received reagents from Roche Diagnostics. I.D. has received an academic trial grant and reagents (given to the institution) from Roche Diagnostics; an academic trial grant from Ferring; speaker's honoraria from Novartis; support for attending meetings from Ferring and Theramex. M.L. has received consulting fees and speaker's honoraria from AstraZeneca, Exact Sciences, Lilly, Novartis, Pfizer, Roche Diagnostics, Roche Pharma and Seagen; speaker's honoraria from Ipsen, Sandoz and Takeda. R.A.A. has received consulting fees from Ferring, NeRRE Therapeutics, Roche Diagnostics and Sojournix Inc; payment from Merck and IBSA for educational events; laboratory materials from Roche Diagnostics. S.M.N. has received grants from CSO, ESHRE and MRC; consulting fees from Access Fertility, Coopers Genomics, Ferring, Merck, Modern Fertility and TFP; speaker's honoraria from Ferring, Merck and Roche Diagnostics; payment for expert testimony from Medical Defence Work; support for attending meetings from Ferring and Merck. S.M.N. has personally invested in TFP.

R.A.A. had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

## References

Acibucu F, Acibucu DO, Akkar OB, Dokmetas HS. Evaluation of ovarian reserve with AMH level in patients with well-differentiated thyroid cancer receiving radioactive iodine ablation treatment. *Exp Clin Endocrinol Diabetes* 2016; **124**:593–596.

- Al-Janabi HT, Al-Taee HA, Alawad AS. The impact of age on anti-müllerian hormone serum level in women attending chemotherapy unit for primary breast cancer. *Middle East Fertil Soc J* 2018;**23**: 126–130.
- Anderson RA, Cameron DA. Pretreatment serum anti-müllerian hormone predicts long-term ovarian function and bone mass after chemotherapy for early breast cancer. *J Clin Endocrinol Metab* 2011;**96**:1336–1343.
- Anderson RA, Mansi J, Coleman RE, Adamson DJA, Leonard RCF. The utility of anti-Müllerian hormone in the diagnosis and prediction of loss of ovarian function following chemotherapy for early breast cancer. *Eur J Cancer* 2017;**87**:58–64.
- Anderson RA, Remedios R, Kirkwood AA, Patrick P, Stevens L, Clifton-Hadley L, Roberts T, Hatton C, Kalakonda N, Milligan DW et al. Determinants of ovarian function after response-adapted therapy in patients with advanced Hodgkin's lymphoma (RATHL): a secondary analysis of a randomised phase 3 trial. *Lancet Oncol* 2018;**19**:1328–1337.
- Anderson RA, Rosendahl M, Kelsey TW, Cameron DA. Pretreatment anti-Müllerian hormone predicts for loss of ovarian function after chemotherapy for early breast cancer. *Eur J Cancer* 2013;**49**:3404–3411.
- Anderson RA, Su HI. The clinical value and interpretation of anti-Müllerian hormone in women with cancer. *Front Endocrinol (Lausanne)* 2020;**11**:574263.
- Bath LE, Wallace WH, Shaw MP, Fitzpatrick C, Anderson RA. Depletion of ovarian reserve in young women after treatment for cancer in childhood: detection by anti-Müllerian hormone, inhibin B and ovarian ultrasound. *Hum Reprod* 2003;**18**:2368–2374.
- Behringer K, Mueller H, Goergen H, Thielen I, Eibl AD, Stumpf V, Wessels C, Wiehlputz M, Rosenbrock J, Halbsguth T et al. Gonadal function and fertility in survivors after Hodgkin lymphoma treatment within the German Hodgkin Study Group HD13 to HD15 trials. *J Clin Oncol* 2013;**31**:231–239.
- Ben-Aharon I, Granot T, Meizner I, Hasky N, Tobar A, Rizel S, Yerushalmi R, Ben-Haroush A, Fisch B, Stemmer SM. Long-term follow-up of chemotherapy-induced ovarian failure in young breast cancer patients: the role of vascular toxicity. *Oncologist* 2015;**20**: 985–991.
- Berjeb KK, Debbabi L, Braham M, Zemni Z, Chtourou S, Hannachi H, Hamdoun M, Ayadi M, Kacem K, Zhioua F et al. Evaluation of ovarian reserve before and after chemotherapy. *J Gynecol Obstet Hum Reprod* 2021;**50**:102035.
- Bi X, Zhang J, Cao D, Sun H, Feng F, Wan X, Xiang Y, Qiu L, Cheng X, Yang J et al. Anti-Müllerian hormone levels in patients with gestational trophoblastic neoplasia treated with different chemotherapy regimens: a prospective cohort study. *Oncotarget* 2017;**8**:113920–113927.
- Brougham MF, Crofton PM, Johnson EJ, Evans N, Anderson RA, Wallace WH. Anti-Müllerian hormone is a marker of gonadotoxicity in pre- and postpubertal girls treated for cancer: a prospective study. *J Clin Endocrinol Metab* 2012;**97**:2059–2067.
- Cameron K, Sammel MD, Prewitt M, Gracia C. Differential rates of change in measures of ovarian reserve in young cancer survivors across the reproductive lifespan. *J Clin Endocrinol Metab* 2019;**104**: 1813–1822.
- Cameron KE, Kole MB, Sammel MD, Ginsberg JP, Gosiengfiao Y, Mersereau JE, Su HI, Gracia CR. Acute menopausal symptoms in young cancer survivors immediately following chemotherapy. *Oncology* 2018;**94**:200–206.
- Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, Hartmann-Boyce J, Ryan R, Shepperd S, Thomas J et al. Synthesis without meta-analysis (SWiM) in systematic reviews: reporting guideline. *BMJ* 2020;**368**:l6890.
- Celebi F, Ordu C, Ilgun S, Ozturk A, Erdogan Iyigun Z, Alco G, Duymaz T, Aktepe F, Soybir G, Baysal B et al.; The effect of systemic chemotherapy on ovarian function: a prospective clinical trial. *Eur J Breast Health* 2020;**16**:177–182.
- Chai J, Howie AF, Cameron DA, Anderson RA. A highly-sensitive anti-Müllerian hormone assay improves analysis of ovarian function following chemotherapy for early breast cancer. *Eur J Cancer* 2014;**50**:2367–2374.
- Charpentier A-M, Chong AL, Gingras-Hill G, Ahmed S, Cigsar C, Gupta AA, Greenblatt E, Hodgson DC. Anti-Müllerian hormone screening to assess ovarian reserve among female survivors of childhood cancer. *J Cancer Surviv* 2014;**8**:548–554.
- Chemersinski A, Cameron K, Sammel M, Ginsberg J, Carlson C, Gracia C. Relationship of menopausal symptoms and ovarian reserve in reproductive-aged cancer survivors. *J Cancer Surviv* 2020;**14**:607–613.
- Dafopoulos A, Dafopoulos K, Georgoulas P, Galazios G, Limberis V, Tsikouras P, Koutlaki N, Maroulis G. Smoking and AMH levels in women with normal reproductive history. *Arch Gynecol Obstet* 2010;**282**:215–219.
- D'Avila ÂM, Biolchi V, Capp E, Corleta HVE. Age, anti-müllerian hormone, antral follicles count to predict amenorrhea or oligomenorrhea after chemotherapy with cyclophosphamide. *J Ovarian Res* 2015;**8**:82.
- D'Avila ÂM, Capp E, Corleta HVE. Antral follicles count and anti-Müllerian hormone levels after gonadotoxic chemotherapy in patients with breast cancer: cohort study. *Rev Bras Ginecol Obstet* 2017;**39**:162–168.
- de Kat AC, van der Schouw YT, Eijkemans MJC, Broer SL, Verschuren WMM, Broekmans FJM. Can menopause prediction be improved with multiple AMH measurements? Results from the prospective Doetinchem cohort study. *J Clin Endocrinol Metab* 2019;**104**:5024–5031.
- Decanter C, Cloquet M, Dassonneville A, D'Orazio E, Mailliez A, Pigny P. Different patterns of ovarian recovery after cancer treatment suggest various individual ovarian susceptibilities to chemotherapy. *Reprod Biomed Online* 2018;**36**:711–718.
- Decanter C, Delepine J, Behal H, Manier S, Bruno B, Barbatti M, Robin C, Labreuche J, Morschhauser F, Pigny P. Longitudinal study of AMH variations in 122 adolescents and young adults (AYA) and non-AYA lymphoma patients to evaluate the chemo-induced ovarian toxicity to further personalise fertility preservation counselling. *Hum Reprod* 2021;**36**:2743–2752.
- Decanter C, Morschhauser F, Pigny P, Lefebvre C, Gallo C, Dewailly D. Anti-Müllerian hormone follow-up in young women treated by chemotherapy for lymphoma: preliminary results. *Reprod Biomed Online* 2010;**20**:280–285.
- Decanter C, Peigne M, Mailliez A, Morschhauser F, Dassonneville A, Dewailly D, Pigny P. Toward a better follow-up of ovarian

- recovery in young women after chemotherapy with a hypersensitive antimüllerian hormone assay. *Fertil Steril* 2014;**102**:483–487.
- Demeestere I, Brice P, Peccatori FA, Kentos A, Dupuis J, Zachee P, Casanovas O, Van Den Neste E, Dechene J, De Maertelaer V et al. No evidence for the benefit of gonadotropin-releasing hormone agonist in preserving ovarian function and fertility in lymphoma survivors treated with chemotherapy: final long-term report of a prospective randomized trial. *J Clin Oncol* 2016;**34**:2568–2574.
- Demeestere I, Racape J, Dechene J, Dupuis J, Morschhauser F, De Wilde V, Lazarovici J, Ghesquieres H, Touati M, Sibon D et al. Gonadal function recovery in patients with advanced Hodgkin lymphoma treated with a PET-adapted regimen: prospective analysis of a randomized phase III trial (AHL2011). *J Clin Oncol* 2021;**39**:3251–3260.
- Dezellus A, Barriere P, Campone M, Lemanski C, Vanlemmens L, Mignot L, Delozier T, Levy C, Bendavid C, Debled M et al. Prospective evaluation of serum anti-Müllerian hormone dynamics in 250 women of reproductive age treated with chemotherapy for breast cancer. *Eur J Cancer* 2017;**79**:72–80.
- Di Paola R, Costantini C, Tecchio C, Salvagno GL, Montemezzi R, Perandini A, Pizzolo G, Zaffagnini S, Franchi M. Anti-Müllerian hormone and antral follicle count reveal a late impairment of ovarian reserve in patients undergoing low-gonadotoxic regimens for hematological malignancies. *Oncologist* 2013;**18**:1307–1314.
- Dillon KE, Sammel MD, Ginsberg JP, Lechtenberg L, Prewitt M, Gracia CR. Pregnancy after cancer: results from a prospective cohort study of cancer survivors. *Pediatr Blood Cancer* 2013a;**60**:2001–2006.
- Dillon KE, Sammel MD, Prewitt M, Ginsberg JP, Walker D, Mersereau JE, Gosiengfiao Y, Gracia CR. Pretreatment antimüllerian hormone levels determine rate of posttherapy ovarian reserve recovery: acute changes in ovarian reserve during and after chemotherapy. *Fertil Steril* 2013b;**99**:477–483.
- Dólleman M, Verschuren WMM, Eijkemans MJC, Dollé MET, Jansen EHJM, Broekmans FJM, van der Schouw YT. Reproductive and lifestyle determinants of anti-Müllerian hormone in a large population-based study. *J Clin Endocrinol Metab* 2013;**98**:2106–2115.
- Elchuri SV, Patterson BC, Brown M, Bedient C, Record E, Wasilewski-Masker K, Mertens AC, Meacham LR. Low anti-Müllerian hormone in pediatric cancer survivors in the early years after gonadotoxic therapy. *J Pediatr Adolesc Gynecol* 2016;**29**:393–399.
- Elgindy EA, El-Haieg DO, Khorshid OM, Ismail El, Abdelgawad M, Sallam HN, Abou-Setta AM. Gonadotrophin suppression to prevent chemotherapy-induced ovarian damage: a randomized controlled trial. *Obstet Gynecol* 2013;**121**:78–86.
- Eltzur S, Frank S, Goshen-Lago T, Barzilai-Birenboim S, Gilad G, Avrahami G, Goldberg T, Litichever N, Masarwa A, Oron G et al. Long-term ovarian reserve and fertility outcomes in female survivors of childhood acute lymphoblastic leukemia. *Leuk Lymphoma* 2021;**62**:2211–2218.
- El-Shalakany AH, Ali MS, Abdelmaksoud AA, Abd El-Ghany S, Hasan EA. Ovarian function in female survivors of childhood malignancies. *Pediatr Hematol Oncol* 2013;**30**:328–335.
- Evrano B, Faki S, Polat SB, Bestepe N, Ersoy R, Cakir B. Effects of radioactive iodine therapy on ovarian reserve: a prospective pilot study. *Thyroid* 2018;**28**:1702–1707.
- Finkelstein JS, Lee H, Karlamangla A, Neer RM, Sluss PM, Burnett-Bowie SM, Darakananda K, Donahoe PK, Harlow SD, Prizand SH et al. Antimüllerian hormone and impending menopause in late reproductive age: the study of women's health across the nation. *J Clin Endocrinol Metab* 2020;**105**:e1862–e1871.
- George SA, Williamson Lewis R, Schirmer DA, Effinger KE, Spencer JB, Mertens AC, Meacham LR. Early detection of ovarian dysfunction by anti-Müllerian hormone in adolescent and young adult-aged survivors of childhood cancer. *J Adolesc Young Adult Oncol* 2019;**8**:18–25.
- Gharwan H, Lai C, Grant C, Dunleavy K, Steinberg SM, Shovlin M, Fojo T, Wilson WH. Female fertility following dose-adjusted EPOCH-R chemotherapy in primary mediastinal B-cell lymphomas. *Leuk Lymphoma* 2016;**57**:1616–1624.
- Goldfarb SB, Turan V, Bedoschi G, Taylan E, Abdo N, Cigler T, Bang H, Patil S, Dickler MN, Oktay KH. Impact of adjuvant chemotherapy or tamoxifen-alone on the ovarian reserve of young women with breast cancer. *Breast Cancer Res Treat* 2021;**185**:165–173.
- Gracia CR, Sammel MD, Freeman E, Prewitt M, Carlson C, Ray A, Vance A, Ginsberg JP. Impact of cancer therapies on ovarian reserve. *Fertil Steril* 2012;**97**:134–140.e131.
- Gupta AA, Lee Chong A, Deveault C, Traubici J, Maloney AM, Knight S, Lorenzo A, Allen L. Anti-Müllerian hormone in female adolescent cancer patients before, during, and after completion of therapy: a pilot feasibility study. *J Pediatr Adolesc Gynecol* 2016;**29**:599–603.
- Hamre H, Kiserud CE, Ruud E, Thorsby PM, Fossa SD. Gonadal function and parenthood 20 years after treatment for childhood lymphoma: a cross-sectional study. *Pediatr Blood Cancer* 2012;**59**:271–277.
- Hamy AS, Porcher R, Eskenazi S, Cuvier C, Giacchetti S, Coussy F, Hocini H, Tournant B, Perret F, Bonfils S et al. Anti-Müllerian hormone in breast cancer patients treated with chemotherapy: a retrospective evaluation of subsequent pregnancies. *Reprod Biomed Online* 2016;**32**:299–307.
- Henry NL, Xia R, Schott AF, McConnell D, Banerjee M, Hayes DF. Prediction of postchemotherapy ovarian function using markers of ovarian reserve. *Oncologist* 2014;**19**:68–74.
- Iwase A, Sugita A, Hirokawa W, Goto M, Yamamoto E, Takikawa S, Nakahara T, Nakamura T, Kondo M, Kikkawa F. Anti-Müllerian hormone as a marker of ovarian reserve following chemotherapy in patients with gestational trophoblastic neoplasia. *Eur J Obstet Gynecol Reprod Biol* 2013;**167**:194–198.
- Jayasinghe YL, Wallace WHB, Anderson RA. Ovarian function, fertility and reproductive lifespan in cancer patients. *Expert Rev Endocrinol Metab* 2018;**13**:125–136.
- Kelsey TW, Wright P, Nelson SM, Anderson RA, Wallace WHB. A validated model of serum anti-Müllerian hormone from conception to menopause. *PLoS One* 2011;**6**:e22024.
- Kim HA, Choi J, Park CS, Seong MK, Hong SE, Kim JS, Park IC, Lee JK, Noh WC; The ASTRRA trial investigators. Post-chemotherapy serum anti-Müllerian hormone level predicts ovarian function recovery. *Endocr Connect* 2018;**7**:949–956.
- Krawczuk-Rybak M, Leszczynska E, Poznanska M, Zelazowska-Rutkowska B, Wysocka J. Anti-Müllerian hormone as a sensitive marker of ovarian function in young cancer survivors. *Int J Endocrinol* 2013a;**2013**:125080.

- Krawczuk-Rybak M, Leszczynska E, Poznanska M, Zelazowska-Rutkowska B, Wysocka J. The progressive reduction in the ovarian reserve in young women after anticancer treatment. *Horm Metab Res* 2013b;**45**:813–819.
- Krawczuk-Rybak M, Płonowski M, Leszczyńska E, Latoch E, Sawicka-Żukowska M, Muszyńska-Roslan K, Skalska-Sadowska J, Wachowiak J, Sga-Pondel D, Kazanowska B et al. The influence of different intensity of treatment on hormonal markers of gonadal function in acute lymphoblastic leukemia survivors. *Hematol Oncol* 2019;**37**:609–616.
- Lambertini M, Olympios N, Lequesne J, Calbrix C, Fontanilles M, Loeb A, Leheurteur M, Demeestere I, Di Fiore F, Perdrix A et al. Impact of taxanes, endocrine therapy, and deleterious germline BRCA mutations on anti-mullerian hormone levels in early breast cancer patients treated with anthracycline- and cyclophosphamide-based chemotherapy. *Front Oncol* 2019;**9**:575.
- Lee DY, Kim JY, Yu J, Kim SW. Prediction of successful ovarian protection using gonadotropin-releasing hormone agonists during chemotherapy in young estrogen receptor-negative breast cancer patients. *Front Oncol* 2020;**10**:863.
- Lee DY, Park YH, Lee JE, Choi D. Prediction of ovarian function recovery in young breast cancer patients after protection with gonadotropin-releasing hormone agonist during chemotherapy. *Breast Cancer Res Treat* 2018;**171**:649–656.
- Leiper A, Houwing M, Davies EG, Rao K, Burns S, Morris E, Laven J, van der Kooi AL, van den Heuvel Eibrink M, Nussey S. Anti-Mullerian hormone and Inhibin B after stem cell transplant in childhood: a comparison of myeloablative, reduced intensity and treosulfan-based chemotherapy regimens. *Bone Marrow Transplant* 2020;**55**:1985–1995.
- Leonard RCF, Adamson DJA, Bertelli G, Mansi J, Yellowlees A, Dunlop J, Thomas GA, Coleman RE, Anderson RA. GnRH agonist for protection against ovarian toxicity during chemotherapy for early breast cancer: the Anglo Celtic Group OPTION trial. *Ann Oncol* 2017;**28**:1811–1816.
- Li HWR, Robertson DM, Burns C, Ledger WL. Challenges in measuring AMH in the clinical setting. *Front Endocrinol (Lausanne)* 2021;**12**:691432.
- Li X, Liu S, Ma L, Chen X, Weng H, Huang R, Yu Y, Zong X. Can anti-Müllerian hormone be a reliable biomarker for assessing ovarian function in women postchemotherapy. *Cancer Manag Res* 2020;**12**:8171–8181.
- Lie Fong S, Laven JSE, Hakvoort-Cammel FGJ, Schipper I, Visser JA, Themmen APN, de Jong FH, van den Heuvel-Eibrink MM. Assessment of ovarian reserve in adult childhood cancer survivors using anti-Mullerian hormone. *Hum Reprod* 2009;**24**:982–990.
- Loubersac S, Dezellus A, Lefebvre T, Reignier A, Barriere P, Masson D, Freour T; RESOVA Investigators Group. Evolution of serum Anti-Mullerian Hormone (AMH) level in young women treated with chemotherapy for breast cancer according to basal AMH level. *Eur J Obstet Gynecol Reprod Biol* 2020;**254**:132–137.
- Lunsford AJ, Whelan K, McCormick K, McLaren JF. Antimullerian hormone as a measure of reproductive function in female childhood cancer survivors. *Fertil Steril* 2014;**101**:227–231.
- Lutchman Singh K, Muttukrishna S, Stein RC, McGarrigle HH, Patel A, Parikh B, Groome NP, Davies MC, Chatterjee R. Predictors of ovarian reserve in young women with breast cancer. *Br J Cancer* 2007;**96**:1808–1816.
- Malisic E, Susnjar S, Milovanovic J, Todorovic-Rakovic N, Kesic V. Assessment of ovarian function after chemotherapy in women with early and locally advanced breast cancer from Serbia. *Arch Gynecol Obstet* 2018;**297**:495–503.
- Martin HL, Ullah S, Abbas N, Scott-Hoy A, Kichenadasse G, Karapetis CS, Roy A, Sukumaran S, Ross DM, Khattak MA et al. Predicting chemotherapy-induced menopause using baseline and post-chemotherapy anti-Müllerian hormone levels: Results of a pilot study. *Cancer Rep* 2021;**4**:e11342.
- Mittica M, Dotto A, Comina M, Teliti M, Monti E, Giusti M. Cross-sectional and prospective study on anti-Mullerian hormone changes in a cohort of pre-menopausal women with a history of differentiated thyroid cancer. *Thyroid Res* 2020;**13**:1.
- Miyoshi Y, Ohta H, Namba N, Tachibana M, Miyamura T, Miyashita E, Hashii Y, Oue T, Isobe A, Tsutsui T et al. Low serum concentrations of anti-Mullerian hormone are common in 53 female childhood cancer survivors. *Horm Res Paediatr* 2013;**79**:17–21.
- Morarji K, McArdle O, Hui K, Gingras-Hill G, Ahmed S, Greenblatt EM, Warner E, Sridhar S, Ali AMF, Azad A et al. Ovarian function after chemotherapy in young breast cancer survivors. *Curr Oncol* 2017;**24**:e494–e502.
- Morse H, Elfving M, Turkiewicz A, Andersen CY, Ora I. Severe gonadotoxic insult manifests early in young girls treated for Ewing sarcoma. *Medicine (Baltimore)* 2016;**95**:e4512.
- Nielsen SN, Andersen AN, Schmidt KT, Rechnitzer C, Schmiegelow K, Bentzen JG, Larsen EC. A 10-year follow up of reproductive function in women treated for childhood cancer. *Reprod Biomed Online* 2013;**27**:192–200.
- Nies M, Cantineau AEP, Arts E, van den Berg MH, van Leeuwen FE, Muller Kobold AC, Klein Hesselink MS, Burgerhof JGM, Brouwers AH, van Dam E et al. Long-term effects of radioiodine treatment on female fertility in survivors of childhood differentiated thyroid carcinoma. *Thyroid* 2020;**30**:1169–1176.
- Nitzschke M, Raddatz J, Bohlmann MK, Stute P, Strowitzki T, von Wolff M. GnRH analogs do not protect ovaries from chemotherapy-induced ultrastructural injury in Hodgkin's lymphoma patients. *Arch Gynecol Obstet* 2010;**282**:83–88.
- Nystrom A, Morse H, Nordlof H, Wiebe K, Artman M, Ora I, Giwercman A, Henic E, Elfving M. Anti-mullerian hormone compared with other ovarian markers after childhood cancer treatment. *Acta Oncol* 2019;**58**:218–224.
- Oktay KH, Bedoschi G, Goldfarb SB, Taylan E, Titus S, Palomaki GE, Cigler T, Robson M, Dickler MN. Increased chemotherapy-induced ovarian reserve loss in women with germline BRCA mutations due to oocyte deoxyribonucleic acid double strand break repair deficiency. *Fertil Steril* 2020;**113**:1251–1260.e1251.
- Palinska-Rudzka KE, Ghobara T, Parsons N, Milner J, Lockwood G, Hartshorne GM. Five-year study assessing the clinical utility of anti-Mullerian hormone measurements in reproductive-age women with cancer. *Reprod Biomed Online* 2019;**39**:712–720.
- Panay N, Anderson RA, Nappi RE, Vincent AJ, Vujovic S, Webber L, Wolfman W. Premature ovarian insufficiency: an International Menopause Society White Paper. *Climacteric* 2020;**23**:426–446.
- Parissone F, Di Paola R, Balter R, Garzon S, Zaffagnini S, Neri M, Vitale V, Tridello G, Cesaro S. Female adolescents and young



- women previously treated for pediatric malignancies: assessment of ovarian reserve and gonadotoxicity risk stratification for early identification of patients at increased infertility risk. *J Pediatr Endocrinol Metab* 2021;**34**:25–33.
- Partridge AH, Ruddy KJ, Gelber S, Schapira L, Abusief M, Meyer M, Ginsburg E. Ovarian reserve in women who remain premenopausal after chemotherapy for early stage breast cancer. *Fertil Steril* 2010;**94**:638–644.
- Passildas J, Collard O, Savoye AM, Dohou J, Ginzac A, Thivat E, Durando X, Kwiatkowski F, Penault-Llorca F, Abrial C et al. Impact of chemotherapy-induced menopause in women of child-bearing age with non-metastatic breast cancer - preliminary results from the MENOCOR study. *Clin Breast Cancer* 2019;**19**:e74–e84.
- Ramezani Tehrani F, Bidhendi Yarandi R, Solaymani-Dodaran M, Tohidi M, Firouzi F, Azizi F. Improving prediction of age at menopause using multiple anti-Mullerian hormone measurements: the Tehran lipid-glucose study. *J Clin Endocrinol Metab* 2020;**105**:dgaa083.
- Rosendahl M, Andersen CY, Ernst E, Westergaard LG, Rasmussen PE, Loft A, Andersen AN. Ovarian function after removal of an entire ovary for cryopreservation of pieces of cortex prior to gonadotoxic treatment: a follow-up study. *Hum Reprod* 2008;**23**:2475–2483.
- Rosendahl M, Andersen CY, la Cour Freiesleben N, Juul A, Lossl K, Andersen AN. Dynamics and mechanisms of chemotherapy-induced ovarian follicular depletion in women of fertile age. *Fertil Steril* 2010;**94**:156–166.
- Roshandel R, van Dijk M, Overbeek A, Kaspers G, Lambalk C, Beerendonk C, Bresters D, van der Heiden-van der Loo M, van den Heuvel-Eibrink M, Kremer L et al.; LATER-VEVO Study Group. Female reproductive function after treatment of childhood acute lymphoblastic leukemia. *Pediatr Blood Cancer* 2021;**68**:e28894.
- Ruddy KJ, O'Neill A, Miller KD, Schneider BP, Baker E, Sparano JA, Dang C, Northfelt DW, Sledge GW Jr, Partridge AH. Biomarker prediction of chemotherapy-related amenorrhea in premenopausal women with breast cancer participating in E5103. *Breast Cancer Res Treat* 2014;**144**:591–597.
- Ruddy KJ, Schaid DJ, Batzler A, Cecchini RS, Partridge AH, Norman A, Fehrenbacher L, Stewart EA, Trabuco E, Ginsburg E et al. Anti-mullerian hormone as a serum biomarker for risk of chemotherapy-induced amenorrhea. *J Natl Cancer Inst* 2021;**113**:1105–1108.
- Shandley LM, Spencer JB, Fothergill A, Mertens AC, Manatunga A, Paplomata E, Howards PP. Impact of tamoxifen therapy on fertility in breast cancer survivors. *Fertil Steril* 2017;**107**:243–252.e245.
- Silva C, Caramelo O, Almeida-Santos T, Ribeiro Rama AC. Factors associated with ovarian function recovery after chemotherapy for breast cancer: a systematic review and meta-analysis. *Hum Reprod* 2016;**31**:2737–2749.
- Silva C, Ribeiro Rama AC, Reis Soares S, Moura-Ramos M, Almeida-Santos T. Adverse reproductive health outcomes in a cohort of young women with breast cancer exposed to systemic treatments. *J Ovarian Res* 2019;**12**:102.
- Spears N, Lopes F, Stefansdottir A, Rossi V, De Felici M, Anderson RA, Klinger FG. Ovarian damage from chemotherapy and current approaches to its protection. *Hum Reprod Update* 2019;**25**:673–693.
- Steiner AZ, Pritchard D, Stanczyk FZ, Kesner JS, Meadows JW, Herring AH, Baird DD. Association between biomarkers of ovarian reserve and infertility among older women of reproductive age. *JAMA* 2017;**318**:1367–1376.
- Sterne JA, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, Henry D, Altman DG, Ansari MT, Boutron I et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016;**355**:i4919.
- Su HI, Kwan B, Whitcomb BW, Shliakhitsava K, Dietz AC, Stark SS, Martinez E, Sluss PM, Sammel MD, Natarajan L. Modeling variation in the reproductive lifespan of female adolescent and young adult cancer survivors using AMH. *J Clin Endocrinol Metab* 2020;**105**:2740–2751.
- Su HI, Sammel MD, Green J, Velders L, Stankiewicz C, Matro J, Freeman EW, Gracia CR, DeMichele A. Antimullerian hormone and inhibin B are hormone measures of ovarian function in late reproductive-aged breast cancer survivors. *Cancer* 2010;**116**:592–599.
- Thomas-Teinturier C, Allodji RS, Svetlova E, Frey MA, Oberlin O, Millischer AE, Epelboin S, Decanter C, Pacquement H, Tabone MD et al. Ovarian reserve after treatment with alkylating agents during childhood. *Hum Reprod* 2015;**30**:1437–1446.
- Trapp E, Steidl J, Rack B, Kupka MS, Andergassen U, Juckstock J, Kurt A, Vilsmaier T, de Gregorio A, de Gregorio N et al. Anti-Mullerian hormone (AMH) levels in premenopausal breast cancer patients treated with taxane-based adjuvant chemotherapy - a translational research project of the SUCCESS A study. *Breast* 2017;**35**:130–135.
- van Beek RD, van den Heuvel-Eibrink MM, Laven JS, de Jong FH, Themmen AP, Hakvoort-Cammel FG, van den Bos C, van den Berg H, Pieters R, de Muinck Keizer-Schrama SM. Anti-Mullerian hormone is a sensitive serum marker for gonadal function in women treated for Hodgkin's lymphoma during childhood. *J Clin Endocrinol Metab* 2007;**92**:3869–3874.
- van den Berg MH, Overbeek A, Lambalk CB, Kaspers GJL, Bresters D, van den Heuvel-Eibrink MM, Kremer LC, Loonen JJ, van der Pal HJ, Ronckers CM et al.; DCOG LATER-VEVO study group. Long-term effects of childhood cancer treatment on hormonal and ultrasound markers of ovarian reserve. *Hum Reprod* 2018;**33**:1474–1488.
- van der Kooi ALF, van den Heuvel-Eibrink MM, van den Berg SAA, van Dorp W, Pluijm SMF, Laven JSE. Changes in anti-Mullerian hormone and inhibin B in children treated for cancer. *J Adolesc Young Adult Oncol* 2019;**8**:281–290.
- van der Kooi ALF, van den Heuvel-Eibrink MM, van Noortwijk A, Neggers SJ, Pluijm SM, van Dulmen-den Broeder E, van Dorp W, Laven JS. Longitudinal follow-up in female Childhood Cancer Survivors: no signs of accelerated ovarian function loss. *Hum Reprod* 2017;**32**:193–200.
- van Dorp W, Haupt R, Anderson RA, Mulder RL, van den Heuvel-Eibrink MM, van Dulmen-den Broeder E, Su HI, Winther JF, Hudson MM, Levine JM et al. Reproductive function and outcomes in female survivors of childhood, adolescent, and young adult cancer: a review. *J Clin Oncol* 2018;**36**:2169–2180.

- van Velsen EFS, Visser WE, van den Berg SAA, Kam BLR, van Ginhoven TM, Massolt ET, Peeters RP. Longitudinal analysis of the effect of radioiodine therapy on ovarian reserve in females with differentiated thyroid cancer. *Thyroid* 2020;**30**:580–587.
- Webber L, Davies M, Anderson R, Bartlett J, Braat D, Cartwright B, Cifkova R, de Muinck Keizer-Schrama S, Hogervorst E, Janse F et al.; European Society for Human Reproduction and Embryology (ESHRE) Guideline Group on POI. ESHRE Guideline: management of women with premature ovarian insufficiency. *Hum Reprod* 2016;**31**:926–937.
- Wenners A, Grambach J, Koss J, Maass N, Jonat W, Schmutzler A, Mundhenke C. Reduced ovarian reserve in young early breast cancer patients: preliminary data from a prospective cohort trial. *BMC Cancer* 2017;**17**:632.
- Yaish I, Azem F, Gutfeld O, Silman Z, Serebro M, Sharon O, Shefer G, Limor R, Stern N, Tordjman KM. A single radioactive iodine treatment has a deleterious effect on ovarian reserve in women with thyroid cancer: results of a prospective pilot study. *Thyroid* 2018;**28**:522–527.
- Yu B, Douglas N, Ferin MJ, Nakhuda GS, Crew K, Lobo RA, Hershman DL. Changes in markers of ovarian reserve and endocrine function in young women with breast cancer undergoing adjuvant chemotherapy. *Cancer* 2010;**116**:2099–2105.
- Zhong Y, Lin Y, Cheng X, Huang X, Zhou Y, Mao F, Wang Y, Guan J, Shen S, Xu Y et al. GnRHa for ovarian protection and the association between AMH and ovarian function during adjuvant chemotherapy for breast cancer. *J Cancer* 2019;**10**:4278–4285.