



Petermann-Rocha, F. et al. (2022) Children who sleep more may have longer telomeres: evidence from a longitudinal population study in Spain. *Pediatric Research*, (doi: [10.1038/s41390-022-02255-w](https://doi.org/10.1038/s41390-022-02255-w)).

This is the Author Accepted Manuscript.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/278181/>

Deposited on: 18 November 2022

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

Children who sleep more may have longer telomeres: findings from the INMA birth cohort study

Fanny Petermann-Rocha^{1,2}, Desirée Valera-Gran^{*3,4}, Paula Fernandes-Pires³, Dries S. Martens⁵, Jordi Júlvez⁶, Cristina Rodríguez-Dehli^{7,8}, Ainara Andiarena^{9,10}, Manuel Lozano^{11,12}, Ana Fernández-Somoano^{8,13,14}, Aitana Lertxundi^{9,13,15}, Sabrina Llop^{12,13}, Mònica Guxens,^{13,16,17,18} Tim S. Nawrot⁵, Eva-María Navarrete-Muñoz^{3,4,19}

¹ Faculty of Medicine, Universidad Diego Portales, Santiago, Chile

² Institute of Cardiovascular Medical Sciences and Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK.

³ Department of Surgery and Pathology, Miguel Hernandez University, 03550 Alicante, Spain.

⁴ Grupo de Investigación en Terapia Ocupacional (InTeO), Miguel Hernández University, 03550 Alicante, Spain.

⁵ Centre for Environmental Sciences, Hasselt University, Hasselt, Belgium

⁶ Institut d'Investigació Sanitària Pere Virgili (IISPV), Hospital Universitari Sant Joan de Reus, Reus, Spain

⁷ Servicio de Pediatría. Hospital San Agustín. Heros Street, 4. 33410, Avilés, Asturias, Spain

⁸ Instituto de Investigación Sanitaria del Principado de Asturias (ISPA). Roma Avenue s/n. 33001, Oviedo. Spain.

⁹ Health Research Institute BIODONOSTIA, Donostia-San Sebastian, Basque Country, Spain.

¹⁰ Faculty of Psychology, University of the Basque Country (UPV/EHU), San Sebastian, Basque Country, Spain.

¹¹ Preventive Medicine and Public Health, Food Sciences, Toxicology and Forensic Medicine Department, Universitat de València, Valencia, Spain

¹² Epidemiology and Environmental Health Joint Research Unit, FISABIO–Universitat Jaume I–Universitat de València, Valencia, Spain

¹³ Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Madrid, Spain

¹⁴ Unidad de Epidemiología Molecular del Cáncer, Instituto Universitario de Oncología del Principado de Asturias (IUOPA)-Departamento de Medicina, Universidad de Oviedo. Julián Clavería Street, s/n 33006, Oviedo, Asturias, Spain

¹⁵Preventive medicine and public health department, University of Basque Country (UPV/EHU), Leioa, Basque Country, Spain

¹⁶ISGlobal, Barcelona, Spain

¹⁷Universitat Pompeu Fabra, Barcelona, Spain

¹⁸Department of Child and Adolescent Psychiatry/Psychology, Erasmus MC, University Medical Centre, Rotterdam, The Netherlands

¹⁹Department of Surgery and Pathology, Miguel Hernandez University, Alicante Institute for Health and Biomedical Research (ISABIAL), 03550, Alicante, Spain.

Corresponding author: Desirée Valera Gran. Department of Surgery and Pathology, Miguel Hernandez University, 03550 Alicante, Spain. Email: dvalera@umh.es. Telephone: +34 96.523.3705

Abstract

Inadequate sleep duration has been suggested as a chronic stressor associated with changes in telomere length. However, the association of these two variables in children is limited. This study aimed to explore the association between sleep duration and telomere length (TL) using the INMA birth cohort study data. A sample of 1014 children was included in this study (cross-sectional: 686; longitudinal: 872). Sleep duration (hours/day [h/day]) was reported by caregivers at age 4 and classified in three categories (7-10 h/day; >10-11 h/day; >11-14 h/day). Leucocyte TL at age 4 and age 7-9 were the two outcomes of interest. They were measured using quantitative PCR methods. Multiple robust linear regression models, through log-level regression models, were used to report the percentage (%) of difference among tertiles of sleep duration. In comparison to children who slept between >10 and 11 h/day, those in the highest category (> 11 h/day) had 8.5% (95% CI: 3.56-13.6) longer telomeres at age 4. In contrast, children in the lowest sleep category showed 2.2% longer leucocyte TL than their counterparts, but this association was non-significant ($p=0.162$). No significant associations were identified between sleep duration at age 4 and TL at age 7-9.

Keywords: Child; Telomere length; Sleep; Lifestyle

Total words: 2923

Introduction

Sleep is a necessary physiological process and has a critical role in promoting balanced health¹. In children, adequate sleep is associated with normal growth, wellbeing, and different development domains such as nutrition, hygiene, communication, and physical contact^{2,3}. Inadequate sleep, instead – defined mainly as the number of hours a child sleeps – negatively impacts cognitive functions, socioemotional domains, early childhood development, and physical health². The American Academy of Sleep Medicine recommends sleeping 10 to 13 hours per day for children between 3 and 5 years old to reach their full developmental potential³. However, not all children meet this recommendation. For instance, 34.9% of American children and adolescents aged 4 months to 17 years reported sleeping less than the recommendations for their age⁴. In Spain, Ruitter et al. estimated that sleep duration in children between 2 and 14 years had decreased by 20 minutes in the last decades and that only 55% of children were sleeping enough hours per day⁵.

In addition to the aforementioned consequences of sleep disturbance, inadequate sleep duration has been suggested as a chronic stressor associated with changes in telomere length⁶⁻⁸. Telomeres are nucleoprotein structures containing repeat sequences of tandem TTAGGG DNA stretches that protect chromosome ends from illicit DNA repair. Naturally, they shorten over time; however, they are susceptible to faster shortening under stressors. Previous studies have identified that shorter telomeres are associated with a higher risk of adverse health outcomes and have been identified as a useful ageing biomarker⁹.

Although studies evaluating telomere length in children are limited, previous works highlighted the association between childhood abuse, early life adversity, childhood socioeconomic status, and maternal factors (such as depression, smoking, and inheritance) with telomere length^{10,11}. Regarding sleep, studies conducted in adults have shown that poor sleep quality is associated with shorter telomere length⁶⁻⁸. In children, two studies evaluated the potential association between sleep duration and telomere length. However, results from both studies are inconclusive since one evidenced a positive association and the other no association between these two variables^{12,13}. Considering that the literature has proposed that the environmental conditions during adulthood might have less impact over telomere length than those during childhood^{14,15} – and the poorly investigated role of

sleep in children – this study aimed to explore the association between sleep duration and telomere length using data from the INMA birth cohort study.

Methods

This study was carried out using data from the INMA (INfancia y Medio Ambiente, Environment and Childhood) birth cohort study. The INMA project's main aim is to investigate the role of environmental factors during pregnancy and early life and their effects on child growth and development. More details about the INMA project can be found online <https://www.proyectoinma.org/> and have been published elsewhere¹⁶.

In brief, pregnant women from the general population were recruited between 2004 and 2008 in four areas of Spain (Asturias, Gipuzkoa, Sabadell and Valencia) using the following inclusion criteria: ≥16 years-old, singleton pregnancy, no assisted conception, intention to deliver at the reference hospital and to have no communication problems. Of the original sample, 1014 children had available data on the exposure (sleep at age 4), at least one of the outcomes (telomere length at 4 years, or at 7-to- 9 years; 'hereafter: 7-9'), and covariates and were, therefore, included in the analyses (Figure 1). Of them, 686 and 875 children had information for telomere length at 4 and 7-9 years, respectively (Figure 1).

Ethics declarations

The regional Ethical Committees approved the INMA birth cohort study. Written informed consent was obtained from all participants. This study complies with the Helsinki declaration for human studies.

Sleep categories – exposure

Caregivers (parents/legal tutors) reported child's sleep time (hours per day [h/day]) in the assessment carried out at age 4 using questionnaires. During the evaluation, the examiner asked 'how many hours does your child sleep during the week (h/day)?' and 'how many hours does your child sleep during the weekend (h/day)?' The average sleep per day was estimated as the sum of hours during the week and weekend divided by seven using these two questions as follows: $((\text{weekday sleep time} \times 5) + (\text{weekend sleep time} \times 2)) / 7$. The American Academy of Sleep Medicine recommends children sleep between 10 and 13 hours/day at age 4³. However, only 7 children were reported as sleeping more than 13

h/day whilst 89 less than 10 h/day, representing a relatively homogeneous sample. Consequently, the average per day was categorized in tertiles following the participants' distribution as follows: i) 7 to 10 h/day, ii) 10.02 to 11 h/day (hereafter, '>10 to 11 h/day'), and iii) 11.03 to 14 h/day (hereafter, '>11 to 14 h/day').

Leucocyte telomere length – outcome

Leucocyte telomere length (LTL) at age 4 was available in Gipuzkoa, Sabadell and Asturias (average age: 4.4 years, standard deviation [SD] 0.2 years; interquartile range 4.4-4.5 years). The cross-sectional analysis was restricted to these participants only (Figure 1). On the other hand, Gipuzkoa, Asturias and Valencia had available telomere data at age 7 while Sabadell at age 9 only (average age: 8.2 years; SD: 0.6 years; interquartile range: 7.7-9.2 years). Therefore, these outcomes were pooled together to create the variable LTL at age 7-9. The longitudinal analysis was restricted to these participants only (Figure 1).

Blood samples were collected during clinical examination and adequately stored in EDTA tubes. At age 4, DNA was extracted from blood using the Flexigen AGKT-WB-640 (Qiagen) kit in Gipuzkoa samples, Chemagen kit (Perkin Elmer) in Sabadell and from buffy coat applying the QIAamp DNA Mini Kit (Qiagen) in Asturias. At age 7-9, DNA was extracted from buffy coats using the aforementioned kits for Gipuzkoa, Sabadell and Asturias. In Valencia, DNA was extracted from buffy coats using the Chemagen kit (Perkin Elmer)¹⁷.

As described in supplementary methods, LTL was determined using quantitative PCR methods. Different single-copy gene primers were used to assess LTL at 9 years in the Sabadell cohort samples¹⁸. Relative Leucocyte telomere length was determined separately for each cohort and normalized separately using qBase software (Biogazelle, Zwijnaarde, Belgium) and expressed as the ratio of telomere copy number to single-copy gene number (T/S) relative to the average T/S ratio of the cohort sample set. The reliability of the applied protocol was assessed by interclass correlation coefficients of triplicate measures (T/S ratios, telomere copy number and single-copy gene number measures).

Covariates

Age (calculated from the date of birth and assessment at 4 years), sex (female or male), the cohort of origin (Gipuzkoa, Sabadell, Asturias, or Valencia), blood extraction date (the day

when telomere information was collected; then codified as the season of extraction), mother's social class, parity (number of previous children, classified as 0 or ≥ 1), adherence to a relative Mediterranean Diet Score (rMED) and television (TV) time (reported by the caregivers regarding the total hours during the week and weekend watching TV/videos, i.e., screen time) were the covariates included in the main analyses. rMED was previously published in children¹⁹ and is based on the Buckland et al. index excluding alcohol consumption²⁰ since this study was restricted to children. The dietary index was calculated using the food intake of a validated food frequency questionnaire of eight components: vegetables (excluding potatoes), fruit (including nuts, seeds, and fruit juices), legumes, cereals (including whole grains and bread), fish (including seafood), meat (including processed meat), dairy products (including low-fat and high-fat products), and olive oil. Each rMED component was calculated in grams per 1000 kcal/day and divided into tertiles of intake.

Statistical analyses

Descriptive characteristics by children's sleep categories are presented as median with their respective interquartile range for quantitative variables. For categorical variables, data are reported as frequencies with their respective percentages. The distribution of the continuous variables was checked using the Lilliefors correction of Kolmogorov–Smirnov test and compared using ANOVA or Kruskal-Wallis, and Chi-square tests, both for main covariates and additional descriptive variables used in the sensitivity analyses.

Associations were initially analyzed using meta-analytic techniques to obtain combined estimates to quantify the heterogeneity among the study cohorts. The heterogeneity was quantified using I^2 statistics in R²¹. Since all I^2 values obtained for the primary outcomes were $< 50\%$; we analyzed adding the cohort variable to the adjustment of all the models (data not shown).

Associations between sleep categories and LTL at age 4 were investigated using multiple robust linear regression models, through log-level regression models, where the LTL was \log_{10} -transformed. Therefore, the results are reported as % difference and their respective 95% CI. Children whose parents/tutors reported sleeping between >10 and 11 h/day were

used as the reference group. Same analyzes were performed when LTL at age 7-9 was used as the outcome of interest.

All analyses were adjusted using three incremental models: Model 1, adjusted for blood date (at age 4 or age 7-9, according to the outcome of interest), cohort, age at sleep assessment and sex of the child. Model 2: as per model 1, but also for social class and parity of the mother at baseline assessment. Model 3: as per model 2, but also for the rMED and TV time. These potential confounder factors were selected based on previous literature and also in those variables with p-values <0.20 in the individual bivariate analyses at age 4 and 7-9 and those that changed the magnitude of the main effect by 10% using a backwards-forward elimination procedure²².

Finally, to investigate whether the association differed by sex, the analyses were repeated and stratified by sex (male and female) using the maximally adjusted model.

R 4.0.5 (packages 'robustbase', 'nortest', 'meta', 'lmtest', 'foreign', 'car', 'gdata') and Stata 17 were used to perform the statistical analyses. A p-value lower than 0.05 was considered statistically significant.

Results

Characteristics of the sample

Cohort characteristics by sleep categories are presented in Table 1. A total of 489 (48.2%) caregivers reported their children slept between 7-10 h/day, while only 132 (13.0%) children were reported sleeping more than 11 hours per day. Overall, and compared to those in the lowest sleep category (7-10 h/day), children who slept more than 11 hours per day were more likely to be male and from Asturias and their mothers were more likely to belong to a lower social class (IV +V). They also tended to have a better rMED and watch fewer TV hours per week. More information regarding the children's characteristics is available in Table 1.

Associations between sleep duration and LTL

Associations between sleep categories and LTL measured as a percentage difference at ages 4 and 7-9 are presented in Table 2. Compared to those in the medium category (sleep between >10 and 11 h/day), children in the highest category (more than 11 h/day) had 6.9%

(95% CI: 1.94-12.1) longer telomeres at age 4 (model 1). After further adjusting the model for other sociodemographic and lifestyle factors (models 2 and 3), the percentage difference at age 4 was even higher in this group (% difference: 8.48 [95% CI:3.56-13.6]). On the other hand, children in the lowest sleep category showed 2.2% longer LTL compared to their counterparts (model 3); however, this association was non-significant ($p=0.162$). Regarding sleep duration at age 4 and LTL at age 7-9, both children in the lowest and highest category showed longer LTL when compared to the models that evaluated LTL at the age of 4 years (Table 2).

Exploring differences in the association between sleep duration and LTL by sex

Finally, when the associations were stratified by sex, similar patterns of associations were identified for sleep at age 4 and telomere at age 4 (Supplementary Table 1). In the cross-sectional analyses, analyses remained significant for both sexes. Yet, boys had longer telomeres at age 4 than their counterparts (% difference _{boys}: 10% [95% CI: 2.97-17.8] and % difference _{girls}: 7.03% [95% CI: 0.24-14.3]). No differences were identified in the longitudinal analysis by sex (Supplementary Table 1).

Discussion

This study showed that, compared with children whose caregivers reported they slept between >10 and 11 h/day, LTL was longer in those in the highest sleep category (>11 h/day) independently of a wide range of confounder factors at age 4. This finding remained consistent by sex. Notwithstanding the above, we did not observe a significant association between sleep duration at age 4 and LTL later in childhood (age 7-9). Other unmeasured confounder factors could also explain the lack of significant association at this age. Yet, as previous authors have proposed, environmental conditions might have a major impact on telomere length during the first years of life¹⁴.

During the first 4 years of life, there is a rapid decline in LTL because of proliferative cells' increased turnover associated with growth¹⁴. Sleep is a unique window of opportunity to restore cellular health²³. Even if some biological mechanisms underlying the association between sleep and LTL have been proposed, they are still unclear and need to be elucidated. For example, sleep is related to changes in the immune system through the sleep-wake cycle, and disruption in this cycle has been associated with higher inflammation.

The latter increases the circulation of proinflammatory cytokines, which may affect the telomere length^{23 24}. In the same line, changes in cortisol secretion and melatonin have also been linked to telomere length variation through higher oxidative stress^{25 26}. Hence, it may be hypothesized that a reduced stress environment, lower inflammation and/or oxidation^{6 9 27} in children who slept more hours per day may be the potential mechanisms that might explain longer telomeres at age 4 in our study. Nonetheless, the complex biology of telomeres – influenced by environmental and genetic factors – makes the investigation in this field very challenging. Therefore, future studies still need to elucidate which biological pathways might explain the association between sleep duration and LTL.

Sleep disturbance and its role in telomere length have been more widely investigated in adults⁶⁻⁸. Thus far, few studies have explored this association in children. James et al. investigated the cross-sectional association between telomere length and sleep duration at age 9 from 1567 children of the Fragile Families and Child Wellbeing Study (a population-based birth cohort of children born between 1998 and 2000 in American cities)¹². According to this study, each hour less sleep was associated with 0.015 log-kilobase shorter telomeres, i.e., children with fewer sleep hours had shorter telomeres than those who slept longer¹². Inconsistently, Nguyen et al. showed no evidence for the association between sleep duration – objectively measured – and telomere length in blood in adolescents of 11-12 years from the Longitudinal Study of Australian Children ($\beta = 0.01$ (95% CI: -0.04-0.06))¹³. The latter might be explained by the age of the participants since, as it has been proposed, the rate of telomere loss becomes more stable later in life compared with the first years¹⁴.

Short sleep duration is a modifiable risk factor contributing to non-communicable diseases such as type 2 diabetes, hypertension, and obesity in children²⁸⁻³⁰. The overuse of technology and screen time has disturbed many children's sleep hygiene, especially nighttime sleep³¹. Sleep routines provide security and help with activity transitions in children and moderate impulsivity^{2 31 32}. A previous systematic review of approaches to assist in sleep hygiene summarized that using positive routines, controlled comforting and gradual extinction or sleep remodeling are some recommended techniques³³. Given the acquaintance sleep benefits, consistent bedtime routines and adequate sleep are encouraged to promote positive child development and may be associated with a longer telomere length, as disclosed in this study.

This study leveraged data from the INMA birth cohort study, a pioneer project in Spain investigating the role of environmental factors during pregnancy and the beginning of life on growth and development. LTL was objectively measured following standard methods by trained professionals. In addition, we were able to adjust our analyses for an extensive range of confounder factors, including data collected during pregnancy, at birth, and during the 4-year follow-up interview. However, this study is not exempt from limitations. Firstly, although children from the INMA project were from different Spain areas, they may not represent the Spanish children population; therefore, estimates should not be fully generalized. Secondly, due to the observational nature of this study, causality cannot be inferred. Nonetheless, the prospective design of the INMA project allows verifying long-term effects in follow-up assessments and identifying potential etiological factors of disturbances of normal child development over time, thereby establishing a temporal sequence of events. Thirdly, recall bias is possible with self-reported data, as it was the sleep data in this study. Nonetheless, any inaccuracy should be understood as non-differential. Fourthly, LTL was measured using PCR, which shows a higher technical variability than, e.g., Terminal Restriction Fragment (TRF) analysis. However, in large cross-sectional settings, as assessed by qPCR, LTL may be in line with TRF estimated LTL³⁴. Yet, among the limitations of the PCR method are that it does not provide absolute LTL measures as well as issues detecting very short telomeres or telomeric losses. Therefore, even if a large amount of epidemiological research has conducted their investigation on LTL using the PCR approach³⁵, findings should be interpreted with caution, and telomere dynamics should be confirmed in future longitudinal-based studies. Finally, unmeasured or residual confounding is possible even if we included a long list of confounder factors. Moreover, traumatic events^{10 11}, a risk factor widely investigated and associated with telomere length, were not included as confounder factors since there was no available information.

In conclusion, children that slept more h/day had a longer LTL at age 4 independently of a wide range of confounder factors. No significant differences were identified in LTL at age 7-9. Therefore, sleep routines are encouraged to promote positive child development. Yet, considering the complex biology of telomere length, future studies still need to elucidate which biological pathways might explain the association between sleep duration and telomere length.

Data availability

The data that support the findings of this study are not available for sharing due to ethical and legal restrictions implemented by the regional Ethical Committees and the Ethical Committee of the General Hospital of Alicante. As stated in the informed consent form from participants, we guaranteed the confidentiality of collected personal information from questionnaires and related data. Requests to access the data should be submitted to the corresponding author. Requests will be reviewed by the research team and will require a data transfer agreement.

Author Contributions

F.P-R, D. V-G and E.N-M. contributed to the conception and design of the study, advised on all statistical aspects, and interpreted the data. F.P-R performed the literature search and the analyses. All authors critically reviewed this and previous drafts. All authors approved the final draft for submission, with final responsibility for publication. E.N-M is the guarantor.

Conflict of Interest Disclosures

None.

Funding/support

This study was funded by Instituto de Salud Carlos III/Agencia Estatal de Investigación, grant number PI18/00825 Project: “Dieta y actividad física en embarazo y tras el nacimiento y longitud del telómero en niños y adolescentes: Proyecto TeloDiPA” and Unión Europea (FEDER) “Una manera de hacer Europa”; Generalitat Valenciana (GVA/2021/191); Dries Martens holds a postdoctoral grant by the Flemish Scientific Fund (FWO grant 12X9620N). In Sabadell was funded by grants from Instituto de Salud Carlos III (Red INMA G03/176; CB06/02/0041; PI041436; PI081151 incl. FEDER funds; PI12/01890 incl. FEDER funds; CP13/00054 incl. FEDER funds, PI15/00118 incl. FEDER funds, CPII18/00018), CIBERESP, Generalitat de Catalunya-CIRIT 1999SGR 00241, Generalitat de Catalunya-AGAUR (2009 SGR 501, 2014 SGR 822), Fundació La marató de TV3 (090430), Spanish Ministry of Economy and Competitiveness (SAF2012-32991 incl. FEDER funds), Agence Nationale de Securite Sanitaire de l’Alimentation de l’Environnement et du Travail (1262C0010), EU Commission (261357,

308333, 603794 and 634453). We acknowledge support from the Spanish Ministry of Science and Innovation and the State Research Agency through the “Centro de Excelencia Severo Ochoa 2019-2023” Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program. In Asturias was funded by ISCIII: PI04/2018, PI09/02311, PI13/02429, PI18/00909 co-funded by FEDER, “A way to make Europe”/“Investing in your future”, Obra Social Cajastur/Fundación Liberbank, and Universidad de Oviedo. In Valencia, this study was supported by grants from Instituto de Salud Carlos III [FIS-FEDER: 13/1944, 16/1288, 19/1338; Miguel Servet-FSE: MS15/0025, MSII20/0006 incl. FEDER funds], Generalitat Valenciana [BEST/2020/059 and AICO/2020/285].

Role of the Funder/Sponsor

The funders had no role in the design of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval and submission of the manuscript.

References

1. Frank MG, Heller HC. The Function(s) of Sleep. *Handb Exp Pharmacol* 2019;253:3-34. doi: 10.1007/164_2018_140 [published Online First: 2019/04/21]
2. Mindell JA, Williamson AA. Benefits of a bedtime routine in young children: Sleep, development, and beyond. *Sleep Medicine Reviews* 2018;40:93-108. doi: <https://doi.org/10.1016/j.smrv.2017.10.007>
3. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended Amount of Sleep for Pediatric Populations: A Consensus Statement of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2016;12(6):785-86. doi: 10.5664/jcsm.5866
4. Wheaton AG, Claussen AH. Short Sleep Duration Among Infants, Children, and Adolescents Aged 4 Months-17 Years - United States, 2016-2018. *MMWR Morb Mortal Wkly Rep* 2021;70(38):1315-21. doi: 10.15585/mmwr.mm7038a1 [published Online First: 2021/09/24]
5. de Ruiter I, Olmedo-Requena R, Sánchez-Cruz J-J, et al. Changes in sleep duration in Spanish children aged 2–14 years from 1987 to 2011. *Sleep Medicine* 2016;21:145-50. doi: <https://doi.org/10.1016/j.sleep.2015.12.021>
6. Carroll JE, Esquivel S, Goldberg A, et al. Insomnia and Telomere Length in Older Adults. *Sleep* 2016;39(3):559-64. doi: 10.5665/sleep.5526 [published Online First: 2015/12/31]
7. Huang P, Zhou J, Chen S, et al. The association between obstructive sleep apnea and shortened telomere length: a systematic review and meta-analysis. *Sleep Med* 2018;48:107-12. doi: 10.1016/j.sleep.2017.09.034 [published Online First: 2018/06/09]
8. Savolainen K, Eriksson JG, Kajantie E, et al. The history of sleep apnea is associated with shorter leukocyte telomere length: the Helsinki Birth Cohort Study. *Sleep Med* 2014;15(2):209-12. doi: 10.1016/j.sleep.2013.11.779 [published Online First: 2013/12/24]
9. Rentscher KE, Carroll JE, Mitchell C. Psychosocial Stressors and Telomere Length: A Current Review of the Science. *Annual Review of Public Health* 2020;41(1):223-45. doi: 10.1146/annurev-publhealth-040119-094239
10. Deighton S, Neville A, Pusch D, et al. Biomarkers of adverse childhood experiences: A scoping review. *Psychiatry Research* 2018;269:719-32. doi: <https://doi.org/10.1016/j.psychres.2018.08.097>
11. Coimbra BM, Carvalho CM, Moretti PN, et al. Stress-related telomere length in children: A systematic review. *Journal of Psychiatric Research* 2017;92:47-54. doi: <https://doi.org/10.1016/j.jpsychires.2017.03.023>
12. James S, McLanahan S, Brooks-Gunn J, et al. Sleep Duration and Telomere Length in Children. *J Pediatr* 2017;187:247-52.e1. doi: 10.1016/j.jpeds.2017.05.014 [published Online First: 2017/06/13]
13. Nguyen MT, Lycett K, Olds T, et al. Objectively measured sleep and telomere length in a population-based cohort of children and midlife adults. *Sleep* 2020;43(1) doi: 10.1093/sleep/zsz200 [published Online First: 2019/11/17]
14. Gorenjak V, Petrelis AM, Stathopoulou MG, et al. Telomere length determinants in childhood. *Clin Chem Lab Med* 2020;58(2):162-77. doi: 10.1515/cclm-2019-0235 [published Online First: 2019/08/30]
15. Benetos A, Kark JD, Susser E, et al. Tracking and fixed ranking of leukocyte telomere length across the adult life course. *Aging Cell* 2013;12(4):615-21. doi: 10.1111/accel.12086 [published Online First: 2013/04/23]
16. Guxens M, Ballester F, Espada M, et al. Cohort Profile: the INMA--Infancia y Medio Ambiente-- (Environment and Childhood) Project. *Int J Epidemiol* 2012;41(4):930-40. doi: 10.1093/ije/dyr054 [published Online First: 2011/04/08]
17. Martens DS, Van Der Stukken C, Derom C, et al. Newborn telomere length predicts later life telomere length: Tracking telomere length from birth to child- and adulthood. *eBioMedicine* 2021;63 doi: 10.1016/j.ebiom.2020.103164

18. Martens DS, Janssen BG, Bijmens EM, et al. Association of Parental Socioeconomic Status and Newborn Telomere Length. *JAMA Netw Open* 2020;3(5):e204057-e57. doi: 10.1001/jamanetworkopen.2020.4057
19. Notario-Barandiaran L, Valera-Gran D, Gonzalez-Palacios S, et al. High adherence to a mediterranean diet at age 4 reduces overweight, obesity and abdominal obesity incidence in children at the age of 8. *Int J Obes (Lond)* 2020;44(9):1906-17. doi: 10.1038/s41366-020-0557-z [published Online First: 2020/03/11]
20. Buckland G, González CA, Agudo A, et al. Adherence to the Mediterranean diet and risk of coronary heart disease in the Spanish EPIC Cohort Study. *Am J Epidemiol* 2009;170(12):1518-29. doi: 10.1093/aje/kwp282 [published Online First: 2009/11/12]
21. Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *Bmj* 2003;327(7414):557-60. doi: 10.1136/bmj.327.7414.557 [published Online First: 2003/09/06]
22. Mickey RM, Greenland S. The impact of confounder selection criteria on effect estimation. *Am J Epidemiol* 1989;129(1):125-37. doi: 10.1093/oxfordjournals.aje.a115101 [published Online First: 1989/01/01]
23. Cribbet MR, Carlisle M, Cawthon RM, et al. Cellular aging and restorative processes: subjective sleep quality and duration moderate the association between age and telomere length in a sample of middle-aged and older adults. *Sleep* 2014;37(1):65-70. doi: 10.5665/sleep.3308 [published Online First: 2014/01/29]
24. Redwine L, Dang J, Irwin M. Cellular adhesion molecule expression, nocturnal sleep, and partial night sleep deprivation. *Brain, Behavior, and Immunity* 2004;18(4):333-40. doi: <https://doi.org/10.1016/j.bbi.2004.01.001>
25. Prather AA, Puterman E, Lin J, et al. Shorter leukocyte telomere length in midlife women with poor sleep quality. *J Aging Res* 2011;2011:721390. doi: 10.4061/2011/721390 [published Online First: 2011/11/03]
26. Liang G, Schernhammer E, Qi L, et al. Associations between rotating night shifts, sleep duration, and telomere length in women. *PLoS One* 2011;6(8):e23462. doi: 10.1371/journal.pone.0023462 [published Online First: 2011/08/20]
27. Prather AA, Gurfein B, Moran P, et al. Tired telomeres: Poor global sleep quality, perceived stress, and telomere length in immune cell subsets in obese men and women. *Brain Behav Immun* 2015;47:155-62. doi: 10.1016/j.bbi.2014.12.011 [published Online First: 2014/12/24]
28. Dutil C, Chaput JP. Inadequate sleep as a contributor to type 2 diabetes in children and adolescents. *Nutr Diabetes* 2017;7(5):e266. doi: 10.1038/nutd.2017.19 [published Online First: 2017/05/10]
29. Sluggett L, Wagner SL, Harris RL. Sleep Duration and Obesity in Children and Adolescents. *Can J Diabetes* 2019;43(2):146-52. doi: 10.1016/j.cjcd.2018.06.006 [published Online First: 2018/09/30]
30. DelRosso LM, Mogavero MP, Ferri R. Effect of Sleep Disorders on Blood Pressure and Hypertension in Children. *Curr Hypertens Rep* 2020;22(11):88. doi: 10.1007/s11906-020-01100-x [published Online First: 2020/09/08]
31. Bathory E, Tomopoulos S. Sleep Regulation, Physiology and Development, Sleep Duration and Patterns, and Sleep Hygiene in Infants, Toddlers, and Preschool-Age Children. *Current Problems in Pediatric and Adolescent Health Care* 2017;47(2):29-42. doi: <https://doi.org/10.1016/j.cppeds.2016.12.001>
32. Galland BC, Taylor BJ, Elder DE, et al. Normal sleep patterns in infants and children: A systematic review of observational studies. *Sleep Medicine Reviews* 2012;16(3):213-22. doi: <https://doi.org/10.1016/j.smrv.2011.06.001>
33. Halal CS, Nunes ML. Education in children's sleep hygiene: which approaches are effective? A systematic review. *J Pediatr (Rio J)* 2014;90(5):449-56. doi: 10.1016/j.jped.2014.05.001 [published Online First: 2014/06/29]

34. Nettle D, Gadalla SM, Lai T-P, et al. Measurement of Telomere Length for Longitudinal Analysis: Implications of Assay Precision. *American Journal of Epidemiology* 2021;190(7):1406-13. doi: 10.1093/aje/kwab025
35. Lai TP, Wright WE, Shay JW. Comparison of telomere length measurement methods. *Philos Trans R Soc Lond B Biol Sci* 2018;373(1741) doi: 10.1098/rstb.2016.0451 [published Online First: 2018/01/18]

Table 1. Cohort characteristics by sleep categories at baseline

	Sleep categories				p-value
	Total participants	7-10 h/day	>10-11 h/day	>11-14 h/day	
Total children, n (%)	1014 (100)	489 (48.2)	393 (38.8)	132 (13.0)	
Child's characteristics					
Age (years), median (IQR)	4.4 (4.3-4.5)	4.4 (4.3-4.5)	4.4 (4.3-4.5)	4.4 (4.3-4.5)	0.860
Sex, n (%)					0.223
Male	528 (52.1)	249 (50.9)	192 (48.8)	78 (59.1)	
Cohort, n (%)					
Asturias	317 (31.3)	138 (28.2)	125 (31.8)	54 (40.9)	
Gipuzkoa	217 (21.4)	102 (20.9)	88 (22.4)	27 (20.5)	<0.001
Sabadell	272 (26.8)	119 (24.3)	118 (30.0)	35 (26.5)	
Valencia	208 (20.5)	130 (26.6)	62 (15.8)	16 (12.1)	
Season of blood extraction at age 4, n (%)*					
Winter	181 (26.4)	73 (24.7)	72 (24.8)	36 (36.0)	
Spring	232 (33.8)	93 (31.4)	108 (37.3)	31 (31.0)	<0.001
Summer	105 (15.3)	52 (17.5)	43 (14.8)	10 (10.0)	
Autumn	168 (24.5)	78 (26.4)	67 (23.1)	23 (23.0)	
Season of blood extraction at age 7-9, n (%)*					
Winter	233 (26.4)	114 (26.0)	91 (26.8)	28 (26.7)	
Spring	227 (25.7)	113 (25.7)	91 (26.8)	23 (21.9)	0.186
Summer	184 (21.9)	99 (22.6)	71 (20.8)	24 (22.8)	
Autumn	230 (26.0)	113 (25.7)	87 (25.6)	30 (28.6)	
Relative Mediterranean diet score, median (IQR)	8.0 (7.0-10.0)	8.0 (6.0-10.0)	9.0 (7.0-11.0)	9.0 (7.0-10.0)	0.002
TV time, (h/week), median (IQR)	9.0 (5.8-13.5)	9.0 (6.5-14.0)	9.0 (5.8-12.0)	7.5 (5.3-11.3)	0.002
Mothers' characteristics					
Parity, n (%)					
≥1	435 (42.9)	208 (42.5)	181 (46.3)	45 (34.1)	0.048
Social class at baseline, n (%)					
I+II	259 (25.5)	135 (27.6)	92 (23.4)	32 (24.2)	0.031
III	258 (25.5)	116 (23.7)	118 (30.0)	24 (18.2)	
IV + V	497 (49.0)	238 (48.7)	183 (46.6)	76 (57.6)	

The distribution of the continuous variables was checked using the Lilliefors correction of Kolmogorov–Smirnov test and compared using ANOVA, Kruskal-Wallis, and Chi-square tests. N, number; h: hour; IQR: interquartile range.

*Available only at age 4 and 7-9, respectively ~ used in sensitivity analyses, available in fewer participants.

Table 2. Associations between sleep categories and telomere length at age 4 and age 7-9.

	Model 1			Model 2		Model 3	
	n	% difference (95% CI)	P-value	% difference (95% CI)	P-value	% difference (95% CI)	P-value
Cross-sectional association							
Sleep categories at 4 – telomere at age 4	686						
Medium (>10-11 h/day)	290	Ref.		Ref.		Ref.	
Low (7-10 h/day)	296	1.49 (-1.57-4.64)	0.343	2.19 (-0.88-5.36)	0.164	2.21 (-0.88-5.38)	0.162
High (>11-14 h/day)	100	6.88 (1.94-12.1)	0.006	8.43 (3.47-13.6)	<0.001	8.48 (3.56-13.6)	<0.001
Longitudinal association							
Sleep categories at 4 – telomere at age 7-9	875						
Medium (>10-11 h/day)	334	Ref.		Ref.		Ref.	
Low (7-10 h/day)	437	1.15 (-1.88-4.28)	0.460	1.51 (-1.57-4.68)	0.340	1.72 (-1.36-4.89)	0.276
High (>11-14 h/d>ay)	104	3.65 (-1.23-8.77)	0.145	4.26 (-0.70-9.47)	0.094	3.86 (-1.19-9.17)	0.136

Data presented as % difference and their respective 95% CI. Model 1: adjusted for blood date, cohort, age at baseline and sex of the child. Model 2: as per model 1, but also for social class at baseline and parity Model 3: as per model 2, but also for relative Mediterranean parameter and TV time.




Figure 1. Diagram of participant included in the study