



Bayes-Genis, A. et al. (2022) Biomarkers in heart failure clinical trials. A review from the Biomarkers Working Group of the Heart Failure Association of the European Society of Cardiology. *European Journal of Heart Failure*, 24(10), pp. 1767-1777. (doi: [10.1002/ejhf.2675](https://doi.org/10.1002/ejhf.2675))

This is the author version of the work. There may be differences between this version and the published version. You are advised to consult the published version if you wish to cite from it:  
<https://doi.org/10.1002/ejhf.2675>

<https://eprints.gla.ac.uk/279208/>

Deposited on 12 September 2022

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

## Biomarkers in Heart Failure Clinical Trials

### A review from the Biomarkers Working Group of the Heart Failure Association of the ESC

Antoni Bayés-Genís<sup>1,2,3</sup>, Alberto Aimo<sup>4,5</sup>, Pardeep Jhund<sup>6</sup>, A Mark Richards<sup>7</sup>, Rudolf A de Boer<sup>8</sup>, Henrike Arfsten<sup>9</sup>, Iacopo Fabiani<sup>5</sup>, Josep Lupón<sup>1</sup>, Stefan D Anker<sup>10</sup>, Arantxa González<sup>3,11</sup>, Vincenzo Castiglione<sup>4</sup>, Marco Metra<sup>12</sup>, Christian Mueller<sup>13</sup>, Julio Núñez<sup>3,14</sup>, Patrick Rossignol<sup>15</sup>, Andrea Barison<sup>5</sup>, Javed Butler<sup>16</sup>, John Teerlink<sup>17</sup>, Gerasimos Filippatos<sup>18</sup>, Piotr Ponikowski<sup>19</sup>, Giuseppe Vergaro<sup>4,5</sup>, Faiez Zannad<sup>20</sup>, Petar Seferovic<sup>21,22</sup>, Giuseppe Rosano<sup>23</sup>, Andrew JS Coats<sup>24</sup>, Michele Emdin<sup>4,5</sup>, James L Januzzi<sup>25</sup>

<sup>1</sup> Institut del Cor, Hospital Universitari Germans Trias i Pujol, Badalona, Barcelona, Spain; <sup>2</sup> Department of Medicine, Universitat Autònoma de Barcelona, Barcelona, Spain; <sup>3</sup> CIBERCV, Carlos III Institute of Health, Madrid, Spain; <sup>4</sup> Scuola Superiore Sant'Anna, Pisa, Italy; <sup>5</sup> Cardiology Division, Fondazione Toscana Gabriele Monasterio, Pisa, Italy; <sup>6</sup> Institute of Cardiovascular & Medical Sciences, University of Glasgow, UK; <sup>7</sup> University of Otago, New Zealand; <sup>8</sup> University Medical Centre Groningen, Department of Cardiology, Groningen, The Netherlands; <sup>9</sup> Clinical Division of Cardiology, Department of Internal Medicine II, Medical University of Vienna, Vienna, Austria; <sup>10</sup> Department of Cardiology (CVK), and Berlin Institute of Health Center for Regenerative Therapy (BCRT), German Center for Cardiovascular Research (DZHK) partner site Berlin, Charité Universitätsmedizin, Berlin, Germany; <sup>11</sup> Program of Cardiovascular Diseases, CIMA Universidad de Navarra and IdiSNA, Navarra Institute for Health Research, Pamplona, Spain; <sup>12</sup> Cardiology Department, ASST Spedali Civili; Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, University of Brescia, Italy; <sup>13</sup> Department of Cardiology, University Hospital, Basel, Switzerland; <sup>14</sup> Hospital Clínico Universitario de Valencia, INCLIVA, Universidad de Valencia, Valencia, Spain; <sup>15</sup> Academic Hospital (CHU), Nancy, France; <sup>16</sup> Department of Medicine, University of Mississippi Medical Center, Jackson, Mississippi, USA; <sup>17</sup> Heart Failure and of the Echocardiography Laboratory at the San Francisco Veterans Affairs Medical Center in San Francisco, California; <sup>18</sup> HF Unit, Attikon University Hospital, Athens, Greece; <sup>19</sup> Centre for Heart Diseases, University Hospital, Wroclaw, Poland; <sup>20</sup> Université de Lorraine, Centre d'Investigations Cliniques-Plurithématique 1433, and Inserm U1116, CHRU Nancy, F-CRIN

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ejhf.2675](https://doi.org/10.1002/ejhf.2675)

INI-CRCT, Nancy, France; <sup>21</sup> Department of Cardiology, University Clinical Center of Serbia, Belgrade, Serbia; <sup>22</sup> Serbian Academy of Sciences and Arts, Belgrade, Serbia; <sup>23</sup> IRCCS San Raffaele, Rome, Italy; <sup>24</sup> University of Warwick, Coventry, UK; <sup>25</sup> Massachusetts General Hospital and Baim Institute for Clinical Research, Boston, Massachusetts, USA.

Short title: Biomarkers in HF trials

Word count: 3,821 (text)

**Address for correspondence:**

Antoni Bayés-Genís

Cardiology Department, Hospital Universitari Germans Trias i Pujol

Carretera de Canyet, s/n, 08916 Badalona, Barcelona, Spain

Email: abayesgenis@gmail.com

Phone: +34 679418491

## Abstract

The approval of new heart failure (HF) therapies has slowed over the past two decades in part due to the high costs of conducting large randomized clinical trials that are needed to adequately power major clinical endpoint studies. Several biomarkers have been identified reflecting different elements of HF pathophysiology, with possible applications in diagnosis, risk stratification, treatment monitoring, and even in the design of clinical trials. Biomarkers could potentially be used to refine study inclusion criteria to enable enrolment of patients who are more likely to respond to a therapeutic intervention, despite being at sufficient risk to meet predetermined study endpoint rates. When there is a close relationship between biomarker levels and clinical endpoints, changes in biomarker levels after a given treatment can act as a surrogate endpoint, potentially reducing the duration and cost of a clinical trial. Natriuretic peptides have been widely used in clinical trials with a variable amount of added value, which such variation being probably due to the absence of a close pathophysiological connection to the study drug. Notable exceptions to this include sacubitril/valsartan and vericiguat. Future studies should seek to adopt unbiased approaches for discovery of true companion diagnostics; with -omics-based tools, biomarkers might be more precisely selected for use in clinical trials to identify responses that closely reflect the biological effects of the drug under investigation. Finally, biomarkers associated with cardiac damage and remodelling, such as cardiac troponin, could be employed as safety endpoints provided that standardization between different assays is achieved.

Word count: 248 (abstract)

Keywords: biomarkers; clinical trials; criteria; natriuretic peptides; inclusion; risk prediction; -omics.

## Abbreviations and acronyms

ACC/AHA/HFSA, American College of Cardiology/American Heart Association/HF Society of America

BNP, B-type natriuretic peptide

CI, confidence interval

cTn, cardiac troponin

ESC, European Society of Cardiology

FDA, Food and Drug Administration

GUIDE-IT, Guiding Evidence Based Therapy Using Biomarker Intensified Treatment in Heart Failure

HF, heart failure

HFpEF, heart failure with preserved ejection fraction

HFREF, heart failure with reduced ejection fraction

HR, hazard ratio

hs-cTn(I/T), high-sensitivity cardiac troponin (I/T)

NP, natriuretic peptide

NT-proBNP, N-terminal pro-B-type natriuretic peptide

PARADIGM-HF, Prospective Comparison of ARNI (Angiotensin Receptor-Neprilysin Inhibitor) with ACEI (Angiotensin-Converting-Enzyme Inhibitor) to Determine Impact on Global Mortality and Morbidity in Heart Failure

PARAGON-HF, Efficacy and Safety of LCZ696 Compared to Valsartan, on Morbidity and Mortality in Heart Failure Patients With Preserved Ejection Fraction

PONTIAC, NT-proBNP selected prevention of cardiac events in a population of diabetic patients without a history of cardiac disease

PROVE-HF, Effects of Sacubitril/Valsartan Therapy on Biomarkers, Myocardial Remodeling and Outcomes

STOP-HF, St Vincent's Screening TO Prevent Heart Failure

TOPCAT, Treatment of Preserved Cardiac Function Heart Failure With an Aldosterone Antagonist Trial

U.S., United States

VICTORIA, Vericiguat Global Study in Patients With Heart Failure and Reduced Ejection Fraction

Cardiovascular disorders remain the leading cause of morbidity and mortality worldwide, but the pace of drug and device development is considerably slower than for other conditions such as neoplastic disorders (1). A possible reason is the high cost of clinical trials adequately powered for hard clinical endpoints. Problems with trial design might be another issue, as 20% of drugs completing the phase 3 stage are not approved by the United States (U.S.) Food and Drug Administration (FDA), and only 6% of all novel cardiovascular drugs entering phase 1 are ultimately approved (2). It has been noted that clinical development programs selecting patients through biomarker-based criteria have higher success rates at each phase of development (3). Indeed, biomarkers might facilitate the enrolment of patients who are most likely to respond to therapeutic intervention, despite still having the sufficient number of events to meet the trial endpoints (1).

A biomarker (from “biological marker”) has been defined as “a characteristic that is objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention” (4). The analysis of circulating substances is the most common form of biomarker measurement, and has become widespread in clinical research (5,6). A 2019 FDA guidance document indicates that biomarkers have potential utility to enrol HF patients with a greater event risk, stratify patients based on their predicted prognosis, and allow early proof of concept and dose selection studies (7). Despite the almost exclusive focus on B-type natriuretic peptide (BNP) and N-terminal pro-BNP (NT-proBNP), many biomarkers reflecting different elements of HF pathophysiology exist (8-11), and the identification of specific biomarkers for each drug has been proposed (1).

Building on the FDA document and some authoritative papers (1,6,12,13), this review article provides specific suggestions for biomarker use in HF clinical trials.

### **Possible applications of heart failure biomarkers**

Many biomarkers have been proposed to guide the diagnosis and management of HF (14) (**Figure 1**). Ibrahim and Januzzi summarized in 5 points the characteristics of an ideal HF biomarker: 1) the

method by which a biomarker is judged should be thorough; 2) the assays used to measure the biomarker should be robust; 3) the biomarker should reflect an important pathophysiologic pathway involved in the HF disease process; 4) the biomarker should provide information other than what is already available by routine physical exam and laboratory evaluation; 5) the biomarker should add to clinical judgment for understanding diagnosis, prognosis, or management of HF (15). Biomarkers can predict the development of signs and symptoms, identify patients with subclinical disease, help diagnose HF, predict disease trajectories, guide therapeutic management, or serve as surrogate endpoints (16,17).

BNP and NT-proBNP >80<sup>th</sup> percentile predict a higher risk of new-onset HF at 5 years in the general population (18). Two randomized controlled trials (St Vincent's Screening TO Prevent Heart Failure [STOP-HF] and NT-proBNP selected prevention of cardiac events in a population of diabetic patients without a history of cardiac disease [PONTIAC]) evaluated a BNP- or NT-proBNP-guided approach in individuals with risk factors for HF. The STOP-HF trial enrolled subjects aged >40 years with at least one risk factor or cardiovascular comorbidity (19), while the PONTIAC trial enrolled diabetic patients (20). The positive results of these trials informed the American College of Cardiology/American Heart Association/HF Society of America (ACC/AHA/HFSA) Guideline recommendation of a NP-based screening in patients at risk for HF (IIa B-R recommendation) (21).

In symptomatic patients, BNP or NT-proBNP are recommended to diagnose chronic HF by both the European Society of Cardiology (ESC; I B recommendation) (22) and ACC/AHA/HFSA (I A recommendation) Guidelines (21). BNP or NT-proBNP are also recommended to diagnose acute HF (21,22). High NP levels are also essential components of the universal definition of HF (23). A large number of biomarkers have proven useful for the prediction of disease trajectories in patients with acute decompensated or ambulatory HF (5,14). The most studied are again NPs, and their use as prognostic tools in outpatient care is recommended by ACC/AHA/HFSA Guidelines (1 A), as well as on admission (1 A) and at discharge (2a B-NR) (21). Additionally, current evidence suggests that sex modulates cardiovascular biomarkers biology. However, the biological



Accepted Article

mechanisms are still poorly defined, with limited translation into clinical practice. In detail, there are limited sex-specific trials, with consistent underrepresentation of women as a standard or special population and limited inclusion of biological sex in research study design. This ultimately results in a general uncertainty regarding different tailoring of clinical care for men and women, starting from specific biomarkers thresholds. Future approaches should include a sex-specific approach, from recruitment to study design and reporting (24).

Regarding biomarkers integration in diagnostic, prognostic and therapeutic setting, circulating biomarkers and imaging techniques provide independent and complementary information to guide management of HF. A recent consensus document by the Heart Failure Association (HFA) of the European Society of Cardiology (ESC) recently presented evidence-based indications relevant to the integration of imaging techniques and biomarkers in HF (25). This includes screening, diagnosis, risk stratification, the guidance of treatment, and monitoring. An individually tailored approach includes a deep characterization of cardiac dysfunction, age, gender, and comorbidities. Circulating biomarkers associated with different pathways are an ideal complement of multi-modal imaging techniques in patient-centered and condition-specific approaches to HF assessment and management.

Several studies have tried to find biomarkers able of guiding therapeutic management in HF, with disappointing results (5). The Guiding Evidence Based Therapy Using Biomarker Intensified Treatment in Heart Failure (GUIDE-IT) trial enrolled patients with LVEF <40%, an HF episode within the prior 12 months, and elevated NPs within the previous month (BNP >400 ng/L or NT-proBNP >2000 ng/L), who were randomized to an NT-proBNP-guided strategy (target NT-proBNP <1000 ng/L) or the standard of care. The study was discontinued prematurely for futility after 894 patients out of the planned 1,100 were enrolled, because there was no significant difference in the risk of cardiovascular death or HF hospitalization over a median of 15 months (hazard ratio [HR] 0.98, 95% confidence interval [CI], 0.79-1.22; p=0.88), with the neutral results most likely caused by similar treatment (26).

Accepted Article

Finally, biomarkers could serve as surrogate endpoints that could be reached in a shorter time than hard endpoints, thus reducing the follow-up time, the size of the study group and overall cost of clinical trials (15). The prerequisite is a close relationship between biomarker change and a clinical endpoint. At present, such relationship has not emerged for any biomarker. In the Prospective Comparison of ARNI (Angiotensin Receptor-Neprilysin Inhibitor) with ACEI (Angiotensin-Converting-Enzyme Inhibitor) to Determine Impact on Global Mortality and Morbidity in Heart Failure (PARADIGM-HF) trial, an early decrease in NT-proBNP predicted fewer clinical end-points, but it did not predict the superior treatment effect of the experimental therapy (27). Nonetheless, the perfect alignment of biology and pharmacology (NT-proBNP changes due to drug effects) led to the FDA utilizing changes in NT-proBNP in paediatric studies of sacubitril/valsartan as a bridge to approval (28).

### **Biomarkers in HF clinical trials**

Biomarkers have many possible applications in HF trials (**Table 1**). BNP and NT-proBNP have been by far the most commonly evaluated biomarkers, and have been employed usually for patient selection or as surrogate endpoints (**Figure 2**).

#### Inclusion criteria

The use of biomarkers as inclusion criteria has become standard practice in HF clinical trials, encouraging enrolment of patients with a level of risk that allows to design adequately powered trials that are still economically viable.

NPs have several characteristics that make their use ideal for patient selection for clinical trials. Most notably, they are inexpensive, reproducible, reliable, are sensitive and specific to diagnose HF, and have strong prognostic value. Given their significant associations with the presence and severity of HF (29,30), NP-based inclusion criteria might allow to accurately identify patients with HF. This is particularly important in patients with HF and preserved ejection fraction (HFpEF) because the

potential for misdiagnosis of HF is greater in this population, which might in turn dilute the observed benefit of the investigational treatment. Most of the major recent trials have used NP-based inclusion criteria (31-34). In the Treatment of Preserved Cardiac Function Heart Failure With an Aldosterone Antagonist Trial (TOPCAT) trial, patients could be enrolled based on clinical suspicion of HF plus either elevated BNP or NT-proBNP or a recent HF hospitalization. While no significant benefit from spironolactone was found for the primary outcome, a significant benefit was seen in patients enrolled based on elevated NPs (HR, 0.65; 95% CI, 0.49-0.87; p=0.003) (33). Furthermore, patients enrolled without meeting the biomarker entry requirement had a lower event rate, suggesting misdiagnosis of HF (35). A recent analysis of the Efficacy and Safety of LCZ696 Compared to Valsartan, on Morbidity and Mortality in Heart Failure Patients With Preserved Ejection Fraction (PARAGON-HF) trial focused on the 193 patients (4.0%) not meeting the final NP-based enrolment criteria (NT-proBNP >300 ng/L for patients in sinus rhythm or >900 ng/L for those in atrial fibrillation/flutter). These patients had lower rates of the primary endpoint of total HF hospitalizations and cardiovascular death as compared with patients meeting final enrolment criteria (8.6 [6.7-11.2] events per 100 patient-years vs. 14.0 [13.4-14.7] events per 100 patient-years; p=0.01). The rate ratio for the treatment effect comparing sacubitril/valsartan with valsartan was 0.85 (95% confidence interval 0.74-0.99; p=0.04) in those who met final criteria (36).

In 2020, a search on the ClinicalTrials.gov dataset identified 3,446 trials, with 10.6% using BNP or NT-proBNP as inclusion criteria (6). Among these trials, 43% used both NPs, 33% used only NT-proBNP, and 24% used only BNP in determining eligibility. Significant variations existed across trials in terms of BNP and NT-proBNP cut-off values used as inclusion criteria (**Supplemental Table 1**). Only some trials (10% in patients with decompensated HF, 20% in ambulatory outpatients with HF) employed adjusted cut-offs for at least some patient populations, although NP levels are known to be lower in black, obese, or patients with HFpEF, and higher in elderly patients or those with atrial fibrillation. Ibrahim et al. went on to propose recommendations for the use of NPs for patient selection in HF clinical trials (**Table 2**) (6).

Elevated NPs predict a higher risk of adverse outcomes, including death and rehospitalization (37,38). The value of NPs for risk prediction seems similar in patients with HFpEF or HF with reduced EF (HFrEF) despite lower concentrations of both BNP or NT-proBNP in patients with HFpEF (39). The possibility to predict future disease trajectories is particularly useful for enriching the study population with patients who have higher event rates to minimize the risk for type II error in underpowered studies. In the PARADIGM-HF trial, patients were required to have a BNP  $\geq 150$  ng/L or NT-proBNP  $\geq 600$  ng/L, or a BNP  $\geq 100$  ng/L or NT-proBNP  $\geq 400$  ng/L if they had been hospitalized for HF within the previous 12 months. The resulting high event rates allowed for more timely completion of this event-driven study (40). Higher concentrations of selected predictive biomarkers may enrich a trial with study participants at high risk, which usually benefit the most from a certain treatment (41-43). On the other hand, an excessively high biomarker threshold could lead to enrol patients who are unlikely to respond to therapy because of their excessive risk. An example of this may be provided by a *post hoc* analysis of the Vericiguat Global Study in Patients With Heart Failure and Reduced Ejection Fraction (VICTORIA) trial, which demonstrated a significant benefit in the primary composite endpoint only in subjects with NT-proBNP levels up to 8,000 ng/L (44).

#### Assessment of response to treatment

In the non-randomized Effects of Sacubitril/Valsartan Therapy on Biomarkers, Myocardial Remodeling and Outcomes (PROVE-HF) study, early changes in NT-proBNP were associated with a subsequent reverse remodelling in patients with HFrEF receiving sacubitril/valsartan (45). The drop in NT-proBNP was reported to precede a decrease in high-sensitivity cardiac troponin T (hs-cTnT), and the combination of these biomarker changes predicted reverse remodelling more accurately than each biomarker alone (46). Besides sacubitril/valsartan, several drug classes, proven to be beneficial in HFrEF, have been shown to reduce NPs. For example, treatment with spironolactone (47) and empagliflozin (48) were associated with a decrease in NT-proBNP. With the notable exception of

sacubitril/valsartan, biomarker changes are usually not a direct effect of therapy, but rather reflect an amelioration of HF severity. In the case of sodium-glucose cotransporter 2 inhibitors, our incomplete understanding of the drug effects on cardiomyocytes does not allow us to define with certainty whether NT-proBNP changes partially represent a direct manifestation of the mechanism of action of the drug (49).

Regardless of the lack of a biological link between specific therapies and natriuretic peptide changes, when reduction in BNP or NT-proBNP occurs after a therapy is applied, it is more often than not accompanied by a temporal shift in prognosis. Often, the prognostic meaning of NP lowering may be significantly greater than other risk markers: for example, despite an unclear association between sodium-glucose cotransporter 2 inhibitor effects and the biology determining NT-proBNP and hs-cTnT release, reduction of both markers were the strongest predictors of outcome in the recent EMPEROR program focused on empagliflozin in HF (50). However, NPs remain surrogate measurements whose relationship with outcomes remains indirect. It must be tested whether they may actually substitute outcome for drug approval whereas outcome, as well as quality of life, remain major endpoints that need direct assessment.

The use of a biomarker to monitor response to therapy in clinical trials need to be carefully assessed for each drug-biomarker combination. In particular, 1) the biomarker should be affected by a specific treatment in a manner that indicates how effective the treatment is; 2) changes in the biomarker levels should reflect the benefit of the therapy; 3) the biomarker should be sampled with an optimal, pre-specified timing and frequency to track treatment response, limiting the fluctuations due to its biological variability; 4) a pre-specified threshold to assess treatment response should be chosen, based either on percent or absolute changes (1). For most biomarkers, these points remain unclear, and more work in this area is needed as it might lead to their use as a surrogate endpoint for expedited regulatory approval of promising therapies. The optimal conditions to choose a surrogate biomarker are summarized in **Table 3**.

### Biomarkers as elements of composite endpoints

Despite their relative strength as indicators of prognosis, changes in biomarker concentrations may nonetheless be insufficient to represent the entire clinical effect of a given treatment (51). Indeed, biomarker changes, as well as symptom resolution, and clinical events should be considered together as measure of treatment benefit. A ranking approach allows to incorporate both events and quantitative measures of functional status (e.g., quality of life assessment, 6-minute walking distance or cardiac biomarkers), and to assess them based on a prespecified hierarchical system (52). A pre-specified hierarchical ranking system can overcome the discrepancies often found between phase 2 and 3 trials, where the phase 2 trial shows an improvement in symptoms or congestion (surrogate endpoints), but these positive findings do not translate into a significant benefit in terms of hard clinical outcomes in the subsequent phase 3 trial. Indeed, the combination of continuous biomarker data and clinical endpoints in a hierarchical system in a phase 2 study might provide a more reliable indicator of the success or failure of the treatment in the following phase 3 study (52). The global rank endpoint of Felker and Maisel is an example of hierarchical approach that is based on the occurrence of death, the relief of dyspnoea at 24 hours, and changes in three biomarkers (cTn, creatinine, and BNP/NT-proBNP) (53). A limitation of this approach is that if we consider the low percentage of in-hospital death and the very high likelihood of meaningful symptom improvement at 24 hours, this endpoint can be simplified as a comparison of biomarkers to test the efficacy of a new drug. However, since there is no established relationship between changes in cTn or BNP and outcomes in acute HF, this approach is not considered a valid surrogate for regulatory review (54). The win ratio is a novel method of incorporating morbidity, mortality, health-related quality of life and NPs in HF studies in a hierarchical fashion (55,56). Nonetheless, the more complex the ranking approach, the less easy it is for physicians to understand the benefit of a certain treatment and to carry out cost-effectiveness analyses (57).

### Subgroup analyses

Biomarkers may be used to provide insight into specific subgroups of patients included in clinical trials, generating novel hypotheses (12). This may provide insight into the mechanism of drug action (8), the benefits of a drug (58), or the identification of subgroups or specific phenotypes of patients that report a differential benefit (8). Notably, although NPs are recognized biomarkers in many different settings, including HFpEF, we should highlight its limitations, as recently shown by Verbrugge et al. (59). In particular, while patients with HFpEF and normal NP display in general mild functional alterations, with more favorable clinical outcomes, there remains an increased residual risk of events, emphasizing the importance of a phenotype potentially poorly tracked by a biomarkers-only based strategy in large clinical trials in this specific setting.

### Surrogates for safety

Numerous types of cardiac toxicity have been revealed throughout drug development, and interest in using biomarkers as an early system for toxicity detection has increased. In preclinical studies, biomarkers may be used to identify hazards, generate hypotheses, and guide patient monitoring in clinical trials. In phase 2 to 4 trials, it is critical to have effective tools to monitor cardiac toxicity for innovative therapies to ensure patient safety. When biomarkers are used to detect potentially toxic effects of HF drugs, it is critical to establish a solid biological link between the biomarker and therapy toxicity, and any risk estimated from the change in biomarker levels over time should be rigorously validated, including knowledge of the minimum amount of change required to predict risk (60-62). One should not assume a rise in biomarker is necessarily a sign of toxicity: the importance of clear understanding of how therapies may cause unexpected rise in biomarkers is illustrated in the increase of NPs following treatment with drugs from the neuregulin class (63) or increase in hs-cTn following myosin activator therapy with omecantiv mecarbil (64). In both cases, no increase in risk was observed with the rise in biomarker levels.

cTn is an indicator of cardiomyocyte damage, and measurement of cTn through hs-assays may be used in HF trials to assess the toxicity of treatment strategies. Nonetheless, several hs-cTnI tests exist, each with its own reference range, and there are no specific guidelines establishing hs-cTn cut-offs above which concentrations would signal an abnormality (65). It is then difficult to say, for example, if a 4 ng/L higher median change in hs-cTnI in patients on a novel drug vs. placebo is clinically relevant (64). Changes in hs-cTn should be interpreted considering the clinical setting and other laboratory data. If hs-cTn is employed in clinical trials as a toxicity monitor without a clear grasp of how to interpret concentration changes, it may jeopardize an otherwise beneficial medication.

### **When might biomarkers not be used?**

The use of biomarkers in the development of cardiovascular therapies is particularly attractive when assessing slowly progressive cardiovascular diseases, patients with little or no symptoms, subclinical cardiovascular diseases, populations with low rates of clinical events, or therapeutic options requiring accelerated approval. Studies on patients with clinical HF usually do not meet these criteria, unless specific aetiologies are considered (for example, hypertrophic cardiomyopathy or cardiac amyloidosis).

Biomarkers are not strictly required when clinically relevant measures can be assessed, such as patient-reported outcomes, physical performance, and mortality (1). Biomarkers should not be utilized when they have not been completely validated, or when they have not been evaluated in the patient demographic or disease state being researched. This is not trivial: intra-individual variability between biomarkers differs greatly (66), and the origin of most biomarkers is not the cardiac muscle but rather other organs such as fatty tissue, liver, and kidney (67). Therefore, changes over time of most biomarkers reflect changes in the entire organism, not changes of the cardiac muscle. Furthermore, short-term biomarker changes in patients with HF treated with a brief treatment exposure should not be viewed as conclusive for a longer-term benefit if the therapy in question does not have long-lasting effects (68,69).



### **Proposals for the use of biomarkers in future clinical trials**

Based on the considerations above, we may formulate several suggestions for the future use of biomarkers in clinical trials (**Table 4**). As a first point, the use of biomarkers in HF clinical trials requires standardization of analytical aspects, which is now less relevant for NPs, but is a significant issue for other biomarkers, such as hs-cTn. Patient enrolment in HF trials is now commonly based on NP levels, but there is a lack of consensus on the specific cut-offs to be employed. In addition, factors affecting NP values (including age, gender, EF, atrial fibrillation, or chronic kidney disease) are not given sufficient consideration. Recently proposed cut-offs (**Table 2**) are in agreement with a previous position paper by the ESC (13), and might represent the first step toward standardized NP-based inclusion criteria. A relevant field of application is represented by valvular heart disease. The recent ESC/EACTS Guidelines (70) emphasize the central role of cardiac biomarkers in valvular patients, in particular NPs (71), especially in asymptomatic patients or to better tailor the follow-up. However, biomarkers use should be adopted selectively and, at present, starting from trials guidance in percutaneous setting (72), their standardization should be almost completely addressed along all trial phases.

Instead of focusing on NPs in every trial, future studies could try to identify specific biomarkers for each drug. These patterns could be employed for risk enrichment and to assess the response to treatment. A panel of biomarkers reflecting different cardiac alterations, such as collagen peptides or sST2 associated with fibrosis, C-reactive protein and interleukins with inflammation, endothelin or adrenomedullin with endothelial dysfunction, antigen carbohydrate 125 or bio-adrenomedullin with fluid overload, urinary sodium for monitoring decongestion, or direct renin for renin-angiotensin-aldosterone system activation could provide additional useful information regarding the pathophysiological impact on cardiac remodelling features (73). In addition, an unbiased -omic approach could be more accurate than the simple evaluation of biomarkers more closely associated with specific disease mechanisms (74-76) (**Figure 4**). The heterogeneous setting of HFpEF seems particularly well-suited for this kind of approach (77).

The main priorities for cardiac safety biomarkers are to define the strengths and limitations of *in vitro* models, and the relative advantages of human- versus animal-derived models for specific questions. Appropriate applications need assessment of study duration (acute vs. chronic effects), resource availability, the value of single versus multiparametric outcomes, and overall translational fidelity. Additionally, unbiased -omics may help identify safety markers before phase 3 studies (78).

Finally, future clinical trials in HF should systematically include a plan for the collection, storage, and management of laboratory samples, to warrant the adequate quality of the aliquots and to ensure the possibility to test novel biomarkers.

**Funding:** This work did not receive any funding.

**Potential conflicts of interest:** The UMCG, which employs Dr. De Boer, has received research grants and/or fees from AstraZeneca, Abbott, Boehringer Ingelheim, Cardior Pharmaceuticals GmbH, Ionis Pharmaceuticals, Inc., Novo Nordisk, and Roche (outside the submitted work). Dr. de Boer received speaker fees from Abbott, AstraZeneca, Bayer, Novartis, and Roche (outside the submitted work). Dr. Núñez reports personal fees from AstraZeneca, Novartis, Boehringer-Ingelheim, Eli Lilly, Rovi, NovoNordisk, and Vifor Pharma (outside the submitted work). The other Authors do not report any conflicts of interest relevant to the present review.

## Figure legends

**Figure 1. Main pathophysiological pathways involved in heart failure and their most representative biomarkers.**

BNP, B-type natriuretic peptide; CA-125, cancer antigen-125; CRP, C-reactive protein; FGF-23, fibroblast growth factor-23; fT3, triiodothyronine; GDF15, growth differentiation factor-15; GFR, glomerular filtration rate; hs-TnI/T, high-sensitivity troponin I/T; KIM-1, kidney injury molecule-1; MMP, matrix metalloprotease; MPO, myeloperoxidase; MR-proADM, mid-regional pro-adrenomedullin; MR-proANP, mid-regional pro-atrial natriuretic peptide; NAG, N-acetyl- $\beta$ -(D)-glucosaminidase; NGAL, neutrophil gelatinase-associated lipocalin; NT-proBNP, N-terminal pro-B-type natriuretic peptide; PRA, plasma renin activity; sST2, soluble suppression of tumorigenesis-2; TIMP, tissue inhibitor of metalloproteinase; TNF- $\alpha$ , tumour necrosis factor alpha. Reprinted with permission from: Castiglione et al., 2022 (5).

**Figure 2. Use of natriuretic peptides in heart failure clinical trials.**

This analysis refers to all studies registered in ClinicalTrials.gov up to July 2019. Reprinted with permission from: Ibrahim et al., 2020 (6).

**Figure 3. Proposed new approach to biomarker-based design of heart failure (HF) clinical trials.**

BNP, B-type natriuretic peptide; NT-proBNP, N-terminal pro-B-type natriuretic peptide.

## Tables

**Table 1. Possible applications of biomarkers in heart failure clinical trials.**

<b>Phase 2</b>	<b>Phase 3</b>	<b>Phase 4</b>
<ul style="list-style-type: none"><li>• Development of mechanistic hypotheses</li><li>• Inclusion criteria</li><li>• Safety monitoring</li></ul>	<ul style="list-style-type: none"><li>• Inclusion criteria</li><li>• Risk enrichment</li><li>• Assessment of response to treatment (surrogate endpoint or part of composite endpoint)</li><li>• Identification of patient subgroups where the drug seems more effective</li><li>• Safety monitoring</li></ul>	<ul style="list-style-type: none"><li>• Inclusion criteria</li><li>• Assessment of treatment benefit (if only surrogate endpoint in phase 3 trial)</li><li>• Safety monitoring</li></ul>

**Table 2. Possible natriuretic peptide-based cut-offs in heart failure (HF) clinical trials.**

<p>If the goal is to exclude HF:</p> <ul style="list-style-type: none"><li>• BNP &lt;100 ng/L or NT-proBNP &lt;300 ng/L to exclude acute HF</li><li>• BNP &lt;35 ng/L or NT-proBNP &lt;125 ng/L to exclude chronic HF</li></ul>
<p>If the goal is to include patients with probable acute HF in the emergency department setting, symptoms of dyspnoea should be present accompanied by the following cut-off values:</p> <ul style="list-style-type: none"><li>• BNP &gt;100 ng/L</li><li>• NT-proBNP &gt;450 ng/L (age &lt;50 years); &gt;900 ng/L (age 50-75 years); &gt;1,800 ng/L (age &gt;75 years)</li></ul>
<p>To enrich a study population for risk in stable chronic HFrEF and HFpEF clinical trials:</p> <ul style="list-style-type: none"><li>• BNP <math>\geq</math>100 ng/L or NT-proBNP <math>\geq</math>360 ng/L for HFpEF trials</li><li>• BNP <math>\geq</math>150 ng/L or NT-proBNP <math>\geq</math>600 ng/L for HFrEF trials</li><li>• It is advisable to remember clinical presentation, including severity of symptoms, LVEF, and comorbidities independently add to the observed risk in patients with elevated BNP or NT-proBNP</li></ul>
<p>If higher event rates are desired, higher concentrations of BNP and NT-proBNP concentrations should be considered. For example:</p> <ul style="list-style-type: none"><li>• BNP &gt;400 ng/L</li><li>• NT-proBNP &gt;900 ng/L</li></ul>
<p>To enrich a study population for risk in advanced/morbid chronic HF clinical trials:</p>

- BNP  $\geq 300$  ng/L or NT-proBNP  $\geq 1000$  ng/L

Special circumstances:

1. Atrial fibrillation

- BNP and NT-proBNP: consider an increase of enrolment threshold by  $\geq 30\%$

2. Black patients

- BNP and NT-proBNP: consider lowering enrolment threshold by  $\geq 30\%$

3. Elderly patients (>75 years)

- BNP: consider raising enrolment threshold by at least 20-30%
- NT-proBNP: >1800 ng/L

4. Chronic kidney disease

- Exclude patients with ESRD or receiving RRT from enrolment by NPs

4. Neprilysin inhibition

- Avoid BNP use for monitoring response to neprilysin inhibition until more data is available regarding response of the various BNP assays to therapy. BNP should be used for study population enrichment, however
- No adjustment needed for NT-proBNP

5. Obesity

- BNP and NT-proBNP: Consider lowering enrolment threshold by at least 20-30% for patients with BMI  $\geq 30$  kg/m<sup>2</sup>

BMI, body mass index; BNP, B-type natriuretic peptide; ESRD, end-stage renal disease; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; LVEF, left ventricular ejection fraction; NP, natriuretic peptide; NT-proBNP, N-terminal pro-B-type natriuretic peptide; RRT, renal replacement therapy. Modified with permission from: Ibrahim et al., 2020 (4).

**Table 3. Optimal conditions to choose a surrogate biomarker.**

An excellent understanding of disease pathophysiology
A strong mechanistic rationale for a relationship between the surrogate endpoint and the clinical outcome(s) of interest
Clinical data supporting the existence of the same relationship
Some evidence of an association between the magnitude of biomarker changes and the clinical effect size



**Table 4. Proposals for the use of biomarkers in future clinical trials.**

Broader and more reasoned use of biomarkers in clinical trials
Standardized BNP/NT-proBNP cut-offs as inclusion criteria, considering study goals and specific patient features (e.g., atrial fibrillation or chronic kidney disease)
More extensive use of biomarkers as surrogate endpoints or elements of composite endpoints
Transition from a NP-centred strategy to biomarker selection via an unbiased -omic approach
Implementation of biomarkers of cardiac damage as safety endpoints
Unbiased -omics to identify safety markers before phase 3 studies
Storage of samples for future biomarker studies

BNP, B-type natriuretic peptide; NT-proBNP, N-terminal pro-B-type natriuretic peptide.

## References

1. Januzzi JL, Jr., Canty JM, Das S, DeFilippi CR, Gintant GA, Gutstein DE, et al. Gaining efficiency in clinical trials with cardiac biomarkers: JACC Review Topic of the Week. *J Am Coll Cardiol*. 2021;77:1922-33.
2. <https://www.statista.com/statistics/1186648/us-clinical-development-success-rate-novel-cardiovascular-drugs/>
3. BIO. Clinical Development Success Rates 2006-2015; June 2016. Available at: <https://www.bio.org/sites/default/files/legacy/bioorg/docs/Clinical%20Development%20Success%20Rates%202006-2015%20-%20BIO,%20Biomedtracker,%20Amplion%202016.pdf>.
4. Biomarkers and surrogate endpoints: preferred definitions and conceptual framework. *Clin Pharmacol Therapeut*. 2001;69:89-95.
5. Castiglione V, Aimo A, Vergaro G, Saccaro L, Passino C, Emdin M. Biomarkers for the diagnosis and management of heart failure. *Heart Fail Rev*. 2022;27:625-43.
6. Ibrahim NE, Burnett JC, Jr., Butler J, Camacho A, Felker GM, Fiuzat M, et al. Natriuretic peptides as inclusion criteria in clinical trials: A JACC: Heart Failure Position Paper. *JACC Heart Fail*. 2020;8:347-58.
7. U.S. Department of Health and Human Services, Food and Drug Administration, Center for Drug Evaluation and Research (CDER), Center for Biologics Evaluation and Research (CBER). FDA Draft Guidance: Treatment for Heart Failure: Endpoints for Drug Development Guidance for Industry, June 2019. Silver Spring, MD: Food and Drug Administration, 2019.
8. Ahmad T, Fiuzat M, Pencina MJ, Geller NL, Zannad F, Cleland JG, et al. Charting a roadmap for heart failure biomarker studies. *JACC Heart Fail*. 2014;2:477-88.
9. van Kimmenade RR, Januzzi JL, Jr. Emerging biomarkers in heart failure. *Clinical Chem*. 2012;58:127-38.

10. Braunwald E. The war against heart failure: the Lancet lecture. *Lancet*. 2015;385:812-24.
11. González A, Richards AM, de Boer RA, Thum T, Arfsten H, Hülsmann M, et al. Cardiac remodelling - Part 1: From cells and tissues to circulating biomarkers. A review from the Study Group on Biomarkers of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail*. 2022.
12. Ibrahim NE, Gaggin HK, Konstam MA, Januzzi JL, Jr. Established and emerging roles of biomarkers in heart failure clinical trials. *Circ Heart Fail*. 2016;9.
13. Mueller C, McDonald K, de Boer RA, Maisel A, Cleland JGF, Kozhuharov N, et al. Heart Failure Association of the European Society of Cardiology practical guidance on the use of natriuretic peptide concentrations. *Eur J Heart Fail*. 2019;21:715-31.
14. Braunwald E. Biomarkers in heart failure. *N Engl J Med*. 2008;358:2148-59.
15. Ibrahim NE, Januzzi JL, Jr. Beyond Natriuretic Peptides for Diagnosis and Management of Heart Failure. *Clin Chem*. 2017;63:211-22.
16. Vasan RS. Biomarkers of cardiovascular disease: molecular basis and practical considerations. *Circulation*. 2006;113:2335-62.
17. Emdin M, Vittorini S, Passino C, Clerico A. Old and new biomarkers of heart failure. *Eur J Heart Fail*. 2009;11:331-5.
18. Wang TJ, Larson MG, Levy D, Benjamin EJ, Leip EP, Omland T, et al. Plasma natriuretic peptide levels and the risk of cardiovascular events and death. *N Engl J Med*. 2004;350:655-63.
19. Ledwidge M, Gallagher J, Conlon C, Tallon E, O'Connell E, Dawkins I, et al. Natriuretic peptide-based screening and collaborative care for heart failure: the STOP-HF randomized trial. *JAMA*. 2013;310:66-74.
20. Hülsmann M, Neuhold S, Resl M, Strunk G, Brath H, Francesconi C, et al. PONTIAC (NT-proBNP selected prevention of cardiac events in a population of diabetic patients without a

history of cardiac disease): a prospective randomized controlled trial. *J Am Coll Cardiol.* 2013;62:1365-72.

21. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2022:101161cir0000000000001063.
22. McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, Burri H, Butler J, Čelutkienė J, Chioncel O, Cleland JGF, Coats AJS, Crespo-Leiro MG, Farmakis D, Gilard M, Heymans S, Hoes AW, Jaarsma T, Jankowska EA, Lainscak M, Lam CSP, Lyon AR, McMurray JJV, Mebazaa A, Mindham R, Muneretto C, Francesco Piepoli M, Price S, Rosano GMC, Ruschitzka F, Kathrine Skibelund A; ESC Scientific Document Group. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J.* 2021;42:3599-3726.
23. Bozkurt B, Coats AJ, Tsutsui H, Abdelhamid M, Adamopoulos S, Albert N, Anker SD, Atherton J, Böhm M, Butler J, Drazner MH, Felker GM, Filippatos G, Fonarow GC, Fiuzat M, Gomez-Mesa JE, Heidenreich P, Imamura T, Januzzi J, Jankowska EA, Khazanie P, Kinugawa K, Lam CSP, Matsue Y, Metra M, Ohtani T, Francesco Piepoli M, Ponikowski P, Rosano GMC, Sakata Y, Seferović P, Starling RC, Teerlink JR, Vardeny O, Yamamoto K, Yancy C, Zhang J, Zieroth S. Universal definition and classification of heart failure: A Report of the Heart Failure Society of America, Heart Failure Association of the European Society of Cardiology, Japanese Heart Failure Society and Writing Committee of the Universal Definition of Heart Failure. *J Card Fail.* 2021 Mar 1:S1071-9164(21)00050-6.
24. Lau ES, Binek A, Parker SJ, Shah SH, Zanni MV, Van Eyk JE, Ho JE. Sexual dimorphism in cardiovascular biomarkers: clinical and research implications. *Circ Res.* 2022;130:578-592.
25. Moura B, Aimo A, Al-Mohammad A, Flammer A, Barberis V, Bayes-Genis A, et al. Integration of imaging and circulating biomarkers in heart failure: a consensus document by the Biomarkers

and Imaging Study Groups of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Fail.* 2021;23:1577-1596.

26. Felker GM, Anstrom KJ, Adams KF, Ezekowitz JA, Fiuzat M, Houston-Miller N, et al. Effect of natriuretic peptide-guided therapy on hospitalization or cardiovascular mortality in high-risk patients with heart failure and reduced ejection fraction: a randomized clinical trial. *JAMA.* 2017;318:713-20.
27. Zile MR, Claggett BL, Prescott MF, McMurray JJ, Packer M, Rouleau JL, et al. Prognostic implications of changes in n-terminal pro-b-type natriuretic peptide in patients with heart failure. *J Am Coll Cardiol.* 2016;68:2425-36.
28. <https://www.medscape.com/viewarticle/919362>, 10.1093/eurheartj/ehr352, <https://clinicaltrials.gov/ct2/show/study/NCT02678312?term=NCT+02678312&rank=1>
29. Januzzi JL, Jr., Rehman S, Mueller T, van Kimmenade RR, Lloyd-Jones DM. Importance of biomarkers for long-term mortality prediction in acutely dyspneic patients. *Clin Chem.* 2010;56:1814-21.
30. Daubert MA, Adams K, Yow E, Barnhart HX, Douglas PS, Rimmer S, et al. NT-proBNP goal achievement is associated with significant reverse remodeling and improved clinical outcomes in HFrEF. *JACC Heart Fail.* 2019;7:158-68.
31. Solomon SD, McMurray JJV, Anand IS, Ge J, Lam CSP, Maggioni AP, et al. Angiotensin-neprilysin inhibition in heart failure with preserved ejection fraction. *N Engl J Med.* 2019;381:1609-20.
32. Anker SD, Butler J, Filippatos G, Ferreira JP, Bocchi E, Böhm M, et al. Empagliflozin in heart failure with a preserved ejection fraction. *N Engl J Med.* 2021;385(16):1451-61.
33. Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med.* 2014;370:1383-92.

34. Solomon SD, de Boer RA, DeMets D, Hernandez AF, Inzucchi SE, Kosiborod MN, et al. Dapagliflozin in heart failure with preserved and mildly reduced ejection fraction: rationale and design of the DELIVER trial. *Eur J Heart Fail.* 2021;23:1217-25.
35. Myhre PL, Vaduganathan M, Claggett BL, Anand IS, Sweitzer NK, Fang JC, et al. Association of Natriuretic Peptides With Cardiovascular Prognosis in Heart Failure With Preserved Ejection Fraction: Secondary Analysis of the TOPCAT Randomized Clinical Trial. *JAMA Cardiol.* 2018;3:1000-5.
36. Pabón MA, Cunningham JW, Claggett BL, Packer M, Zile M, Pfeffer MA, et al. Natriuretic peptide-based inclusion criteria in heart failure with preserved ejection fraction clinical trials: insights from PARAGON-HF. *Eur J Heart Fail.* 2022.
37. Januzzi JL, van Kimmenade R, Lainchbury J, Bayes-Genis A, Ordonez-Llanos J, Santalo-Bel M, et al. NT-proBNP testing for diagnosis and short-term prognosis in acute destabilized heart failure: an international pooled analysis of 1256 patients: the International Collaborative of NT-proBNP Study. *Eur Heart J.* 2006;27:330-7.
38. Rehman SU, Martinez-Rumayor A, Mueller T, Januzzi JL, Jr. Independent and incremental prognostic value of multimarker testing in acute dyspnea: results from the ProBNP Investigation of Dyspnea in the Emergency Department (PRIDE) study. *Clin Chim Acta.* 2008;392:41-5.
39. van Veldhuisen DJ, Linssen GC, Jaarsma T, van Gilst WH, Hoes AW, Tijssen JG, et al. B-type natriuretic peptide and prognosis in heart failure patients with preserved and reduced ejection fraction. *J Am Coll Cardiol.* 2013;61:1498-506.
40. McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med.* 2014;371:993-1004.
41. Packer M, Fowler MB, Roecker EB, Coats AJ, Katus HA, Krum H, et al. Effect of carvedilol on the morbidity of patients with severe chronic heart failure: results of the carvedilol

prospective randomized cumulative survival (COPERNICUS) study. *Circulation*. 2002;106:2194-9.

42. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet*. 1999;353:2001-7.
43. Motiwala SR, Szymonifka J, Belcher A, Weiner RB, Baggish AL, Sluss P, et al. Serial measurement of galectin-3 in patients with chronic heart failure: results from the ProBNP Outpatient Tailored Chronic Heart Failure Therapy (PROTECT) study. *Eur J Heart Fail*. 2013;15:1157-63.
44. Ezekowitz JA, O'Connor CM, Troughton RW, Alemayehu WG, Westerhout CM, Voors AA, et al. N-terminal pro-B-type natriuretic peptide and clinical outcomes: vericiguat heart failure with reduced ejection fraction study. *JACC Heart Fail*. 2020;8:931-9.
45. Januzzi JL, Jr., Prescott MF, Butler J, Felker GM, Maisel AS, McCague K, et al. Association of change in n-terminal pro-b-type natriuretic peptide following initiation of sacubitril-valsartan treatment with cardiac structure and function in patients with heart failure with reduced ejection fraction. *JAMA*. 2019;322:1085-95.
46. Murphy SP, Prescott MF, Maisel AS, Butler J, Piña IL, Felker GM, et al. Association between angiotensin receptor-neprilysin inhibition, cardiovascular biomarkers, and cardiac remodeling in heart failure with reduced ejection fraction. *Circ Heart Fail*. 2021;14:e008410.
47. Ozkara A, Turgut F, Selcoki Y, Karanfil A, Metin MR, Kanbay M, et al. Probrain natriuretic peptide for assessment of efficacy in heart failure treatment. *Adv Ther*. 2007;24:1233-9.
48. Packer M, Anker SD, Butler J, Filippatos G, Pocock SJ, Carson P, et al. Cardiovascular and renal outcomes with empagliflozin in heart failure. *N Engl J Med*. 2020;383:1413-24.
49. Vallon V, Verma S. Effects of SGLT2 inhibitors on kidney and cardiovascular function. *Ann Rev Physiol*. 2021;83:503-28.

50. Pocock SJ, Ferreira JP, Gregson J, Anker SD, Butler J, Filippatos G, et al. Novel biomarker-driven prognostic models to predict morbidity and mortality in chronic heart failure: the EMPEROR-Reduced trial. *Eur Heart J*. 2021;42:4455-64.
51. Brown PM, Ezekowitz JA. Composite end points in clinical trials of heart failure therapy: how do we measure the effect size? *Circ Heart Fail*. 2017;10.
52. Packer M. Development and evolution of a hierarchical clinical composite end point for the evaluation of drugs and devices for acute and chronic heart failure: a 20-year perspective. *Circulation*. 2016;134:1664-78.
53. Felker GM, Maisel AS. A global rank end point for clinical trials in acute heart failure. *Circ Heart Fail*. 2010;3:643-6.
54. Zannad F, Garcia AA, Anker SD, Armstrong PW, Calvo G, Cleland JG, et al. Clinical outcome endpoints in heart failure trials: a European Society of Cardiology Heart Failure Association consensus document. *Eur J Heart Fail*. 2013;15:1082-94.
55. Voors AA, Angermann CE, Teerlink JR, Collins SP, Kosiborod M, Biegus J, et al. The SGLT2 inhibitor empagliflozin in patients hospitalized for acute heart failure: a multinational randomized trial. *Nat Med*. 2022;28:568-74.
56. Ferreira JP, Jhund PS, Duarte K, Claggett BL, Solomon SD, Pocock S, et al. Use of the win ratio in cardiovascular trials. *JACC Heart Fail*. 2020;8:441-50.
57. Pocock SJ, Ariti CA, Collier TJ, Wang D. The win ratio: a new approach to the analysis of composite endpoints in clinical trials based on clinical priorities. *Eur Heart J*. 2012;33:176-82.
58. Braunwald E. Heart failure. *JACC Heart Fail*. 2013;1:1-20.
59. Verbrugge FH, Omote K, Reddy YNV, Sorimachi H, Obokata M, Borlaug BA. Heart failure with preserved ejection fraction in patients with normal natriuretic peptide levels is associated with increased morbidity and mortality. *Eur Heart J*. 2022;43:1941-1951.




60. Curigliano G, Cardinale D, Suter T, Plataniotis G, de Azambuja E, Sandri MT, et al. Cardiovascular toxicity induced by chemotherapy, targeted agents and radiotherapy: ESMO Clinical Practice Guidelines. *Ann Oncol*. 2012;23 Suppl 7:vii155-66.
61. Yeh ET, Tong AT, Lenihan DJ, Yusuf SW, Swafford J, Champion C, et al. Cardiovascular complications of cancer therapy: diagnosis, pathogenesis, and management. *Circulation*. 2004;109:3122-31.
62. Sawaya H, Sebag IA, Plana JC, Januzzi JL, Ky B, Cohen V, et al. Early detection and prediction of cardiotoxicity in chemotherapy-treated patients. *Am J Cardiol*. 2011;107:1375-80.
63. Jabbour A, Hayward CS, Keogh AM, Kotlyar E, McCrohon JA, England JF, et al. Parenteral administration of recombinant human neuregulin-1 to patients with stable chronic heart failure produces favourable acute and chronic haemodynamic responses. *Eur J Heart Fail*. 2011;13:83-92.
64. Teerlink JR, Diaz R, Felker GM, McMurray JJV, Metra M, Solomon SD, et al. Cardiac myosin activation with omecamtiv mecarbil in systolic heart failure. *N Engl J Med*. 2021;384:105-16.
65. Sherwood MW, Kristin Newby L. High-sensitivity troponin assays: evidence, indications, and reasonable use. *JAHA*. 2014;3:e000403.
66. Meijers WC, van der Velde AR, Muller Kobold AC, Dijck-Brouwer J, Wu AH, Jaffe A, et al. Variability of biomarkers in patients with chronic heart failure and healthy controls. *Eur J Heart Fail*. 2017;19:357-65.
67. Du W, Piek A, Schouten EM, van de Kolk CWA, Mueller C, Mebazaa A, et al. Plasma levels of heart failure biomarkers are primarily a reflection of extracardiac production. *Theranostics*. 2018;8:4155-69.
68. Cohen-Solal A, Logeart D, Huang B, Cai D, Nieminen MS, Mebazaa A. Lowered B-type natriuretic peptide in response to levosimendan or dobutamine treatment is associated with improved survival in patients with severe acutely decompensated heart failure. *J Am Coll Cardiol*. 2009;53:2343-8.

69. Packer M, O'Connor C, McMurray JJV, Wittes J, Abraham WT, Anker SD, et al. Effect of ularitide on cardiovascular mortality in acute heart failure. *N Engl J Med.* 2017;376:1956-64.
70. Vahanian A, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J.* 2022;43:561-632.
71. Gardezi SK, Coffey S, Prendergast BD, Myerson SG. Serum biomarkers in valvular heart disease. *Heart.* 2018;104:349-358.
72. Stone GW, Vahanian AS, Adams DH, Abraham WT, Borer JS, Bax JJ, et al. Clinical trial design principles and endpoint definitions for transcatheter mitral valve repair and replacement: part 1: clinical trial design principles: a consensus document from the Mitral Valve Academic Research Consortium. *J Am Coll Cardiol.* 2015;66:278-307.
73. Aimo A, Vergaro G, González A, Barison A, Lupón J, Delgado V, et al. Cardiac remodelling part 2: clinical, imaging and laboratory findings. A review from the Biomarkers Working Group of the Heart Failure Association of the ESC. *Eur J Heart Fail.* 2022.
74. Michelhaugh SA, Januzzi JL, Jr. Finding a needle in a haystack: proteomics in heart failure. *JACC Basic Transl Sci.* 2020;5:1043-53.
75. Michelhaugh SA, Camacho A, Ibrahim NE, Gaggin H, D'Alessandro D, Coglianese E, et al. Proteomic signatures during treatment in different stages of heart failure. *Circulation Heart Fail.* 2020;13:e006794.
76. Bayes-Genis A, Liu PP, Lanfear DE, de Boer RA, González A, Thum T, et al. Omics phenotyping in heart failure: the next frontier. *Eur Heart J.* 2020;41:3477-84.
77. Adamo L, Yu J, Rocha-Resende C, Javaheri A, Head RD, Mann DL. Proteomic signatures of heart failure in relation to left ventricular ejection fraction. *J Am Coll Cardiol.* 2020;76:1982-94.
78. Meijers WC, Bayes-Genis A, Mebazaa A, Bauersachs J, Cleland JGF, Coats AJS, et al. Circulating heart failure biomarkers beyond natriuretic peptides: review from the Biomarker


Study Group of the Heart Failure Association (HFA), European Society of Cardiology (ESC).

Eur J Heart Fail. 2021;23:1610-32.


Accepted Article



EJHF\_2675\_Figure 1.png



EJHF\_2675\_Figure 2.tif



EJHF\_2675\_Figure 3.png

Permission note: all permissions to reproduce original material have been obtained; permissions will be shown if needed.

Accepted Article