

A large-scale genomic snapshot of *Klebsiella* spp. isolates in Northern Italy reveals limited transmission between clinical and non-clinical settings

Received: 19 July 2021

Accepted: 10 October 2022

Published online: 21 November 2022

 Check for updates

Harry A. Thorpe¹, Ross Booton², Teemu Kallonen³, Marjorie J. Gibbon⁴, Natacha Couto⁴, Virginie Passet⁵, Sebastián López-Fernández⁵, Carla Rodrigues⁵, Louise Matthews⁶, Sonia Mitchell⁶, Richard Reeve⁶, Sophia David⁷, Cristina Merla⁸, Marta Corbella⁸, Carolina Ferrari⁸, Francesco Comandatore⁹, Piero Marone⁸, Sylvain Brisse⁵, Davide Sasserà^{10,12}, Jukka Corander^{1,7,11,12} & Edward J. Feil^{4,12} ✉

The *Klebsiella* group, found in humans, livestock, plants, soil, water and wild animals, is genetically and ecologically diverse. Many species are opportunistic pathogens and can harbour diverse classes of antimicrobial resistance genes. Healthcare-associated *Klebsiella pneumoniae* clones that are non-susceptible to carbapenems can spread rapidly, representing a high public health burden. Here we report an analysis of 3,482 genome sequences representing 15 *Klebsiella* species sampled over a 17-month period from a wide range of clinical, community, animal and environmental settings in and around the Italian city of Pavia. Northern Italy is a hotspot for hospital-acquired carbapenem non-susceptible *Klebsiella* and thus a pertinent setting to examine the overlap between isolates in clinical and non-clinical settings. We found no genotypic or phenotypic evidence for non-susceptibility to carbapenems outside the clinical environment. Although we noted occasional transmission between clinical and non-clinical settings, our data point to a limited role of animal and environmental reservoirs in the human acquisition of *Klebsiella* spp. We also provide a detailed genus-wide view of genomic diversity and population structure, including the identification of new groups.

The *Klebsiella* genus is a member of the Enterobacteriaceae family. The most well studied species is *Klebsiella pneumoniae*, which the World Health Organization has recognized as a critical priority healthcare-associated pathogen¹. Antibiotic resistance has spread rapidly within *K. pneumoniae* and other members of the genus since the early 1980s²; the plasmid-mediated spread of genes encoding

carbapenemases over the last two decades is of particular concern^{3,4}. Widespread clones of *K. pneumoniae* and other *Klebsiella* species associated with these genes are spread through the healthcare network⁵; in common with other key antimicrobial resistance (AMR) determinants, genes encoding carbapenemases have also been detected in multiple non-clinical settings including livestock and wastewater^{6–8}.

A full list of affiliations appears at the end of the paper. ✉ e-mail: e.feil@bath.ac.uk




Fig. 1 | Summary of the sampling effort. a, Geographical summary of the whole sampling area. **b**, More detail of the region around the city of Pavia as highlighted by the red box in **a**. The size of each point indicates the number of samples and the colours represent the source. **c**, Timeline of the sampling effort broken down by source. Further details are hidden to preserve anonymity.

The potential public health risks posed by these non-clinical reservoirs of antibiotic resistance has led to a widening adoption of the One Health framework for AMR management⁹. This integrative approach is underpinned by a synthesis of antibiotic stewardship and AMR surveillance within clinical, community, agricultural and environmental settings. However, existing data on the abundance and distribution of AMR strains and genes in the environment does not provide a full picture and our current understanding of their maintenance and spread within and between ecological settings is fragmentary¹⁰. Given the urgent requirement for policy priorities informed by robust risk assessments, this represents a key knowledge gap.

A powerful way to infer pathogen transmission dynamics is to use whole-genome sequencing (WGS) combined with phylogenetic and other statistical analyses. WGS has been applied successfully within healthcare settings^{11,12} and played a key role in the management of the severe acute respiratory syndrome coronavirus 2 pandemic; however, capturing transmission pathways within complex non-clinical settings is more challenging. Large contemporaneous samples from well-defined regions are required to control for the confounding effects of divergence time and geographical variation. Nevertheless, a picture has begun to emerge that the risk of transmission of AMR genes and strains from environmental or agricultural settings into the clinic may be rather low, at least in well-resourced regions; this is a view seemingly at odds with the prevailing One Health *cri de coeur*¹³. Recent reports suggested that transmission of AMR strains and/or genes between humans and agricultural animals is limited in *Escherichia coli*¹⁴, *Enterococcus faecium*¹⁵ and *K. pneumoniae*^{16,17}. The evidence, however, is equivocal¹⁸; a compelling counter-example is the study on colistin

resistance dissemination in humans in China, which was largely driven by aquaculture activities¹⁹.

The sequencing of large and carefully sampled collections of isolates holds the promise to inform on the rate of transmission between settings²⁰ and shed light on the relevant biological and ecological factors underpinning transmission barriers. While commonalities of gene and strain profiles between settings point to ample opportunities for mixing, the risks to public health of environmental reservoirs of AMR are difficult to gauge in the absence of this broad framework^{10,21,22}. Recent advances in bioinformatics tools and analytical approaches provide the means to extract critical added value from genome data, enabling a more nuanced view of how microbes and mobile elements move through complex ecosystems.

In this study, we report a large-scale One Health study based on WGS data for 3,482 sequenced isolates, recovered from 6,548 samples. The sequenced genomes represent 15 *Klebsiella* species (including *Raoultella* species²³) and approximately half ($n = 1,705$) are *K. pneumoniae*. These data informed a large-scale transmission analysis to identify environmental reservoirs likely to pose the greatest risk to public health. Samples from multiple clinical, community, veterinary, agricultural and environmental sources were taken within a 17-month period around a single city, Pavia, in Northern Italy. This represents an unprecedented contemporaneous sampling and sequencing effort within a restricted geographical area that is a known hotspot for healthcare-associated multidrug-resistant *K. pneumoniae*⁵. We describe the distribution of species, strains, AMR and virulence genes in different settings and provide detail on the phylogeny and diversity of the *Klebsiella* genus, including the identification of new lineages of potential species status.

Results

Sequencing, species assignments and phylogenetic analysis

After quality control, 3,483 high-quality read sets and assemblies were retained: 2,796 from diverse sources recovered using Simmons Citrate Agar with Inositol (SCAI) media and 687 from ongoing clinical surveillance projects (summaries in Extended Data Fig. 1 and Supplementary Tables 1 and 2). Full isolate metadata, including species and lineage assignments, source, genotypic and phenotypic resistance data and phylogenetic trees are available for download from the Microreact project (<https://microreact.org/project/KLEBPAVIA>). A summary of the main metadata fields used in the Microreact project is provided in Supplementary Table 3. Figure 1 provides a summary of the sampling. A summary of the species assignments and sources of the 3,482 sequenced *Klebsiella* isolates and phylogenetic trees are given in Fig. 2. We used three-letter species abbreviations throughout this article; these are provided in the main text and the legend to Fig. 2.

We inferred a neighbour-joining tree of all 3,483 isolates using Mash²⁴ distances (Fig. 2b) and generated a more statistically robust RAxML²⁵ tree (GTR+ Γ) based on a representative subset of 703 isolates (Fig. 2a). These trees were consistent with each other and with Kleborate²⁶ species assignments, except for those cases where clusters were not present in the Kleborate database. We identified 15 recognized *Klebsiella* species, including *Klebsiella pasteurii* (*K. pas*) and *Klebsiella spallanzanii* (*K. spa*), which were first isolated during the course of this study²⁷, and 8 isolates of *Klebsiella huaxiensis* (*K. hua*), which was previously only recovered from a urine sample in China²⁸. Our data also resolved a new cluster of six isolates, to which we have assigned the label *Klebsiella quasiterrigena* (*K. qte*) (Extended Data Fig. 2) and two isolates from a hospital carriage that are positioned approximately equidistantly from *Klebsiella grimontii* (*K. gri*) and *K. pas* (labelled NA; Extended Data Fig. 3). A single isolate recovered from the surface of an automated teller machine (ATM) was assigned as a new species belonging to the genus *Superficiebacter*, designated *Superficiebacter maynardsmithii*²⁹, and was retained as a convenient outgroup. Previous WGS studies did not support the assignment of the *Raoultella* species as a separate genus^{7,23,30}, which is consistent with our data. Hence, we refer to these species as *Klebsiella*. Phylogenetic analyses revealed four higher-order clusters, which we have referred to as species complexes (SPECs), extending those used in²⁶. These were named according to the canonical species in each group: *K. pneumoniae* (*K. pne* SPEC), *Klebsiella oxytoca* (*K. oxy* SPEC), *Klebsiella ornithocolytica* (*K. orn* SPEC (*Raoultella*)) and *Klebsiella aerogenes* (*K. aer* SPEC) (Extended Data Figs. 2–4).

K. pneumoniae (*K. pne*) was by far the most commonly sampled species, accounting for approximately half of the isolates ($n = 1,705$). This proportion was inflated by the inclusion of the 687 diagnostic isolates, 676 of which were sampled from healthcare settings. Of these isolates, 571 were *K. pne* (84%), confirming its dominance as a cause of human infection, with the opportunistic pathogen *K. oxy* being the second most common ($n = 40$; 6%) (Supplementary Table 4).

Species clonality and population structures

We compared the population structures of the different species by delineating SCs using PopPunk³¹ (Extended Data Fig. 5 and Fig. 3). This revealed high levels of diversity in all species, as previously described for *K. pne*^{32,33}. In total, we identified 1,168 SCs across all species, of which only 41 (3.5%) were represented by more than 10 isolates and 50% of all

isolates corresponded to SCs that were observed no more than 6 times. The most common SC within each species represented between 3 and 10% of the population (Fig. 3a) and the SC accumulation curves were not close to saturation (Fig. 3b). *K. orn* showed particularly high diversity; 147 SCs were identified from 258 isolates and the most common SC accounted for 3% of the isolates. Pairwise divergences tended to be distributed around a modal average of approximately 1% (Fig. 3c) and each lineage was roughly equidistant to every other lineage (Extended Data Fig. 6). In some cases (for example, *K. pne*, *K. gri*), a much smaller peak was also evident at a much lower divergence, reflecting expansion of individual SCs. *Klebsiella michiganensis* (*K. mic*), *K. hua*, *K. spa*, *Klebsiella terrigena* (*K. ter*) and *K. aer* also showed more diverged modal peaks, with core genome distances up to 3%; this reflects the presence of deep subdivisions within these species, which is consistent with nascent speciation; this was also evident in the individual species trees (Extended Data Fig. 6 and Supplementary Table 5).

Species are distributed non-randomly across different sources


We explored the prevalence and distribution of the 15 recognized *Klebsiella* species and *K. qte* across different epidemiological and ecological sources (Figs. 2 and 4). Most of the disease isolates from hospital patients were recovered from diagnostic plates; thus, it was not valid to compare source prevalence between the diagnostic and SCAI samples. Therefore, the analysis presented in Fig. 4 was restricted to the 2,795 *Klebsiella* isolates recovered using the SCAI sampling strategy from any of the major source categories ($n = 23$ isolates excluded). Considering all sources, prevalence (calculated as the percentage of samples that were positive for at least 1 species) was highest for water samples (river, 100%; environmental, 85.2%; farm 86.1%) and turtles (82.6%), most of which are riverine. The source with the next highest prevalence was humans (hospital carriage, 58.5%; community carriage, 62.9%) and livestock (cows, 59.6%; pigs, 49.4%). The prevalence from soil was 44.6% and from plants 26.8%. While a high prevalence was observed from farm surfaces (53.1%), the prevalence from environmental and hospital surfaces was much lower (15.9%). Most species can be isolated from most sources; 20 sources harboured at least 7 species and 11 sources harboured at least 10 species.

We used a permutation test to gauge whether different species were non-randomly distributed between sources (Fig. 4). *K. pne* was significantly overrepresented in hospital carriage and in livestock (cows and pigs), as expected^{16,34}, but was underrepresented in sheep, water (and turtles), invertebrates and soil/plants. Species within the *K. orn* SPEC were significantly overrepresented in soil and plants and underrepresented in hospital carriage, which is also consistent with previous work³⁴. Other distributions were more surprising; for example, we did not find any evidence that *Klebsiella varicicola* (*K. var*) is associated with plants, contrary to its original description³⁵ and species from the *K. oxy* SPEC tended to be overrepresented in invertebrates. While this is consistent with a previous report of a symbiotic relationship between houseflies and *K. oxy*³⁶, to our knowledge this specialism has not been described before in other species of this SPEC. These data also point to a significant overrepresentation of *K. mic* in hospital carriage and we note that a small but consequential proportion (17 out of 600; 2.8%) of the diagnostic isolates from hospital disease correspond to this species.

An important caveat with this analysis is that statistical association can result from clonality rather than ecological adaptation. For

Fig. 2 | Phylogenetic tree with metadata and sample and source distributions. **a**, Maximum-likelihood phylogenetic tree constructed from core genes, coloured by species, with the SPECs shown. Only one isolate from each species is shown as this tree is intended to show the distances between species. **b**, Neighbour-joining phylogenetic tree constructed from pairwise Mash distances between all isolates, coloured by species, with the SPECs shown. The metadata rings show sources (inner rings) and resistance and virulence scores (outer rings). **c**, Bar plot showing

the number of sequenced samples from each species. The dark bars show samples from SCAI media and the transparent ones show diagnostic samples. **d**, Bar plot showing the number of sequenced samples from each high-level source. The dark bars show samples from SCAI media and the transparent ones show diagnostic samples. With the following exceptions, the three-letter species abbreviations used are explained in the main text: *Klebsiella quasipneumoniae* subsp. *similipneumoniae* (*K. qps*); *Klebsiella planticola* (*K. pla*).






Fig. 3 | Clonality and population structure. **a**, Composition of the eight most common species as determined by SC frequencies. For each species, the isolates were grouped by SC and the SCs were ranked by their frequencies as a proportion of the dataset (top 30 SCs shown). **b**, The number of unique SCs as isolates were sampled. Accumulation curves were produced by randomizing the order of the

isolates and counting the SCs, and then repeating this 100 times (mean values plotted). The dashed grey line indicates the $x = y$ line. **c**, Distribution of pairwise core genome distances for each species. The distances were estimated using PopPunk and the points were arranged in the x direction by density to show their distributions.

example, the apparent overrepresentation of *K. oxy* in turtles is due to the clonal expansion of a single lineage (*K. oxy* SC1) within a population of turtles in a pond at a botanical garden. However, we did not find evidence for clonal expansion of *K. mic* within hospital settings nor for certain *K. mic* lineages being more strongly associated with humans than others.

Distribution of resistance genes

Kleborate²⁶ assigns isolates to 1 of 4 resistance scores: 0 = low level resistance; 1 = extended-spectrum beta-lactamase (ESBL)-positive; 2 = carbapenemase-positive; and 3 = carbapenemase plus colistin-positive. The distribution of species according to these categories and to each source is shown in Figs. 2 and 5a; a full breakdown




Fig. 4 | The distribution of species according to source. Only *Klebsiella* samples from the SCAI dataset ($n = 2,795$) are shown and 23 of these samples were removed either because they were from very poorly sampled sources (21) or could not be confidently assigned to a species (2). The rows represent species delimited according to SPECTs and the columns represent sources delimited according to source categories. The grey shaded rows at the bottom of the table give the total number of positive samples for the corresponding source, and below, the total number of samples for that source. The grey shading reflects the percentage prevalence from each source. The number of positive samples are

shown for each species from each source and a blank cell indicates zero positive samples. The red shading shows the relative enrichment of each species from each source, given the overall prevalence from that source and assuming a null hypothesis whereby all species would be equally likely to be observed from any given source. The dark red and blue borders show those categories where the number of samples is significantly higher or lower than expected, respectively, as determined by a permutation test. The bar plot to the right shows the number of samples from each species and the total sampling effort.

of resistance classes is shown in Extended Data Fig. 7; 82.4% (2,870 out of 3,482) of the isolates were category 0, with those scoring 1–3 being either *K. pne* from multiple sources or isolates of other species from hospital patients (exceptions are discussed below). None of the isolates, including *K. pne*, recovered from outside a hospital setting harboured a carbapenemase gene or showed phenotypic non-susceptibility to carbapenems.

Only three isolates of species other than *K. pne* from outside the hospital setting harboured an ESBL gene; in each case, the gene in question was *bla_{SHV-12}*. These were a *K. orn* isolate recovered from a fly caught within a hospital (SPARK_2923_C1), a *K. orn* isolate from environmental water (SPARK_1613_C1) and a *Klebsiella quasivariicola* (*K. qva*) isolate from a pig (SPARK_1906_C1). Excluding *K. pne*, there were 9 isolates from other species recovered from hospital patients that harboured ESBLs (*bla_{CTX-M-15}*, $n = 4$; *bla_{SHV-12}*, $n = 5$). Of note are a pair of clonally related isolates (SPARK_1773_C1, SPARK_2031_C1) belonging to clone K.qpq_SC_11_ST571, which harboured *bla_{CTX-M-15}* plus the virulence factors *ybt*, *iro* and *rmpA*. These isolates were recovered from urine samples from two inpatients at the same hospital in April 2018. This is consistent with hospital transmission of a new *Klebsiella quasipneumoniae* subsp. *quasipneumoniae* (*K. qpq*) clone exhibiting both resistance and virulence genes.

Excluding *K. pne*, only three isolates from other species harboured a carbapenemase gene; these were all isolated from the hospital environment and carried *bla_{VIM-1}*. Two of these (*K. mic* SPARK_1816_C1 and *K. gri* SPARK_1652_C1) presented nearly identical genotypic and phenotypic resistance profiles to each other and to five isolates of *K. pne*. This resistance profile is characterized by the presence of the *bla_{SHV-12}*, *bla_{VIM-1}*, *mph(A)* and *qnrS* genes, harboured by a class 1 integron (GenBank accession no. MN783743) associated with the conjugative IncA plasmid pR210-2-VIM³⁷. This plasmid is known to circulate in multiple Enterobacteriaceae species in Italy³⁸ and the re-emergence of VIM-1 in this region is thought to reflect the increased use of ceftazidime-avibactam against *K. pneumoniae* carbapenemase (KPC)-producing bacteria. Closer analysis revealed the presence of this plasmid in distinct *K. pne* clones within a single patient and in other *Klebsiella* species (Extended Data Fig. 8).

Regarding the 1,705 isolates of *K. pne*, 1,105 (64.8%) exhibited a low level of resistance (category 0), 411 (24.1%) carried an ESBL (category 1), 175 (10.3%) carried a carbapenemase gene (category 2) and only 14 (0.8%) carried a carbapenemase gene and colistin resistance (category 3). Two ESBL genes were dominant; *bla_{CTX-M-15}* and variants of *bla_{SHV-27}*, which together accounted for 83.5% of all ESBL genes. These were distributed non-randomly between sources; 238 out of 256 (93%) of




Fig. 5 | Distribution of resistance and virulence genes according to species and source. a. Resistance genes were identified and grouped into levels 0–3 by Kleborate. **b.** Virulence genes were identified and grouped into levels 0–5 by Kleborate. The area of the circles is proportional to the number of isolates

and the text shows the number of isolates. The shading shows the proportion of isolates from a given species and source, which correspond to a given resistance or virulence level.

the *K. pne* isolates bearing *bla*_{CTX-M-15} were from humans, the exceptions being from hospital surfaces and companion animals. In contrast, only 51 out of 170 (30%) of the *K. pne* isolates bearing *bla*_{SHV-27} variants were from human sources compared to 87 out of 170 (51%) from cows. Of the 175 *K. pne* isolates harbouring carbapenemase genes, all were isolated from the hospital environment and the majority ($n = 161$; 92%) carried *bla*_{KPC} and corresponded to the healthcare-associated clones ST258/512 or ST307.

*K. pne*_ST307_SC1 was the most abundant clone in the dataset and was isolated from hospital surfaces and companion animals as well as hospital patients, although none of the ST307 isolates from non-human sources harboured *bla*_{KPC}. Eleven *K. pne* isolates harboured *bla*_{VIM-17}, including those discussed above, and 3 *K. pne* isolates harboured *bla*_{OXA-48}. Of the 192 isolates with a carbapenemase gene for which phenotypic resistance data were also available, 91% showed phenotypic resistance to ertapenem, 71.7% to imipenem and 77.7% to meropenem. In contrast, the values were 0.8% (27 isolates), 0.18% (6 isolates) and 0.28% (9 isolates), respectively, for isolates (from all species) without a carbapenemase gene; these exceptions are likely due to changes in membrane permeability³⁹. Consistent with the genotypic data, there was no evidence for any phenotypic resistance to carbapenems outside of the hospital environment.

There were 14 isolates in the highest resistance category; these were all *K. pne* isolates from hospitals and all harboured the carbapenemase gene *bla*_{KPC} plus a mutated *mgrB* gene known to confer colistin resistance. All except one of these isolates belong to the common healthcare-associated clone ST258/512, with the exception of a single ST307 isolate. Phylogenetic analysis of the 95 ST258/512 isolates suggested at least 5 acquisitions of the *mgrB* chromosomal mutation into this clone (Supplementary Fig. 1). The available phenotypic data confirmed resistance to colistin in 13 out of 14 of these isolates, 1 of which (SPARK_1222_C1) was originally assigned as sensitive using the BD Phoenix 100 automated system but was subsequently found to be resistant using the Sensititre platform. Phenotypic sensitivity to the other isolate containing *mgrB* (SPARK_372_C2) could not be confirmed because this isolate lost viability. In total, phenotypic resistance to colistin was observed in 46 *K. pne* isolates, 41 of which were from humans. Besides the 12 phenotypically resistant isolates harbouring an *mgrB* mutation, Kleborate did not detect a mechanism for colistin resistance in the other cases, including three *K. pne* isolates from pigs and a single *K. aer* isolate from a goat. This was not unexpected since many *mcr* variants are not included in the Kleborate database and colistin resistance can also be conferred through mutations responsible for membrane synthesis⁴⁰. The final non-human colistin-resistant isolate was a single *K. pne* isolate from a cow that harboured *mcr-1*.

Distribution of virulence genes

Like the genotypic resistance profiles, all isolates were assigned to 1 of 6 categories based on the presence of genes encoding the known virulence factors yersiniabactin (*ybt*), aerobactin (*iuc*), salmochelin (*iro*) and colibactin (*clb*), as identified by Kleborate (Figs. 2 and 5b); 2,749 out of 3,482 (78.9%) of all isolates and 1233 out of 1705 (72.3%) of *K. pne* isolates were in the lowest virulence category and 669 out of 3,483 (19.2%) of all isolates corresponded to virulence category 1, reflecting the presence of *ybt*, but the frequency of this locus varied markedly between species: *ybt* was present in 410 out of 1706 (24%) of the *K. pne* isolates, 249 out of 258 (96.5%) of the *K. orn* isolates, 6 out of 279 (2.1%) of the *K. var* isolates and 2 out of 171 (1.1%) of the *K. aer* isolates. The *ybt* locus in *K. orn* was assigned as an 'unknown' type by Kleborate, chromosomally located close to an transfer RNA-Asparagine site (with no evidence for an associated integrative conjugative element) and was phylogenetically distinct from the *ybt* locus in *K. pne*^{41,42}. Despite being a core locus in *K. orn*, this distinct *ybt* variant was not found in any other species, including related species from *K. orn* SPEC.

While only 7 isolates corresponded to virulence category 2 (*ybt* + *clb*), 45 *K. pne* isolates and 1 *K. oxy* isolate were assigned as virulence category 3. These isolates harboured the *iuc* locus that encodes the siderophore *iuc* and 38 out of 46 were recovered from pigs. In total, 42 out of 87 (48%) of the pig isolates harboured *iuc* and in 40 out of 42 (95%) cases harboured *iuc3*. Three *K. pne* isolates and one *K. oxy* isolate from the farm environment (water and surfaces) also harboured *iuc3*; a similar association between *iuc3* and pig isolates has recently been described in Germany⁴³. The high frequency of *iuc3* in porcine isolates contrasts with clinical isolates, in which *iuc1* and *iuc2* are more common⁴⁴. The porcine *iuc3* was observed on multiple sequence types (STs) and from different farms; hence, it is not a simple consequence of clonal spread. Preliminary analysis also suggests that *iuc3* is carried by diverse plasmids (Extended Data Fig. 9).

Twelve isolates were predicted to show a high level of virulence (categories 4 and 5). The two category 5 isolates corresponded to the hypervirulent lineage *K. pne* ST23 and contained all five virulence loci. These isolates were from patients in different hospitals and were sufficiently diverged to rule out epidemiological linkage. One of these ST23 isolates, SPARK_1158_C1, isolated from the urine of a hospital inpatient, had also acquired the resistance genes *qnrS1* and *bla*_{TEM} and exhibited phenotypic resistance to ciprofloxacin and levofloxacin. Of the ten *K. pne* isolates corresponding to category 4, four were from cases of hospital disease, four from pigs (all containing *iuc3*) and two from dogs. The two isolates from dogs, representing STs 5 and 25, harboured *ybt*, *iuc*, *iro* and *rmpA*.

The distribution of sublineages below the species level

We examined the distribution across sources of subspecies SCs as defined using PopPunk, using the same permutation test used to examine species distributions (Supplementary Figs. 2–17). This analysis revealed that different *K. pne* lineages were associated with either cows or humans (Supplementary Fig. 2) and this was also borne out by phylogenetic analysis (Supplementary Fig. 18). The lineages SC1_ST307, SC2_ST17, SC3_ST512, SC4_ST45 and SC11_ST392 were mostly associated with humans, although these varied in the degree to which they were associated with hospital carriage versus hospital disease. For example, 66% of the SC1_ST307 isolates were associated with hospital disease and 28% with hospital carriage. These figures contrasted with *K. pne*_SC2_ST17, the second most common lineage in our dataset ($n = 128$), for which the equivalent figures were 20% and 57%, respectively. Other *K. pne* SCs were associated with cows rather than humans (for example, SC5_ST661, SC9_ST3068, SC10_ST2703, SC17_ST3345). Some intermingling occurred, particularly in SC5_ST661, which contained clonal expansions of both bovine and human isolates. This lineage was previously observed from both human and bovine sources¹⁶ and may represent a more generalist clone that is adapted to and able to transmit between both cows and humans.

Fewer statistically significant SC enrichments were apparent for other species, likely due to smaller sample sizes. However, a number of observations were notable. As discussed, *K. mic* was enriched within hospital carriage (Supplementary Table 4) but this was not due to the expansion of a single SC. Twenty-five of the 30 most common SCs of this species were present in hospital carriage samples but no single SC was significantly more commonly associated with hospital carriage relative to the others (Supplementary Fig. 3). In contrast, the association of *K. gri* with invertebrates was largely driven by *K. gri* SC1 (Supplementary Fig. 7). This is unlikely to reflect clonal expansion or sampling bias since *K. gri* SC1 was associated with different invertebrate hosts (a cockroach, fly, wasp and an unspecified bug) sampled in different locations. This clone, which has no notable resistance or virulence attributes, was also recovered from a cockroach caught in a hospital environment; an isolate very closely related to this clone was recovered from an outpatient of the same hospital (Supplementary Fig. 19).




Fig. 6 | Transmission heatmaps and networks. a, b, Heatmaps showing the number of transmission events between each pair of sources, as determined by SNP thresholds of 1 (**a**) and 10 (**b**). The shading is proportional to the number of events and does not account for the number of samples from each source. **c, d,** Transmission networks showing the number of transmission events between

each pair of sources, using the same data as in the heatmaps in **a** (**c**) and **b** (**d**), except that within-source events are not shown. The nodes represent the sources and the area of the node is proportional to the number of samples from that source. The edges show the number of transmission events and the thickness of the edge is proportional to the number of events between the two sources.

Quantifying transmission

To quantify and compare transmission events between different settings, we used a single-nucleotide polymorphism (SNP) threshold-based approach (thresholds: 0, 1, 2, 5, 10, 20). It was clear from the resulting transmission matrices and networks (Fig. 6 and Supplementary Table 6) that most of the transmission occurred within a single source and, most importantly, that acquisition by humans almost always originated from other humans rather than from animals or the environment. In particular, our analysis further reinforces the view that transmission of *K. pne*, and other species, between cows and humans, which are the two most deeply sampled sources, is limited. Despite this, we note that sporadic transmission events occur relatively commonly between humans and companion animals and very occasionally between humans and other sources, including river water and invertebrates.

Discussion

The One Health framework is integral to AMR research programmes aiming to mitigate the risks posed by non-clinical reservoirs of AMR through careful surveillance and stewardship. These risks exist on multiple levels, with the simplest being sporadic transmission events, for example, from livestock to farmers. If these events are epidemiologically distinct, and onward human-to-human transmission is unlikely,

then the public health risk posed by each event is likely to be low. However, the recovery of isolates from dogs that are predicted to be hypervirulent, and the widespread healthcare-associated clone ST307 from both cats and dogs, suggest that transmission of high-risk clones between companion animals and humans may warrant close attention. We note that ST307 and ST15, the latter lineage being the most common *K. pne* lineage recovered from dogs in our data, have both previously been associated with companion animals⁴⁵.

A more complex question relates to the emergence of high-risk lineages that combine heightened virulence and/or resistance attributes with the ability to move between, and spread within, different settings. A full understanding of the emergence of such clones requires a consideration of likely anthropogenic drivers, such as inappropriate use of antibiotics or other environmental stresses, but also of the underlying ecology since free transmission will be less likely if the selective landscape is characterized by local adaptation and specialization.

Our data revealed low levels of resistance and virulence genes outside of clinical settings and within species other than *K. pne*. Given the caveat that this may in part reflect biases in the database used by Kleborate towards well-characterized genes common in *K. pne*, this suggests that the emergence and subsequent spread of highly virulent and/or resistant lineages within the environment is a rare event.

The acquisition of a variant *ybt* by *K. orn* as a core locus provides a possible exception, although the functional and epidemiological significance of this is unknown and it has not been observed in any other species. An additional example is the high frequency of porcine *K. pne* isolates that harbour the virulence locus *iuc3*. Although none of the pigs harbouring *iuc3* isolates showed signs of disease, it is likely that this *iuc* lineage is adaptive in the pig host. While much is unclear about the function and significance of the *iuc3* locus in *K. pne* isolates from pigs, and the near universal presence of *ybt* in *K. orn*, their close surveillance is warranted.

In contrast to the evidence from environmental and animal settings, our data provide evidence for the emergence of new and potentially high-risk lineages within the hospital setting, even in species other than *K. pne*. The new lineage *K. qpq* ST571, associated with hospital disease, is a worrying example since this clone harbours both resistance (*bla*_{CTX-M-15}) and virulence (*ybt*, *iro* and *rmpA*) genes. Although only two isolates of this lineage were observed, they were both isolated from urine from different patients in the same hospital and are likely to be epidemiologically linked. A second example is the interspecies transfer of the plasmid pR210-2-VIM-like carrying *bla*_{VIM-1} from *K. pne* to *K. mic* and *K. gri* isolates, again within the hospital environment, as well as the inpatient transfer of this plasmid between different *K. pne* lineages. Our data also revealed a surprisingly high rate of *K. mic* within hospital carriage, although in this case there was no evidence for the hospital spread of high-risk *K. mic* lineages. The high frequency of this species in hospital carriage, combined with the recovery of a *K. mic* strain harbouring the *bla*_{VIM-1}, as well as previous reports of strains of this species harbouring carbapenemase genes⁴⁶, urges heightened clinical awareness of this species.

We note that species and SCs within species are non-randomly distributed, with the clearest example being the distinct sets of *K. pne* SCs of human and bovine origin, which is consistent with previous studies¹⁶. Moreover, our analysis revealed that transmission is much more common within, than between settings, and that most cases of acquisition of *Klebsiella* by humans is likely to be from other humans, which has been suggested earlier for *E. coli*⁴⁷. Thus, our data and analyses broadly challenge the view that bacterial strains and the resistance genes they carry can 'flow' unimpeded between different settings and we argue that local adaptation plays a role in mitigating transmission. The ecological and phylogenetic distribution of virulence and resistance genes also points to barriers to transmission between the clinical environment, animals and the environment. High levels of virulence and/or resistance tends to be rare in species other than in *K. pne* and outside the hospital environment. The complete absence of genotypic or phenotypic evidence for carbapenem non-susceptibility outside of healthcare settings is particularly noteworthy.

In conclusion, in this study we described WGS data incorporating multiple species of the *Klebsiella* genus from diverse sources. Our findings broadly corroborate recent research indicating hospitals as the hubs of *K. pne* resistance dissemination in Europe⁵ and justify a continued focus on breaking the transmission chains throughout the healthcare network. Moreover, our analysis suggested that new lineages of heightened virulence and/or resistance are most likely to emerge within hospital settings rather than in the environment or animals, although this possibility cannot be discounted. We contend that the One Health perspective is pertinent for restricting sporadic transmission events and that transmission dynamics will vary according to the region and the pathogen under study. For example, contact between humans and animals may be much more common in many low-resource regions⁴⁸. We also recognize that the transmission dynamics of strains can be distinct from the plasmids they carry. Finally, we acknowledge limitations in our study with respect to wastewater and food-borne transmission, which may play important roles in the transmission cycle between humans, animals and the environment.

Methods

Sampling

A summary of the sampling and subsequent methodology is provided in Extended Data Fig. 1. In this study, two strategies were used to obtain samples: 5,900 samples were collected in the city of Pavia (Northern Italy) and the surrounding province between June 2017 and November 2018. Information on the samples collected is given in Supplementary Table 1. To summarize, the following types of samples were collected: stool and rectal swabs from hospital inpatients and outpatients (four different hospitals) and from a nursing home; stool from healthy volunteers; stool and rectal swabs from companion animals, farm animals and animals admitted to veterinary clinics (dogs, cats, cattle, pigs, poultry, turtles, rabbits and wild birds); invertebrates; samples of edible and ornamental plants, both wild and purchased from grocery shops, garden centres and wholesale distribution; soil samples; samples of drinking water (drinking fountains) and surface water (rivers and irrigation ditches); surface swabs from hospital, anthropic surfaces (including ATM keypads, ticket machines, buses, benches, supermarket trolleys) and farm surfaces (including enclosure, buckets and milking machines). Seven hundred and twenty-two *Klebsiella* isolates obtained from the laboratory diagnostic routine from urine, wound swabs, respiratory samples and blood cultures of hospital patients with infections were also processed. All sampling metadata was stored in Microsoft Excel 365 spreadsheets.

Metadata

Detailed metadata on all the sequenced isolates is given in Supplementary Table 2 and the Microreact project. For guidance, a summary of the main metadata fields used in the Microreact project is provided in Supplementary Table 3 and instructions on the use of this platform are available at <https://microreact.org/>. Maps showing the locations of the sequenced isolates and a sampling timeline (sequenced isolates only) is given in Fig. 1.

Sample processing

Stool and rectal swab samples (fecal swabs; Copan), both from human and animals, were enriched in Luria Bertani (LB) broth with amoxicillin (10 mg ml⁻¹) at 36 ± 1 °C for 24 h. Invertebrates were frozen for at least 24 h after sampling and surviving bacteria were recovered from the surface of the animals as well as the gut. Freezing was carried out for practical and sanitary reasons, although this procedure was not optimal due to a loss of sensitivity and potentially some sampling bias resulting from bacterial cell death. For the surface, each insect was washed with sterile water for 2 min and an aliquot of the washing was enriched in LB broth with amoxicillin (10 mg ml⁻¹). For the gut, the insect's surface was washed with ethanol 70% for 5 min (once the surface had been sampled) and then air-dried. Small insects were ground with a pestle, while larger ones were dissected with a sterile scalpel to separate the gut. The gut was then used to inoculate LB broth with amoxicillin (10 mg ml⁻¹), which was left to incubate at 36 ± 1 °C for 24 h.

For plants, each sample was divided into four portions: rhizosphere; rhizoplane; epiphyte; endophyte. The portions corresponding to the rhizosphere, rhizoplane and epiphytes were washed once with PBS (pH 7.2). The buffer used for washing was then added to the LB broth with amoxicillin (10 mg ml⁻¹) at 36 ± 1 °C for 24 h. Endophytes were washed with ethanol 70% for 2 min (once the surface had been sampled) and rinsed with sterile water before being washed with a sodium hypochlorite 2% and Triton X-100 1% solution and incubated for 2 min before washing with sterile water. Endophytes were ground once in PBS (pH 7.2) with pestle and mortar. An aliquot of 1 ml was enriched in LB broth with amoxicillin (10 mg ml⁻¹) at 36 ± 1 °C for 24 h.

Soil samples (5 g) were washed in PBS (pH 7.2), which was then added to the LB broth with amoxicillin (10 mg ml⁻¹) at 36 ± 1 °C for 24 h. Water samples (1 l for both drinking and river water) were filtered through a sterile filter unit (pore size 0.2 µm; Thermo Fisher

Scientific) and the membranes were enriched in LB broth with amoxicillin (10 mg ml^{-1}) at $36 \pm 1^\circ \text{C}$ for 24 h. For environmental water (mainly from ditches and ponds), we sampled and filtered at least 50 ml of water (higher volumes when possible) and then proceeded in the same way as the drinking and river waters. Surface swabbing was performed on areas of 10 cm^2 for each point by using a swab rinse kit (Copan). After the collection, the swab and its medium were enriched in LB broth with amoxicillin (10 mg ml^{-1}) at $36 \pm 1^\circ \text{C}$ for 24 h. For each of the above samples, $1 \mu\text{l}$ of each enrichment was plated on SCAI^{50,51} medium and the plates were incubated at $36 \pm 1^\circ \text{C}$ for 48 h.

Diagnostic isolates

In addition to the *Klebsiella* isolates recovered using selective SCAI media from diverse sources, we also assembled an additional 722 diagnostic *Klebsiella* isolates recovered from hospital patients and veterinary clinics as part of ongoing surveillance programmes within the Pavia catchment area, of which 687 were successfully sequenced: 676 (98%) of the isolates were from human clinical cases, with 484 (70%) being from a single hospital, 600 (89%) were from hospital disease and 76 (11%) were from hospital carriage. Most of the human clinical isolates were recovered from urine ($n = 417$; 62%), the other sample types being blood ($n = 74$; 11%), rectal swab ($n = 70$; 10%), bronchial ($n = 45$; 7%), sputum ($n = 20$; 3%) and other minor sources.

Species identification and antimicrobial susceptibility testing

Yellow and mucoid colonies on SCAI plates suspected to belong to the *Klebsiella* genus were identified at the species level through matrix-assisted laser desorption/ionization (MALDI)–time-of-flight (TOF) mass spectrometry (Microflex LT/SH; Bruker Daltonik GmbH) equipped with the Bruker biotyper 3.1 software. Once confirmed to be members of this genus, the isolates were subcultured on MacConkey agar for antibiotic susceptibility testing and DNA extraction. Antibiotic susceptibility was tested for all the isolates using the BD Phoenix 100 automated system and the dedicated panels NMIC-402 for all the diagnostic routine samples and NMIC-417 for all the other samples. The antibiotics and range of antibiotic concentrations present in the two panels are listed in Supplementary Table 7. We also used broth microdilution (Sensititre panel DKMGN; Thermo Fisher Diagnostics) on a subset of isolates to confirm phenotypic resistance to colistin in those cases where there were discrepancies between the genotypic and phenotypic data.

DNA extraction, sequencing and bioinformatics

For all positive samples, genomic DNA was extracted from at least one colony using a QIASymphony instrument (QIAGEN) and a dedicated kit (QIASymphony DSP Virus/Pathogen; QIAGEN). All the extracts were stored at -80°C . Genomic DNA libraries were prepared using the Nextera XT Library Prep Kit (Illumina) according to the manufacturer's protocol. Illumina sequencing was performed at 3 centres: Wellcome Trust Sanger Institute, HiSeq X10, 150 base pair (bp), paired-end reads ($n = 3,418$); University of Bath, MiSeq, 250 bp paired-end reads ($n = 110$); MicrobesNG (Birmingham), HiSeq, 200 bp paired-end reads ($n = 15$), resulting in 3,543 isolates in total, of which 3,482 were confirmed as *Klebsiella* spp. and to be of high quality. The Illumina sequence reads were trimmed with Trimmomatic v.0.33 (ref. ⁵²). SPAdes v.3.9.0 (ref. ⁵³) was used to generate de novo assemblies from the trimmed sequence reads using k -mer sizes of 41, 49, 57, 65, 77, 85 and 93 and with the `--cov-cutoff` flag set to 'auto'. The assemblies were annotated using Prokka v.1.12 (ref. ⁵⁴). Kleborate v.2.0.0 (ref. ²⁶) was used to group the isolates into *Klebsiella* species using Mash distances and assign multilocus sequence typing (MLST) types to several species. It was also used to detect antimicrobial resistance and virulence genes in the assemblies. Mob-Suite v.2.1.0 (ref. ⁵⁵) and Prokka v.1.4.5 were also used to identify and annotate plasmids; Abricate v.1.0.0 (<https://github.com/tseemann/abricate>) was used to assign

resistance genes to the plasmid contigs (PlasmidFinder and ResFinder databases downloaded on 21 May 2021 from https://bitbucket.org/genomicepidemiology/plasmidfinder_db and https://bitbucket.org/genomicepidemiology/resfinder_db, respectively). Two phylogenetic trees were generated to show species relationships across the whole genus. The first was a neighbour-joining tree of all isolates generated by RapidNJ from Mash v.2.0 (ref. ²⁴) distances. The second was a maximum-likelihood tree of a subset of 703 isolates generated by RAxML v.8.2.8 (ref. ²⁵) (GTR+ Γ substitution model) from an alignment of core genes generated by Roary v.3.12.0 (ref. ⁵⁶). Individual species trees were generated by PopPunk v.2.0.2 (ref. ³¹) core distances using the neighbour-joining method.

MALDI–TOF

All of the sequenced isolates were initially assigned to 1 of 7 *Klebsiella* species by MALDI–TOF and, apart from 1 exception, subsequently assigned to 1 of 15 species on the basis of the genome sequences using Kleborate and phylogenetics. The accuracy of the MALDI–TOF assignments varied according to species; 88.4% of the isolates assigned as *K. pneumoniae* by MALDI–TOF were confirmed by WGS, whereas only 30% of the isolates assigned as *K. oxytoca* were confirmed as this species. These discrepancies reflect the fact that reference databases are currently unable to distinguish *K. oxytoca* from closely related species (Supplementary Table 8).

Lineage assignment at the subspecies level

Kleborate was used to assign STs to the isolates but this was only possible for those species for which MLST schemes have been previously established. Therefore, we carried out intraspecies lineage assignments into SCs for all species using PopPunk v.2.0.2. For each species, the number of components to fit in the mixture model (k) was chosen based on the scatter plot of core and accessory distances. The model was then fitted and the boundary refined using an iterative process of moving the boundary and reassessing the network features. In all cases, the core boundary was used to define the clusters. For most species, 2 components provided the best fit, the exceptions being *K. aer* ($k = 6$), *K. mic*, *K. qppq* and *K. ter* ($k = 3$). For two species, there were outlying isolates (SPARK_1532_C1, SPARK_1532_C2; SPARK_1553_C1, SPARK_871_C1), which were removed to fit the model and then re-assigned as query sequences. Plots of the final fits are shown in Extended Data Fig. 5. The SCs defined by PopPunk were named according to their relative abundance within each species, with SC1 being the most abundant, followed by SC2 and so on. For those species where STs could also be identified by Kleborate, the SCs defined by PopPunk matched closely with STs (Rand Index > 0.98), although SCs tended to be slightly more inclusive groupings. For ease of reference, we also used a compound identifier that combined SC with the corresponding canonical ST (for example, the most common group in *K. pne* was designated *K.pne*_SC1_ST307).

Inferring transmission events

To quantify the frequency of transmission events within and between different sources, we aggregated data over all *Klebsiella* species and identified transmission events using a threshold-based approach based on SNP distances using a mapping procedure. Briefly, for each SC in the dataset we used the 'sample_n' function from the dplyr v.0.8.3 package in R v.3.6.1 to randomly choose a reference isolate to represent the SC. This was an objective and practical solution, given the high diversity of the dataset (1,168 SCs across 15 species). We mapped all isolates from a given SC to this random SC-specific reference using Snippy v.4.6.0 (<https://github.com/tseemann/snippy>), with variant calling by freebayes v.1.3.2 and the parameters `--mapqual 60`, `--basequal 13`, `--mincov 10`, `--minfrac AUTO`, `--minqual 100`, `--maxsoft 10`. We then used snp-dists v.0.7.0 (<https://github.com/tseemann/snp-dists>) to count the number of SNPs between each pair of isolates from that SC.

For each SC, we created a network where all isolates were connected to all other isolates and then pruned this network into 'putative transmission clusters' by removing all the links between isolates where the SNP distance was greater than the chosen threshold SNP distance (an example is given in Extended Data Fig. 10). For each of these putative transmission clusters, we recorded every unique source pair (including same source pairs) connected by a pair of isolates differing by an SNP distance \leq threshold. We excluded pairs of samples from the same individual hosts. Each unique source pair (not isolate pair) within each cluster, which satisfied these criteria, was recorded as a single transmission event. This approach is conservative because multiple transmission events occurring within a cluster between any given source pair are only counted once, but this avoids double counting of single transmission events. To obtain transmission frequencies, we normalized the count of transmission events for each source pair by the total number of isolates in each pair of sources. We carried out this analysis using 6 different SNP thresholds: 0, 1, 2, 5, 10 and 20 SNPs. The upper threshold (20 SNPs) was informed by a recent analysis of *K. pne* transmission within and between healthcare settings⁵.

Statistics and plotting

All statistics and plotting were performed with R v.3.6.1. The permutation tests were done by randomly permuting the source categories and cluster labels (either species or SC depending on the dataset) 10,000 times and then deriving a *P* value by dividing the number of permutations that were at least as extreme as the observed data by 10,000. The test was two-tailed; therefore, a significance level of 0.025 was used at each tail. The *P* values were corrected using the Benjamini–Hochberg method from the *p.adjust* function in R. Other tests of association were done using the chi-squared test. All the plotting was done in ggplot2 v.3.3.5 and the multi-panel plots were produced using cowplot v.1.1.0.

Ethical approval

This study was approved by the ethical committee of the San Matteo Hospital in Pavia under no. 20170001787 on 25 May 2017. The proceeding number is 2017000759 and the internal code of the project is 0890170117. The ethical procedure includes written informed consent from all the patients participating in the study.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

All raw Illumina sequence data are available from the European Nucleotide Archive under accession no. [PRJEB27342](https://www.ebi.ac.uk/ena/record/PRJEB27342). The individual accession numbers for all sequenced isolates are in Supplementary Table 9. The metadata and phylogenetic analysis are available for download from the Microreact project <https://microreact.org/project/KLEB-PAVIA>; additional metadata is provided in Supplementary Tables 1 and 2. The PlasmidFinder and ResFinder databases were downloaded on 21 May 2021 from https://bitbucket.org/genomicpidemiology/plasmidfinder_db and https://bitbucket.org/genomicpidemiology/resfinder_db, respectively. Map tiles by Stamen Design, under CC BY 3.0. Data by OpenStreetMap, under ODbL.

Code availability

The R code for the transmission analysis and permutation tests is available from https://github.com/harry-thorpe/SPARK_code.

References

- Taconelli, E. et al. Discovery, research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis. *Lancet Infect. Dis.* **18**, 318–327 (2018).
- Knothe, H., Shah, P., Krcmery, V., Antal, M. & Mitsuhashi, S. Transferable resistance to cefotaxime, ceftiofur, cefamandole and cefuroxime in clinical isolates of *Klebsiella pneumoniae* and *Serratia marcescens*. *Infection* **11**, 315–317 (1983).
- Yigit, H. et al. Novel carbapenem-hydrolyzing β -lactamase, KPC-1, from a carbapenem-resistant strain of *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* **45**, 1151–1161 (2001).
- Wilson, H. & Török, M. E. Extended-spectrum β -lactamase-producing and carbapenemase-producing Enterobacteriaceae. *Microb. Genom.* **4**, e000197 (2018).
- David, S. et al. Epidemic of carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by nosocomial spread. *Nat. Microbiol.* **4**, 1919–1929 (2019).
- Köck, R. et al. Carbapenem-resistant Enterobacteriaceae in wildlife, food-producing, and companion animals: a systematic review. *Clin. Microbiol. Infect.* **24**, 1241–1250 (2018).
- Gibbon, M. et al. A high prevalence of *bla*_{OXA-48} in *Klebsiella (Raoultella) ornithinolytica* and related species in hospital wastewater in South West England. *Microb. Genom.* **7**, mgen000509 (2021).
- Mills, M. C. & Lee, J. The threat of carbapenem-resistant bacteria in the environment: evidence of widespread contamination of reservoirs at a global scale. *Environ. Pollut.* **255**, 113143 (2019).
- Walsh, T. R. A one-health approach to antimicrobial resistance. *Nat. Microbiol.* **3**, 854–855 (2018).
- Baquero, F., Coque, T. M., Martínez, J.-L., Aracil-Gisbert, S. & Lanza, V. F. Gene transmission in the One Health microbiosphere and the channels of antimicrobial resistance. *Front. Microbiol.* **10**, 2892 (2019).
- Balloux, F. et al. From theory to practice: translating whole-genome sequencing (WGS) into the clinic. *Trends Microbiol.* **26**, 1035–1048 (2018).
- Gouliouris, T. et al. Quantifying acquisition and transmission of *Enterococcus faecium* using genomic surveillance. *Nat. Microbiol.* **6**, 103–111 (2021).
- Robinson, T. P. et al. Antibiotic resistance is the quintessential One Health issue. *Trans. R. Soc. Trop. Med. Hyg.* **110**, 377–380 (2016).
- Ludden, C. et al. One Health genomic surveillance of *Escherichia coli* demonstrates distinct lineages and mobile genetic elements in isolates from humans versus livestock. *mBio* **10**, e02693–18 (2019).
- Gouliouris, T. et al. Genomic surveillance of *Enterococcus faecium* reveals limited sharing of strains and resistance genes between livestock and humans in the United Kingdom. *mBio* **9**, e01780–18 (2018).
- Ludden, C. et al. A One Health study of the genetic relatedness of *Klebsiella pneumoniae* and their mobile elements in the East of England. *Clin. Infect. Dis.* **70**, 219–226 (2020).
- Schubert, H. et al. Reduced antibacterial drug resistance and *bla*_{CTX-M} β -lactamase gene carriage in cattle-associated *Escherichia coli* at low temperatures, at sites dominated by older animals, and on pastureland: implications for surveillance. *Appl. Environ. Microbiol.* **87**, e01468–20 (2021).
- Hanage, W. P. Two health or not two health? That is the question. *mBio* **10**, e00550–19 (2019).
- Shen, Y. et al. Anthropogenic and environmental factors associated with high incidence of *mcr-1* carriage in humans across China. *Nat. Microbiol.* **3**, 1054–1062 (2018).
- Boon, R. D. et al. One Health drivers of antibacterial resistance: quantifying the relative impacts of human, animal and environmental use and transmission. *One Health* **12**, 100220 (2021).
- Stanton, I. C., Bethel, A., Leonard, A. F. C., Gaze, W. H. & Garside, R. What is the research evidence for antibiotic resistance exposure and transmission to humans from the environment? A systematic map protocol. *Environ. Evid.* **9**, 12 (2020).

22. Holmes, A. H. et al. Understanding the mechanisms and drivers of antimicrobial resistance. *Lancet* **387**, 176–187 (2016).
23. Ma, Y. et al. Proposal for reunification of the genus *Raoultella* with the genus *Klebsiella* and reclassification of *Raoultella electrica* as *Klebsiella electrica* comb. nov. *Res. Microbiol.* **172**, 103851 (2021).
24. Ondov, B. D. et al. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol.* **17**, 132 (2016).
25. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
26. Lam, M. M. C. et al. A genomic surveillance framework and genotyping tool for *Klebsiella pneumoniae* and its related species complex. *Nat. Commun.* **12**, 4188 (2021).
27. Merla, C. et al. Description of *Klebsiella spallanzanii* sp. nov. and of *Klebsiella pasteurii* sp. nov. *Front. Microbiol.* **10**, 2360 (2019).
28. Hu, Y., Wei, L., Feng, Y., Xie, Y. & Zong, Z. *Klebsiella huaxiensis* sp. nov., recovered from human urine. *Int. J. Syst. Evol. Microbiol.* **69**, 333–336 (2019).
29. Batisti Biffignandi, G. et al. Genome of *Superficieibacter maynardsmithii*, a novel, antibiotic susceptible representative of Enterobacteriaceae. *G3 (Bethesda)* **11**, jkab019 (2021).
30. Wyres, K. L., Lam, M. M. C. & Holt, K. E. Population genomics of *Klebsiella pneumoniae*. *Nat. Rev. Microbiol.* **18**, 344–359 (2020).
31. Lees, J. A. et al. Fast and flexible bacterial genomic epidemiology with PopPUNK. *Genome Res.* **29**, 304–316 (2019).
32. Huynh, B.-T. et al. *Klebsiella pneumoniae* carriage in low-income countries: antimicrobial resistance, genomic diversity and risk factors. *Gut Microbes* **11**, 1287–1299 (2020).
33. Holt, K. E. et al. Genomic analysis of diversity, population structure, virulence, and antimicrobial resistance in *Klebsiella pneumoniae*, an urgent threat to public health. *Proc. Natl Acad. Sci. USA* **112**, E3574–E3581 (2015).
34. Bagley, S. T. Habitat association of *Klebsiella* species. *Infect. Control* **6**, 52–58 (1985).
35. Rosenblueth, M., Martínez, L., Silva, J. & Martínez-Romero, E. *Klebsiella varicola*, a novel species with clinical and plant-associated isolates. *Syst. Appl. Microbiol.* **27**, 27–35 (2004).
36. Lam, K., Thu, K., Tsang, M., Moore, M. & Gries, G. Bacteria on housefly eggs, *Musca domestica*, suppress fungal growth in chicken manure through nutrient depletion or antifungal metabolites. *Naturwissenschaften* **96**, 1127–1132 (2009).
37. Dong, N. et al. Evolution of carbapenem-resistant serotype K1 hypervirulent *Klebsiella pneumoniae* by acquisition of *bla_{VIM-1}*-bearing plasmid. *Antimicrob. Agents Chemother.* **63**, e01056–19 (2019).
38. Arcari, G. et al. A multispecies cluster of VIM-1 carbapenemase-producing *Enterobacterales* linked by a novel, highly conjugative, and broad-host-range IncA plasmid forebodes the reemergence of VIM-1. *Antimicrob. Agents Chemother.* **64**, e02435–19 (2020).
39. Navon-Venezia, S., Kondratyeva, K. & Carattoli, A. *Klebsiella pneumoniae*: a major worldwide source and shuttle for antibiotic resistance. *FEMS Microbiol. Rev.* **41**, 252–275 (2017).
40. Ayoub Moubareck, C. Polymyxins and bacterial membranes: a review of antibacterial activity and mechanisms of resistance. *Membranes (Basel)* **10**, 181 (2020).
41. Wang, M. et al. Genomic insights into evolution of pathogenicity and resistance of multidrug-resistant *Raoultella ornithinolytica* WM1. *Ann. N. Y. Acad. Sci.* **1497**, 74–90 (2021).
42. Lam, M. M. C. et al. Genetic diversity, mobilisation and spread of the yersiniabactin-encoding mobile element ICEKp in *Klebsiella pneumoniae* populations. *Microb. Genom.* **4**, e000196 (2018).
43. Klaper, K., Hammerl, J. A., Rau, J., Pfeifer, Y. & Werner, G. Genome-based analysis of *Klebsiella* spp. isolates from animals and food products in Germany, 2013–2017. *Pathogens* **10**, 573 (2021).
44. Lam, M. M. C. et al. Tracking key virulence loci encoding aerobactin and salmochelin siderophore synthesis in *Klebsiella pneumoniae*. *Genome Med.* **10**, 77 (2018).
45. Garcia-Fierro, R. et al. Comparative phylogenomics of ESBL-, AmpC- and carbapenemase-producing *Klebsiella pneumoniae* originating from companion animals and humans. *J. Antimicrob. Chemother.* **77**, 1263–1271 (2022).
46. Seiffert, S. N. et al. First clinical case of KPC-3-producing *Klebsiella michiganensis* in Europe. *New Microbes New Infect.* **29**, 100516 (2019).
47. Mughini-Gras, L. et al. Attributable sources of community-acquired carriage of *Escherichia coli* containing β -lactam antibiotic resistance genes: a population-based modelling study. *Lancet Planet. Health* **3**, e357–e369 (2019).
48. Osei Sekyere, J. & Reta, M. Genomic and resistance epidemiology of gram-negative bacteria in Africa: a systematic review and phylogenomic analyses from a One Health perspective. *mSystems* **5**, e00897–20 (2020).
49. Connor, T. R. et al. CLIMB (the Cloud Infrastructure for Microbial Bioinformatics): an online resource for the medical microbiology community. *Microb. Genom.* **2**, e000086 (2016).
50. Van Kregten, E., Westerdal, N. A. & Willers, J. M. New, simple medium for selective recovery of *Klebsiella pneumoniae* and *Klebsiella oxytoca* from human feces. *J. Clin. Microbiol.* **20**, 936–941 (1984).
51. Passet, V. & Brisse, S. Association of tellurite resistance with hypervirulent clonal groups of *Klebsiella pneumoniae*. *J. Clin. Microbiol.* **53**, 1380–1382 (2015).
52. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).
53. Bankevich, A. et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* **19**, 455–477 (2012).
54. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).
55. Robertson, J. & Nash, J. H. E. MOB-suite: software tools for clustering, reconstruction and typing of plasmids from draft assemblies. *Microb. Genom.* **4**, e000206 (2018).
56. Page, A. J. et al. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* **31**, 3691–3693 (2015).

Acknowledgements

This work was funded by the SpARK project, awarded to E.J.F., ‘The rates and routes of transmission of multidrug-resistant *Klebsiella* clones and genes into the clinic from environmental sources’, which has received funding under the 2016 Joint Programming Initiative on Antimicrobial Resistance call ‘Transmission dynamics’ (medical research council (MRC) reference no. MR/R00241X/1) and by the French Government’s Investissement d’Avenir program Laboratoire d’Excellence ‘Integrative Biology of Emerging Infectious Diseases’ (no. ANR-10-LABX-62-IBEID). J.C. and H.A.T. were funded by the European Research Council grant no. 742158. J.C. and T.K. were funded by the Norwegian Research Council grant no. 271162. The use of the MRC Cloud Infrastructure for Microbial Bioinformatics⁴⁹ was critical for the computational aspects of this work. We thank R. Zadoks and A. McNally for advice during the course of the project.

Author contributions

H.A.T. carried out the extensive bioinformatics and statistical analysis and helped write the paper and design the study. R.B. developed and implemented the methods with input from L.M. and R.R. T.K., M.J.G., N.C., V.P., S.L.F., C.R., S.M., S.D. and F.C. helped with data analysis and manuscript preparation. C.M., M.C. and C.F. carried out the sampling,

microbiology and susceptibility testing. P.M. helped with the clinical sampling logistics. S.B., D.S., J.C. and E.J.F. designed the study, contributed to the analysis and prepared the paper.

Competing interests

The authors declare no competing interests.

Additional information

Extended data is available for this paper at <https://doi.org/10.1038/s41564-022-01263-0>.

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41564-022-01263-0>.

Correspondence and requests for materials should be addressed to Edward J. Feil.

Peer review information *Nature Microbiology* thanks the anonymous reviewers for their contribution to the peer review of this work.


Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.


Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022


¹Department of Biostatistics, University of Oslo, Oslo, Norway. ²Bristol Veterinary School, University of Bristol, Bristol, UK. ³Department of Clinical Microbiology, Turku University Hospital, Turku, Finland. ⁴The Milner Centre for Evolution, Department of Life Sciences, University of Bath, Bath, UK. ⁵Institut Pasteur, Université Paris Cité, Biodiversity and Epidemiology of Bacterial Pathogens, Paris, France. ⁶Boyd Orr Centre for Population and Ecosystem Health, School of Biodiversity, One Health and Veterinary Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, UK. ⁷Parasites and Microbes, Wellcome Sanger Institute, Cambridge, UK. ⁸Microbiology and Virology Unit, Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo, Pavia, Italy. ⁹Romeo ed Enrica Invernizzi Pediatric Research Center, Department of Biomedical and Clinical Sciences Luigi Sacco, Università di Milano, Milan, Italy. ¹⁰Department of Biology and Biotechnology, Università di Pavia, Pavia, Italy. ¹¹Department of Mathematics and Statistics, Helsinki Institute of Information Technology, University of Helsinki, Helsinki, Finland. ¹²These authors contributed equally: Davide Sasseria, Jukka Corander, Edward J. Feil. ✉e-mail: e.feil@bath.ac.uk




Extended Data Fig. 1 | Summary of the sampling and subsequent methodology. Flowchart showing the number of initial samples taken, and the number of remaining samples after each processing step.




Extended Data Fig. 2 | *K. orn* SPEC phylogeny. Circular phylogenetic tree for *K. ornitholytica* and related species within the species complex *K. orn* SPEC. The tree was generated from Mash distances using the neighbour-joining method.



Extended Data Fig. 3 | *K.oxy* SPEC phylogeny. Circular phylogenetic tree for *K. oxytoca* and related species within the species complex *K.oxy* SPEC. The tree was generated from Mash distances using the neighbour-joining method.




Extended Data Fig. 4 | *K.pne* SPEC phylogeny. Circular phylogenetic tree for *K. pneumoniae* and related species within the species complex *K.pne* SPEC. The tree was generated from Mash distances using the neighbour-joining method.




Extended Data Fig. 5 | Lineage assignment with PopPunk. PopPunk was used to divide each species into sequence clusters (SCs). For each species, the number of components to fit in the mixture model (k) was chosen based on the scatter plot of core and accessory distances. The model was then fit, and the boundary

refined using an iterative process of moving the boundary and reassessing the network features. In all cases the core boundary was used to define the clusters. The number of isolates, number of SCs, model K , and core distance boundaries are shown for each species.




Extended Data Fig. 6 | Phylogenies of individual species. Core genome phylogenetic trees for each species were generated using the neighbour-joining method.




Extended Data Fig. 7 | The distribution of resistance genes according to species and source. Detailed breakdown of the distribution of all resistance gene classes identified by Kleborate by species and source. The area of the circles

is proportional to the number of isolates, and the text shows the number of isolates. The shading shows the proportion of isolates from a given species and source which harbour at least one gene from that class.




Extended Data Fig. 8 | Alignment of *bla*_{VIM} plasmids. Alignment of the MN783743 and CP034084 reference plasmids with the contigs identified in our data as carrying the *bla*_{VIM} gene in Inca/C plasmids. Arrows represent the *bla*_{VIM} gene (turquoise), conjugal transfer system (green), resistance genes (red), mobile elements (yellow), hypothetical proteins (plum), replication protein *repA* (orange), integrases (dark blue) and the mercury resistance operon (pink). The three bracketed SPARK_315 SCAI carriage isolates were obtained from the same faecal sample from an outpatient in Hospital I (H1). The plasmid was found in two hospitals (H1 and H2). Two of these (SPARK_315_C1 and SPARK_315_C2) were members of the same clonal lineage (SC152_ST2418), whilst SPARK_315_C3 was an unrelated isolate related to the hypervirulent lineage ST23 (SC299_ST23-2LV). The plasmids are however essentially identical within all three isolates (which

is identical to CP034084), indicating within-patient transfer. SPARK_355_C1 is a diagnostic carriage sample from a rectal swab from a different inpatient in Hospital I corresponding to lineage *K.pne*_SC66_ST253 and also contains the same VIM plasmid. SPARK_1466_C1 also contains a very similar plasmid. This isolate corresponds to a different lineage (*K.pne*_SC195_ST606) and is a diagnostic isolate from a urine sample from inpatient in Hospital 2. SPARK_1816_C1 is a *K.mic* diagnostic isolate (SC14) from a rectal swab from inpatient in Hospital I and contains also contains this plasmid. Finally, SPARK_1652_C1 is a *K.gri* diagnostic isolate (SC44), also from a urine sample from an outpatient in hospital I. This isolate contains the same plasmid, but with a deletion near the resistance cassette.



Extended Data Fig. 9 | Comparison of *iuc3*-containing contigs from *K.pne*. Comparison of short-read contigs from *K.pne* isolates from pigs harbouring *iuc3* in this study. The ST and position of *iuc3* are shown. Contigs were compared with the Artemis Comparison Tool (ACT); sections of similarity of >500 bp are shown.

K.pne_SC5_ST661



Extended Data Fig. 10 | Transmission clusters. Networks were used to identify transmission clusters using multiple SNP thresholds. To ensure that the estimates were conservative, only one event was counted for each pair of sources that satisfied the threshold in each component of the network. These events are denoted by the red edges. Only one SC is shown as an example of the method.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

- n/a Confirmed
- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
 - A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
 - The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
 - A description of all covariates tested
 - A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
 - A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
 - For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value
Noted. Give P values as exact values whenever suitable.
 - For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
 - For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
 - Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection Excel 365 was used to compile and record the list of samples as they were being generated

Data analysis Bruker Biotyper 3.1 was used to interpret the data from Maldi-ToF for species ID
 Trimmomatic v. 0.33 was used to trim the Illumina reads
 SPAdes v3.9.0 was used to generate de novo assemblies
 Prokka v1.12 was used to annotate the genome assemblies, and Prokka v1.4.5 was used to annotate the plasmids
 Kleborate v2.0.0 was used to assign species and ST (and call resistance and virulence genes) from the assemblies
 PopPunk v2.0.2 was used to delimit sequence clusters (SCs)
 Snippy v4.6.0 was used for mapping
 Abricate v1.0.0 was used to assign AMR genes to contigs
 RapidNJ, (no version information available), and RaXML v8.2.8 were used for phylogenetic analysis
 R code for the transmission analysis and permutation tests has been deposited in github:
https://github.com/harry-thorpe/SPARK_code

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres [to our policy](#)

Data availability: Short-read data available under accession numbers ERR3412430 to ERR3412448;ERR3440341 to ERR3440427;ERR3448863 to ERR3449598;ERR3469775 to ERR3469909;ERR3479903 to ERR3480717;ERR3844616 to ERR3844777;ERR3904469 to ERR3904709;ERR3931787 to ERR3932313;ERR3967745 to ERR3967936;ERR4022833 to ERR4023150;ERR4139181 to ERR4139191;ERR4374646 to ERR4374837. The metadata and tree are available to download from the Microreact project at <https://microreact.org/project/KLEBPAVIA>. PlasmidFinder and Resfinder databases downloaded on 21/05/2021 from https://bitbucket.org/genomicpidemiology/plasmidfinder_db and https://bitbucket.org/genomicpidemiology/resfinder_db, respectively. Map tiles by Stamen Design, under CC BY 3.0. Data by OpenStreetMap, under ODbL.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	The study focussed on the characterisation -by whole-genome sequencing- of a large collection of bacteria within to the genus <i>Klebsiella</i> . The aim was to examine the likely role of one-health reservoirs of <i>Klebsiella</i> in the emergence and spread of antimicrobial resistance.
Research sample	The <i>Klebsiella</i> isolates were sampled from multiple clinical, community, animal and environmental settings. Animal samples were from swabs only.
Sampling strategy	Most isolates were recovered by culturing on selective media (SCAI). Clinical isolates were acquired as part of ongoing surveillance projects in the Pavia region. The final sample size (3482 complete genomes) is one of the largest studies of its kind and is sufficient to draw inferences regarding transmission frequencies between different settings. No size calculation was carried out as we were not aiming to capture all the diversity within these populations.
Data collection	Metadata (place, time, source information) pertaining to all the samples (both positive and negative) were recorded by members of the team in Pavia. The sequence data was generated on the Illumina platform at the Wellcome Trust Sanger Institute in Cambridge, UK.
Timing and spatial scale	The data were collected in and around the Northern Italian city of Pavia between June 2017 and November 2018.
Data exclusions	A small number of genome sequences were excluded on the basis that they did not pass QC.
Reproducibility	Multiple samples were taken from each source
Randomization	The samples were grouped by source (eg human, cow, river water, rhizosphere etc). These are discrete categories.
Blinding	The isolates were cultured without selecting for distinct species or resistance profile. Species ID by WGS was derived independently from species ID by Maldi-ToF.
Did the study involve field work?	<input checked="" type="checkbox"/> Yes <input type="checkbox"/> No

Field work, collection and transport

Field conditions	Environmental parameters such as temperature and rainfall were not recorded.
Location	In and around the Northern Italian city of Pavia
Access & import/export	Permission was granted for access to farms, veterinary clinics and for clinical samples. All appropriate procedures were followed when transporting the isolates.
Disturbance	NA

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input checked="" type="checkbox"/>	<input type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input type="checkbox"/>	<input checked="" type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern

Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Human research participants

Policy information about [studies involving human research participants](#)

Population characteristics	The study was focused on bacterial isolates from stools and rectal swabs from humans (and other non-human biological materials). Full anonymity was provided to the enrolled persons, with only the collection of limited metadata useful for the study (these are available in a linked microreact project and separately in a supplementary table). No additional data from the patients are available to the authors, no follow up was performed.
Recruitment	<p>The samples included stool and rectal swabs from hospital inpatients and outpatients (four different hospitals) and from a nursing home; stool from healthy volunteers.</p> <p>For patients, no selection was performed, and all the samples obtained by the research group through the clinical laboratory of the San Matteo Hospital in Pavia was analyzed, up to the monthly analytical capacity of the research Group.</p> <p>For healthy volunteers, no selection was performed. The study was publicized by word of mouth and through the channels of the Hospital and of the University of Pavia. All the volunteers that contacted us were enrolled. Limited metadata was required (see above) and full anonymity was guaranteed.</p>
Ethics oversight	This study was approved by the Ethical Committee of the San Matteo Hospital in Pavia under number 20170001787 in date 25/05/2017. The proceeding number is 2017000759 and the internal code of the project is 0890170117. The ethical procedure includes written informed consent from all the patients participating in the study.

Note that full information on the approval of the study protocol must also be provided in the manuscript.