

Anthropometric measures and adverse outcomes in heart failure with reduced ejection fraction: revisiting the obesity paradox

Jawad H. Butt^{1,2}, Mark C Petrie ¹, Pardeep S. Jhund¹, Naveed Sattar ¹, Akshay S. Desai³, Lars Køber², Jean L. Rouleau⁴, Karl Swedberg⁵, Michael R. Zile⁶, Scott D. Solomon³, Milton Packer ⁷, and John J.V. McMurray ^{1*}

¹British Heart Foundation Cardiovascular Research Centre, University of Glasgow, Glasgow, UK; ²Department of Cardiology, Copenhagen University Hospital—Rigshospitalet, Copenhagen, Denmark; ³Division of Cardiovascular Medicine, Brigham and Women's Hospital, Boston, MA, USA; ⁴Institut de Cardiologie de Montréal, Université de Montréal, Montréal, QC, Canada; ⁵Department of Molecular and Clinical Medicine, University of Gothenburg, Gothenburg, Sweden; ⁶Department of Medicine, Medical University of South Carolina and Ralph H. Johnson Veterans Administration Medical Center, Charleston, South Carolina, USA; and ⁷Baylor Heart and Vascular Institute, Baylor University Medical Center, Dallas, TX, USA

Received 13 September 2022; revised 9 January 2023; accepted 2 February 2023

See the editorial comment for this article 'Revisiting the obesity paradox in heart failure: what is the best anthropometric index to gauge obesity?', by R. Sato and S. von Haehling, <https://doi.org/10.1093/eurheartj/ehad079>.

Abstract

Aims

Although body mass index (BMI) is the most commonly used anthropometric measure, newer indices such as the waist-to-height ratio, better reflect the location and amount of ectopic fat, as well as the weight of the skeleton, and may be more useful.

Methods and results

The prognostic value of several newer anthropometric indices was compared with that of BMI in patients with heart failure (HF) and reduced ejection fraction (HFrEF) enrolled in prospective comparison of ARNI with ACEI to determine impact on global mortality and morbidity in heart failure. The primary outcome was HF hospitalization or cardiovascular death. The association between anthropometric indices and outcomes were comprehensively adjusted for other prognostic variables, including natriuretic peptides. An 'obesity-survival paradox' related to lower mortality risk in those with BMI ≥ 25 kg/m² (compared with normal weight) was identified but this was eliminated by adjustment for other prognostic variables. This paradox was less evident for waist-to-height ratio (as an exemplar of indices not incorporating weight) and eliminated by adjustment: the adjusted hazard ratio (aHR) for all-cause mortality, for quintile 5 vs. quintile 1, was 1.10 [95% confidence interval (CI) 0.87–1.39]. However, both BMI and waist-to-height ratio showed that greater adiposity was associated with a higher risk of the primary outcome and HF hospitalization; this was more evident for waist-to-height ratio and persisted after adjustment e.g. the aHR for HF hospitalization for quintile 5 vs. quintile 1 of waist-to-height ratio was 1.39 (95% CI 1.06–1.81).

Conclusion

In patients with HFrEF, alternative anthropometric measurements showed no evidence for an 'obesity-survival paradox'. Newer indices that do not incorporate weight showed that greater adiposity was clearly associated with a higher risk of HF hospitalization.

* Corresponding author. Tel: +44 141 3303479, Fax: +44 141 330 6955, Email: john.mcmurray@glasgow.ac.uk

© The Author(s) 2023. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Structured Graphical Abstract

Key Question


This study assesses newer anthropometric indices in patients with heart failure and reduced ejection fraction (HF_rEF) enrolled in PARADIGM-HF, focusing on their prognostic value and on the “obesity paradox”

Key Finding

The “obesity survival paradox” related to body mass index was eliminated by comprehensive adjustment for prognostic variables. Alternative anthropometric measurements showed less evidence of this paradox, and two indices that incorporate waist circumference and height, but not weight, showed a clearer association between greater adiposity and a higher risk of heart failure (HF) hospitalization.

Take Home Message

These findings collectively suggest a need to test the potential benefits of intentional weight loss in patients living with obesity and HF_rEF.



The upper part of the figure describes the calculation of each of the anthropometric measures. The lower part of the figure shows the risk of outcomes according to continuous body mass index (left panel) and waist-to-height ratio (right panel). The solid line represents the hazard ratio and the shaded area the 95% confidence interval. The blue spline is adjusted for treatment and region. The red spline is adjusted for treatment, age, sex, region, systolic blood pressure, heart rate, estimated glomerular filtration rate, left ventricular ejection fraction, log of *n*-terminal pro-B-type natriuretic peptide, body mass index (only in the waist-to-height ratio analyses), New York Heart Association functional class, heart failure aetiology, duration of heart failure, prior heart failure hospitalization, a history of diabetes, and atrial fibrillation. BMI, body mass index; BRI, body roundness index; BSA, body surface area; BSI, body shape index; CI, confidence interval; HF, heart failure; HR, hazard ratio; RFM, relative fat mass; WHR, waist-to-hip ratio; WHtR, waist-to-height ratio; WWI, weight-adjusted-waist index.

Keywords

Heart failure with reduced ejection fraction • Obesity • Body mass index • Angiotensin receptor-neprilysin inhibitor • Clinical trial

Introduction

Although obesity has been repeatedly shown to be an independent risk factor for the development of heart failure (HF),^{1–3} its prognostic importance in established HF, especially HF with reduced ejection fraction (HFrEF), is less clear and an 'obesity-survival paradox' has been described in patients with HFrEF.^{4–8} However, the associations between obesity and outcomes in HFrEF have generally been based on body mass index (BMI), calculated as weight in kilograms divided as height in meters squared, which has many limitations as a measure of adiposity. BMI does not take into account the location of body fat or its amount, relative to muscle, or the weight of the skeleton, which may differ according to sex, age, and race.^{9–12} In HF specifically, there is also the contribution of retained fluid to body weight. Consequently, alternative anthropometric indices have been proposed such as waist circumference, waist-to-hip ratio, and weight-adjusted-weight index, which may better reflect intra-abdominal fat ('central obesity') and body shape index, body roundness index, and relative fat mass, which have been suggested to better reflect the distribution of body fat and total fat mass.^{13–17} Recently, some of these were found to be better predictors of incident HF in the general population than BMI,^{18–20} but they are not commonly measured in clinical practice.

Waist-to-height ratio is of particular interest as it should, to some extent, take account of sex- and race-based differences in stature and the distribution of body fat, and the National Institute for Health and Care Excellence in the United Kingdom has recently suggested that waist-to-height ratio should replace BMI in the evaluation of adiposity.^{21,22}

To complicate matters further, the association between BMI (and potentially any other anthropometric index) and outcome in patients with HFrEF is also confounded by the relationship between adiposity and natriuretic peptide levels. Higher BMI is associated with lower natriuretic peptide levels, possibly due to increased clearance or potentially because patients with obesity present with symptoms at an earlier stage in the development of HF.^{23,24} Although natriuretic peptide level is one of the most powerful prognostic variables in HFrEF,²⁵ few analyses of the association between BMI (or other anthropometric indices) and outcome have accounted for this.^{4–8,26–34} Therefore, we have examined the newer anthropometric indices described above in patients with HFrEF, focusing on their prognostic value and whether an 'obesity-survival paradox' is observed, as has been reported for BMI. We have also adjusted all analyses for natriuretic peptide levels. We carried out these analyses in a global population enrolled in the prospective comparison of angiotensin receptor-neprilysin inhibitor with angiotensin-converting enzyme inhibitor (ACE-i) to determine impact on global mortality and morbidity in heart failure trial (prospective comparison of ARNI with ACEi to determine impact on global mortality and morbidity in heart failure [PARADIGM-HF]) which included 1832 women and 6567 men with HFrEF enrolled in 47 countries on six continents.³⁵

Methods

PARADIGM-HF was a randomized, double-blind, placebo-controlled trial in patients with chronic HFrEF, evaluating the efficacy and safety of the angiotensin receptor-neprilysin inhibitor sacubitril/valsartan compared with enalapril, added to standard care. The design and primary results of PARADIGM-HF have been reported previously.^{35,36} The institutional review boards of all participating institutions approved the protocol, and all patients gave written informed consent.

Patients and study procedures

Key inclusion criteria were age ≥ 18 years, New York Heart Association (NYHA) functional class II–IV, left ventricular ejection fraction (LVEF) of $\leq 35\%$ (changed

from $\leq 40\%$ by a protocol amendment), elevated natriuretic peptide levels, and treatment with a stable dose of an ACE-i or angiotensin receptor blocker (ARB) equivalent to enalapril 10 mg/day for at least 4 weeks before the screening visit. Key exclusion criteria were symptomatic hypotension or systolic blood pressure < 95 mmHg at randomization (100 mmHg at screening), estimated glomerular filtration rate (eGFR) < 30 mL/min/1.73 m² at randomization (or screening), potassium > 5.4 mmol/L at randomization (> 5.2 mmol/L at screening), a history of angioedema, and intolerance to ACE-i or ARB.³⁶

On trial entry, ongoing therapy with ACE-i or ARB was stopped, and patients received enalapril 10 mg twice daily for 2 weeks followed by sacubitril/valsartan, up-titrated from 100 mg twice daily to 200 mg twice daily, for additional 4–6 weeks. Patients tolerating both drugs at the target doses were then randomly assigned to double-blind therapy with sacubitril/valsartan or enalapril in a 1:1 ratio.³⁶

Anthropometric measures

Data on anthropometric measures were obtained at the randomization visit. The calculation of each of these measures is described in [Figure 1](#).

In the analyses using BMI, patients were divided according to the World Health Organization (WHO) BMI categories, i.e. underweight (< 18.5 kg/m²); normal weight (18.5–24.9 kg/m²); overweight (25.0–29.9 kg/m²); obesity class I (30.0–34.9 kg/m²); obesity class II (35.0–39.9 kg/m²) and obesity class III (≥ 40 kg/m²). The choice of waist-to-height ratio as the exemplar anthropometric index that does not include weight was prespecified as it is easiest to calculate. We analyzed this index by quintile as the lowest quintile identified patients with a waist-to-height ratio in the range 0.150–0.520, the upper end of which approximates the < 0.5 regarded as healthy.

Outcomes

The primary outcome in PARADIGM-HF was the composite of HF hospitalization or cardiovascular death. In the present analysis, we also examined each of the components of the primary outcome, non-cardiovascular death, and death from any cause.

Statistical analyses

Baseline characteristics were summarized as frequencies with percentages, means with standard deviation (SD), or medians with interquartile ranges. Differences in baseline characteristics were tested using the Cochran–Armitage trend test for binary variables, the Cochran–Mantel–Haenszel test for categorical variables, and the Jonckheere–Terpstra test and linear regression for non-normal and normally distributed continuous variables, respectively.



Time-to-event data, regardless of treatment allocation, were evaluated using Cox proportional-hazards models, adjusted for treatment-group assignment and geographic region, and hazard ratios (HR) with 95% confidence intervals (CIs) were reported. In addition, HRs adjusted for treatment-group assignment, age, sex, race, geographic region, systolic blood pressure, heart rate, eGFR, LVEF, BMI (not included when analyzing outcomes according to BMI), log of *n*-terminal pro-B-type natriuretic peptide (NT-proBNP), NYHA functional class, HF aetiology, duration of HF, prior HF hospitalization, and a history of diabetes and atrial fibrillation were reported. The relationship between anthropometric measurements as continuous variables and the risk of outcomes was also examined in restricted cubic spline analyses (with the median value as reference).

To compare the effects of sacubitril/valsartan vs. enalapril, time-to-event data were evaluated with Cox proportional-hazards models, with geographical region and treatment-group assignment as fixed-effect factors.

All analyses were conducted using SAS version 9.4 (SAS Institute, Cary, NC) and STATA version 17.0 (College Station, TX). A *P*-value of .05 was considered statistically significant.

Results

Of the 8399 patients randomized in PARADIGM-HF, data on BMI were available in 99.9% (as were data on body surface area), and data on



waist-to-height ratio were available in 98.6% (as were data on waist circumference, relative fat mass, body roundness index, body shape index, and weight-adjusted-weight index although our focus was on waist-to-height ratio). Waist-to-hip ratio was available in 97.7% of randomized patients. Median duration of follow-up was 27 months (25th–75th percentile, 19–36 months).

Patient characteristics

Body mass index

Median BMI was 27.5 kg/m² (25th–75th percentile, 24.5–31.0 kg/m²) and 27.6 kg/m² (25th–75th percentile, 24.0–32.0 kg/m²) in men and women, respectively (Figure 2A). In total, 153 patients had a BMI <18.5 kg/m²; 2268 patients between 18.5–24.9 kg/m²; 3249 patients between 25–29.9 kg/m²; 1810 patients between 30–34.9 kg/m²; and 909 patients ≥35 kg/m². Detailed baseline characteristics according to these BMI categories are presented in Table 1a. Compared with patients with normal weight, those with higher BMI were younger, more often female, and white. Patients with higher BMI had more comorbidity, longer duration HF, more prior hospitalization for HF, and worse NYHA functional class, and Kansas City Cardiomyopathy Questionnaire (KCCQ) scores. However, they had higher LVEF and were more likely to have a non-ischemic aetiology (although patients

with a BMI <18.5 kg/m² were more likely than the other BMI categories to have a non-ischemic aetiology). Patients with higher BMI had lower natriuretic peptide levels (irrespective of the presence of atrial fibrillation) but a higher urinary cGMP/BNP ratio (as a marker of tissue responsiveness to natriuretic peptides). Patients with higher BMI had higher aldosterone, hemoglobin, uric acid, and blood urea nitrogen, levels, as well as neutrophil/lymphocyte ratio, compared with patients with normal weight, but a lower eGFR (Table 1a).

Waist-to-height ratio

Median waist-to-height ratio was 0.58 (25th–75th percentile, 0.54–0.64) and 0.59 (25th–75th percentile, 0.53–0.66) in men and women, respectively (Figure 2B). Baseline characteristics of patients according to quintiles of waist-to-height ratio are presented in Table 1b. Overall, the differences were similar to those described above and in Table 1a.

Overlap between body mass index categories and waist-to-height ratio quintiles

The overlap between BMI categories and waist-to-height ratio quintiles is shown in Table 2. Of the 2397 patients not overweight or obese,

Table 1a Baseline characteristics according to body mass index

	Underweight BMI <18.5 n = 153	Normal weight BMI 18.5–24.9 n = 2268	Overweight BMI 25–29.9 n = 3249	Obesity class I BMI 30–34.9 n = 1810	Obesity class II/III BMI ≥35 n = 909	P-value
Age (years), mean (SD)	60.9 ± 13.9	64.5 ± 12.2	64.6 ± 11.1	63.4 ± 10.7	60.3 ± 10.4	<.001
Male sex, n (%)	89 (58.2)	1776 (78.3)	2635 (81.1)	1417 (78.3)	643 (70.7)	.04
Geographic region, n (%)						<.001
North America	3 (2.0)	98 (4.3)	208 (6.4)	158 (8.7)	135 (14.9)	
Latin America	24 (15.7)	420 (18.5)	601 (18.5)	295 (16.3)	91 (10.0)	
Western Europe and other	16 (10.5)	427 (18.8)	875 (26.9)	489 (27.0)	238 (26.2)	
Central Europe	12 (7.8)	541 (23.9)	1090 (33.5)	755 (41.7)	428 (47.1)	
Asia-Pacific	98 (64.1)	782 (34.5)	475 (14.6)	113 (6.2)	17 (1.9)	
Race, n (%)						<.001
White	30 (19.6)	1092 (48.1)	2238 (68.9)	1419 (78.4)	758 (83.4)	
Black	8 (5.2)	134 (5.9)	127 (3.9)	83 (4.6)	76 (8.4)	
Asian	99 (64.7)	785 (34.6)	488 (15.0)	115 (6.4)	20 (2.2)	
Other	16 (10.5)	257 (11.3)	396 (12.2)	193 (10.7)	55 (6.1)	
Physiological measures, median (IQR) or mean (SD)						
Systolic blood pressure (mmHg)	118.4 ± 15.5	118.7 ± 14.8	121.4 ± 15.0	123.4 ± 15.6	124.7 ± 15.8	<.001
Heart rate (bpm)	76.1 ± 11.3	71.8 ± 11.6	71.8 ± 11.9	72.8 ± 12.4	74.4 ± 12.6	<.001
Body mass index	17.5 (16.9–18.0)	23.0 (21.5–24.1)	27.2 (26.1–28.5)	31.9 (30.8–33.1)	37.7 (36.1–40.4)	<.001
Waist-to-height ratio	0.46 (0.42–0.49)	0.52 (0.48–0.55)	0.58 (0.55–0.61)	0.64 (0.61–0.67)	0.71 (0.67–0.76)	<.001
Waist circumference	75 (68–81)	87 (81–93)	99 (92–104)	109 (103–115)	121 (113–130)	<.001
Waist-to-hip ratio	0.91 (0.84–0.94)	0.93 (0.88–0.98)	0.96 (0.92–1.02)	0.99 (0.94–1.05)	1.00 (0.94–1.06)	<.001
Relative fat mass	23.8 (18.9–31.4)	26.7 (23.6–30.0)	30.3 (28.1–33.1)	33.4 (31.7–36.1)	37.2 (35.0–44.7)	<.001
Weight-adjusted-waist index	11.0 (10.1–12.0)	11.0 (10.3–11.5)	11.1 (10.6–11.7)	11.3 (10.8–11.9)	11.5 (10.9–12.1)	<.001
Body shape index	0.088 (0.080–0.095)	0.084 (0.080–0.088)	0.084 (0.080–0.087)	0.083 (0.079–0.087)	0.082 (0.078–0.086)	<.001
Body roundness index	2.6 (1.9–3.3)	3.8 (3.1–4.4)	5.0 (4.3–5.7)	6.4 (5.7–7.2)	8.2 (7.1–9.5)	<.001
Current smoker, n (%)	26 (17.0)	380 (16.8)	455 (14.0)	231 (12.8)	116 (12.8)	<.001
Ischemic cause of HF, n (%)	69 (45.1)	1371 (60.4)	2006 (61.7)	1079 (59.6)	504 (55.4)	.22
Duration of HF, n (%)						<.001
<=1 year	87 (56.9)	814 (35.9)	929 (28.6)	475 (26.2)	215 (23.7)	

Continued

Table 1a Continued

	Underweight BMI < 18.5 n = 153	Normal weight BMI 18.5–24.9 n = 2268	Overweight BMI 25–29.9 n = 3249	Obesity class I BMI 30–34.9 n = 1810	Obesity class II/III BMI ≥ 35 n = 909	P-value
1–5 years	53 (34.6)	878 (38.7)	1270 (39.1)	667 (36.9)	361 (39.7)	
> 5 years	13 (8.5)	576 (25.4)	1050 (32.3)	668 (36.9)	333 (36.6)	
LVEF, mean (SD)	26.5 ± 6.4	28.8 ± 6.3	29.5 ± 6.1	30.0 ± 6.1	30.5 ± 6.1	<.001
NYHA class at randomization, n (%)						
I	11 (7.2)	130 (5.7)	150 (4.6)	72 (4.0)	25 (2.8)	
II	122 (79.7)	1667 (73.6)	2325 (71.7)	1256 (69.5)	541 (59.6)	
III	20 (13.1)	455 (20.1)	745 (23.0)	462 (25.6)	335 (36.9)	
IV	0 (0.0)	14 (0.6)	24 (0.7)	16 (0.9)	6 (0.7)	
KCCQ-OSS, mean (SD)	74.7 ± 18.6	75.0 ± 18.5	74.6 ± 18.6	71.2 ± 20.3	65.5 ± 21.1	<.001
KCCQ-CSS, mean (SD)	79.6 ± 17.1	78.9 ± 18.1	77.5 ± 18.4	73.8 ± 20.2	67.8 ± 20.8	<.001
Medical history, n (%)						
Hospitalization for HF	85 (55.6)	1343 (59.2)	2053 (63.2)	1165 (64.4)	623 (68.5)	<.001
Hypertension	65 (42.5)	1360 (60.0)	2290 (70.5)	1427 (78.8)	792 (87.1)	<.001
Diabetes	23 (15.0)	579 (25.5)	1083 (33.3)	747 (41.3)	471 (51.8)	<.001
History of atrial fibrillation	17 (11.1)	640 (28.2)	1191 (36.7)	779 (43.0)	459 (50.5)	<.001
Atrial fibrillation on electrocardiogram	10 (6.6)	398 (17.9)	784 (24.6)	524 (29.2)	318 (35.5)	<.001
Previous myocardial infarction	30 (19.6)	934 (41.2)	1525 (46.9)	785 (43.4)	355 (39.1)	.32
Previous stroke	8 (5.2)	189 (8.3)	279 (8.6)	154 (8.5)	95 (10.5)	.06
Chronic obstructive pulmonary disease	22 (14.4)	275 (12.1)	389 (12.0)	240 (13.3)	154 (16.9)	.002
Cancer	6 (3.9)	91 (4.0)	175 (5.4)	91 (5.0)	49 (5.4)	.08
Treatment, n (%)						
Beta-blocker	134 (87.6)	2076 (91.5)	3006 (92.5)	1720 (95.0)	865 (95.2)	<.001
Mineralocorticoid-receptor antagonist	84 (54.9)	1244 (54.9)	1786 (55.0)	1030 (56.9)	522 (57.4)	.09
Diuretic	123 (80.4)	1720 (75.8)	2564 (78.9)	1518 (83.9)	804 (88.4)	<.001
Digitalis	62 (40.5)	817 (36.0)	906 (27.9)	476 (26.3)	274 (30.1)	<.001
Antiplatelet	87 (56.9)	1323 (58.3)	1902 (58.5)	961 (53.1)	459 (50.5)	<.001
Oral anticoagulant	11 (7.2)	536 (23.6)	1028 (31.6)	706 (39.0)	400 (44.0)	<.001
Statin	67 (43.8)	1142 (50.4)	1907 (58.7)	1079 (59.6)	525 (57.8)	<.001

Continued

Table 1a Continued

	Underweight BMI <18.5 n = 153	Normal weight BMI 18.5–24.9 n = 2268	Overweight BMI 25–29.9 n = 3249	Obesity class I BMI 30–34.9 n = 1810	Obesity class II/III BMI ≥35 n = 909	P-value
Implantable cardioverter-defibrillator	5 (3.3)	225 (9.9)	519 (16.0)	329 (18.2)	163 (17.9)	<.001
Cardiac resynchronization therapy	1 (0.7)	129 (5.7)	245 (7.5)	122 (6.7)	75 (8.3)	.002
Biomarkers, median (IQR) or mean (SD) ^a						
NT-proBNP (pg/mL)						
No atrial fibrillation on electrocardiogram	2510 (1225–6807)	1964 (991–4371)	1450 (831–2836)	1229 (728–2196)	1072 (672–1938)	<.001
Atrial fibrillation on electrocardiogram	3515 (1605–10835)	2893 (1632–5715)	2102 (1203–4053)	1748 (1113–3231)	1508 (876–2468)	<.001
BNP (pg/mL)						
No atrial fibrillation on electrocardiogram	428 (159–763)	309 (167–633)	254 (151–448)	223 (141–391)	208 (132–346)	<.001
Atrial fibrillation on electrocardiogram	578 (215–1167)	334 (195–664)	264 (173–471)	221 (146–396)	194 (143–290)	<.001
ucGMP/BNP ratio (nmol/pg)						
No atrial fibrillation on electrocardiogram	1.5 (1.0–3.5)	3.2 (1.6–6.7)	4.6 (2.6–7.9)	4.8 (2.6–7.6)	5.2 (2.7–10.2)	<.001
Atrial fibrillation on electrocardiogram	2.8 (0.9–3.4)	5.1 (2.9–8.4)	5.1 (2.6–10.0)	5.7 (3.6–9.0)	7.5 (4.1–11.1)	.001
Aldosterone (pmol/L)						
No mineralocorticoid-receptor antagonist	246 (179–566)	180 (126–269)	212 (138–289)	221 (136–327)	260 (165–355)	<.001
Mineralocorticoid-receptor antagonist	322 (98–430)	262 (155–394)	261 (158–466)	327 (190–535)	325 (196–597)	.001
Troponin T (ng/L)	19 (8–26)	15 (10–24)	15 (10–23)	15 (10–23)	14 (9–22)	.59
ST2 (ng/mL)	22 (18–37)	33 (26–43)	32 (25–41)	33 (27–42)	32 (25–40)	.90
MMP-9 (ng/mL)	119 (53–173)	58 (35–121)	63 (38–119)	64 (41–126)	78 (44–161)	.005
MMP-2 (ng/mL)	147 (133–173)	133 (114–157)	134 (115–154)	135 (118–159)	137 (119–158)	.13
TIMP1 (ng/mL)	158 (137–199)	127 (104–158)	122 (104–149)	123 (105–148)	129 (106–153)	.96
Galectin 3 (ng/mL)	24 (21–30)	17 (14–23)	17 (14–21)	17 (14–21)	18 (15–22)	.11
GDF-15 (ng/L)	2238 (1983–4798)	1805 (1176–2536)	1590 (1153–2298)	1695 (1219–2466)	1598 (1074–2404)	.20
Hemoglobin (g/L)	128 (119–139)	135 (125–146)	140 (130–151)	143 (132–153)	143 (132–154)	<.001
Hemoglobin A1c (%)	6.0 (5.7–6.4)	6.1 (5.7–6.6)	6.2 (5.8–6.8)	6.3 (5.9–7.0)	6.5 (6.0–7.5)	<.001
White blood cell count (10 ⁹ /L)	6.4 (5.6–8.1)	6.5 (5.4–7.8)	6.7 (5.7–8.0)	6.8 (5.8–8.1)	7.2 (6.0–8.6)	<.001
Neutrophil count (10 ⁹ /L)	3.9 (3.1–5.1)	4.0 (3.1–5.0)	4.2 (3.3–5.2)	4.3 (3.5–5.3)	4.5 (3.6–5.6)	<.001
Neutrophil/lymphocyte ratio	2.1 (1.6–3.1)	2.3 (1.7–3.1)	2.4 (1.7–3.2)	2.4 (1.8–3.2)	2.4 (1.8–3.4)	<.001
Monocyte count (10 ⁹ /L)	0.4 (0.3–0.5)	0.4 (0.3–0.6)	0.5 (0.4–0.6)	0.5 (0.4–0.6)	0.5 (0.4–0.6)	<.001

Continued

Table 1a Continued

	Underweight BMI <18.5 n = 153	Normal weight BMI 18.5–24.9 n = 2268	Overweight BMI 25–29.9 n = 3249	Obesity class I BMI 30–34.9 n = 1810	Obesity class II/III BMI ≥35 n = 909	P-value
Lymphocyte count (10 ⁹ /L)	1.8 (1.4–2.2)	1.7 (1.3–2.2)	1.8 (1.4–2.2)	1.8 (1.4–2.2)	1.8 (1.5–2.3)	<.001
eGFR (mL/min/1.73 m ²)	69 (56–86)	68 (55–82)	66 (54–79)	65 (53–78)	66 (53–79)	<.001
Creatinine (μmol/L)	87 (74–101)	93 (80–111)	96 (82–114)	96 (82–116)	96 (80–114)	<.001
Blood urea nitrogen (mmol/L)	5.4 (3.9–8.0)	6.4 (5.0–8.6)	6.8 (5.4–8.9)	6.8 (5.7–8.6)	6.8 (5.4–8.9)	<.001
Cystatin (mg/L)	1.4 (1.2–1.7)	1.2 (1.0–1.5)	1.2 (1.0–1.4)	1.1 (1.0–1.4)	1.2 (1.0–1.4)	.40
Urine albumin-creatinine ratio (mg/mmol)						
No diabetes	3.9 (0.9–8.1)	0.8 (0.2–2.3)	0.9 (0.3–2.6)	0.8 (0.2–2.8)	1.1 (0.3–3.2)	.22
Diabetes	4.3 (2.7–5.8)	1.5 (0.5–6.1)	1.7 (0.6–7.1)	2.1 (0.7–6.3)	1.9 (0.7–8.1)	.17
KIM-1 (pg/mL)	260 (165–318)	124 (84–185)	120 (81–179)	134 (92–201)	149 (99–220)	<.001
Uric acid (μmol/L)	357 (297–434)	381 (321–464)	399 (327–482)	404 (339–488)	428 (351–512)	<.001
ALT (IU/L)	16 (11–21)	17 (13–22)	18 (14–24)	19 (14–25)	19 (15–25)	<.001
AST (IU/L)	23 (19–27)	21 (18–26)	21 (17–25)	20 (17–25)	20 (17–24)	<.001
Alkaline phosphatase (IU/L)	79 (65–97)	74 (60–92)	71 (58–88)	70 (57–87)	71 (58–88)	<.001
Total bilirubin (μmol/L)	9 (7–14)	9 (7–14)	9 (7–14)	9 (7–12)	9 (7–12)	<.001
Albumin (g/L)	42 (39–44)	43 (41–45)	43 (41–45)	43 (41–45)	42 (40–44)	.44

BNP, B-type natriuretic peptide; CSS, clinical summary score; eGFR, estimated glomerular filtration rate; HF, heart failure; KCCQ, Kansas City Cardiomyopathy Questionnaire; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; NT-proBNP, n-terminal pro-B-type natriuretic peptide; OSS, overall summary score; SD, standard deviation; ucGMP, urinary cyclic guanosine monophosphate.
^aRandomization visit, except ST2, MMP2, MMP9, TIMP1, GDF-15, and KIM-1, which were collected at the screening visit. Number of patients with missing biomarkers are: NT-proBNP, n = 14; BNP, n = 59; ucGMP/BNP ratio, n = 6375; aldosterone, n = 6381; troponin T, n = 6345; ST2, n = 6803; MMP-9, n = 6999; MMP-2, n = 6998; TIMP1, n = 6983; galectin 3, n = 6354; GDF-15, n = 6850; Hemoglobin, n = 274; hemoglobin A1c, n = 180; white blood cell count, n = 294; neutrophil count, n = 421; neutrophil/lymphocyte ratio, n = 421; lymphocyte count, n = 421; monocyte count, n = 421; creatinine, n = 0; blood urea nitrogen, n = 161; Cystatin, n = 6343; urine albumin-creatinine ratio, n = 6416; KIM-1, n = 6846; uric acid, n = 186; ALT, n = 219; AST, n = 199; total bilirubin, n = 199; albumin, n = 179.

Table 1b Baseline characteristics according to quintile of waist-to-height ratio

	Quintile 1 n = 1635	Quintile 2 n = 1659	Quintile 3 n = 1658	Quintile 4 n = 1655	Quintile 5 n = 1674	P-value
Age (years), mean (SD)	62.3 ± 12.9	64.1 ± 11.3	64.3 ± 11.2	64.9 ± 10.7	63.4 ± 10.7	.001
Male sex, n (%)	1250 (76.5)	1346 (81.1)	1370 (82.6)	1314 (79.4)	1193 (71.3)	<.001
Geographic region, n (%)						<.001
North America	103 (6.3)	94 (5.7)	115 (6.9)	119 (7.2)	161 (9.6)	
Latin America	239 (14.6)	285 (17.2)	304 (18.3)	305 (18.4)	275 (16.4)	
Western Europe and other	243 (14.9)	386 (23.3)	415 (25.0)	458 (27.7)	479 (28.6)	
Central Europe	446 (27.3)	507 (30.6)	572 (34.5)	616 (37.2)	679 (40.6)	
Asia-Pacific	604 (36.9)	387 (23.3)	252 (15.2)	157 (9.5)	80 (4.8)	
Race, n (%)						<.001
White	769 (47.0)	1006 (60.6)	1108 (66.8)	1232 (74.4)	1332 (79.6)	
Black	118 (7.2)	87 (5.2)	64 (3.9)	70 (4.2)	82 (4.9)	
Asian	606 (37.1)	389 (23.4)	261 (15.7)	161 (9.7)	85 (5.1)	
Other	142 (8.7)	177 (10.7)	225 (13.6)	192 (11.6)	175 (10.5)	
Physiological measures, median (IQR) or mean (SD)						
Systolic blood pressure (mmHg)	118.7 ± 14.5	119.5 ± 14.8	121.8 ± 15.2	122.5 ± 15.2	124.3 ± 16.0	<.001
Heart rate (bpm)	72.3 ± 11.6	71.8 ± 12.1	71.9 ± 11.6	72.1 ± 12.3	73.8 ± 12.3	<.001
Body mass index	22.7 (20.6–24.7)	25.2 (23.6–26.9)	27.4 (25.6–29.4)	30.0 (27.9–32.1)	34.4 (31.2–37.7)	<.001
Waist-to-height ratio	0.48 (0.46–0.51)	0.55 (0.53–0.56)	0.59 (0.58–0.59)	0.63 (0.62–0.64)	0.70 (0.67–0.74)	<.001
Waist circumference	81 (75–86)	92 (89–96)	99 (96–103)	107 (102–110)	118 (112–125)	<.001
Waist-to-hip ratio	0.90 (0.85–0.95)	0.94 (0.90–0.98)	0.97 (0.93–1.01)	0.99 (0.95–1.04)	1.02 (0.97–1.08)	<.001
Relative fat mass	23.8 (21.1–25.4)	27.7 (26.7–28.5)	30.0 (29.4–30.7)	32.5 (31.7–33.3)	36.4 (34.6–46.0)	<.001
Weight-adjusted-waist index	10.1 (9.6–10.6)	10.8 (10.5–11.2)	11.2 (10.8–11.6)	11.5 (11.1–11.9)	12.0 (11.6–12.5)	<.001
Body shape index	0.078 (0.073–0.083)	0.083 (0.079–0.086)	0.084 (0.081–0.087)	0.085 (0.081–0.088)	0.086 (0.083–0.090)	<.001
Body roundness index	3.1 (2.6–3.5)	4.3 (4.0–4.5)	5.1 (4.9–5.3)	6.1 (5.8–6.4)	7.8 (7.2–8.9)	<.001
Current smoker, n (%)	292 (17.9)	254 (15.3)	240 (14.5)	196 (11.8)	208 (12.4)	<.001
Ischemic cause of HF, n (%)	906 (55.4)	1000 (60.3)	1036 (62.5)	1039 (62.8)	984 (58.8)	.02
Duration of HF, n (%)						<.001
≤1 year	624 (38.2)	528 (31.8)	470 (28.3)	430 (26.0)	436 (26.0)	
1–5 years	613 (37.5)	656 (39.5)	647 (39.0)	656 (39.6)	623 (37.2)	

Continued

Table 1b Continued

	Quintile 1 n = 1635	Quintile 2 n = 1659	Quintile 3 n = 1658	Quintile 4 n = 1655	Quintile 5 n = 1674	P-value
> 5 years	398 (24.3)	475 (28.6)	541 (32.6)	569 (34.4)	615 (36.7)	
LVEF, mean (SD)	28.4 ± 6.2	29.2 ± 6.4	29.5 ± 6.2	30.0 ± 6.1	30.4 ± 6.1	<.001
NYHA class at randomization, n (%)						
I	85 (5.2)	104 (6.3)	70 (4.2)	63 (3.8)	55 (3.3)	
II	1172 (71.8)	1188 (71.7)	1185 (71.5)	1210 (73.3)	1080 (64.6)	
III	366 (22.4)	356 (21.5)	396 (23.9)	364 (22.0)	517 (30.9)	
IV	10 (0.6)	10 (0.6)	6 (0.4)	14 (0.8)	20 (1.2)	
KCCQ-OSS, mean (SD)	74.3 ± 18.6	75.2 ± 18.0	74.6 ± 18.9	73.4 ± 19.1	67.4 ± 21.1	<.001
KCCQ-CSS, mean (SD)	78.5 ± 18.3	78.5 ± 17.9	77.6 ± 18.4	75.9 ± 19.1	69.7 ± 21.0	<.001
Medical history, n (%)						
Hospitalization for HF	991 (60.6)	995 (60.0)	1054 (63.6)	1045 (63.1)	1121 (67.0)	<.001
Hypertension	935 (57.2)	1087 (65.5)	1164 (70.2)	1297 (78.4)	1382 (82.6)	<.001
Diabetes	362 (22.1)	498 (30.0)	551 (33.2)	640 (38.7)	815 (48.7)	<.001
History of atrial fibrillation	472 (28.9)	561 (33.8)	591 (35.6)	668 (40.4)	761 (45.5)	<.001
Atrial fibrillation on electrocardiogram	304 (19.0)	385 (23.5)	370 (22.8)	440 (26.8)	513 (31.0)	<.001
Previous myocardial infarction	625 (38.2)	715 (43.1)	763 (46.0)	785 (47.4)	698 (41.7)	.004
Previous stroke	143 (8.7)	141 (8.5)	148 (8.9)	123 (7.4)	166 (9.9)	.55
Chronic obstructive pulmonary disease	193 (11.8)	180 (10.8)	209 (12.6)	221 (13.4)	264 (15.8)	<.001
Cancer	64 (3.9)	82 (4.9)	82 (4.9)	87 (5.3)	91 (5.4)	.046
Treatment, n (%)						
Beta-blocker	1497 (91.6)	1526 (92.0)	1559 (94.0)	1539 (93.0)	1586 (94.7)	<.001
Mineralocorticoid-receptor antagonist	938 (57.4)	914 (55.1)	911 (54.9)	924 (55.8)	929 (55.5)	.44
Diuretic	1247 (76.3)	1291 (77.8)	1304 (78.6)	1349 (81.5)	1458 (87.1)	<.001
Digitalis	635 (38.8)	561 (33.8)	502 (30.3)	409 (24.7)	407 (24.3)	<.001
Antiplatelet	906 (55.4)	933 (56.2)	995 (60.0)	933 (56.4)	908 (54.2)	.55
Oral anticoagulant	380 (23.2)	511 (30.8)	509 (30.7)	588 (35.5)	651 (38.9)	<.001
Statin	740 (45.3)	903 (54.4)	1019 (61.5)	1031 (62.3)	966 (57.7)	<.001
Implantable cardioverter-defibrillator	160 (9.8)	237 (14.3)	241 (14.5)	285 (17.2)	286 (17.1)	<.001
Cardiac resynchronization therapy	99 (6.1)	107 (6.4)	109 (6.6)	107 (6.5)	136 (8.1)	.03

Continued

Table 1b Continued

	Quintile 1 n = 1635	Quintile 2 n = 1659	Quintile 3 n = 1658	Quintile 4 n = 1655	Quintile 5 n = 1674	P-value
Biomarkers, median (IQR) or mean (SD) ^a						
NT-proBNP (pg/mL)						
No atrial fibrillation on electrocardiogram	1934 (972–4360)	1644 (898–3546)	1481 (838–3003)	1306 (744–2425)	1213 (727–2212)	<.001
Atrial fibrillation on electrocardiogram	2883 (1480–5583)	2109 (1331–4093)	2212 (1189–4145)	1803 (1107–3346)	1672 (1034–2859)	<.001
BNP (pg/mL)						
No atrial fibrillation on electrocardiogram	299 (160–637)	287 (167–548)	251 (155–465)	230 (143–392)	219 (136–390)	<.001
Atrial fibrillation on electrocardiogram	329 (186–665)	277 (172–488)	262 (170–446)	230 (156–412)	215 (143–354)	<.001
ucGMP/BNP ratio (nmol/pg)						
No atrial fibrillation on electrocardiogram	3.4 (1.6–7.1)	4.2 (2.5–8.0)	4.7 (2.5–7.9)	4.6 (2.4–8.2)	4.6 (2.5–7.7)	.02
Atrial fibrillation on electrocardiogram	3.8 (1.5–6.6)	6.7 (3.2–10.2)	5.6 (2.8–10.0)	5.9 (2.8–9.8)	6.3 (3.8–10.5)	.02
Aldosterone (pmol/L)						
No mineralocorticoid-receptor antagonist	208 (125–327)	199 (135–283)	190 (131–275)	228 (141–322)	231 (161–347)	.003
Mineralocorticoid-receptor antagonist	305 (171–522)	279 (156–458)	250 (178–483)	294 (164–469)	330 (208–558)	.04
Troponin T (ng/L)	16 (9–25)	13 (9–22)	15 (10–23)	14 (10–22)	15 (10–25)	.32
ST2 (ng/mL)	32 (24–43)	32 (27–42)	32 (25–41)	32 (26–42)	32 (25–41)	.72
MMP-9 (ng/mL)	56 (35–107)	62 (39–123)	63 (39–117)	59 (38–132)	75 (42–139)	.008
MMP-2 (ng/mL)	137 (115–162)	132 (113–151)	134 (117–152)	133 (115–157)	136 (118–159)	.22
TIMP1 (ng/mL)	130 (104–161)	122 (102–149)	122 (103–146)	122 (104–150)	128 (107–151)	.27
Galectin 3 (ng/mL)	17 (14–22)	17 (14–22)	17 (14–21)	17 (14–21)	18 (15–22)	.007
GDF-15 (ng/L)	1780 (1215–2434)	1620 (1130–2373)	1517 (1104–2246)	1622 (1200–2304)	1763 (1222–2573)	.14
Hemoglobin (g/L)	135 (125–147)	139 (129–150)	141 (130–151)	141 (130–152)	141 (130–152)	<.001
Hemoglobin A1c (%)	6.0 (5.7–6.5)	6.1 (5.7–6.7)	6.2 (5.8–6.8)	6.2 (5.8–6.9)	6.4 (5.9–7.4)	<.001
White blood cell count (10 ⁹ /L)	6.5 (5.3–7.8)	6.6 (5.4–7.8)	6.8 (5.7–8.1)	6.8 (5.7–8.1)	7.1 (6.0–8.4)	<.001
Neutrophil count (10 ⁹ /L)	3.9 (3.1–5.0)	4.1 (3.2–5.0)	4.2 (3.4–5.2)	4.3 (3.4–5.3)	4.5 (3.6–5.5)	<.001
Neutrophil/lymphocyte ratio	2.2 (1.6–3.1)	2.4 (1.7–3.3)	2.4 (1.8–3.1)	2.4 (1.8–3.2)	2.5 (1.8–3.4)	<.001
Monocyte count (10 ⁹ /L)	0.4 (0.3–0.6)	0.4 (0.3–0.6)	0.5 (0.4–0.6)	0.5 (0.4–0.6)	0.5 (0.4–0.6)	<.001
Lymphocyte count (10 ⁹ /L)	1.7 (1.4–2.2)	1.7 (1.3–2.2)	1.8 (1.4–2.2)	1.8 (1.4–2.3)	1.8 (1.4–2.3)	<.001
eGFR (mL/min/1.73m ²)	69 (56–83)	67 (54–79)	65 (53–78)	65 (53–79)	65 (52–78)	<.001
Creatinine (μmol/L)	92 (79–109)	95 (81–113)	96 (83–114)	95 (81–115)	95 (80–116)	<.001

Continued

Table 1b Continued

	Quintile 1 n = 1635	Quintile 2 n = 1659	Quintile 3 n = 1658	Quintile 4 n = 1655	Quintile 5 n = 1674	P-value
Blood urea nitrogen (mmol/L)	6.4 (5.0–8.2)	6.8 (5.4–8.6)	6.8 (5.4–8.6)	6.8 (5.7–8.6)	7.1 (5.7–9.3)	<.001
Cystatin (mg/L)	1.2 (1.0–1.4)	1.1 (1.0–1.4)	1.1 (1.0–1.4)	1.2 (1.0–1.4)	1.2 (1.0–1.5)	.01
Urine albumin-creatinine ratio (mg/mmol)						
No diabetes	0.9 (0.0–2.8)	0.8 (0.2–2.1)	1.0 (0.3–2.8)	0.8 (0.3–2.9)	1.1 (0.3–2.9)	.19
Diabetes	1.6 (0.3–9.6)	1.5 (0.7–4.9)	1.8 (0.8–7.1)	1.7 (0.6–7.3)	2.1 (0.7–7.8)	.08
KIM-1 (pg/mL)	126 (82–192)	115 (78–176)	123 (80–181)	125 (88–197)	151 (99–213)	<.001
Uric acid (µmol/L)	375 (315–458)	393 (327–476)	404 (333–482)	404 (339–488)	410 (333–500)	<.001
ALT (IU/L)	17 (13–23)	18 (14–24)	18 (14–24)	18 (14–24)	18 (14–24)	<.001
AST (IU/L)	22 (18–27)	21 (18–26)	21 (17–25)	20 (17–25)	20 (17–24)	<.001
Alkaline phosphatase (IU/L)	73 (59–91)	71 (58–89)	72 (57–90)	70 (57–86)	73 (59–90)	<.001
Total bilirubin (µmol/L)	9 (7–14)	10 (7–14)	9 (7–12)	9 (7–12)	9 (7–12)	<.001
Albumin (g/L)	43 (41–45)	43 (41–45)	43 (41–45)	43 (41–45)	43 (41–44)	<.001

BNP, B-type natriuretic peptide; CSS, clinical summary score; eGFR, estimated glomerular filtration rate; HF, heart failure; KCCQ, Kansas City Cardiomyopathy Questionnaire; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; NT-proBNP, n-terminal pro-B-type natriuretic peptide; OSS, overall summary score; SD, standard deviation; ucGMP, urinary cyclic guanosine monophosphate.
^aRandomization visit, except ST2, MMP2, MMP9, TIMP1, GDF-15, and KIM-1, which were collected at the screening visit. Number of patients with missing biomarkers are: NT-proBNP, n = 14; BNP, n = 59; ucGMP/BNP ratio, n = 6375; Aldosterone, n = 6381; troponin T, n = 6345; ST2, n = 6803; MMP-9, n = 6999; MMP-2, n = 6998; TIMP1, n = 6983; galectin 3, n = 6354; GDF-15, n = 6850; Hemoglobin A1c, n = 180; White blood cell count, n = 294; Neutrophil count, n = 421; neutrophil/lymphocyte ratio, n = 421; monocyte count, n = 421; lymphocyte count, n = 421; eGFR, n = 0; creatinine, n = 0; blood urea nitrogen, n = 161; cystatin, n = 6343; urine albumin-creatinine ratio, n = 6416; KIM-1, n = 6846; uric acid, n = 186; ALT, n = 219; AST, n = 236; alkaline phosphatase, n = 199; total bilirubin, n = 199; albumin, n = 179.

according to BMI, only 1253 (52%) were in waist-to-height ratio quintile 1 (covering the range of waist-to-height ratio thought to be healthy). The remaining 1144 participants (48%) not overweight or obese, according to BMI, were in waist-to-height ratio quintiles 2–5 (mainly quintiles 2 and 3).

There was less reclassification of patients classified as obese by BMI; of the 2680 patients categorized as in obesity class I or class II/III, 2249 (84%) were in the fourth or fifth quintile of waist-to-height ratio.

Outcomes according to anthropometric measures

Body mass index

In the analyses of BMI by WHO category, the risk of the primary outcome did not differ significantly between patients with obesity and those with normal weight in the analysis adjusted for only region and treatment. However, the risk of death (whether due to cardiovascular or all causes), was significantly lower in overweight and obese patients; conversely, the risk of heart failure hospitalization was higher among individuals in obesity class II/III (Table 3a). After adjustment for prognostic variables, including NT-proBNP, the association between higher BMI and lower risk of death was eliminated and the association between higher BMI and higher risk of HF hospitalization was accentuated (Table 3a).

When BMI was examined as a continuous variable, a similar picture was evident although the small 'underweight' BMI category appeared to be associated with a higher risk of death from non-cardiovascular causes (Figure 3).

Waist-to-height ratio

In the analyses of waist-to-height ratio by quintile, the risk of the primary outcome did not differ significantly between patients in quintile 5, compared to those in quintile 1, in the analysis adjusted for only region and treatment (Table 3b). However, the risk of death (whether due to cardiovascular or all causes), tended to be lower in patients in quintile 5, although this was not as clear as for BMI. Conversely, the risk of HF hospitalization was significantly higher in individuals in quintile 5, compared with quintile 1. After adjustment for prognostic variables, the association between higher waist-to-height ratio and lower risk of death was eliminated and the association between higher waist-to-height ratio and higher risk of HF hospitalization persisted (Table 3).

When waist-to-height ratio was examined as a continuous variable, a similar picture was evident although a low waist-to-height ratio did not

appear to be associated with a higher risk of death from non-cardiovascular causes (Figure 4).

Other anthropometric measures

The association between the other anthropometric measures examined as continuous variables and outcomes are shown in Supplementary material online, Figures S1–S7. The association between waist-to-hip ratio examined as a continuous variable and outcomes in men and women, respectively, are shown in Supplementary material online, Figures S8 and S9. After adjustment for prognostic variables, greater adiposity, as assessed by body roundness index and relative fat mass, was associated with a higher risk of HF hospitalization. None of the anthropometric measures (i.e. waist circumference, waist-to-hip ratio, body shape index, weight-adjusted-waist index, and body surface area) was associated with cardiovascular death.

Effects of sacubitril/valsartan according to anthropometric measures

Body mass index

Compared with enalapril, sacubitril/valsartan reduced the risk of HF hospitalization or cardiovascular death across BMI categories (P for interaction = 0.50) (Table 4). The beneficial effect of sacubitril/valsartan was consistent across BMI categories for HF hospitalization (P for interaction = .41), cardiovascular death (P for interaction = .81), and death from any cause (P for interaction = .97) (Table 4).

Waist-to-height ratio

The beneficial effect of sacubitril/valsartan was also consistent for the primary and secondary endpoints according to quintiles of waist-to-height ratio (see Supplementary material online, Table S1).

Discussion

Among the patients with HFrEF in PARADIGM-HF, there was no longer evidence of a BMI-related 'obesity-survival paradox' after comprehensive adjustment for other prognostic variables. Moreover, the counterintuitive epidemiologic observation of a lower risk of death in patients with greater adiposity was less apparent with the newer anthropometric indices. All anthropometric indices examined showed that greater adiposity was associated with a higher risk of HF hospitalization as this was more evident with the newer indices (Structured Graphical Abstract).

Table 2 Correlation between body mass index categories and waist-to-height ratio quintiles

	Underweight BMI <18.5 n = 151	Normal weight BMI 18.5–24.9 n = 2246	Overweight BMI 25–29.9 n = 3203	Obesity class I BMI 30–34.9 n = 1779	Obesity II/III BMI ≥35 n = 901
WHtR quintile 1	129	1124	347	30	5
WHtR quintile 2	14	746	819	69	11
WHtR quintile 3	4	282	1055	295	21
WHtR quintile 4	2	71	745	715	122
WHtR quintile 5	2	23	237	670	742

Spearman correlation coefficient: 0.74.

Table 3a Outcomes according to body mass index

	Underweight BMI <18.5 n = 153	Normal weight BMI 18.5–24.9 n = 2268	Overweight BMI 25–29.9 n = 3249	Obesity class I BMI 30–34.9 n = 1810	Obesity class II/III BMI ≥35 n = 909
HF hospitalization or cardiovascular death					
n (%)	38 (24.8)	556 (24.5)	764 (23.5)	432 (23.9)	239 (26.3)
Event rate per 100 person-years (95% CI)	12.5 (9.1–17.1)	12.1 (11.1–13.1)	11.2 (10.4–12.0)	11.3 (10.3–12.4)	12.6 (11.1–14.3)
HR (95% CI) ^a	1.02 (0.73–1.42)	Reference	0.94 (0.84–1.06)	0.94 (0.83–1.08)	1.04 (0.88–1.21)
HR (95% CI) ^b	1.20 (0.86–1.67)	Reference	0.90 (0.81–1.01)	0.86 (0.75–0.99)	0.92 (0.78–1.09)
HR (95% CI) ^c	1.06 (0.76–1.47)	Reference	1.03 (0.92–1.16)	1.09 (0.95–1.25)	1.24 (1.05–1.48)
HF hospitalization					
n (%)	18 (11.8)	289 (12.7)	451 (13.9)	273 (15.1)	163 (17.9)
Event rate per 100 person-years (95% CI)	5.9 (3.7–9.4)	6.3 (5.6–7.0)	6.6 (6.0–7.3)	7.1 (6.3–8.0)	8.6 (7.4–10.0)
HR (95% CI) ^a	0.96 (0.60–1.55)	Reference	1.04 (0.90–1.22)	1.10 (0.93–1.31)	1.26 (1.03–1.54)
HR (95% CI) ^b	1.16 (0.72–1.88)	Reference	0.98 (0.84–1.14)	0.96 (0.81–1.15)	1.06 (0.86–1.32)
HR (95% CI) ^c	1.03 (0.63–1.67)	Reference	1.12 (0.96–1.31)	1.22 (1.01–1.46)	1.43 (1.15–1.78)
Cardiovascular death					
n (%)	30 (19.6)	387 (17.1)	474 (14.6)	242 (13.4)	117 (12.9)
Event rate per 100 person-years (95% CI)	9.3 (6.5–13.3)	7.9 (7.1–8.7)	6.5 (5.9–7.1)	5.8 (5.1–6.6)	5.6 (4.6–6.7)
HR (95% CI) ^a	1.14 (0.78–1.65)	Reference	0.85 (0.74–0.98)	0.77 (0.65–0.91)	0.75 (0.60–0.93)
HR (95% CI) ^b	1.27 (0.87–1.86)	Reference	0.83 (0.73–0.96)	0.75 (0.63–0.89)	0.72 (0.58–0.91)
HR (95% CI) ^c	1.11 (0.76–1.63)	Reference	0.96 (0.83–1.11)	0.96 (0.80–1.15)	1.00 (0.79–1.26)
Non-cardiovascular death					
n (%)	6 (3.9)	79 (3.5)	119 (3.7)	65 (3.6)	24 (2.6)
Event rate per 100 person-years (95% CI)	1.9 (0.8–4.1)	1.6 (1.3–2.0)	1.6 (1.4–1.9)	1.6 (1.2–2.0)	1.1 (0.8–1.7)
HR (95% CI) ^a	1.47 (0.64–3.39)	Reference	0.88 (0.66–1.17)	0.81 (0.58–1.13)	0.57 (0.36–0.91)
HR (95% CI) ^b	1.75 (0.75–4.08)	Reference	0.87 (0.65–1.16)	0.83 (0.59–1.18)	0.64 (0.39–1.06)
HR (95% CI) ^c	1.70 (0.73–3.98)	Reference	0.91 (0.68–1.23)	0.91 (0.64–1.30)	0.72 (0.44–1.19)
All-cause death					
n (%)	36 (23.5)	466 (20.5)	593 (18.3)	307 (17.0)	141 (15.5)
Event rate per 100 person-years (95% CI)	11.2 (8.1–15.5)	9.5 (8.7–10.4)	8.1 (7.5–8.8)	7.4 (6.6–8.3)	6.7 (5.7–7.9)
HR (95% CI) ^a	1.18 (0.84–1.67)	Reference	0.86 (0.75–0.97)	0.78 (0.67–0.90)	0.71 (0.58–0.86)
HR (95% CI) ^b	1.33 (0.94–1.88)	Reference	0.84 (0.74–0.95)	0.76 (0.65–0.89)	0.71 (0.57–0.87)
HR (95% CI) ^c	1.19 (0.84–1.69)	Reference	0.95 (0.84–1.08)	0.95 (0.81–1.11)	0.94 (0.76–1.16)

BMI, body mass index; CI, confidence interval; HF, heart failure; HR, hazard ratio.

^aAdjusted for treatment and region.

^bAdjusted for treatment, age, sex, region, systolic blood pressure, heart rate, estimated glomerular filtration rate, left ventricular ejection fraction, NYHA functional class, HF etiology, duration of HF, prior HF hospitalization, a history of diabetes, and atrial fibrillation.

^cAdjusted for the above-mentioned variables and log of NT-proBNP.

In the minimally adjusted analyses (adjusted for only region and treatment assignment), overweight and obesity, defined using conventional BMI categories, were associated with a lower risk of death from any cause, and cardiovascular causes, compared with normal weight, as has been reported previously.^{4–8,37–40} However, most prior studies did not adjust for other prognostic variables which vary greatly across BMI categories, particularly

the most powerful of these, i.e. natriuretic peptides.^{4–8,26–34,37–40} We were able to adjust for a broad range of prognostic variables, including NT-proBNP, and thereby minimize the impact of any potential residual confounding. After this adjustment, the 'survival paradox' related to high BMI was eliminated. Furthermore, none of the newer anthropometric measures showed the same association between greater adiposity and a lower risk of




Figure 3 Outcomes according to body mass index. This figure shows the risk of heart failure hospitalization or cardiovascular death, its components, and all-cause death, according to continuous body mass index. The solid line represents the hazard ratio and the shaded area the 95% CI. The reference is a body mass index of 25 kg/m². The blue spline is adjusted for treatment and region. The red spline is adjusted for treatment, age, sex, region, systolic blood pressure, heart rate, estimated glomerular filtration rate, left ventricular ejection fraction, log of *n*-terminal pro-B-type natriuretic peptide, New York Heart Association functional class, heart failure aetiology, duration of heart failure, prior heart failure hospitalization, a history of diabetes, and atrial fibrillation. CI, confidence interval; HF, heart failure; HR, hazard ratio.

cardiovascular and all-cause death as was seen with BMI when adjusted for conventional risk variables (but not NT-proBNP); additional adjustment for NT-proBNP eliminated any suggestion of an ‘obesity-survival paradox’ with BMI. Indeed, when each of these indices was analyzed as a continuous variable their relationship with death was entirely flat. It is not certain why the newer indices showed a weaker relationship between adiposity and fatal outcomes, compared with BMI. However, an inspection of patient characteristics and biomarkers suggested a somewhat steeper gradient in age, atrial fibrillation, and NYHA class, as well as natriuretic peptide, aldosterone, and hemoglobin levels across BMI categories (normal to obesity class II/III) compared with the other indices (lowest to highest quintile). This may also explain why adjustment for other prognostic variables changed the association between higher adiposity and risk of death more for BMI than the other anthropometric indices.

Importantly, three of the newer anthropometric indices (waist-to-height ratio, relative fat mass, and body roundness index) demonstrated a significantly higher risk of HF hospitalization (and the composite of HF hospitalization or death from cardiovascular causes) in patients with greater adiposity, an association that was less apparent with BMI. These newer indices incorporate waist circumference and height, with the former better reflecting intra-abdominal fat (‘central obesity’) and the latter accounting for sex- and race-based differences in stature and skeletal weight. Notably, neither newer index included measured overall weight, the interpretation of which in patients with HF may be confounded by fluid retention or unintentional weight loss

due to other illness. Interestingly, despite the steeper gradient in prognostic variables across BMI categories, compared to waist-to-height ratio quintiles, described above, the association between higher BMI and HF hospitalization was not as strong as that between higher waist-to-height ratio and HF hospitalization. This suggests that these new anthropometric indices identify pathophysiologic processes not reflected by conventional prognostic variables e.g. related to the distribution of body fat.

Different associations with outcomes were seen at the lower end of the range of adiposity and these also varied between the anthropometric indices. When considered as a continuous variable, a very low BMI was associated with a higher risk of death, which has been shown previously. However, we demonstrated that this was explained by a significant excess of non-cardiovascular death rather than cardiovascular death. The excess risk persisted after adjustment for other prognostic variables. This relationship was not as clearly evident with waist-to-height ratio (or the other newer anthropometric indices).

The relationship between adiposity and health-related quality of life (measured with the Kansas City Cardiomyopathy Questionnaire) was consistent between BMI and waist-to-height ratio. Both showed a steep decline in health-related quality of life with increasing adiposity, a relationship which was consistent with the relationship between waist-to-height ratio and HF hospitalization but opposite to that for BMI and mortality (in an unadjusted analysis).

Collectively these data show that greater adiposity in HFrEF is associated with a higher symptom burden, worse quality of life, and a

Table 3b Outcomes according to quintile of waist-to-height ratio

	Quintile 1 n = 1635	Quintile 2 n = 1659	Quintile 3 n = 1658	Quintile 4 n = 1655	Quintile 5 n = 1674
HF hospitalization or cardiovascular death					
n (%)	378 (23.1)	378 (22.8)	433 (26.1)	390 (23.6)	425 (25.4)
Event rate per 100 person-years (95% CI)	11.4 (10.3–12.6)	10.9 (9.8–12.0)	12.4 (11.3–13.6)	11.2 (10.1–12.4)	12.3 (11.1–13.5)
HR (95% CI) ^a	Reference	0.98 (0.85–1.13)	1.13 (0.98–1.30)	1.02 (0.88–1.18)	1.12 (0.97–1.29)
HR (95% CI) ^b	Reference	1.00 (0.86–1.16)	1.18 (1.01–1.39)	1.12 (0.93–1.33)	1.18 (0.96–1.44)
HR (95% CI) ^c	Reference	1.02 (0.88–1.19)	1.24 (1.06–1.46)	1.18 (0.99–1.41)	1.27 (1.03–1.55)
HF hospitalization					
n (%)	198 (12.1)	208 (12.5)	257 (15.5)	230 (13.9)	284 (17.0)
Event rate per 100 person-years (95% CI)	6.0 (5.2–6.9)	6.0 (5.2–6.9)	7.4 (6.5–8.3)	6.6 (5.8–7.5)	8.2 (7.3–9.2)
HR (95% CI) ^a	Reference	1.02 (0.84–1.24)	1.26 (1.04–1.52)	1.12 (0.92–1.37)	1.37 (1.13–1.65)
HR (95% CI) ^b	Reference	1.01 (0.83–1.24)	1.24 (1.00–1.54)	1.13 (0.89–1.43)	1.30 (1.00–1.69)
HR (95% CI) ^c	Reference	1.04 (0.85–1.28)	1.31 (1.05–1.62)	1.19 (0.93–1.51)	1.39 (1.06–1.81)
Cardiovascular death					
n (%)	263 (16.1)	248 (14.9)	263 (15.9)	240 (14.5)	224 (13.4)
Event rate per 100 person-years (95% CI)	7.5 (6.6–8.4)	6.7 (5.9–7.6)	7.0 (6.2–7.9)	6.4 (5.6–7.3)	5.9 (5.2–6.7)
HR (95% CI) ^a	Reference	0.93 (0.78–1.10)	0.98 (0.82–1.17)	0.91 (0.76–1.09)	0.85 (0.70–1.02)
HR (95% CI) ^b	Reference	0.98 (0.82–1.18)	1.09 (0.90–1.33)	1.11 (0.89–1.39)	1.06 (0.82–1.38)
HR (95% CI) ^c	Reference	1.01 (0.84–1.21)	1.16 (0.95–1.41)	1.19 (0.95–1.49)	1.15 (0.88–1.49)
Non-cardiovascular death					
n (%)	58 (3.5)	55 (3.3)	58 (3.5)	60 (3.6)	55 (3.3)
Event rate per 100 person-years (95% CI)	1.6 (1.3–2.1)	1.5 (1.1–1.9)	1.5 (1.2–2.0)	1.6 (1.2–2.1)	1.4 (1.1–1.9)
HR (95% CI) ^a	Reference	0.82 (0.56–1.19)	0.80 (0.55–1.16)	0.82 (0.56–1.18)	0.71 (0.49–1.04)
HR (95% CI) ^b	Reference	0.82 (0.56–1.21)	0.81 (0.53–1.23)	0.86 (0.55–1.36)	0.87 (0.52–1.48)
HR (95% CI) ^c	Reference	0.83 (0.57–1.22)	0.82 (0.54–1.25)	0.88 (0.56–1.39)	0.90 (0.53–1.52)
All-cause death					
n (%)	321 (19.6)	303 (18.3)	321 (19.4)	300 (18.1)	279 (16.7)
Event rate per 100 person-years (95% CI)	9.1 (8.2–10.2)	8.2 (7.3–9.2)	8.5 (7.6–9.5)	8.0 (7.2–9.0)	7.3 (6.5–8.2)
HR (95% CI) ^a	Reference	0.91 (0.77–1.06)	0.94 (0.81–1.11)	0.89 (0.76–1.05)	0.82 (0.69–0.97)
HR (95% CI) ^b	Reference	0.95 (0.81–1.12)	1.04 (0.87–1.24)	1.06 (0.87–1.30)	1.03 (0.81–1.30)
HR (95% CI) ^c	Reference	0.97 (0.83–1.15)	1.09 (0.91–1.31)	1.13 (0.92–1.38)	1.10 (0.87–1.39)

CI, confidence interval; HF, heart failure; HR, hazard ratio.

^aAdjusted for treatment and region.

^bAdjusted for treatment, age, sex, region, systolic blood pressure, heart rate, estimated glomerular filtration rate, left ventricular ejection fraction, body mass index, NYHA functional class, HF etiology, duration of HF, prior HF hospitalization, a history of diabetes, and atrial fibrillation.

^cAdjusted for the above-mentioned variables and log of NT-proBNP.

greater risk of HF hospitalization. Other studies have shown that obese patients also have a higher risk of developing diabetes, atrial fibrillation, sleep apnea, and other comorbid conditions, compared with non-obese patients.^{26,34,41–44} Also, BMI > 35 kg/m² is a contraindication to heart transplantation.⁴⁵ Therefore, there is a strong rationale for promoting weight loss in obese patients especially as the 'obesity-survival paradox' seems to be an artifact of unadjusted analyses of

BMI. Unfortunately, few randomized controlled trials using dietary and exercise intervention, bariatric surgery, or novel pharmacological therapies have been conducted in patients with HFrEF, although the latter are being investigated in individuals with HFpEF.^{46–52} The 2021 European Society of Cardiology and 2022 American College of Cardiology/American Heart Association guidelines do not provide any recommendation regarding weight management in HFrEF.^{45,53}




Figure 4 Outcomes according to waist-to-height ratio. This figure shows the risk of heart failure hospitalization or cardiovascular death, its components, and all-cause death, according to continuous waist-to-height ratio. The solid line represents the hazard ratio and the shaded area the 95% CI. The reference is the median waist-to-height ratio (0.58). The blue spline is adjusted for treatment and region. The red spline is adjusted for treatment, age, sex, region, systolic blood pressure, heart rate, estimated glomerular filtration rate, left ventricular ejection fraction, log of *n*-terminal pro-B-type natriuretic peptide, body mass index, New York Heart Association functional class, heart failure aetiology, duration of heart failure, prior heart failure hospitalization, a history of diabetes, and atrial fibrillation. CI, confidence interval; HF, heart failure; HR, hazard ratio.

Table 4 Effects of sacubitril/valsartan compared with enalapril according to body mass index

	Underweight BMI <18.5 <i>n</i> = 153 HR (95% CI)	Normal weight BMI 18.5–24.9 <i>n</i> = 2268 HR (95% CI)	Overweight BMI 25–29.9 <i>n</i> = 3249 HR (95% CI)	Obesity class I BMI 30–34.9 <i>n</i> = 1810 HR (95% CI)	Obesity class II/III BMI ≥35 <i>n</i> = 909 HR (95% CI)	P-value for interaction
HF hospitalization or cardiovascular death	0.80 (0.42–1.53)	0.91 (0.77–1.07)	0.79 (0.68–0.91)	0.75 (0.62–0.90)	0.70 (0.54–0.91)	.50
HF hospitalization	0.93 (0.37–2.38)	0.97 (0.77–1.22)	0.75 (0.62–0.90)	0.74 (0.59–0.95)	0.71 (0.52–0.97)	.41
Cardiovascular death	0.79 (0.38–1.66)	0.88 (0.72–1.07)	0.80 (0.67–0.96)	0.72 (0.55–0.92)	0.73 (0.50–1.06)	.81
All-cause death	0.81 (0.41–1.60)	0.90 (0.75–1.08)	0.82 (0.70–0.96)	0.83 (0.66–1.04)	0.81 (0.58–1.13)	.97

BMI, body mass index; CI, confidence interval; HF, heart failure; HR hazard ratio. Models adjusted for region.

Efforts to find effective and safe approaches to reducing weight in patients with HF_rEF are therefore warranted.

Effect of sacubitril/valsartan according to anthropometric measures

The effect of HF therapies according to anthropometric measures is also of interest because of the hypothesis that the ‘obesity-survival

paradox’ could, to some extent, reflect a greater effect of treatment in obese patients. Support for this was provided by a *post hoc* analysis of the Eplerenone in Mild Patients Hospitalization and Survival Study in Heart Failure (EMPHASIS-HF) trial, in which the benefit of the mineralocorticoid receptor antagonist, eplerenone, appeared to be greater in patients with larger waist circumference.⁵⁴ However, such a relationship has not been established for other effective therapies in HF_rEF, and a recent analysis of the dapagliflozin and prevention of adverse outcomes in

heart failure (DAPA-HF) trial demonstrated that the benefit of dapagliflozin was consistent across the spectrum of BMI.³⁴ In the present analysis, the beneficial effect of sacubitril/valsartan, compared with enalapril, was consistent for all outcomes across the spectrum of BMI and waist-to-height ratio. Specifically, there was no evidence of a diminished benefit in underweight patients or a larger benefit in obese patients.

Study limitations

This study has some limitations. Due to the observational nature of this study, the possibility of unmeasured confounding, despite adjustment for known prognostic variables, remains. Abdominal anthropometric measurements, such as waist circumference, are associated with higher measurement error than BMI, especially when these measurements are performed by different individuals.⁵⁵ In addition, the analyses on the association between anthropometric measures at randomization and clinical adverse outcomes did not account for any change in e.g. weight or waist circumference during follow-up. Previous studies have suggested that the 'obesity-survival paradox' in HF may be affected by the level of cardiorespiratory fitness.^{56–59} Although it would have been interesting to examine the association between anthropometric measures and outcomes according to cardiorespiratory fitness levels, these data were not available. Finally, only 153 patients had a BMI <18.5 kg/m² (the 'underweight' BMI category) and 171 patients a waist-to-height ratio <0.40, and our findings clearly cannot be extrapolated to patients with a low BMI or waist-to-height ratio.

Conclusion

In a large cohort of patients with HFrEF, the 'obesity-survival paradox' related to BMI was eliminated by comprehensive adjustment for prognostic variables. Importantly, alternative anthropometric measurements showed less evidence of this paradox, and two indices that incorporate waist circumference and height, but not weight, showed a clearer association between greater adiposity and a higher risk of HF hospitalization. Greater adiposity was associated with worse symptoms and health-related quality of life, irrespective of the anthropometric index used. These findings collectively suggest a need to test the potential benefits of intentional weight loss in patients living with obesity and HFrEF.

Author contributions

Michael Zile, MD (Writing – review & editing: Equal), Karl Swedberg, MD (Writing – review & editing: Equal), Jean L Rouleau, MD (Writing – review & editing: Equal), John J.J.V. McMurray, MD, Milton Packer, MD (Writing – review & editing: Equal), Scott D. Solomon, MD (Writing – review & editing: Equal), Pardeep S. Jhund (Writing – review & editing: Equal), Mark C. Petrie, MBChB (Writing – review & editing: Equal), Jawad H. Butt, MD (Writing – original draft: Lead), Lars Kober, MD (Writing – review & editing: Equal), Akshay S Desai, MD (Writing – review & editing: Equal), and Naveed Sattar, MD (Writing – review & editing: Equal).

Acknowledgements

None.

Supplementary data

Supplementary data is available at *European Heart Journal* online.

Data availability

Trial data will be made available by the sponsor, Novartis, in accordance with their data sharing policy.

Conflict of interest

Dr. Butt reports advisory board honoraria from Bayer.

Dr. Petrie has received research grants or consultancy fees from SQ Innovations, AstraZeneca, Roche, Boehringer Ingelheim, Pharmacosmos, Eli Lilly, Napp Pharmaceuticals, Novartis, and Novo Nordisk and has served on clinical events committees for AbbVie, Alnylam, AstraZeneca, Bayer, Boehringer Ingelheim, GlaxoSmithKline, Resverlogix, and Novo Nordisk.

Dr. Jhund has received consulting fees, advisory board fees, and lecture fees from Novartis; advisory board fees from Cytokinetics; and grant support from Boehringer Ingelheim.

Dr. Sattar has consulted for or received lecture fees from Afimmune, Amgen, AstraZeneca, Boehringer Ingelheim, Eli Lilly, Hanmi Pharmaceuticals, Merck Sharp & Dohme, Novartis, Novo Nordisk, Pfizer, and Sanofi. He has received grant support from AstraZeneca, Boehringer Ingelheim, Novartis, and Roche Diagnostics through his institution, the University of Glasgow.

Dr. Desai reported receiving personal fees from Abbott, Biofourmis, Boston Scientific, Boehringer Ingelheim, Merck, Regeneron, and Relpsa and grants and personal fees from AstraZeneca, Alnylam, and Novartis outside the submitted work.

Dr. Kober reports personal fees from speaker honoraria from Novartis, AstraZeneca, Novo Nordisk, and Boehringer Ingelheim.

Dr. Rouleau has received grants and consulting fees from Novartis and consulting fees from Abbott, AstraZeneca, MyoKardia, and Sanofi.

Dr. Swedberg reports honoraria from AstraZeneca, Boehringer 12 Ingelheim, and Novartis.

Dr. Zile has received research funding from Novartis and has been a consultant for Novartis, Abbott, Boston Scientific, CVRx, EBR, Endotronics, Ironwood, Merck, Medtronic, and Myokardia V Wave.

Dr. Solomon has received research grants from Actelion, Alnylam, Amgen, AstraZeneca, Bellerophon, Bayer, BMS, Celladon, Cytokinetics, Eidos, Gilead, GSK, Ionis, Lilly, Lone Star Heart, Mesoblast, MyoKardia, NIH/NHLBI, Neurotronik, Novartis, NovoNordisk, Respicardia, Sanofi Pasteur, Theracos, and has consulted for Abbott, Action Akros, Alnylam, Amgen, Arena, AstraZeneca, Bayer, Boeringer-Ingelheim, BMS, Cardior, Cardurion, Corvia, Cytokinetics, Daiichi-Sankyo, Gilead, GSK, Ironwood, Lilly, Merck, Myokardia, Novartis, Roche, Takeda, Theracos, Quantum Genetics, Cardurion, AoBiome, Janssen, Cardiac Dimensions, Tenaya, Sanofi-Pasteur, Dinaqor, Tremeau, CellProThera, Moderna, and American Regent.

Dr. Packer has received consulting fees from AbbVie, Akcea, Actavis, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, Cardiorentis, Daiichi Sankyo, Gilead, Johnson & Johnson, Novo Nordisk, Pfizer, Relpsa, Sanofi, Synthetic Biologics, and Theravance.

Dr. McMurray reports payments to his employer, Glasgow University, for work on clinical trials, consulting, lecturing and other activities: Alnylam, Amgen, AstraZeneca, Bayer, Boehringer Ingelheim, BMS, Cardurion, Cytokinetics, Dal-Cor, GSK, Ionis, KBP Biosciences, Novartis, Pfizer, and Theracos. Personal lecture fees: Abbott, Hikma, Sun Pharmaceuticals, Servier, Theracos; and personal payments from Abbott, Hikma, Ionis, Sun Pharmaceuticals, Servier. The other authors report no conflicts.

Funding

The PARADIGM-HF trial was funded by Novartis. Profs. McMurray and Jhund are supported by British Heart Foundation Centre of Research Excellence Grant RE/18/6/34217.

References

- Aune D, Sen A, Norat T, Janszky I, Romundstad P, Tonstad S, et al. Body mass index, abdominal fatness, and heart failure incidence and mortality: a systematic review and dose-response meta-analysis of prospective studies. *Circulation* 2016;**133**:639–649. <https://doi.org/10.1161/CIRCULATIONAHA.115.016801>
- Kenchaiah S, Evans JC, Levy D, Wilson PVF, Benjamin EJ, Larson MG, et al. Obesity and the risk of heart failure. *N Engl J Med* 2002;**347**:305–313. <https://doi.org/10.1056/NEJMoa020245>
- Pandey A, LaMonte M, Klein L, Ayers C, Psaty BM, Eaton CB, et al. Relationship between physical activity, body mass index, and risk of heart failure. *J Am Coll Cardiol* 2017;**69**:1129–1142. <https://doi.org/10.1016/j.jacc.2016.11.081>
- Zhang J, Begley A, Jackson R, Harrison M, Pellicori P, Clark AL, et al. Body mass index and all-cause mortality in heart failure patients with normal and reduced ventricular ejection fraction: a dose–response meta-analysis. *Clin Res Cardiol* 2019;**108**:119–132. <https://doi.org/10.1007/s00392-018-1302-7>
- Padwal R, McAlister FA, McMurray JJV, Cowie MR, Rich M, Pocock S, et al. The obesity paradox in heart failure patients with preserved versus reduced ejection fraction: a meta-analysis of individual patient data. *Int J Obes* 2014;**38**:1110–1114. <https://doi.org/10.1038/ijo.2013.203>
- Oga EA, Eseyin OR. The obesity paradox and heart failure: a systematic review of a decade of evidence. *J Obes* 2016;**2016**:1–9. <https://doi.org/10.1155/2016/9040248>
- Mahajan R, Stokes M, Elliott A, Munawar DA, Khokhar KB, Thiyagarajah A, et al. Complex interaction of obesity, intentional weight loss and heart failure: a systematic review and meta-analysis. *Heart* 2020;**106**:58–68. <https://doi.org/10.1136/heartjnl-2019-314770>
- Horwich TB, Fonarow GC, Clark AL. Obesity and the obesity paradox in heart failure. *Prog Cardiovasc Dis* 2018;**61**:151–156. <https://doi.org/10.1016/j.pcad.2018.05.005>
- Heymsfield SB, Peterson CM, Thomas DM, Heo M, Schuna JM. Why are there race/ethnic differences in adult body mass index-adiposity relationships? A quantitative critical review. *Obes Rev* 2016;**17**:262–275. <https://doi.org/10.1111/obr.12358>
- Okorodudu DO, Jumean MF, Montori VM, Romero-Corral A, Somers VK, Erwin PJ, et al. Diagnostic performance of body mass index to identify obesity as defined by body adiposity: a systematic review and meta-analysis. *Int J Obes* 2010;**34**:791–799. <https://doi.org/10.1038/ijo.2010.5>
- Piché ME, Poirier P, Lemieux I, Després JP. Overview of epidemiology and contribution of obesity and body fat distribution to cardiovascular disease: an update. *Prog Cardiovasc Dis* 2018;**61**:103–113. <https://doi.org/10.1016/j.pcad.2018.06.004>
- Rothman KJ. BMI-related errors in the measurement of obesity. *Int J Obes* 2008;**32**:S56–S59. <https://doi.org/10.1038/ijo.2008.87>
- Krakauer NY, Krakauer JC. A new body shape index predicts mortality hazard independently of body mass index. *PLoS One* 2012;**7**:e39504. <https://doi.org/10.1371/journal.pone.0039504>
- Woolcott OO, Bergman RN. Relative fat mass (RFM) as a new estimator of whole-body fat percentage — A cross-sectional study in American adult individuals. *Sci Rep* 2018;**8**:10980. <https://doi.org/10.1038/s41598-018-29362-1>
- Park Y, Kim NH, Kwon TY, Kim SG. A novel adiposity index as an integrated predictor of cardiometabolic disease morbidity and mortality. *Sci Rep* 2018;**8**:16753. <https://doi.org/10.1038/s41598-018-35073-4>
- Thomas DM, Bredlau C, Bosty-Westphal A, Mueller M, Shen W, Gallagher D, et al. Relationships between body roundness with body fat and visceral adipose tissue emerging from a new geometrical model. *Obesity* 2013;**21**:2264–2271. <https://doi.org/10.1002/oby.20408>
- Zhang C, Rexrode KM, Van Dam RM, Li TY, Hu FB. Abdominal obesity and the risk of all-cause, cardiovascular, and cancer mortality: sixteen years of follow-up in US women. *Circulation* 2008;**117**:1658–1667. <https://doi.org/10.1161/CIRCULATIONAHA.107.739714>
- Loehr LR, Rosamond WD, Poole C, McNeill AM, Chang PP, Folsom AR, et al. Association of multiple anthropometrics of overweight and obesity with incident heart failure: the atherosclerosis risk in communities study. *Circ Heart Fail* 2009;**2**:18–24. <https://doi.org/10.1161/CIRCHEARTFAILURE.108.813782>
- Gelber RP, Gaziano JM, Orav EJ, Manson JAE, Buring JE, Kurth T. Measures of obesity and cardiovascular risk among men and women. *J Am Coll Cardiol* 2008;**52**:605–615. <https://doi.org/10.1016/j.jacc.2008.03.066>
- Suthahar N, Meems LMG, Withaar C, Gorter TM, Kieneker LM, Gansevoort RT, et al. Relative fat mass, a new index of adiposity, is strongly associated with incident heart failure: data from PREVEND. *Sci Rep* 2022;**12**:147. <https://doi.org/10.1038/s41598-021-02409-6>
- Ashwell M, Gibson S. Waist-to-height ratio as an indicator of early health risk: simpler and more predictive than using a matrix based on BMI and waist circumference. *BMJ Open* 2016;**6**:e010159. <https://doi.org/10.1136/bmjopen-2015-010159>
- National Institute for Health and Care Excellence. Obesity: identification, assessment and management. <https://www.nice.org.uk/guidance/cg189/resources/obesity-identification-assessment-and-management-pdf-35109821097925> (11 December 2022)
- Horwich TB, Hamilton MA, Fonarow GC. B-type natriuretic peptide levels in obese patients with advanced heart failure. *J Am Coll Cardiol* 2006;**47**:85–90. <https://doi.org/10.1016/j.jacc.2005.08.050>
- Mehra MR, Uber PA, Park MH, Scott RL, Ventura HO, Harris BC, et al. Obesity and suppressed B-type natriuretic peptide levels in heart failure. *J Am Coll Cardiol* 2004;**43**:1590–1595. <https://doi.org/10.1016/j.jacc.2003.10.066>
- Savarese G, Orsini N, Hage C, Vedin O, Cosentino F, Rosano GMC, et al. Utilizing NT-proBNP for eligibility and enrichment in trials in HFpEF, HFmrEF, and HFrfEF. *JACC Heart Fail* 2018;**6**:246–256. <https://doi.org/10.1016/j.jchf.2017.12.014>
- Streng KW, Voors AA, Hillege HL, Anker SD, Cleland JG, Dickstein K, et al. Waist-to-hip ratio and mortality in heart failure. *Eur J Heart Fail* 2018;**20**:1269–1277. <https://doi.org/10.1002/ehf.1244>
- Piepoli MF, Corrà U, Veglia F, Bonomi A, Salvioni E, Cattadori G, et al. Exercise tolerance can explain the obesity paradox in patients with systolic heart failure: data from the MECKI score research group. *Eur J Heart Fail* 2016;**18**:545–553. <https://doi.org/10.1002/ehf.534>
- Chandramouli C, Tay WT, Bamadhaj NS, Tromp J, Teng THK, Yap JLL, et al. Association of obesity with heart failure outcomes in 11 Asian regions: a cohort study. *PLoS Med* 2019;**16**:e1002916. <https://doi.org/10.1371/journal.pmed.1002916>
- Zhang Y, Zhang Y, Shi Y, Dong W, Mu Y, Wang J, et al. Influence of Waist-to-hip ratio on the prognosis of heart failure patients with revascularized coronary heart disease. *Front Cardiovasc Med* 2021;**8**:732200. <https://doi.org/10.3389/fcvm.2021.732200>
- Davos CH, Doehner W, Rauchhaus M, Ciccoira M, Francis DP, Coats AJS, et al. Body mass and survival in patients with chronic heart failure without cachexia: the importance of obesity. *J Card Fail* 2003;**9**:29–35. <https://doi.org/10.1054/jcaf.2003.4>
- Gustafsson F, Kragelund CB, Torp-Pedersen C, Seibæk M, Burchardt H, Akkan D, et al. Effect of obesity and being overweight on long-term mortality in congestive heart failure: influence of left ventricular systolic function. *Eur Heart J* 2005;**26**:58–64. <https://doi.org/10.1093/eurheartj/ehi022>
- Kistorp C, Faber J, Galatius S, Gustafsson F, Frystyk J, Flyvbjerg A, et al. Plasma adiponectin, body mass index, and mortality in patients with chronic heart failure. *Circulation* 2005;**112**:1756–1762. <https://doi.org/10.1161/CIRCULATIONAHA.104.530972>
- Krishnamoorthy A, Greiner MA, Bertoni AG, Eapen ZJ, O'Brien EC, Curtis LH, et al. The obesity and heart failure epidemics among African Americans: insights from the Jackson heart study. *J Card Fail* 2016;**22**:589–597. <https://doi.org/10.1016/j.cardfail.2016.03.004>
- Adamson C, Jhund PS, Docherty KF, Böhlövák J, Chiang CE, Diez M, et al. Efficacy of dapagliflozin in heart failure with reduced ejection fraction according to body mass index. *Eur J Heart Fail* 2021;**23**:1662–1672. <https://doi.org/10.1002/ehf.2308>
- McMurray JJV, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med* 2014;**371**:993–1004. <https://doi.org/10.1056/NEJMoa1409077>
- McMurray JJV, Packer M, Desai AS, Gong J, Lefkowitz MP, Rizkala AR, et al. Dual angiotensin receptor and neprilysin inhibition as an alternative to angiotensin-converting enzyme inhibition in patients with chronic systolic heart failure: rationale for and design of the prospective comparison of ARNI with ACEI to determine impact. *Eur J Heart Fail* 2013;**15**:1062–1073. <https://doi.org/10.1093/eurjhf/hft052>
- Oreopoulos A, Padwal R, Kalantar-Zadeh K, Fonarow GC, Norris CM, McAlister FA. Body mass index and mortality in heart failure: a meta-analysis. *Am Heart J* 2008;**156**:13–22. <https://doi.org/10.1016/j.ahj.2008.02.014>
- Sharma A, Lavie CJ, Borer JS, Vallakati A, Goel S, Lopez-Jimenez F, et al. Meta-analysis of the relation of body mass index to all-cause and cardiovascular mortality and hospitalization in patients with chronic heart failure. *Am J Cardiol* 2015;**115**:1428–1434. <https://doi.org/10.1016/j.amjcard.2015.02.024>
- Lavie CJ, Osman AF, Milani RV, Mehra MR. Body composition and prognosis in chronic systolic heart failure: the obesity paradox. *Am J Cardiol* 2003;**91**:891–894. [https://doi.org/10.1016/S0002-9149\(03\)00031-6](https://doi.org/10.1016/S0002-9149(03)00031-6)
- Horwich TB, Fonarow GC, Hamilton MA, MacLellan WR, Woo MA, Tillich JH. The relationship between obesity and mortality in patients with heart failure. *J Am Coll Cardiol* 2001;**38**:789–795. [https://doi.org/10.1016/S0735-1097\(01\)01448-6](https://doi.org/10.1016/S0735-1097(01)01448-6)
- Cornier M-A, Després J-P, Davis N, Grossniklaus DA, Klein S, Lamarche B, et al. Assessing adiposity: a scientific statement from the American heart association. *Circulation* 2011;**124**:1996–2019. <https://doi.org/10.1161/CIR.0b013e318233bc6a>
- Parcha V, Patel N, Kalra R, Suri SS, Arora G, Wang TJ, et al. Obesity and serial nt-probnp levels in guided medical therapy for heart failure with reduced ejection fraction: insights from the guide-it trial. *J Am Heart Assoc* 2021;**10**:e018689. <https://doi.org/10.1161/JAHA.120.018689>
- Bozkurt B, Aguilar D, Deswal A, Dunbar SB, Francis GS, Horwich T, et al. Contributory risk and management of comorbidities of hypertension, obesity, diabetes Mellitus, hyperlipidemia, and metabolic syndrome in chronic heart failure: a scientific statement from the American heart association. *Circulation* 2016;**134**:e535–e578. <https://doi.org/10.1161/CIR.0000000000000450>
- Powell-Wiley TM, Poirier P, Burke LE, Després JP, Gordon-Larsen P, Lavie CJ, et al. Obesity and cardiovascular disease: a scientific statement from the American heart association. *Circulation* 2021;**143**:E984–E1010. <https://doi.org/10.1161/CIR.0000000000000973>
- McDonagh TA, Metra M, Adamo M, Gardner RS, Baumbach A, Böhm M, et al. 2021 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J* 2021;**42**:3599–3726. <https://doi.org/10.1093/eurheartj/ehab368>

46. McCloskey CA, Ramani GV, Mathier MA, Schauer PR, Eid GM, Mattar SG, et al. Bariatric surgery improves cardiac function in morbidly obese patients with severe cardiomyopathy. *Surg Obes Relat Dis* 2007;**3**:503–507. <https://doi.org/10.1016/j.soard.2007.05.006>
47. Evangelista LS, Heber D, Li Z, Bowerman S, Hamilton MA, Fonarow GC. Reduced body weight and adiposity with a high-protein diet improves functional Status, lipid profiles, glycemic control, and quality of life in patients with heart failure: a feasibility study. *J Cardiovasc Nurs* 2009;**24**:207–215. <https://doi.org/10.1097/JCN.0b013e31819846b9>
48. Pritchett AM, Deswal A, Aguilar D. Lifestyle modification with diet and exercise in obese patients with heart failure—a pilot study. *J Obes Weight Loss Ther* 2012;**02**:1–8. <https://doi.org/10.4172/2165-7904.1000118>
49. Horwich TB, Broderick S, Chen L, McCullough PA, Strzelczyk T, Kitzman DW, et al. Relation among body mass index, exercise training, and outcomes in chronic systolic heart failure. *Am J Cardiol* 2011;**108**:1754–1759. <https://doi.org/10.1016/j.amjcard.2011.07.051>
50. Flynn KE, Piña IL, Whellan DJ, Lin L, Blumenthal JA, Ellis SJ, et al. Effects of exercise training on health status in patients with chronic heart failure HF-ACTION randomized controlled trial. *JAMA* 2009;**301**:1451–1459. <https://doi.org/10.1001/jama.2009.457>
51. Research Study to Investigate How Well Semaglutide Works in People Living With Heart Failure and Obesity (STEP-HFpEF). <https://clinicaltrials.gov/ct2/show/NCT04788511>.
52. A Study of Tirzepatide (LY3298176) in Participants With Heart Failure With Preserved Ejection Fraction and Obesity (SUMMIT). <https://clinicaltrials.gov/ct2/show/NCT04847557>.
53. Heidenreich PA, Bozkurt B, Aguilar D, Allen LA, Byun JJ, Colvin MM, et al. 2022 AHA/ACC/HFSA guideline for the management of heart failure: a report of the American college of cardiology/American heart association joint committee on clinical practice guidelines. *Circulation*;2022:e895–e1032.
54. Olivier A, Pitt B, Girerd N, Lamiral Z, Machu JL, McMurray JJV, et al. Effect of eplerenone in patients with heart failure and reduced ejection fraction: potential effect modification by abdominal obesity. Insight from the EMPHASIS-HF trial. *Eur J Heart Fail* 2017;**19**:1186–1197. <https://doi.org/10.1002/ejhf.792>
55. Verweij LM, Terwee CB, Proper KI, Hulshof CTJ, van Mechelen W. Measurement error of waist circumference: gaps in knowledge. *Public Health Nutr* 2013;**16**:281–288. <https://doi.org/10.1017/S1368980012002741>
56. McAuley PA, Keteyian SJ, Brawner CA, Dardari ZA, Al Rifai M, Ehrman JK, et al. Exercise capacity and the obesity paradox in heart failure: the FIT (henry ford exercise testing) project. *Mayo Clin Proc* 2018;**93**:701–708. <https://doi.org/10.1016/j.mayocp.2018.01.026>
57. Lavie CJ, Cahalin LP, Chase P, Myers J, Bensimhon D, Peberdy MA, et al. Impact of cardiorespiratory fitness on the obesity paradox in patients with heart failure. *Mayo Clin Proc* 2013;**88**:251–258. <https://doi.org/10.1016/j.mayocp.2012.11.020>
58. Clark AL, Fonarow GC, Horwich TB. Impact of cardiorespiratory fitness on the obesity paradox in patients with systolic heart failure. *Am J Cardiol* 2015;**115**:209–213. <https://doi.org/10.1016/j.amjcard.2014.10.023>
59. Elagizi A, Carbone S, Lavie CJ, Mehra MR, Ventura HO. Implications of obesity across the heart failure continuum. *Prog Cardiovasc Dis* 2020;**63**:561–569. <https://doi.org/10.1016/j.pcad.2020.09.005>