



Observation of gauge boson joint-polarisation states in $W^\pm Z$ production from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector



The ATLAS Collaboration*

ARTICLE INFO

Article history:

Received 18 November 2022

Received in revised form 13 March 2023

Accepted 2 April 2023

Available online 7 June 2023

Editor: M. Doser

Dataset link: <https://hepdata.cedar.ac.uk>

ABSTRACT

Measurements of joint-polarisation states of W and Z gauge bosons in $W^\pm Z$ production are presented. The data set used corresponds to an integrated luminosity of 139 fb^{-1} of proton-proton collisions at a centre-of-mass energy of 13 TeV recorded by the ATLAS detector at the CERN Large Hadron Collider. The $W^\pm Z$ candidate events are reconstructed using leptonic decay modes of the gauge bosons into electrons and muons. The simultaneous pair-production of longitudinally polarised vector bosons is measured for the first time with a significance of 7.1 standard deviations. The measured joint helicity fractions integrated over the fiducial region are $f_{00} = 0.067 \pm 0.010$, $f_{0\Gamma} = 0.110 \pm 0.029$, $f_{\Gamma 0} = 0.179 \pm 0.023$ and $f_{\Gamma\Gamma} = 0.644 \pm 0.032$, in agreement with the next-to-leading-order Standard Model predictions. Individual helicity fractions of the W and Z bosons are also measured and found to be consistent with joint helicity fractions within the expected amounts of correlation. Both the joint and individual helicity fractions are also measured separately in W^+Z and W^-Z events. Inclusive and differential cross sections for several kinematic observables sensitive to polarisation are presented.

© 2023 Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The study of weak boson polarisation provides a way to directly probe the Standard Model (SM) gauge and Higgs sectors and to discriminate between SM and beyond the Standard Model (BSM) signals. The existence of the longitudinally polarised state of weak bosons is a consequence of the non-vanishing mass of the bosons generated by the electroweak (EW) symmetry breaking mechanism. The measurement of the polarisation in diboson production therefore tests both the SM's gauge symmetry structure, through the existence of the triple gauge coupling, and the particular way this symmetry is spontaneously broken, via the longitudinal helicity state. Angular observables can be used to look for new interactions that can lead to different polarisation behaviour than predicted by the SM, to which the $W^\pm Z$ final state would be particularly sensitive [1,2].

At hadron colliders, the polarisation of the W boson was previously measured in the decay of the top quark by the CDF and DØ [3–5] Collaborations and by the ATLAS [6] and CMS [7] Collaborations, as well as in single W production by ATLAS [8] and CMS [9]. Polarisation and several other related angular coefficient measurements of singly produced Z bosons were published by the CDF [10], CMS [11] and ATLAS [12] Collaborations. The polarisation of W bosons was also measured in ep collisions by the H1 Collaboration [13]. Finally, for dibosons, first measurements of the W polarisation were performed by LEP experiments in W pair production in e^+e^- collisions [14,15] and were used to set limits on anomalous triple gauge couplings in Ref. [16]. Correlations between polarisation states of W particles were also probed experimentally at LEP [17–19]. Helicity fractions of pair-produced vector bosons were measured independently of each other and for the first time in hadronic collisions at a centre-of-mass energy of $\sqrt{s} = 13 \text{ TeV}$ by the ATLAS Collaboration [20], followed by the CMS Collaboration [21], using integrated luminosities of 36 fb^{-1} and 137 fb^{-1} , respectively. Benefiting from the full ATLAS data set available at $\sqrt{s} = 13 \text{ TeV}$, combined with novel analysis techniques, observables more difficult to extract from data, such as the correlations between helicity states of the vector bosons, can now be accessed.

This Letter reports on an observation and measurement of joint polarisation states of weak bosons in $W^\pm Z$ production, extending the pioneering work of Ref. [20]. The W and Z bosons are reconstructed using their decay modes into electrons or muons. The pp collision data were collected with the ATLAS detector from 2015 to 2018 at a centre-of-mass energy of $\sqrt{s} = 13 \text{ TeV}$ and correspond to an integrated luminosity of 139 fb^{-1} . The individual and joint boson polarisation state fractions are measured consistently in the detector fiducial region and their relationships are studied. The reported measurements are compared to the recent fixed-order theory

* E-mail address: atlas.publications@cern.ch.

calculation of polarised W and Z boson pair production at next-to-leading-order (NLO) accuracy in quantum chromodynamics (QCD) [22, 23]. Additionally, the $W^\pm Z$ production cross section is measured within a fiducial phase space both inclusively and differentially as a function of several angular variables related to the helicity states of the $W^\pm Z$ system.

2. Theoretical framework and measurement phase space

A complete description of the polarisation states of the produced W and Z bosons is given in terms of the two-particle joint spin density matrix $\rho_{\lambda_W \lambda'_W \lambda_Z \lambda'_Z}$ expressed in the basis of joint-helicity states $(\lambda_W \lambda_Z)$ where λ_W and λ_Z are the helicities of the W and Z bosons, respectively. In terms of the $W^\pm Z$ production helicity amplitudes, $F_{\lambda_W \lambda_Z}^{(\mu_q \mu_{\bar{q}})}$, the spin density matrix elements are defined by

$$\rho_{\lambda_W \lambda'_W \lambda_Z \lambda'_Z} \equiv \frac{1}{C} \times \sum_{\mu_q \mu_{\bar{q}}} F_{\lambda_W \lambda_Z}^{(\mu_q \mu_{\bar{q}})} F_{\lambda'_W \lambda'_Z}^{(\mu_q \mu_{\bar{q}})*}, \text{ where } C = \sum_{\mu_q \mu_{\bar{q}} \lambda_W \lambda_Z} \left| F_{\lambda_W \lambda_Z}^{(\mu_q \mu_{\bar{q}})} \right|^2.$$

The initial-state helicity is averaged in the numerator and denominator with sums running over $\mu_q, \mu_{\bar{q}} = \pm 1/2$. In the denominator the W and Z bosons helicities are summed over λ_i , where $\lambda_i = \pm 1, 0$, to normalise the trace of the matrix to unity. The complete 9×9 joint spin density matrix $\rho_{\lambda_W \lambda'_W \lambda_Z \lambda'_Z}$ cannot yet be measured because of the limited available data sample. Instead, the following four linear combinations of matrix elements are considered

$$\begin{aligned} f_{00} &= \rho_{0000}, \\ f_{TT} &= \rho_{++--} + \rho_{--++} + \rho_{----} + \rho_{++++}, \\ f_{0T} &= \rho_{00--} + \rho_{00++}, \\ f_{T0} &= \rho_{--00} + \rho_{++00}. \end{aligned}$$

The quantities f_{00} , f_{TT} , f_{0T} and f_{T0} are composed of diagonal elements of the full matrix and can be interpreted as probabilities of correlated helicity states of the two W and Z bosons. The distinction between the two transverse helicities of the bosons (right-handed, R, and left-handed, L, also indicated with + and -, respectively) is not made.

The single-boson spin density matrix $\rho_{\lambda \lambda'}$ is derived from the joint spin density matrix $\rho_{\lambda_W \lambda'_W \lambda_Z \lambda'_Z}$ by summing over indices of one of the bosons. It provides information about the helicity of one of the bosons regardless of the state of the other. Diagonal elements of the single-boson spin density matrix are the individual helicity fractions f_0 , f_L and f_R for the longitudinal, transverse left-handed and transverse right-handed helicity states, respectively. The normalisation is defined such that $f_0 + f_L + f_R = 1$.

Helicity fractions are not Lorentz invariant and hence depend on a chosen reference frame. In diboson production a natural frame choice for single-boson helicity fraction measurement is to fix the reference z-axis to the direction of the corresponding boson in the diboson centre-of-mass frame [23]. This definition is called the modified helicity coordinate system and is used for this measurement. The back-to-back kinematics of the bosons in the $W^\pm Z$ centre-of-mass frame implies that the polarisation vectors of the two bosons are defined with respect to the same reference axis, but in opposite directions. This choice also maximises the decorrelation between 00 and TT helicity states.

The relative contributions of joint helicity states can be determined from the absolute value of the cosine of the scattering angle of the W or Z boson in the WZ centre-of-mass frame, $|\cos \theta_V|$, calculated with respect to the beam axis [24, 25]. Events with 00 helicity have $|\cos \theta_V|$ values closer to 0 while events with TT helicity have $|\cos \theta_V|$ values closer to 1.

The polarisation of an individual gauge boson can be determined from the angular distribution of its decay products. At the Born level, the expected angular distribution for massless fermions in the rest frame of the parent W boson is given by [26–29]

$$\frac{1}{\sigma_{W^\pm Z}} \frac{d\sigma_{W^\pm Z}}{d \cos \theta_{\ell^W}^*} = \frac{3}{8} f_L [(1 - q_W \cdot \cos \theta_{\ell^W}^*)^2] + \frac{3}{8} f_R [(1 + q_W \cdot \cos \theta_{\ell^W}^*)^2] + \frac{3}{4} f_0 \sin^2 \theta_{\ell^W}^*, \quad (1)$$

where q_W is the charge of the W boson and $\theta_{\ell^W}^*$ is defined using the modified helicity frame as the decay angle of the charged lepton in the W rest frame relative to the W direction in the $W^\pm Z$ centre-of-mass frame. All dependencies on the azimuthal angle are integrated over. The expected angular distribution of the lepton decay products of the Z boson is described by the generalisation of Equation (1) [26–28]

$$\frac{1}{\sigma_{W^\pm Z}} \frac{d\sigma_{W^\pm Z}}{d \cos \theta_{\ell^Z}^*} = \frac{3}{8} f_L (1 + 2\alpha \cos \theta_{\ell^Z}^* + \cos^2 \theta_{\ell^Z}^*) + \frac{3}{8} f_R (1 - 2\alpha \cos \theta_{\ell^Z}^* + \cos^2 \theta_{\ell^Z}^*) + \frac{3}{4} f_0 \sin^2 \theta_{\ell^Z}^*, \quad (2)$$

where $\theta_{\ell^Z}^*$ is defined using the modified helicity frame as the decay angle of the negatively charged lepton in the Z rest frame relative to the Z direction in the $W^\pm Z$ centre-of-mass frame. The parameter $\alpha = (2c_v c_a)/(c_v^2 + c_a^2) = 0.147$ is expressed in terms of the vector $c_v = -\frac{1}{2} + 2 \sin^2 \theta_W^{\text{eff}}$ and axial-vector $c_a = -\frac{1}{2}$ couplings of the Z boson to leptons, where the effective value of the Weinberg angle $\sin^2 \theta_W^{\text{eff}} = 0.23152$ [30] is used. Equation (2) also holds for the contribution from γ^* and its interference with the Z boson, with appropriate c_v and c_a coefficients.

The reported helicity fractions and cross sections are measured in a fiducial phase space chosen to closely follow the event selection criteria described in Section 5. It is based on the kinematics of particle-level objects as defined in Ref. [31]. These are final-state prompt¹ leptons associated with the W and Z boson decays. The kinematics of final-state prompt leptons is defined before QED final-state radiation (FSR), i.e. Born leptons.

The Born charged leptons from Z and W boson decay must have $|\eta| < 2.5$ and transverse momentum p_T above 15 GeV and 20 GeV, respectively; the invariant mass of the two leptons from the Z boson decay differs by at most 10 GeV from the world average value of

¹ A prompt lepton is a lepton that is not produced in the decay of a hadron, a τ -lepton, or their descendants.

the Z boson mass $m_Z^{\text{PDG}} = 91.1876$ GeV [30]. The W transverse mass, defined as $m_W^T = \sqrt{2 \cdot p_T^\nu \cdot p_T^\ell \cdot [1 - \cos \Delta\phi(\ell, \nu)]}$, where $\Delta\phi(\ell, \nu)$ is the angle between the lepton and the neutrino in the transverse plane, and p_T^ℓ and p_T^ν are the transverse momenta of the lepton from W boson decay and of the neutrino, respectively, must be greater than 30 GeV. In addition, it is required that the angular distance ΔR between the charged leptons from the W and Z decay is larger than 0.3, and that ΔR between the two leptons from the Z decay is larger than 0.2.

The contribution from off-shell effects is minimised by the tight invariant mass window of ± 10 GeV around the nominal Z boson mass used for the definition of the fiducial phase space. These contributions are estimated from NLO QCD calculations to amount to 2% at the integrated level and in most of the differential distributions [23]. However, all the helicity fractions presented here are effective fractions and contain the small non-resonant contributions present in the fiducial phase space, including γ^* events. Interferences among helicity states are also expected to be small, amounting to 0.6% according to NLO QCD+EW calculations [32]. Their effect is included in the measured helicity fractions.

A total phase space is also defined by requiring only the invariant mass of the lepton pair associated with the Z boson to be in the range $66 < m_{\ell\ell} < 116$ GeV. This serves for the generation of polarised Monte Carlo (MC) event templates described in Section 7 for which no restrictions should be applied to the phase space of the decay leptons from W and Z bosons.

3. ATLAS detector

The ATLAS experiment [33] at the LHC is a multipurpose particle detector with a forward–backward symmetric cylindrical geometry and a near 4π coverage in solid angle.² It consists of an inner tracking detector (ID) surrounded by a thin superconducting solenoid providing a 2 T axial magnetic field, electromagnetic and hadron calorimeters and a muon spectrometer (MS). The inner tracking detector covers the pseudorapidity range $|\eta| < 2.5$. It consists of silicon pixel, silicon microstrip and transition radiation tracking detectors. Lead/liquid-argon (LAr) sampling calorimeters provide electromagnetic (EM) energy measurements with high granularity. A steel/scintillator-tile hadron calorimeter covers the central pseudorapidity range ($|\eta| < 1.7$). The endcap and forward regions are instrumented with LAr calorimeters for both the EM and hadronic energy measurements up to $|\eta| = 4.9$. The muon spectrometer surrounds the calorimeters and is based on three large superconducting air-core toroidal magnets with eight coils each. The field integral of the toroids ranges between 2.0 and 6.0 Tm across most of the detector. The muon spectrometer includes a system of precision tracking chambers and fast detectors for triggering. A two-level trigger system is used to select events. The first-level trigger is implemented in hardware and uses a subset of the detector information to accept events at a rate below 100 kHz. This is followed by a software-based trigger that reduces the accepted event rate to 1 kHz on average depending on the data-taking conditions. An extensive software suite [34] is used in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

4. Signal and background simulation

MC simulation is used to model signal and most background processes.

An inclusive sample of simulated $W^\pm Z$ events is used to correct the signal yield for detector effects and to compare the measurements with the theoretical predictions. The production of $W^\pm Z$ pairs and the subsequent leptonic decays of the vector bosons were simulated at NLO in QCD and LO in QED using the Powheg-Box v2 [35–38] generator, interfaced to PYTHIA 8.210 [39] for simulation of parton showering, hadronisation and the underlying event. Final-state radiation resulting from QED interactions was simulated using PYTHIA 8.210 with the AZNLO [40] set of tuned parameters. The CT10NLO [41] parton distribution function (PDF) set was used for the hard-scattering process, while the CTEQ6L1 [42] PDF set was used for the parton shower. The sample was generated with dynamic renormalisation and factorisation QCD scales, μ_r and μ_f , equal to $m_{WZ}/2$, where m_{WZ} is the invariant mass of the WZ system [20].

For systematic uncertainties studies, two alternative inclusive $W^\pm Z$ signal samples were generated at NLO in QCD. The first alternative sample was generated using MADGRAPH5_AMC@NLO 2.6.5 [43]. Matrix elements containing three leptons, one neutrino and up to two jets in the final state were calculated at NLO QCD and merged with the parton shower from PYTHIA 8.210 using the FxFx scheme [44]. The NNPDF3.0NLO [45] PDF set was used for the hard-scattering process. The default dynamic renormalisation and factorisation scale set by MADGRAPH5_AMC@NLO was used [46]. This MC sample is referred to as MADGRAPH5_AMC@NLO+PYTHIA.

The second alternative $W^\pm Z$ signal sample was generated using the SHERPA 2.2.2 generator [47]. Matrix elements contain all diagrams with four electroweak vertices. They were calculated for up to one parton at NLO and up to three partons at LO using Comix [48] and OPENLOOPS [49], and merged with the SHERPA parton shower [50] according to the ME+PS@NLO prescription [51]. The NNPDF3.0NNLO PDF set was used along with the associated set of tuned parton-shower parameters developed by the SHERPA authors.

Polarised $W^\pm Z$ events were generated at LO in QCD using MADGRAPH5_AMC@NLO 2.7.3 [52] interfaced to PYTHIA 8.244 for simulation of parton showering, hadronisation and the underlying event. The generation of polarised events is not possible at the full NLO QCD accuracy. To account for the real part of NLO corrections, events were simulated with either no jets or one jet in the matrix element at LO, and merged with PYTHIA parton shower using the CKKW-L scheme [53,54]. The NNPDF3.0NLO PDF set was used for the hard-scattering process. Four separate MC samples were generated corresponding to the four joint helicity states 00, 0T, TO and TT, respectively. An inclusive MC sample was created by adding the four polarised samples. These MC sets are referred to as MADGRAPH o,ij@LO.

The background sources in this analysis include processes with two or more electroweak gauge bosons, namely ZZ , WW and VVV ($V = W, Z$); processes with top quarks, such as $t\bar{t}$ and $t\bar{t}V$, single top and tZj ; and processes with gauge bosons associated with jets or photons ($Z + j$ and $Z\gamma$). Electroweak $W^\pm Z$ production, $WZjj$ –EW, arising at the order α_E^6 is also considered as a background and not part of the measured signal. MC simulation is used to estimate the contribution from background processes with three or more prompt

² ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

leptons. Background processes with at least one misidentified lepton are evaluated using data-driven techniques and simulated events are used to assess the systematic uncertainties of these backgrounds.

The SHERPA 2.2.2 event generator was used to simulate both the $q\bar{q}$ and gg -initiated ZZ processes, including $H \rightarrow ZZ$ production, using the NNPDF3.0NNLO PDF set. It provides a matrix element calculation accurate to NLO in α_s for 0- and 1-jet final states, and LO accuracy for 2- and 3-jet final states. Triboson events were simulated by the SHERPA 2.2.2 event generator at NLO accuracy with 0 additional partons and at LO accuracy with one and two additional partons and using the NNPDF3.0NNLO PDF set.

Both the $t\bar{t}Z$ and $t\bar{t}W$ processes were generated at NLO with the MADGRAPH5_AMC@NLOMC generator, using the NNPDF3.0NLO PDF set, and interfaced to the PYTHIA 8.186 [55] parton shower model.

Finally, the tZj and $WZjj$ -EW processes were generated at LO by the MADGRAPH5_AMC@NLOMC generator interfaced with PYTHIA 8.244 for the modelling of the parton shower, hadronisation and underlying event. The parton distribution function set was NNPDF3.0NLO. In the simulation of the parton shower the NNPDF2.3Lo [56] PDF set was used.

All generated MC events were passed through the ATLAS detector simulation [57], based on GEANT4 [58], and processed using the same reconstruction software as used for the data. The event samples include the simulation of additional pp interactions (pile-up) generated with PYTHIA 8.186 using the NNPDF2.3Lo PDF set and the A3 [59] set of tuned parameters for the underlying event and parton fragmentation. Simulated events were reweighted to match the pile-up conditions observed in the data. Scale factors were applied to simulated events to correct for small differences in the efficiencies observed in data and predicted by MC simulation for the trigger, reconstruction, identification, isolation and impact parameter requirements of electrons and muons [60,61]. Furthermore, the electron energy and muon momentum in simulated events were smeared to account for small differences in resolution between data and MC events [60,62].

5. Event reconstruction and selection

Only data recorded with stable beam conditions and with all relevant detector subsystems operational are considered. Candidate events are selected using triggers [63–65] that require at least one electron or muon. These triggers require leptons to pass different transverse momentum threshold and isolation criteria, which depend on the data-taking run period and the instantaneous luminosity. The combined efficiency of these triggers is 99% for $W^\pm Z$ events satisfying the offline selection criteria. Events are required to have a primary vertex compatible with the LHC luminous region inside the ATLAS detector. The primary vertex is defined as the reconstructed vertex with at least two charged-particle tracks that has the largest sum of the p_T^2 for the associated tracks.

Muon candidates are identified by tracks reconstructed in the MS and matched to tracks reconstructed in the ID. Muons are required to pass a ‘medium’ identification selection, which is based on requirements on the number of hits in the ID and the MS [61,62]. The efficiency of this selection averaged over p_T and η is larger than 98%. The muon momentum is measured by combining the MS measurement, corrected for the energy deposited in the calorimeters, and the ID measurement. The p_T of the muon must be greater than 15 GeV and its pseudorapidity must satisfy $|\eta| < 2.5$.

Electron candidates are reconstructed from energy clusters in the electromagnetic calorimeter matched to ID tracks. Electrons are identified using a discriminant that is the value of a likelihood function constructed with information about the shape of the electromagnetic showers in the calorimeter, the track properties and the quality of the track-to-cluster matching for the candidate [60]. Electrons must satisfy a ‘medium’ likelihood requirement, which provides an overall identification efficiency of 90%. The electron momentum is computed from the cluster energy and the direction of the track. The p_T of the electron must be greater than 15 GeV and the pseudorapidity of the cluster must satisfy $|\eta| < 1.37$ or $1.52 < |\eta| < 2.47$.

Electron and muon candidates are required to originate from the primary vertex and to be isolated from other particles using both calorimeter-cluster and ID-track information. The isolation efficiency is at least 90% for electrons with $p_T > 25$ GeV and muons with $p_T > 15$ GeV and at least 99% for electrons and muons with $p_T > 60$ GeV [60,61].

Jets of hadrons are reconstructed using a particle-flow algorithm based on noise-suppressed positive-energy topological clusters in the calorimeter [66]. Energy deposited in the calorimeter by charged particles is subtracted and replaced by the momenta of tracks which are matched to those topological clusters. The jets are clustered using the anti- k_t algorithm [67,68] with a radius parameter $R = 0.4$. They are calibrated according to in situ measurements of the jet energy scale [69]. All jets must have $p_T > 25$ GeV and be reconstructed in the pseudorapidity range $|\eta| < 4.5$. Jets with p_T below 60 GeV and with $|\eta| < 2.4$ have to pass a requirement on the jet vertex tagger [70], a likelihood discriminant that uses a combination of track and vertex information to suppress jets originating from pile-up activity. Jets with $|\eta| < 2.5$ containing a b -hadron are identified with a deep-learning neural network (NN) [71–73] which uses distinctive features of b -hadron decays in terms of the impact parameters of the tracks and the displaced vertices reconstructed in the inner detector. Jets initiated by b -quarks are selected by setting the algorithm’s output threshold such that an 85% b -jet selection efficiency is achieved in simulated $t\bar{t}$ events, with a rejection factor of 40 against light-flavour jets.

The transverse momentum of the neutrino is estimated from the missing transverse momentum in the event, E_T^{miss} , calculated as the negative vector sum of the transverse momentum of all identified hard physics objects (electrons, muons, jets), with a contribution from an additional soft term. This soft term is calculated from ID tracks matched to the primary vertex and not assigned to any of the hard objects [74].

Events are required to contain exactly three lepton candidates satisfying the selection criteria described above. To ensure that the trigger efficiency is well determined, at least one of the candidate leptons is required to have $p_T > 25$ GeV for 2015 and $p_T > 27$ GeV for 2016–2018 data, as well as being geometrically matched to a lepton that was selected by the trigger. No requirement on the number of jets is applied.

In order to suppress background processes with at least four prompt leptons, events with a fourth lepton candidate satisfying looser selection criteria are rejected. For this looser selection, the lepton p_T requirement is lowered to $p_T > 5$ GeV, electrons are allowed to be reconstructed in the region $1.37 < |\eta| < 1.52$ and ‘loose’ identification requirements [60,61] are used for both the electrons and muons. A less stringent requirement is applied for electron isolation and is based only on ID track information. No dedicated identification algorithm is used to suppress events with electrons and muons originating from the decay of τ -leptons.

Candidate events are required to have at least one pair of leptons with the same flavour and opposite charge, with an invariant mass that is consistent with m_Z^{PDG} to within 10 GeV. This pair is considered to be the Z boson candidate. If more than one pair can be formed, the pair whose invariant mass is closest to the nominal Z boson mass is taken as the Z boson candidate. The remaining third lepton is assigned to the W boson decay. The transverse mass of the W candidate, computed using E_T^{miss} and the p_T of the associated lepton, is required to be greater than 30 GeV.

Backgrounds originating from misidentified leptons are suppressed by requiring the lepton associated with the W boson to satisfy more stringent selection criteria. The transverse momentum of this lepton is therefore required to be greater than 20 GeV. This lepton is also required to pass the ‘tight’ identification requirements, which results in an efficiency between 90% and 98% for muons and an overall efficiency of 85% for electrons.

The main challenge in the reconstruction of the full kinematics of the selected events arises for the W boson from incomplete knowledge of the neutrino momentum. In the previous ATLAS publication [20] the neutrino longitudinal momentum p_z^ν was reconstructed analytically using the W mass constraint. In this analysis a neural network regression is developed to reconstruct p_z^ν . Input variables to the NN are p_T and p_z of the charged lepton from the W boson decay, the E_T^{miss} components perpendicular and parallel to the direction of this charged lepton and the p_z^ν reconstructed analytically. The NN is trained using $W^\pm Z$ Powheg+Pythia MC events at detector level to reconstruct event-by-event the true p_z^ν value in the MC simulation. The compatibility of the reconstructed W boson mass m_W with its generated value is ensured by using as the loss function the sum of the root-mean-square (RMS) of the deviations of m_W and p_z^ν reconstructed values from their generated values. The regression method provides a reasonable estimate of p_z^ν for events for which the analytical method fails³ and reduces the p_z^ν error RMS by 10% thanks to the improved resolution at high p_z^ν . This improvement comes from fact that the NN is trained to account for the full width of the generated W boson, whereas the analytical method imposes a single value of the W boson mass, fixed to its pole mass, for all events. Additionally, the regression method improves the m_W resolution from 41% obtained with the analytical method to 31%. This in turn improves the reconstruction of angular observables in the modified helicity frame.

6. Background estimation

The background sources are classified into two groups: events where at least one of the candidate leptons is not a prompt lepton (reducible background) and events where all candidates are prompt leptons or are produced in the decay of a τ -lepton (irreducible background). Candidates that are not prompt leptons are also called ‘misidentified’ leptons.

Events in the first group mostly originate from $Z + j$, $Z\gamma$, $t\bar{t}$, and WW production processes and constitute about 5% of all selected events. This reducible background is estimated with a data-driven method based on the inversion of a matrix containing the efficiencies and misidentification probabilities for prompt and misidentified leptons [20,75–77]. The method exploits the classification of the leptons as loose or tight candidates and the probability that a non-prompt lepton is misidentified as a loose or tight lepton. Tight leptons are leptons passing the selection criteria described in Section 5. Loose leptons are leptons that do not meet the isolation and identification criteria for signal leptons but satisfy only looser criteria. The misidentification probabilities for misidentified leptons are determined from data using dedicated control samples each enriched in misidentified leptons from light- or heavy-flavour jets and from photon conversions, respectively. The lepton efficiencies for prompt leptons are determined as detailed in Refs. [60–62]. The lepton efficiencies and misidentification probabilities are combined with event rates in data samples of $W^\pm Z$ candidate events where at least one and up to three of the leptons are loose. Then, solving a system of linear equations, the number of events with at least one misidentified lepton is obtained, which represents the amount of reducible background in the $W^\pm Z$ sample. About 2% of this background contribution arises from events with two misidentified leptons. The background from events with three misidentified leptons, e.g. from multijet processes, is negligible. The method allows the shape of any kinematic distribution of reducible background events to be estimated. Another independent method of assessing the reducible background was also considered. This method estimates the amount of reducible background using MC simulations scaled to data by process-dependent factors determined from the data-to-MC comparison in dedicated control regions. Agreement within 20% with the matrix method estimate is obtained in both yield and shape of the distributions of irreducible background events.

The events contributing to the second group of background processes represent $\sim 18\%$ of all selected events. They originate from ZZ , $t\bar{t} + V$, VVV (where $V = Z$ or W) and tZj events. Events from $W^\pm Z$ production with at least one τ -lepton decay or from electroweak $WZjj$ –EW production are also considered as backgrounds and not part of the measured signal. The amount of irreducible background is estimated using MC simulations. The dominant contribution in this second group is from ZZ production, where one of the leptons from the ZZ decay falls outside the detector acceptance. It represents about 7.5% of all selected events. The MC-based estimates of the ZZ and $t\bar{t} + V$ backgrounds are validated by comparing the MC expectations with the event yield and several kinematic distributions of data samples enriched in ZZ and $t\bar{t} + V$ events, respectively. The ZZ control sample is selected from events with a Z candidate which meets all of the analysis selection criteria and which is accompanied by two additional leptons, satisfying the lepton criteria described in Section 5. The ZZ MC expectation is rescaled to match the data by combining the $W^\pm Z$ signal region and a single bin containing all of the events from the ZZ control region in the fit detailed in Section 7 and used to extract the helicity fractions. A scaling factor of 1.13 is obtained from the fit for the ZZ MC contribution, in agreement with the ZZ cross-section measurements performed at $\sqrt{s} = 13$ TeV [78]. The $t\bar{t} + V$ control sample is selected by requiring selected $W^\pm Z$ events to have at least two reconstructed b -jets. The observed data event yield in this validation region is matching the $t\bar{t} + V$ MC prediction rescaled by a factor of 1.3. This scaling factor relative to predictions is in line with the $t\bar{t} + Z$ cross-section measurements performed at $\sqrt{s} = 13$ TeV [79]. The contribution of $t\bar{t} + V$ events amounts to $\sim 4\%$ of all selected events. Systematic uncertainties affecting the rescaling of ZZ and $t\bar{t} + V$ MC predictions are discussed in Section 8. Finally, some $W^\pm \gamma^*$ events produced outside of the total phase space can be selected at detector level. The contribution of such events is estimated from the $W^\pm Z$ Powheg+Pythia MC simulation to amount to 0.5% and is subtracted from data.

³ The analytical method relies on a second-order equation which has complex solutions for roughly 30% of the events. In those cases, its imaginary part is set to zero.

7. Measurement methodology

7.1. Joint boson polarisation

In the Born-level fiducial phase space, the $|\cos\theta_V|$, $q_W \cdot \cos\theta_{\ell^W}^*$ and $\cos\theta_{\ell^Z}^*$ angular observables can be used together to separate the joint helicity states from an inclusive MC sample and determine their respective generated fractions. However, when using the same angular observables at detector level, the separation of the different joint helicity states is not sufficient for the measurement of their respective fractions with an expected precision of 5σ . Therefore, a multivariate discriminant exploiting more kinematic observables is used to better separate the four joint helicity states. A deep neural network (DNN) classifier is implemented to exploit kinematic differences between polarisation states in different observables. Keras v2.2.4-tf [80] with the Tensorflow v1.3 [81] backend are used. The DNN is trained and optimised using simulated MADGRAPH 0,1j@LO polarised MC samples. A total of eight variables are used as inputs to the DNN: the transverse momenta of the three leptons and of the neutrino, $p_T^{\ell^W}, p_T^{\ell^Z_1}, p_T^{\ell^Z_2}$ and E_T^{miss} , angular variables as the absolute difference between the rapidities of the Z boson and the lepton from the decay of the W boson, $|y_Z - y_{\ell^W}|$, the azimuthal angle difference between the two leptons of each W and Z -boson decay, $\Delta\phi(\ell^W, \nu)$, $\Delta\phi(\ell^Z_1, \ell^Z_2)$, respectively, and the transverse momentum of the WZ system p_T^{WZ} . The DNN classifier is trained such that the components of the vector output, one for each of the four joint-polarisation states, are combined in a single output node in its last layer, used in the analysis for the separation. The optimisation of the combination is part of the DNN training procedure and a sigmoid activation function is applied to the combination output. The output is therefore a continuous score, labelled ‘DNN score,’ ranging from 0 to 1 that takes typical values peaking around 0, 0.5, and 1 for TT, mixed OT and T0, and 00 helicity states, respectively. The DNN classifier achieves integrals of the receiver operating characteristic (ROC) curves between 0.8 and 0.85 for separating the 00 state from each of the three others joint polarisation states. The sensitivity of the analysis in disentangling the mixed helicity states OT and T0 is achieved by categorising events depending on the $\cos\theta_{\ell^W}^*$ and $\cos\theta_{\ell^Z}^*$ observables. Four categories are defined as ($|\cos\theta_{\ell^W}^*| < 0.5, |\cos\theta_{\ell^Z}^*| < 0.5$), ($|\cos\theta_{\ell^W}^*| > 0.5, |\cos\theta_{\ell^Z}^*| < 0.5$), ($|\cos\theta_{\ell^W}^*| < 0.5, |\cos\theta_{\ell^Z}^*| > 0.5$) and ($|\cos\theta_{\ell^W}^*| > 0.5, |\cos\theta_{\ell^Z}^*| > 0.5$). A one-dimensional representation of the DNN score variable in the four ($|\cos\theta_{\ell^W}^*|, |\cos\theta_{\ell^Z}^*|$) categories, shown in Fig. 1 and referred to as the 4-category DNN score, is built.

To measure joint helicity fractions a binned maximum-likelihood fit [82] to data of the four helicity states templates is performed. The binned likelihood function used in the fit consists of a product of Poisson probability terms over bins of the 4-category DNN score distribution in the $W^\pm Z$ signal region and a single-bin integrating over the ZZ control region. Each source of systematic uncertainty is implemented in the likelihood function as a nuisance parameter with a Gaussian constraint, except for the NLO QCD modelling systematic uncertainty described in section 8. Independent parameters of the fit are the particle level helicity fractions f_{00}, f_{T0}, f_{TT} and the integrated fiducial cross section. The fourth joint helicity fraction f_{OT} is obtained using the constraint $f_{00} + f_{OT} + f_{T0} + f_{TT} = 1$. Differences between detector-level and particle-level fractions due to detector efficiencies and QED final-state radiation effects are accounted for in the fit using independent correction factors applied to each helicity fraction. The factors are obtained from simulation and range from 0.95 to 1.01 depending on the helicity state.

As observed in theory calculations [23], a strong relationship exists between NLO QCD corrections and polarisation effects. Therefore, templates for helicity states generated at LO are insufficient. The MADGRAPH 0,1j@LO MC simulation corrects for the real part of NLO QCD effects but misses virtual corrections that are also important for polarisation measurements. In order to verify that the shapes of the polarised templates are valid, a closure test is performed. Templates are used to fit pseudo-data generated using inclusive MC simulations at NLO QCD accuracy, such as with POWHEG+PYTHIA or MADGRAPH5_AMC@NLO+PYTHIA. The fit is performed on detector-level distributions. Polarisation fractions extracted from the fit are compared with the generated polarisation fractions of the NLO MC samples. Differences from 10% to 50% depending on the polarisation state are observed between extracted and built-in fractions when using polarised templates from the MADGRAPH 0,1j@LO generation. This demonstrates the need for using polarised templates at NLO QCD accuracy for polarisation measurements.

Templates better approaching the NLO QCD accuracy are built using a DNN-based event-by-event reweighting procedure [83]. Four DNNs are trained and each of them is specialised to reweight at particle-level the inclusive MADGRAPH 0,1j@LO events to one of the four joint polarised states. Input variables of the DNNs are those of the polarisation DNN classifier augmented by the invariant mass of the WZ system m_{WZ} , and the angular variables $\cos\theta_{\ell^W}^*$, $\cos\theta_{\ell^Z}^*$ and $|\cos\theta_V|$. The four DNNs are then applied to reweight the inclusive POWHEG+PYTHIA MC events, creating four polarised MC templates with NLO-like accuracy in QCD. This method provides the best fit closure, with no bias on the extracted polarisation fractions visible within the statistical precision of the closure test.

A second method is used to create NLO QCD accurate polarised templates, based on the available fixed-order parton-level theory predictions [23]. Predicted distributions, including the output score of the DNN classifier, are calculated in the parton-level fiducial phase space [23]. Corrections for parton-shower (PS) effects on the shape of the templates are incorporated by binned factors. The correction factors are calculated per helicity state by comparing MADGRAPH 0,1j@LO predictions at particle and parton level. The templates obtained at particle level are transformed to detector level using a folding procedure to account for detector effects. This method provides a good fit closure, reducing the bias up to $\sim 10\%$. This template creation method is used to estimate systematic uncertainties in the template shapes.

Interference among polarisation states manifests itself as a shape difference between the sum of polarised templates and the inclusive distribution. This difference is expected to be at most 4% for the DNN score observable [23]. The fit procedure adjusts fitted polarisation fractions such that the sum of polarised templates matches the inclusive distribution of data, with increased fit uncertainties. The measured polarisation fractions therefore include interference effects.

7.2. Individual boson polarisation

For individual W and Z boson polarisation states, the helicity parameters f_0 and $f_L - f_R$ are measured using a fit of templates for the three f_0 , f_L and f_R helicity states to the $q_W \cdot \cos\theta_{\ell^W}^*$ and $\cos\theta_{\ell^Z}^*$ distributions. The equation $f_0 + f_R + f_L = 1$ is used to constrain the independent parameters of the fit to determine f_0 , $f_L - f_R$ and the integrated fiducial cross section. Templates with NLO QCD accuracy

are also required for the $q_W \cdot \cos\theta_{\ell W}^*$ and $\cos\theta_{\ell Z}^*$ distributions. Neither such templates at NLO QCD accuracy nor templates for left- and right-handed helicity states of the vector bosons can be produced by the `MADGRAPH5_AMC@NLO` 2.7.3 MC event generator. The templates for the $q_W \cdot \cos\theta_{\ell W}^*$ and $\cos\theta_{\ell Z}^*$ distributions for each of the three individual helicity states of the W and Z bosons are therefore created from the inclusive PowHEG+PYTHIA MC sample using a reweighting technique [20]. For each of the gauge bosons the predicted helicity fractions of PowHEG+PYTHIA MC events are determined using the moment method [12,26] in the total phase space. Following the method detailed in Ref. [20], the predicted fractions are used to reweight PowHEG+PYTHIA MC events to represent each of the longitudinal, left- and right-handed states of the W or Z bosons. This creates polarised MC templates at detector level.

7.3. Differential cross section

Independently of polarisation fractions, the inclusive $W^\pm Z$ production cross section is measured differentially within the fiducial phase space defined at particle level and as a function of the variables the most sensitive to polarisation of the vector bosons. Similarly to measurements in Ref. [20], the differential detector-level distributions of background-subtracted data are corrected for detector resolution, efficiencies and QED final-state radiation effects using simulated signal events and an iterative Bayesian unfolding method [84], as implemented in the `RooUnfold` toolkit [85]. The number of iterations ranges from two to four, depending on the resolution in the unfolded variable and chosen in order to minimise the sum of the expected statistical and unfolding uncertainties in the unfolded distribution. The width of the bins in each distribution is chosen according to the experimental resolution and to the statistical significance of the expected number of events in each bin. The fraction of signal MC events generated in a bin which are reconstructed in the same bin is always greater than 35%. The detector-level distributions are unfolded with a response matrix computed using the PowHEG+PYTHIA MC signal sample that includes all four measured decay channels (eee , $ee\mu$, $\mu\mu e$ and $\mu\mu\mu$) and is divided by four, such that cross sections refer to final states where the W and Z bosons decay in a single leptonic channel with muons or electrons.

8. Systematic uncertainties

Systematic uncertainties affecting the shape and normalisation of the 4-category DNN score distribution for the background contributions, as well as the acceptance of the helicity components and the shape of their templates, are considered.

Uncertainties due to theoretical modelling in the helicity templates for higher order QCD corrections are evaluated by varying in the PowHEG+PYTHIA MC simulation the renormalisation and factorisation scales independently by factors of two and one-half, removing combinations where the variations differ by a factor of four. The uncertainties due to the PDF and the α_s value used in the PDF determination are evaluated using the PDF4LHC prescription [86].

For joint polarisation measurements an uncertainty due to the modelling of NLO QCD is estimated by comparing predictions of the 4-category DNN score distributions for each of the joint-helicity states obtained from the parton-level NLO calculation with the DNN reweighting of the PowHEG+PYTHIA MC simulation. The difference between the predicted shapes from the two methods is considered as an uncertainty. The associated nuisance parameter in the fit is parametrised by a flat constraint in the range between the two template shapes from each of the method, with steeply vanishing constraints outside of this range, defined using a double Fermi distribution. A specific uncertainty is attributed to the DNN reweighting method. It is estimated by applying the reweighting method to inclusive `MADGRAPH 0,1j@LO` MC events and comparing the obtained distributions with those of polarised `MADGRAPH 0,1j@LO` MC events. The difference between the shapes is treated as an uncertainty.

For individual boson polarisation measurements a modelling uncertainty in the helicity templates is estimated by comparing predictions of the $q_W \cdot \cos\theta_{\ell W}^*$ and $\cos\theta_{\ell Z}^*$ distributions from the PowHEG+PYTHIA event generator with those of SHERPA. The difference between the predicted shapes from the two generators is considered as an uncertainty. An uncertainty is attributed to the reweighting method used to create templates with individually polarised bosons from inclusive PowHEG+PYTHIA MC events. It is estimated in the same way as for the joint polarisation measurement.

Uncertainties in differential cross-section measurements arising from imperfect description of the data by the PowHEG+PYTHIA MC simulation are evaluated using a data-driven method [87]. Each MC differential distribution is corrected to match the data distribution and the resulting weighted MC distribution at detector level is unfolded with the response matrix used in the actual data unfolding. The new unfolded distribution is compared with the weighted MC distribution at particle level and the difference is taken as the systematic uncertainty. An additional uncertainty is derived to account for more subtle differences between the PowHEG+PYTHIA and `MADGRAPH5_AMC@NLO+PYTHIA` generators (e.g. hadronisation models, additional soft objects, mis-modelling in other kinematic variables). The `MADGRAPH5_AMC@NLO+PYTHIA` generator is used to unfold the data and deviation from the nominal result is taken as the uncertainty. In order to remove effects already accounted for in the data-driven method, the `MADGRAPH5_AMC@NLO+PYTHIA` distributions were first reweighed to match PowHEG+PYTHIA distributions at particle level.

Systematic uncertainties affecting the reconstruction and energy calibration of electrons, muons and jets are propagated through the analysis. The uncertainties due to lepton reconstruction, identification, isolation requirements and trigger efficiencies as well as in the lepton momentum scale and resolution are assessed using tag-and-probe methods in $Z \rightarrow \ell\ell$ events [60–62]. The uncertainties in the jet energy scale and resolution are based on their respective measurements in data [88]. The uncertainty in E_T^{miss} is estimated by propagating the uncertainties in the transverse momenta of reconstructed objects and by applying momentum scale and resolution uncertainties to the track-based soft term [74]. A variation in the pile-up reweighting of MC events is included in order to cover the uncertainty in the ratio of the predicted and measured pp inelastic cross sections [89]. For the measurements of the W charge-dependent polarisation fractions, an uncertainty arising from the charge misidentification of electrons is also considered.

The uncertainty in the amount of background from misidentified leptons takes into account the limited number of events in the control regions as well as the difference in background composition between the control region used to determine the lepton misidentification rate and the control regions used to estimate the yield in the signal region. This results in an uncertainty of about 25% in the total misidentified-lepton background yield and in the shape of the differential distributions of the reducible background events.

Uncertainties due to the theoretical modelling of ZZ generated events are considered. They arise from higher-order QCD corrections and the PDFs and are evaluated in the same way as for $W^\pm Z$ events. The uncertainty due to irreducible background sources other than




Fig. 1. Post-fit distribution of the 4-category DNN score in $W^\pm Z$ events at detector level in the signal region (SR). The 4-category DNN score is defined for each event as the sum of the DNN score and of the category number ranging from 0 to 3. Respective contributions of the $W^\pm Z$ joint-helicity states 00, 0T, T0 and TT and backgrounds are normalised to the expected number of events after the fit. The sum of background events containing misidentified leptons is labelled ‘Misid. leptons’. The sum of VVV , $WZjj$ –EW and tZj background events is labelled ‘Others’. The uncertainty band around the expectation includes all systematic uncertainties as obtained from the fit.

Table 1

Expected and observed numbers of events in the $W^\pm Z$ signal region (left) and in the ZZ control region (right). Numbers are presented before and after a fit to the 4-category DNN score distribution in $W^\pm Z$ events. The sum of background events containing misidentified leptons is labelled ‘Misid. leptons’. The sum of VVV , $WZjj$ –EW and tZj background events is labelled ‘Others’. The pre-fit uncertainties quoted are of statistical and systematic origin but do not take into account correlations between parameters of the fit. The post-fit uncertainties quoted are of statistical and systematic origin and take into account the correlations between the various parameters of the fit. For the $W^\pm Z$ signal templates, the correlations arising from the relationship $f_{00} + f_{0T} + f_{T0} + f_{TT} = 1$ explain the larger post-fit uncertainties compared to the pre-fit ones.

	Signal Region		ZZ Control Region	
	Pre-fit	Post-fit	Pre-fit	Post-fit
WZ in τ	620 ± 60	630 ± 60	WZ unpol.	35.6 ± 1.9
ZZ	1420 ± 120	1630 ± 50	ZZ	2030 ± 150
$t\bar{t} + V$	870 ± 130	820 ± 120	$t\bar{t} + V$	153 ± 23
Misid. leptons	1170 ± 230	1010 ± 220	Misid. leptons	14 ± 4
Others	800 ± 90	780 ± 90	Others	32 ± 8
W_0Z_0	920 ± 40	1190 ± 160	Total MC	2260 ± 150
W_0Z_T	2670 ± 50	1900 ± 500	Data	—
W_TZ_0	2670 ± 60	3100 ± 400		2554
W_TZ_T	10200 ± 230	10900 ± 600		
Total MC	21400 ± 500	21950 ± 170		
Data	—	21936		

ZZ is evaluated by propagating the uncertainty in their MC cross sections. These are 30% for VVV [90], 25% for $WZjj$ –EW [91] and 15% for tZj [92] and $t\bar{t} + V$ [79,93]. The contribution of events from $W^\pm Z$ production with at least one τ -lepton decay is linked in the fit to the measured $W^\pm Z$ production cross section for electron and muon decays. A normalisation uncertainty of 10% is attributed to this contribution to account for a possible mis-modelling in the MC simulation of the efficiency to select $W^\pm Z$ events with a τ -lepton decay.

The uncertainty in the combined 2015–2018 integrated luminosity is 1.7% [94], obtained using the LUCID-2 detector [95] for the primary luminosity measurements. It is applied to the signal normalisation as well as to all background contributions that are estimated using only MC simulations.

The effect of the systematic uncertainties on the final results after the maximum-likelihood fit is shown in Table 3 where the breakdown of the contributions to the uncertainties in the measured joint polarisation fractions is presented.

9. Results

9.1. Joint boson polarisation measurements

The distribution of the 4-category DNN score in the $W^\pm Z$ signal region with background normalisations, signal normalisation and nuisance parameters adjusted by the profile-likelihood fit is shown in Fig. 1. The corresponding pre-fit and post-fit yields in the signal region and in the ZZ control region are detailed in Table 1.

The measurements of f_{00} , f_{0T} , f_{T0} and f_{TT} are summarised in Table 2. The hypotheses of having no events in each of the joint helicity states are tested. In $W^\pm Z$ events, the presence of a pair of W and Z bosons with a simultaneous longitudinal polarisation (f_{00}) is observed with a significance of 7.1σ , compared to 6.2σ expected. The other joint helicity fractions f_{0T} , f_{T0} and f_{TT} are also

Table 2

Measured joint helicity fractions f_{00} , f_{0T} , f_{T0} and f_{TT} in the fiducial phase space, for $W^\pm Z$, W^+Z and W^-Z events. The total uncertainties in the measurements are reported. The measurements are compared with predictions from Powheg+Pythia and from NLO QCD fixed-order calculations [23]. The uncertainties in the Powheg+Pythia prediction include statistical, PDF and QCD scale uncertainties; the uncertainties in the NLO QCD fixed-order prediction include QCD scale uncertainties.

	Data	Powheg+Pythia	NLO QCD
$W^\pm Z$			
f_{00}	0.067 ± 0.010	0.0590 ± 0.0009	0.058 ± 0.002
f_{0T}	0.110 ± 0.029	0.1515 ± 0.0017	0.159 ± 0.003
f_{T0}	0.179 ± 0.023	0.1465 ± 0.0017	0.149 ± 0.003
f_{TT}	0.644 ± 0.032	0.6431 ± 0.0021	0.628 ± 0.004
W^+Z			
f_{00}	0.072 ± 0.016	0.0583 ± 0.0012	0.057 ± 0.002
f_{0T}	0.119 ± 0.034	0.1484 ± 0.0022	0.155 ± 0.003
f_{T0}	0.152 ± 0.033	0.1461 ± 0.0022	0.147 ± 0.003
f_{TT}	0.66 ± 0.04	0.6472 ± 0.0026	0.635 ± 0.004
W^-Z			
f_{00}	0.063 ± 0.016	0.0600 ± 0.0014	0.059 ± 0.002
f_{0T}	0.11 ± 0.04	0.1560 ± 0.0027	0.166 ± 0.003
f_{T0}	0.21 ± 0.04	0.1470 ± 0.0027	0.152 ± 0.003
f_{TT}	0.62 ± 0.05	0.6370 ± 0.0033	0.618 ± 0.004

Table 3

Summary of the relative uncertainties in the joint helicity fractions f_{00} , f_{0T} , f_{T0} , and f_{TT} measured in $W^\pm Z$ events. The uncertainties are reported as percentages.

	f_{00}	f_{0T}	f_{T0}	f_{TT}
Relative uncertainty [%]				
e energy scale and id. efficiency	0.34	0.6	0.8	0.31
μ energy scale and id. efficiency	0.8	0.23	0.23	0.13
E_T^{miss} and jets	3.3	1.3	1.2	0.4
Pile-up	0.6	0.17	0.4	0.15
Misidentified lepton background	2.3	1.6	0.8	0.26
$Z Z$ background	0.9	0.17	0.32	0.07
Other backgrounds	3.0	1.6	1.3	0.4
Parton Distribution Function	0.5	1.8	0.09	0.5
QCD scale	0.19	8	0.9	2.0
Modelling	9	4	2.9	1.2
Total systematic uncertainty	14	15	8	4
Luminosity	0.35	0.24	0.15	0.05
Statistical uncertainty	13	10	12	3.0
Total	19	18	14	5

measured with observed (expected) significances of 3.4σ (5.4σ), 7.1σ (6.6σ) and 11σ (9.7σ), respectively. The joint helicity fractions are also measured in W^+Z and W^-Z events separately, with observed significances for f_{00} of 6.9σ and 4.1σ , respectively. Table 3 shows the main sources of uncertainty in the measurement of the joint helicity fractions. Uncertainties related to data statistics and to QCD effects, namely the QCD scale and modelling uncertainties, make similar contributions to the precision of the measurements. The modelling uncertainty is dominated by the uncertainties in the shape of the distributions of polarised $W^\pm Z$ events due to NLO QCD corrections. Among them, the uncertainty from the DNN reweighting method is dominant, except for f_{00} where the uncertainty from the template generation method is of equal size. Different correlation schemes between bins and between helicity states are tested for this latter modelling uncertainty. All yield consistent results. Uncertainties in the energy scale and reconstruction of jets are dominant among detector uncertainties, showing how observables for joint polarisation are linked to jet emissions. The values of f_{00} , f_{0T} , f_{T0} and f_{TT} measured in $W^\pm Z$ events are shown in Fig. 2. The global level of agreement between the four measured joint helicity fractions with the fixed-order predictions at NLO QCD [23] is observed to be 1.4σ . A similar level of agreement is observed with respect to Powheg+Pythia predictions. The impact of EW corrections on the joint helicity fractions is negligible, smaller than 3% [32].

9.2. Individual boson polarisation measurements

The distributions of $q_W \cdot \cos\theta_{\ell^W}^*$ and $\cos\theta_{\ell^Z}^*$ with background normalisations, signal normalisation and nuisance parameters, adjusted by the profile-likelihood fit, are shown in Fig. 3. The post-fit event yields in the $W^\pm Z$ signal region and in the $Z Z$ control region are compatible with those obtained in the fit to the 4-category DNN score distribution.

The measurements of f_0 and $f_L - f_R$ are summarised in Table 4 where they are compared with the predictions from Powheg+Pythia and, for f_0 , from an NLO QCD fixed-order calculation [23]. Good agreement of the measured helicity fractions of both the W and Z bosons with the predictions from Powheg+Pythia is observed. Measured f_0 values agree within 1σ with the prediction, while $f_L - f_R$ values agree within 1.5σ , except for W^-Z events where the level of agreement deteriorates to 2.3σ . Table 5 shows the main sources of uncertainty in the measurement of the individual helicity fractions. The measurements are dominated by statistical uncertainties, except




Fig. 2. Measured joint helicity fractions f_{00} , f_{0T} , f_{T0} and f_{TT} of the W and Z bosons in $W^\pm Z$ events, compared with NLO QCD fixed-order predictions [23] (upward pointing triangle) and to MC predictions from Powheg+Pythia (downward pointing triangle). The effect of PDF and QCD scale uncertainties on the Powheg+Pythia and NLO QCD fixed-order predictions is of the same size as the respective markers. The full and dashed ellipses around the data points correspond to one and two standard deviations, respectively.




Fig. 3. Post-fit distributions of (a) $q_W \cdot \cos \theta_{\ell^*}^{*W}$ and (b) $\cos \theta_{\ell^*}^{*Z}$ in $W^\pm Z$ events at detector level in the signal region (SR). Respective contributions of the $W^\pm Z$ individual helicity states 0, L and R of the W and Z bosons and backgrounds are normalised to the expected number of events after the fit. The sum of background events containing misidentified leptons is labelled ‘Misid. leptons’. The sum of VVV , $WZjj$ –EW and tZj background events is labelled ‘Others’. The uncertainty band around the expectation includes all systematic uncertainties as obtained from the fit.

for the f_0 measurement of the W boson where the modelling uncertainty is equally important. The values of f_0 and $f_L - f_R$ measured in $W^\pm Z$ events are shown in Fig. 4 for the W and Z bosons.

The compatibility of the measured joint and individual helicity fractions is tested. Ignoring the interference between polarisations, which should represent only 0.6%–0.8% of the events according to NLO QCD and NLO QCD+EW fixed-order predictions [23,32], the following relationships ensue

$$f_{0T} = f_0^W - f_{00}, \quad f_{T0} = f_0^Z - f_{00} \quad \text{and} \quad f_{TT} = 1 + f_{00} - f_0^W - f_0^Z.$$

Using these relationships, joint f_{00} and individual f_0^W , f_0^Z helicity fractions can be measured simultaneously from a fit to the 4-category DNN score observable. The fractions f_0^W and f_0^Z measured in this procedure agree within less than 1σ of their own un-

Table 4

Individual helicity fractions f_0 and $f_L - f_R$ of the W and Z bosons measured in the fiducial phase space for $W^\pm Z$, W^+Z and W^-Z events. The total uncertainties in the measurements are reported. The measurements are compared with predictions from POWHEG+PYTHIA and, for f_0 , from NLO QCD fixed-order calculations [23]. The uncertainties in the POWHEG+PYTHIA prediction include statistical, PDF and QCD scale uncertainties; the uncertainties in the NLO QCD fixed-order prediction include QCD scale uncertainties.

f_0			$f_L - f_R$	
Data	POWHEG+PYTHIA	NLO QCD	Data	POWHEG+PYTHIA
W in W^+Z	0.23 ± 0.05	0.2044 ± 0.0024	0.211 ± 0.002	0.071 ± 0.023
W in W^-Z	0.19 ± 0.05	0.217 ± 0.004	0.225 ± 0.001	0.026 ± 0.027
W in $W^\pm Z$	0.21 ± 0.04	0.2094 ± 0.0016	0.217 ± 0.001	0.059 ± 0.016
Z in W^+Z	0.223 ± 0.025	0.1971 ± 0.0019	0.206 ± 0.002	-0.20 ± 0.10
Z in W^-Z	0.241 ± 0.029	0.2065 ± 0.0023	0.211 ± 0.001	0.10 ± 0.13
Z in $W^\pm Z$	0.231 ± 0.019	0.2009 ± 0.0014	0.208 ± 0.001	-0.10 ± 0.08

Table 5

Summary of the relative uncertainties in the polarisation fractions f_0 and $f_L - f_R$ measured in $W^\pm Z$ events for W and Z bosons. The uncertainties are reported as percentages.

	W^\pm in $W^\pm Z$		Z in $W^\pm Z$	
	f_0	$f_L - f_R$	f_0	$f_L - f_R$
Relative uncertainty [%]				
e energy scale and id. efficiency	1.4	0.8	1.3	0.7
μ energy scale and id. efficiency	2.1	5	0.8	0.5
E_T^{miss} and jets	1.9	2.8	0.28	3.0
Pile-up	1.4	4	1.2	3.1
Misidentified lepton background	3.4	0.8	1.6	1.2
ZZ background	0.7	0.6	0.6	2.5
Other backgrounds	0.9	1.3	0.7	1.3
Parton Distribution Function	0.5	2.9	0.05	0.5
QCD scale	6	6	0.22	5
Modelling	12	3.1	2.2	19
Total systematic uncertainty	14	11	3.5	21
Luminosity	0.25	0.09	0.06	0.19
Statistical uncertainty	13	40	9	90
Total	19	40	10	90




Fig. 4. Individual helicity fractions f_0 and $f_L - f_R$ measured for the (a) W and (b) Z bosons in $W^\pm Z$ events, compared with predictions from POWHEG+PYTHIA with $\sin^2 \theta_W = 0.23152$ (downward pointing triangle). The effect of PDF and QCD scale uncertainties on the POWHEG+PYTHIA prediction is of the same size as the triangle marker. The full and dashed ellipses around the data points correspond to one and two standard deviations, respectively.

certainties alone with the f_0^W and f_0^Z values obtained via individual $\cos \theta_{\ell W}^*$ and $\cos \theta_{\ell Z}^*$ distributions. This represents a consistency test of the joint helicity measurement procedure.

The spin-correlation between longitudinal W and Z bosons is tested by comparing joint and individual helicity fractions and measuring the ratio $R_c = f_{00}/(f_0^W f_0^Z)$. In the absence of spin correlations, neglecting interference effects, R_c is equal to 1. A value of $R_c = 1.3$ is predicted by NLO QCD fixed-order calculations, indicating the presence of correlation between polarisation states of the bosons. The ratio R_c is extracted from data by a simultaneous fit to the 4-category DNN score observable of f_{00} , f_0^W and f_0^Z . It is measured in $W^\pm Z$ events to be $R_c = 1.54 \pm 0.35$. The observed significance relative to the no spin-correlations hypothesis, i.e. $R_c = 1$, is 1.6σ .




Fig. 5. The measured $W^\pm Z$ differential cross section in the fiducial phase space as a function of (a) $q_W \cdot \cos \theta_{\ell^*W}$, (b) $\cos \theta_{\ell^*Z}^*$, (c) $|\cos \theta_V|$ and (d) DNN score. The total cross section integrated over each bin and divided by the bin width is represented. The inner and outer error bars on the data points represent the statistical and total uncertainties, respectively. The measurements are compared with the NLO prediction from POWHEG+PYTHIA (solid line). The dashed band shows how the sum of QCD scale and PDF uncertainties affects the POWHEG+PYTHIA NLO prediction. The predictions from the MADGRAPH 0,j@LO and MADGRAPH5_AMC@NLO+PYTHIA MC generators are also indicated by dotted-dashed and dashed lines, respectively. All predictions have been scaled to the NNLO QCD integrated cross section predicted by MATRIX.

9.3. Differential cross-section measurements

The inclusive cross section of $W^\pm Z$ production in the fiducial region for leptonic decay modes (electrons or muons) is determined simultaneously to the joint helicity fractions. It is measured to be $\sigma_{W^\pm Z \rightarrow \ell' v \ell \ell}^{\text{fid.}} = 64.6 \pm 0.5 \text{ (stat.)} \pm 1.8 \text{ (syst.)} \pm 1.1 \text{ (lumi.) fb}$, where the uncertainties correspond to statistical uncertainties, experimental and modelling systematic uncertainties, and luminosity uncertainties. The corresponding SM NNLO QCD prediction from MATRIX [96] is $64.0^{+1.5}_{-1.3}$ fb, where the uncertainty corresponds to the QCD scale uncertainty.

The $W^\pm Z$ production cross section is measured as a function of several variables sensitive to the individual and joint boson polarisation: $q_W \cdot \cos \theta_{\ell^*W}^*$, $\cos \theta_{\ell^*Z}^*$, $|\cos \theta_V|$ and the DNN score. The measured differential cross sections in Fig. 5 are compared with the Born-level predictions at NLO in QCD from the POWHEG+PYTHIA, MADGRAPH5_AMC@NLO+PYTHIA MC generators and to the sum of the polarised MADGRAPH 0,j@LO LO predictions. The Born-level DNN score prediction is evaluated using the same DNN as trained at detector level and evaluated using Born-level input observables. The predicted integrated fiducial cross sections of the three MC generators are rescaled to the NNLO cross section from MATRIX [96], as indicated in Fig. 5. Good agreement of the shapes of the measured distributions with the NLO QCD predictions is observed. The MADGRAPH 0,j@LO LO prediction exhibits larger differences from the measured cross section for the $|\cos \theta_V|$ distribution. This is expected as the $|\cos \theta_V|$ distribution is particularly sensitive to NLO QCD corrections [23] and the MAD-

GRAPH $0,1j@LO$ LO prediction only includes their real parts. The measured cross sections and the Rivet [97] routine for this measurement, including for the evaluation of the DNN classifier score, are published on HEPData [98].⁴

10. Conclusion

Measurements of gauge boson polarisation states from $W^\pm Z$ production in $\sqrt{s} = 13$ TeV pp collisions at the LHC are presented. The data analysed were collected with the ATLAS detector during the years 2015–2018 and correspond to an integrated luminosity of 139 fb^{-1} . The measurements use leptonic decay modes of the gauge bosons to electrons or muons and are performed in a fiducial phase space closely matching the detector acceptance.

Using neural-network-based advanced analysis techniques, joint helicity fractions of pair-produced vector bosons are measured. For the first time, the presence of a W boson and a Z boson with a simultaneous longitudinal polarisation is measured with observed and expected significances of 7.1 and 6.2 standard deviations, respectively. Integrated over the fiducial region, the fractions of the W and Z bosons both longitudinally or transversely polarised are measured in $W^\pm Z$ events to be $f_{00} = 0.067 \pm 0.010$ and $f_{TT} = 0.644 \pm 0.032$, respectively, in agreement with the SM predictions at NLO in QCD. The impact of EW corrections is negligible. The measured fractions of mixed polarisation modes are $f_{0T} = 0.110 \pm 0.029$ and $f_{T0} = 0.179 \pm 0.023$. Joint helicity fractions are also measured separately for W^+Z and W^-Z events.

Individual helicity fractions f_0 and $f_L - f_R$ of the W and Z bosons are measured. The measured values agree with the SM predictions and are consistent with the measured joint helicity fractions when neglecting interference among polarisation states, which from NLO QCD+EW calculations are expected to be smaller than the uncertainty of the present measurement.

Furthermore, the inclusive cross section of $W^\pm Z$ production in the fiducial region for leptonic decay modes is measured to be $\sigma_{W^\pm Z \rightarrow \ell' \nu \ell \ell}^{\text{fid.}} = 64.6 \pm 2.1\text{ fb}$, in agreement with the NNLO SM expectation of $64.0^{+1.5}_{-1.3}\text{ fb}$. The $W^\pm Z$ production cross section is measured as a function of several kinematic variables sensitive to polarisation and compared with SM predictions at NLO in QCD from the POWHEG+PYTHIA and MADGRAPH5_AMC@NLO+PYTHIA MC event generators. In particular, the variable related to the output of the deep neural network classifier used for the joint helicity measurement is unfolded in order to remove detector effects. The differential cross-section distributions are well described by the NLO MC predictions.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

The data for this manuscript are not available. The values in the plots and tables associated to this article are stored in HEPDATA (<https://hepdata.cedar.ac.uk>).

Acknowledgements

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; SSTD, Belarus; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benoziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; JINR; MES of Russia and NRC KI, Russian Federation; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DS/ NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, Canarie, Compute Canada and CRC, Canada; COST, ERC, ERDF, Horizon 2020 and Marie Skłodowska-Curie Actions, European Union; Investissements d'Avenir Labex, Investissements d'Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and GIF, Israel; Norwegian Financial Mechanism 2014–2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelser, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (UK) and BNL (USA), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in Ref. [99].

References

- [1] A. Azatov, J. Elias-Miro, Y. Reyimuaji, E. Venturini, Novel measurements of anomalous triple gauge couplings for the LHC, *J. High Energy Phys.* **10** (2017) 027, arXiv: 1707.08060 [hep-ph].
- [2] G. Panico, F. Riva, A. Wulzer, Diboson interference resurrection, *Phys. Lett. B* **776** (2018) 473, arXiv:1708.07823 [hep-ph].
- [3] CDF D0 Collaborations, Combination of CDF and D0 measurements of the W boson helicity in top quark decays, *Phys. Rev. D* **85** (2012) 071106, arXiv:1202.5272 [hep-ex].

⁴ In the HEPData entry, two sets of cross section measurements are provided, determined using both dressed and Born leptons, respectively.

- [4] CDF Collaboration, Measurement of W-boson polarization in top-quark decay in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys. Rev. Lett. 105 (2010) 042002, arXiv:1003.0224 [hep-ex].
- [5] D0 Collaboration, Measurement of the W boson helicity in top quark decays using 5.4 fb^{-1} of $p\bar{p}$ collision data, Phys. Rev. D 83 (2011) 032009, arXiv:1011.6549 [hep-ex].
- [6] ATLAS Collaboration, Measurement of the W boson polarization in top quark decays with the ATLAS detector, J. High Energy Phys. 06 (2012) 088, arXiv:1205.2484 [hep-ex].
- [7] CMS Collaboration, Measurement of the W boson helicity fractions in the decays of top quark pairs to lepton + jets final states produced in pp collisions at $\sqrt{s} = 8$ TeV, Phys. Lett. B 762 (2016) 512, arXiv:1605.09047 [hep-ex].
- [8] ATLAS Collaboration, Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment, Eur. Phys. J. C 72 (2012) 2001, arXiv:1203.2165 [hep-ex].
- [9] CMS Collaboration, Measurement of the polarization of W bosons with large transverse momenta in W+jets events at the LHC, Phys. Rev. Lett. 107 (2011) 021802, arXiv:1104.3829 [hep-ex].
- [10] CDF Collaboration, First measurement of the angular coefficients of Drell-Yan e^+e^- pairs in the Z mass region from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys. Rev. Lett. 106 (2011) 241801, arXiv:1103.5699 [hep-ex].
- [11] CMS Collaboration, Angular coefficients of Z bosons produced in pp collisions at $\sqrt{s} = 8$ TeV and decaying to $\mu^+\mu^-$ as a function of transverse momentum and rapidity, Phys. Lett. B 750 (2015) 154, arXiv:1504.03512 [hep-ex].
- [12] ATLAS Collaboration, Measurement of the angular coefficients in Z-boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector, J. High Energy Phys. 08 (2016) 159, arXiv:1606.00689 [hep-ex].
- [13] H1 Collaboration, Events with isolated leptons and missing transverse momentum and measurement of W production at HERA, Eur. Phys. J. C 64 (2009) 251, arXiv:0901.0488 [hep-ex].
- [14] L3 Collaboration, Measurement of W polarisation at LEP, Phys. Lett. B 557 (2003) 147, arXiv:hep-ex/0301027.
- [15] OPAL Collaboration, W boson polarization at LEP2, Phys. Lett. B 585 (2004) 223, arXiv:hep-ex/0312047.
- [16] DELPHI Collaboration, Study of W-boson polarisations and triple gauge boson couplings in the reaction $e^+e^- \rightarrow W^+W^-$ at LEP 2, Eur. Phys. J. C 54 (2008) 345, arXiv:0801.1235 [hep-ex].
- [17] OPAL Collaboration, Measurement of W boson polarizations and CP violating triple gauge couplings from W^+W^- production at LEP, Eur. Phys. J. C 19 (2001) 229, arXiv:hep-ex/0009021.
- [18] L3 Collaboration, Study of spin and decay-plane correlations of W bosons in the $e^+e^- \rightarrow W^+W^-$ process at LEP, Eur. Phys. J. C 40 (2005) 333, arXiv:hep-ex/0501036.
- [19] DELPHI Collaboration, Correlations between polarisation states of W particles in the reaction $e^-e^+ \rightarrow W^-W^+$ at LEP2 energies 189-GeV - 209-GeV, Eur. Phys. J. C 63 (2009) 611, arXiv:0908.1023 [hep-ex].
- [20] ATLAS Collaboration, Measurement of $W^\pm Z$ production cross sections and gauge boson polarisation in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Eur. Phys. J. C 79 (2019) 535, arXiv:1902.05759 [hep-ex].
- [21] CMS Collaboration, Measurement of the inclusive and differential WZ production cross sections, polarization angles, and triple gauge couplings in pp collisions at $\sqrt{s} = 13$ TeV, arXiv:2110.11231 [hep-ex], 2021.
- [22] A. Denner, G. Pelliccioli, Polarized electroweak bosons in W^+W^- production at the LHC including NLO QCD effects, J. High Energy Phys. 09 (2020) 164, arXiv:2206.14867 [hep-ph].
- [23] A. Denner, G. Pelliccioli, NLO QCD predictions for doubly-polarized WZ production at the LHC, Phys. Lett. B 814 (2020) 136107, arXiv:2010.07149 (We would like to thank Ansgar Denner and Giovanni Pelliccioli for fruitful discussions and for providing us with the NLO QCD predictions for the specific deep-neural network classifier output used in the analysis and for both W^+Z and W^-Z events).
- [24] U. Baur, T. Han, J. Ohnemus, Amplitude zeros in $W^{+-}Z$ production, Phys. Rev. Lett. 72 (1994) 3941, arXiv:hep-ph/9403248.
- [25] R. Franceschini, G. Panico, A. Pomarol, F. Riva, A. Wulzer, Electroweak precision tests in high-energy diboson processes, J. High Energy Phys. 02 (2018) 111, arXiv:1712.01310 [hep-ph].
- [26] E. Mirkes, J. Ohnemus, W and Z polarization effects in hadronic collisions, Phys. Rev. D 50 (1994) 5692, arXiv:hep-ph/9406381.
- [27] S. Groote, J.G. Korner, P. Tuvike, $O(\alpha_s)$ corrections to the decays of polarized W^\pm and Z bosons into massive quark pairs, Eur. Phys. J. C 72 (2012) 2177, arXiv:1204.5295 [hep-ph].
- [28] W.J. Stirling, E. Vryonidou, Electroweak gauge boson polarisation at the LHC, J. High Energy Phys. 07 (2012) 124, arXiv:1204.6427 [hep-ph].
- [29] G. Gounaris, J. Layssac, G. Moustaka, F.M. Renard, Analytic expressions of cross-sections, asymmetries and W density matrices for $e^-e^+ \rightarrow W^+W^-$ with general three boson couplings, Int. J. Mod. Phys. A 8 (1993) 3285.
- [30] Particle Data Group, Review of particle physics, Phys. Rev. D 98 (2018) 030001.
- [31] ATLAS Collaboration, Proposal for truth particle observable definitions in physics measurements, Tech. rep. ATL-PHYS-PUB-2015-013, CERN, 2015, <https://cds.cern.ch/record/2022743>.
- [32] D.N. Le, J. Baglio, Doubly-polarized WZ hadronic cross sections at NLO QCD+EW accuracy, arXiv:2203.01470 [hep-ph], 2022.
- [33] ATLAS Collaboration, The ATLAS experiment at the CERN large hadron collider, J. Instrum. 3 (2008) S08003.
- [34] ATLAS Collaboration, The ATLAS collaboration software and firmware, ATL-SOFT-PUB-2021-001, <https://cds.cern.ch/record/2767187>, 2021.
- [35] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, J. High Energy Phys. 11 (2004) 040, arXiv:hep-ph/0409146.
- [36] S. Frixione, P. Nason, C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method, J. High Energy Phys. 11 (2007) 070, arXiv:0709.2092 [hep-ph].
- [37] S. Alioli, P. Nason, C. Oleari, E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, J. High Energy Phys. 06 (2010) 043, arXiv:1002.2581 [hep-ph].
- [38] T. Melia, P. Nason, R. Röntsch, G. Zanderighi, W^+W^- , WZ and ZZ production in the POWHEG BOX, J. High Energy Phys. 11 (2011) 078, arXiv:1107.5051 [hep-ph].
- [39] T. Sjöstrand, et al., An introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159, arXiv:1410.3012 [hep-ph].
- [40] ATLAS Collaboration, Measurement of the Z/ γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, J. High Energy Phys. 09 (2014) 145, arXiv:1406.3660 [hep-ex].
- [41] H.-L. Lai, et al., New parton distributions for collider physics, Phys. Rev. D 82 (2010) 074024, arXiv:1007.2241 [hep-ph].
- [42] J. Pumpin, et al., New generation of parton distributions with uncertainties from global QCD analysis, J. High Energy Phys. 07 (2002) 012, arXiv:hep-ph/0201195.
- [43] J. Alwall, et al., The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, J. High Energy Phys. 07 (2014) 079, arXiv:1405.0301 [hep-ph].
- [44] R. Frederix, S. Frixione, Merging meets matching in MC@NLO, J. High Energy Phys. 12 (2012) 061, arXiv:1209.6215 [hep-ph].
- [45] R.D. Ball, et al., Parton distributions for the LHC Run II, J. High Energy Phys. 04 (2015) 040, arXiv:1410.8849 [hep-ph].
- [46] V. Hirschi, O. Mattelaer, Automated event generation for loop-induced processes, J. High Energy Phys. 10 (2015) 146, arXiv:1507.00020 [hep-ph].
- [47] E. Bothmann, et al., Event generation with Sherpa 2.2, SciPost Phys. 7 (2019) 034, arXiv:1905.09127 [hep-ph].
- [48] T. Gleisberg, S. Hoeche, Comix, a new matrix element generator, J. High Energy Phys. 12 (2008) 039, arXiv:0808.3674 [hep-ph].
- [49] F. Cascioli, P. Maierhöfer, S. Pozzorini, Scattering amplitudes with open loops, Phys. Rev. Lett. 108 (2012) 111601, arXiv:1111.5206 [hep-ph].
- [50] S. Schumann, F. Krauss, A Parton shower algorithm based on Catani-Seymour dipole factorisation, J. High Energy Phys. 03 (2008) 038, arXiv:0709.1027 [hep-ph].
- [51] S. Hoeche, F. Krauss, M. Schonherr, F. Siegert, QCD matrix elements + parton showers: the NLO case, J. High Energy Phys. 04 (2013) 027, arXiv:1207.5030 [hep-ph].
- [52] D. Buarque Franzosi, O. Mattelaer, R. Ruiz, S. Shil, Automated predictions from polarized matrix elements, J. High Energy Phys. 04 (2020) 082, arXiv:1912.01725 [hep-ph].
- [53] L. Lönnblad, Correcting the Colour-Dipole Cascade Model with fixed order matrix elements, J. High Energy Phys. 05 (2002) 046, arXiv:hep-ph/0112284.
- [54] L. Lönnblad, S. Prestel, Matching tree-level matrix elements with interleaved showers, J. High Energy Phys. 03 (2012) 019, arXiv:1109.4829 [hep-ph].
- [55] T. Sjöstrand, S. Mrenna, P. Skands, A brief introduction to PYTHIA 8.1, Comput. Phys. Commun. 178 (2008) 852, arXiv:0710.3820 [hep-ph].
- [56] R.D. Ball, et al., Parton distributions with LHC data, Nucl. Phys. B 867 (2013) 244, arXiv:1207.1303 [hep-ph].
- [57] ATLAS Collaboration, The ATLAS simulation infrastructure, Eur. Phys. J. C 70 (2010) 823, arXiv:1005.4568 [physics.ins-det].
- [58] S. Agostinelli, et al., GEANT4 – a simulation toolkit, Nucl. Instrum. Methods Phys. Res., Sect. A 506 (2003) 250.

- [59] ATLAS Collaboration, The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie-Landshoff diffractive model, Tech. rep., CERN, 2016, <https://cds.cern.ch/record/2206965>.
- [60] ATLAS Collaboration, Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data, *J. Instrum.* **14** (2019) P12006, arXiv:1908.00005 [hep-ex].
- [61] ATLAS Collaboration, Muon reconstruction and identification efficiency in ATLAS using the full Run 2 pp collision data set at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **81** (2021) 578, arXiv:2012.00578 [hep-ex].
- [62] ATLAS Collaboration, Muon reconstruction performance of the ATLAS detector in proton–proton collision data at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **76** (2016) 292, arXiv:1603.05598 [hep-ex].
- [63] ATLAS Collaboration, Performance of the ATLAS trigger system in 2015, *Eur. Phys. J. C* **77** (2017) 317, arXiv:1611.09661 [hep-ex].
- [64] ATLAS Collaboration, Performance of electron and photon triggers in ATLAS during LHC Run 2, *Eur. Phys. J. C* **80** (2020) 47, arXiv:1909.00761 [hep-ex].
- [65] ATLAS Collaboration, Performance of the ATLAS muon triggers in Run 2, *J. Instrum.* **15** (2020) P09015, arXiv:2004.13447 [physics.ins-det].
- [66] ATLAS Collaboration, Jet reconstruction and performance using particle flow with the ATLAS Detector, *Eur. Phys. J. C* **77** (2017) 466, arXiv:1703.10485 [hep-ex].
- [67] M. Cacciari, G.P. Salam, G. Soyez, The anti- k_t jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063, arXiv:0802.1189 [hep-ph].
- [68] M. Cacciari, G.P. Salam, G. Soyez, Fastjet user manual, *Eur. Phys. J. C* **72** (2012) 1896, arXiv:1111.6097 [hep-ph].
- [69] ATLAS Collaboration, Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* **81** (2021) 689, arXiv:2007.02645 [hep-ex].
- [70] ATLAS Collaboration, Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector, *Eur. Phys. J. C* **76** (2016) 581, arXiv:1510.03823 [hep-ex].
- [71] ATLAS Collaboration, ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **79** (2019) 970, arXiv:1907.05120 [hep-ex].
- [72] ATLAS Collaboration, Identification of Jets Containing b -Hadrons with Recurrent Neural Networks at the ATLAS Experiment, Tech. rep. ATL-PHYS-PUB-2017-003, CERN, 2017, <https://cds.cern.ch/record/2255226>.
- [73] ATLAS Collaboration, Optimisation and performance studies of the ATLAS b -tagging algorithms for the 2017–18 LHC run, Tech. rep. ATL-PHYS-PUB-2017-013, CERN, 2017, <https://cds.cern.ch/record/2273281>.
- [74] ATLAS Collaboration, Performance of missing transverse momentum reconstruction with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **78** (2018) 903, arXiv:1802.08168 [hep-ex].
- [75] ATLAS Collaboration, Measurements of $W^\pm Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and limits on anomalous gauge boson self-couplings, *Phys. Rev. D* **93** (2016) 092004, arXiv:1603.02151 [hep-ex].
- [76] ATLAS Collaboration, Search for supersymmetry at $\sqrt{s}=8$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector, *J. High Energy Phys.* **06** (2014) 035, arXiv:1404.2500 [hep-ex].
- [77] ATLAS Collaboration, Tools for estimating fake/non-prompt lepton backgrounds with the ATLAS detector at the LHC, *J. Instrum.* (2022), <https://doi.org/10.48550/arXiv.2211.16178>, submitted for publication, arXiv:2211.16178 [hep-ex].
- [78] ATLAS Collaboration, $ZZ \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ cross-section measurements and search for anomalous triple gauge couplings in 13 TeV pp collisions with the ATLAS detector, *Phys. Rev. D* **97** (2018) 032005, arXiv:1709.07703 [hep-ex].
- [79] ATLAS Collaboration, Measurements of the inclusive and differential production cross sections of a top-quark–antiquark pair in association with a Z boson at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* **81** (2021) 737, arXiv:2103.12603 [hep-ex].
- [80] F. Chollet, et al., <https://github.com/fchollet/keras>, 2015.
- [81] Martin Abadi, et al., TensorFlow: large-scale machine learning on heterogeneous systems, Software available from tensorflow.org, arXiv:1603.04467, <https://www.tensorflow.org/>, 2015.
- [82] K. Cranmer, G. Lewis, L. Moneta, A. Shibata, W. Verkerke, HistFactory: a tool for creating statistical models for use with RooFit and RooStats, Tech. rep., New York, 2012, <https://cds.cern.ch/record/1456844>.
- [83] A. Andreassen, B. Nachman, Neural networks for full phase-space reweighting and parameter tuning, *Phys. Rev. D* **101** (2020) 091901, arXiv:1907.08209 [hep-ph].
- [84] G. D’Agostini, A multidimensional unfolding method based on Bayes’ theorem, *Nucl. Instrum. Methods Phys. Res., Sect. A* **362** (1995) 487.
- [85] T. Adye, Unfolding algorithms and tests using RooUnfold, p. 313, arXiv:1105.1160 [physics.data-an], 2011.
- [86] J. Butterworth, et al., PDF4LHC recommendations for LHC Run II, *J. Phys. G* **43** (2016) 023001, arXiv:1510.03865 [hep-ph].
- [87] B. Malaescu, An iterative, dynamically stabilized (IDS) method of data unfolding, in: Proceedings of the PHYSTAT 2011 Workshop, CERN, Geneva, Switzerland, 2011, p. 271, arXiv:1106.3107 [physics.data-an].
- [88] ATLAS Collaboration, Jet energy scale measurements and their systematic uncertainties in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **96** (2017) 072002, arXiv:1703.09665 [hep-ex].
- [89] ATLAS Collaboration, Measurement of the inelastic proton–proton cross section at $\sqrt{s} = 13$ TeV with the ATLAS detector at the LHC, *Phys. Rev. Lett.* **117** (2016) 182002, arXiv:1606.02625 [hep-ex].
- [90] ATLAS Collaboration, Multi-boson simulation for 13 TeV ATLAS analyses, ATL-PHYS-PUB-2016-002, <https://cds.cern.ch/record/2119986>, 2016.
- [91] ATLAS Collaboration, Observation of electroweak $W^\pm Z$ boson pair production in association with two jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Lett. B* **793** (2019) 469, arXiv:1812.09740 [hep-ex].
- [92] ATLAS Collaboration, Observation of the associated production of a top quark and a Z boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **07** (2020) 124, arXiv:2002.07546 [hep-ex].
- [93] ATLAS Collaboration, Modelling of the $t\bar{t}H$ and $t\bar{t}V$ ($V = W, Z$) processes for $\sqrt{s} = 13$ TeV ATLAS analyses, Tech. rep. ATL-PHYS-PUB-2016-005, CERN, 2016, <https://cds.cern.ch/record/2120826>.
- [94] ATLAS Collaboration, Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC, Tech. rep. ATLAS-CONF-2019-021, CERN, 2019, <https://cds.cern.ch/record/2677054>.
- [95] G. Avoni, et al., The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS, *J. Instrum.* **13** (2018) P07017.
- [96] M. Grazzini, S. Kallweit, D. Rathlev, M. Wiesemann, $W^\pm Z$ production at the LHC: fiducial cross sections and distributions in NNLO QCD, *J. High Energy Phys.* **05** (2017) 139, arXiv:1703.09065 [hep-ph].
- [97] C. Bierlich, et al., Robust independent validation of experiment and theory: Rivet version 3, *SciPost Phys.* **8** (2020) 026, arXiv:1912.05451 [hep-ph].
- [98] E. Maguire, L. Heinrich, G. Watt, HEPData: a repository for high energy physics data, in: R. Mount, C. Tull (Eds.), *J. Phys. Conf. Ser.* **898** (2017) 102006, arXiv:1704.05473 [hep-ex].
- [99] ATLAS Collaboration, ATLAS Computing Acknowledgements, ATL-SOFT-PUB-2021-003, <https://cds.cern.ch/record/2776662>, 2021.

The ATLAS Collaboration

G. Aad ¹⁰¹, B. Abbott ¹¹⁹, D.C. Abbott ¹⁰², K. Abeling ⁵⁵, S.H. Abidi ²⁹, A. Aboulhorma ^{35e}, H. Abramowicz ¹⁵⁰, H. Abreu ¹⁴⁹, Y. Abulaiti ¹¹⁶, A.C. Abusleme Hoffman ^{136a}, B.S. Acharya ^{68a,68b,p}, B. Achkar ⁵⁵, C. Adam Bourdarios ⁴, L. Adamczyk ^{84a}, L. Adamek ¹⁵⁴, S.V. Addepalli ²⁶, J. Adelman ¹¹⁴, A. Adiguzel ^{21c}, S. Adorni ⁵⁶, T. Adye ¹³³, A.A. Affolder ¹³⁵, Y. Afik ³⁶, M.N. Agaras ¹³, J. Agarwala ^{72a,72b}, A. Aggarwal ⁹⁹, C. Agheorghiesei ^{27c}, J.A. Aguilar-Saavedra ^{129f}, A. Ahmad ³⁶, F. Ahmadov ^{38,y}, W.S. Ahmed ¹⁰³, S. Ahuja ⁹⁴, X. Ai ⁴⁸, G. Aielli ^{75a,75b}, I. Aizenberg ¹⁶⁸, M. Akbiyik ⁹⁹, T.P.A. Åkesson ⁹⁷, A.V. Akimov ³⁷, K. Al Khoury ⁴¹, G.L. Alberghi ^{23b}, J. Albert ¹⁶⁴, P. Albicocco ⁵³, S. Alderweireldt ⁵²,

- M. Alekse³⁶, I.N. Aleksandrov³⁸, C. Alexa^{27b}, T. Alexopoulos¹⁰, A. Alfonsi¹¹³, F. Alfonsi^{23b}, M. Alhroob¹¹⁹, B. Ali¹³¹, S. Ali¹⁴⁷, M. Aliev³⁷, G. Alimonti^{70a}, W. Alkakhi⁵⁵, C. Allaire⁶⁶, B.M.M. Allbrooke¹⁴⁵, P.P. Allport²⁰, A. Aloisio^{71a,71b}, F. Alonso⁸⁹, C. Alpigiani¹³⁷, E. Alunno Camelia^{75a,75b}, M. Alvarez Estevez⁹⁸, M.G. Alviggi^{71a,71b}, M. Aly¹⁰⁰, Y. Amaral Coutinho^{81b}, A. Ambler¹⁰³, C. Amelung³⁶, M. Amerl¹, C.G. Ames¹⁰⁸, D. Amidei¹⁰⁵, S.P. Amor Dos Santos^{129a}, S. Amoroso⁴⁸, K.R. Amos¹⁶², V. Ananiev¹²⁴, C. Anastopoulos¹³⁸, T. Andeen¹¹, J.K. Anders³⁶, S.Y. Andrean^{47a,47b}, A. Andreazza^{70a,70b}, S. Angelidakis⁹, A. Angerami^{41,aa}, A.V. Anisenkov³⁷, A. Annovi^{73a}, C. Antel⁵⁶, M.T. Anthony¹³⁸, E. Antipov¹²⁰, M. Antonelli⁵³, D.J.A. Antrim^{17a}, F. Anulli^{74a}, M. Aoki⁸², T. Aoki¹⁵², J.A. Aparisi Pozo¹⁶², M.A. Aparo¹⁴⁵, L. Aperio Bella⁴⁸, C. Appelt¹⁸, N. Aranzabal³⁶, V. Araujo Ferraz^{81a}, C. Arcangeletti⁵³, A.T.H. Arce⁵¹, E. Arena⁹¹, J-F. Arguin¹⁰⁷, S. Argyropoulos⁵⁴, J.-H. Arling⁴⁸, A.J. Armbruster³⁶, O. Arnaez¹⁵⁴, H. Arnold¹¹³, Z.P. Arrubarrena Tame¹⁰⁸, G. Artoni^{74a,74b}, H. Asada¹¹⁰, K. Asai¹¹⁷, S. Asai¹⁵², N.A. Asbah⁶¹, J. Assahsah^{35d}, K. Assamagan²⁹, R. Astalos^{28a}, R.J. Atkin^{33a}, M. Atkinson¹⁶¹, N.B. Atlay¹⁸, H. Atmani^{62b}, P.A. Atmasiddha¹⁰⁵, K. Augsten¹³¹, S. Auricchio^{71a,71b}, A.D. Auriol²⁰, V.A. Austrup¹⁷⁰, G. Avner¹⁴⁹, G. Avolio³⁶, K. Axiotis⁵⁶, M.K. Ayoub^{14c}, G. Azuelos^{107,ad}, D. Babal^{28a}, H. Bachacou¹³⁴, K. Bachas^{151,r}, A. Bachiu³⁴, F. Backman^{47a,47b}, A. Badea⁶¹, P. Bagnaia^{74a,74b}, M. Bahmani¹⁸, A.J. Bailey¹⁶², V.R. Bailey¹⁶¹, J.T. Baines¹³³, C. Bakalis¹⁰, O.K. Baker¹⁷¹, P.J. Bakker¹¹³, E. Bakos¹⁵, D. Bakshi Gupta⁸, S. Balaji¹⁴⁶, R. Balasubramanian¹¹³, E.M. Baldin³⁷, P. Balek¹³², E. Ballabene^{70a,70b}, F. Balli¹³⁴, L.M. Baltes^{63a}, W.K. Balunas³², J. Balz⁹⁹, E. Banas⁸⁵, M. Bandieramonte¹²⁸, A. Bandyopadhyay²⁴, S. Bansal²⁴, L. Barak¹⁵⁰, E.L. Barberio¹⁰⁴, D. Barberis^{57b,57a}, M. Barbero¹⁰¹, G. Barbour⁹⁵, K.N. Barends^{33a}, T. Barillari¹⁰⁹, M-S. Barisits³⁶, T. Barklow¹⁴², R.M. Barnett^{17a}, P. Baron¹²¹, D.A. Baron Moreno¹⁰⁰, A. Baroncelli^{62a}, G. Barone²⁹, A.J. Barr¹²⁵, L. Barranco Navarro^{47a,47b}, F. Barreiro⁹⁸, J. Barreiro Guimarães da Costa^{14a}, U. Barron¹⁵⁰, M.G. Barros Teixeira^{129a}, S. Barsov³⁷, F. Bartels^{63a}, R. Bartoldus¹⁴², A.E. Barton⁹⁰, P. Bartos^{28a}, A. Basalaev⁴⁸, A. Basan⁹⁹, M. Baselga⁴⁹, I. Bashta^{76a,76b}, A. Bassalat^{66,b}, M.J. Basso¹⁵⁴, C.R. Basson¹⁰⁰, R.L. Bates⁵⁹, S. Batlamous^{35e}, J.R. Batley³², B. Batool¹⁴⁰, M. Battaglia¹³⁵, D. Battulga¹⁸, M. Bauce^{74a,74b}, P. Bauer²⁴, A. Bayirli^{21a}, J.B. Beacham⁵¹, T. Beau¹²⁶, P.H. Beauchemin¹⁵⁷, F. Becherer⁵⁴, P. Bechtle²⁴, H.P. Beck^{19,q}, K. Becker¹⁶⁶, A.J. Beddall^{21d}, V.A. Bednyakov³⁸, C.P. Bee¹⁴⁴, L.J. Beemster¹⁵, T.A. Beermann³⁶, M. Begalli^{81d}, M. Begel²⁹, A. Behera¹⁴⁴, J.K. Behr⁴⁸, C. Beirao Da Cruz E Silva³⁶, J.F. Beirer^{55,36}, F. Beisiegel²⁴, M. Belfkir¹⁵⁸, G. Bella¹⁵⁰, L. Bellagamba^{23b}, A. Bellerive³⁴, P. Bellos²⁰, K. Beloborodov³⁷, K. Belotskiy³⁷, N.L. Belyaev³⁷, D. Benchekroun^{35a}, F. Bendebba^{35a}, Y. Benhammou¹⁵⁰, D.P. Benjamin²⁹, M. Benoit²⁹, J.R. Bensinger²⁶, S. Bentvelsen¹¹³, L. Beresford³⁶, M. Beretta⁵³, D. Berge¹⁸, E. Bergeaas Kuutmann¹⁶⁰, N. Berger⁴, B. Bergmann¹³¹, J. Beringer^{17a}, S. Berlendis⁷, G. Bernardi⁵, C. Bernius¹⁴², F.U. Bernlochner²⁴, T. Berry⁹⁴, P. Berta¹³², A. Berthold⁵⁰, I.A. Bertram⁹⁰, S. Bethke¹⁰⁹, A. Betti^{74a,74b}, A.J. Bevan⁹³, M. Bhamjee^{33c}, S. Bhatta¹⁴⁴, D.S. Bhattacharya¹⁶⁵, P. Bhattarai²⁶, V.S. Bhopatkar¹²⁰, R. Bi^{29,ag}, R.M. Bianchi¹²⁸, O. Biebel¹⁰⁸, R. Bielski¹²², M. Biglietti^{76a}, T.R.V. Billoud¹³¹, M. Bindi⁵⁵, A. Bingul^{21b}, C. Bini^{74a,74b}, S. Biondi^{23b,23a}, A. Biondini⁹¹, C.J. Birch-sykes¹⁰⁰, G.A. Bird^{20,133}, M. Birman¹⁶⁸, T. Bisanz³⁶, E. Bisceglie^{43b,43a}, D. Biswas^{169,l}, A. Bitadze¹⁰⁰, K. Bjørke¹²⁴, I. Bloch⁴⁸, C. Blocker²⁶, A. Blue⁵⁹, U. Blumenschein⁹³, J. Blumenthal⁹⁹, G.J. Bobbink¹¹³, V.S. Bobrovnikov³⁷, M. Boehler⁵⁴, D. Bogavac³⁶, A.G. Bogdanchikov³⁷, C. Bohm^{47a}, V. Boisvert⁹⁴, P. Bokan⁴⁸, T. Bold^{84a}, M. Bomben⁵, M. Bona⁹³, M. Boonekamp¹³⁴, C.D. Booth⁹⁴, A.G. Borbély⁵⁹, H.M. Borecka-Bielska¹⁰⁷, L.S. Borgna⁹⁵, G. Borissov⁹⁰, D. Bortoletto¹²⁵, D. Boscherini^{23b}, M. Bosman¹³, J.D. Bossio Sola³⁶, K. Bouaouda^{35a}, N. Bouchhar¹⁶², J. Boudreau¹²⁸, E.V. Bouhova-Thacker⁹⁰, D. Boumediene⁴⁰, R. Bouquet⁵, A. Boveia¹¹⁸, J. Boyd³⁶, D. Boye²⁹, I.R. Boyko³⁸, J. Bracinik²⁰, N. Brahimi^{62d}, G. Brandt¹⁷⁰, O. Brandt³², F. Braren⁴⁸, B. Brau¹⁰², J.E. Brau¹²², K. Brendlinger⁴⁸, R. Brener¹⁶⁸, L. Brenner¹¹³, R. Brenner¹⁶⁰, S. Bressler¹⁶⁸, B. Brickwedde⁹⁹, D. Britton⁵⁹, D. Britzger¹⁰⁹, I. Brock²⁴, G. Brooijmans⁴¹, W.K. Brooks^{136f}, E. Brost²⁹, T.L. Bruckler¹²⁵, P.A. Bruckman de Renstrom⁸⁵, B. Brüers⁴⁸, D. Bruncko^{28b,*}, A. Bruni^{23b}, G. Bruni^{23b}, M. Bruschi^{23b}, N. Bruscino^{74a,74b}, L. Bryngemark¹⁴², T. Buanes¹⁶, Q. Buat¹³⁷, P. Buchholz¹⁴⁰, A.G. Buckley⁵⁹, I.A. Budagov^{38,*}, M.K. Bugge¹²⁴, O. Bulekov³⁷, B.A. Bullard⁶¹, S. Burdin⁹¹, C.D. Burgard⁴⁸, A.M. Burger⁴⁰, B. Burghgrave⁸, J.T.P. Burr³², C.D. Burton¹¹, J.C. Burzynski¹⁴¹, E.L. Busch⁴¹, V. Büscher⁹⁹, P.J. Bussey⁵⁹, J.M. Butler²⁵, C.M. Buttar⁵⁹, J.M. Butterworth⁹⁵, W. Buttlinger¹³³, C.J. Buxo Vazquez¹⁰⁶, A.R. Buzykaev³⁷, G. Cabras^{23b}, S. Cabrera Urbán¹⁶², D. Caforio⁵⁸, H. Cai¹²⁸, Y. Cai^{14a,14d}, V.M.M. Cairo³⁶, O. Cakir^{3a}, N. Calace³⁶, P. Calafiura^{17a}, G. Calderini¹²⁶, P. Calfayan⁶⁷, G. Callea⁵⁹,

- L.P. Caloba ^{81b}, D. Calvet ⁴⁰, S. Calvet ⁴⁰, T.P. Calvet ¹⁰¹, M. Calvetti ^{73a,73b}, R. Camacho Toro ¹²⁶,
 S. Camarda ³⁶, D. Camarero Munoz ²⁶, P. Camarri ^{75a,75b}, M.T. Camerlingo ^{76a,76b}, D. Cameron ¹²⁴,
 C. Camincher ¹⁶⁴, M. Campanelli ⁹⁵, A. Camplani ⁴², V. Canale ^{71a,71b}, A. Canesse ¹⁰³, M. Cano Bret ⁷⁹,
 J. Cantero ¹⁶², Y. Cao ¹⁶¹, F. Capocasa ²⁶, M. Capua ^{43b,43a}, A. Carbone ^{70a,70b}, R. Cardarelli ^{75a},
 J.C.J. Cardenas ⁸, F. Cardillo ¹⁶², T. Carli ³⁶, G. Carlino ^{71a}, J.I. Carlotto ¹³, B.T. Carlson ^{128,s},
 E.M. Carlson ^{164,155a}, L. Carminati ^{70a,70b}, M. Carnesale ^{74a,74b}, S. Caron ¹¹², E. Carquin ^{136f}, S. Carrá ^{70a,70b},
 G. Carratta ^{23b,23a}, F. Carrio Argos ^{33g}, J.W.S. Carter ¹⁵⁴, T.M. Carter ⁵², M.P. Casado ^{13,i}, A.F. Casha ¹⁵⁴,
 E.G. Castiglia ¹⁷¹, F.L. Castillo ^{63a}, L. Castillo Garcia ¹³, V. Castillo Gimenez ¹⁶², N.F. Castro ^{129a,129e},
 A. Catinaccio ³⁶, J.R. Catmore ¹²⁴, V. Cavaliere ²⁹, N. Cavalli ^{23b,23a}, V. Cavasinni ^{73a,73b}, E. Celebi ^{21a},
 F. Celli ¹²⁵, M.S. Centonze ^{69a,69b}, K. Cerny ¹²¹, A.S. Cerqueira ^{81a}, A. Cerri ¹⁴⁵, L. Cerrito ^{75a,75b},
 F. Cerutti ^{17a}, A. Cervelli ^{23b}, S.A. Cetin ^{21d}, Z. Chadi ^{35a}, D. Chakraborty ¹¹⁴, M. Chala ^{129f}, J. Chan ¹⁶⁹,
 W.Y. Chan ¹⁵², J.D. Chapman ³², B. Chargeishvili ^{148b}, D.G. Charlton ²⁰, T.P. Charman ⁹³, M. Chatterjee ¹⁹,
 S. Chekanov ⁶, S.V. Chekulaev ^{155a}, G.A. Chelkov ^{38,a}, A. Chen ¹⁰⁵, B. Chen ¹⁵⁰, B. Chen ¹⁶⁴, H. Chen ^{14c},
 H. Chen ²⁹, J. Chen ^{62c}, J. Chen ²⁶, S. Chen ¹⁵², S.J. Chen ^{14c}, X. Chen ^{62c}, X. Chen ^{14b,ac}, Y. Chen ^{62a},
 C.L. Cheng ¹⁶⁹, H.C. Cheng ^{64a}, S. Cheong ¹⁴², A. Cheplakov ³⁸, E. Cheremushkina ⁴⁸, E. Cherepanova ¹¹³,
 R. Cherkaoui El Moursli ^{35e}, E. Cheu ⁷, K. Cheung ⁶⁵, L. Chevalier ¹³⁴, V. Chiarella ⁵³, G. Chiarelli ^{73a},
 N. Chiedde ¹⁰¹, G. Chiodini ^{69a}, A.S. Chisholm ²⁰, A. Chitan ^{27b}, M. Chitishvili ¹⁶², Y.H. Chiu ¹⁶⁴,
 M.V. Chizhov ³⁸, K. Choi ¹¹, A.R. Chomont ^{74a,74b}, Y. Chou ¹⁰², E.Y.S. Chow ¹¹³, T. Chowdhury ^{33g},
 L.D. Christopher ^{33g}, K.L. Chu ^{64a}, M.C. Chu ^{64a}, X. Chu ^{14a,14d}, J. Chudoba ¹³⁰, J.J. Chwastowski ⁸⁵,
 D. Cieri ¹⁰⁹, K.M. Ciesla ^{84a}, V. Cindro ⁹², A. Ciocio ^{17a}, F. Cirotto ^{71a,71b}, Z.H. Citron ^{168,m}, M. Citterio ^{70a},
 D.A. Ciubotaru ^{27b}, B.M. Ciungu ¹⁵⁴, A. Clark ⁵⁶, P.J. Clark ⁵², J.M. Clavijo Columbie ⁴⁸, S.E. Clawson ¹⁰⁰,
 C. Clement ^{47a,47b}, J. Clercx ⁴⁸, L. Clissa ^{23b,23a}, Y. Coadou ¹⁰¹, M. Cobal ^{68a,68c}, A. Coccaro ^{57b},
 R.F. Coelho Barrue ^{129a}, R. Coelho Lopes De Sa ¹⁰², S. Coelli ^{70a}, H. Cohen ¹⁵⁰, A.E.C. Coimbra ^{70a,70b},
 B. Cole ⁴¹, J. Collot ⁶⁰, P. Conde Muiño ^{129a,129g}, M.P. Connell ^{33c}, S.H. Connell ^{33c}, I.A. Connelly ⁵⁹,
 E.I. Conroy ¹²⁵, F. Conventi ^{71a,ae}, H.G. Cooke ²⁰, A.M. Cooper-Sarkar ¹²⁵, F. Cormier ¹⁶³, L.D. Corpe ³⁶,
 M. Corradi ^{74a,74b}, E.E. Corrigan ⁹⁷, F. Corriveau ^{103,w}, A. Cortes-Gonzalez ¹⁸, M.J. Costa ¹⁶², F. Costanza ⁴,
 D. Costanzo ¹³⁸, B.M. Cote ¹¹⁸, G. Cowan ⁹⁴, J.W. Cowley ³², K. Cranmer ¹¹⁶, S. Crépé-Renaudin ⁶⁰,
 F. Crescioli ¹²⁶, M. Cristinziani ¹⁴⁰, M. Cristoforetti ^{77a,77b,d}, V. Croft ¹⁵⁷, G. Crosetti ^{43b,43a}, A. Cueto ³⁶,
 T. Cuhadar Donszelmann ¹⁵⁹, H. Cui ^{14a,14d}, Z. Cui ⁷, A.R. Cukierman ¹⁴², W.R. Cunningham ⁵⁹,
 F. Curcio ^{43b,43a}, P. Czodrowski ³⁶, M.M. Czurylo ^{63b}, M.J. Da Cunha Sargedas De Sousa ^{62a},
 J.V. Da Fonseca Pinto ^{81b}, C. Da Via ¹⁰⁰, W. Dabrowski ^{84a}, T. Dado ⁴⁹, S. Dahbi ^{33g}, T. Dai ¹⁰⁵,
 C. Dallapiccola ¹⁰², M. Dam ⁴², G. D'amen ²⁹, V. D'Amico ¹⁰⁸, J. Damp ⁹⁹, J.R. Dandoy ¹²⁷, M.F. Daneri ³⁰,
 M. Danner ¹⁴¹, V. Dao ³⁶, G. Darbo ^{57b}, S. Darmora ⁶, S.J. Das ²⁹, S. D'Auria ^{70a,70b}, C. David ^{155b},
 T. Davidek ¹³², D.R. Davis ⁵¹, B. Davis-Purcell ³⁴, I. Dawson ⁹³, K. De ⁸, R. De Asmundis ^{71a}, M. De Beurs ¹¹³,
 N. De Biase ⁴⁸, S. De Castro ^{23b,23a}, N. De Groot ¹¹², P. de Jong ¹¹³, H. De la Torre ¹⁰⁶, A. De Maria ^{14c},
 A. De Salvo ^{74a}, U. De Sanctis ^{75a,75b}, A. De Santo ¹⁴⁵, J.B. De Vivie De Regie ⁶⁰, D.V. Dedovich ³⁸,
 J. Degens ¹¹³, A.M. Deiana ⁴⁴, F. Del Corso ^{23b,23a}, J. Del Peso ⁹⁸, F. Del Rio ^{63a}, F. Deliot ¹³⁴,
 C.M. Delitzsch ⁴⁹, M. Della Pietra ^{71a,71b}, D. Della Volpe ⁵⁶, A. Dell'Acqua ³⁶, L. Dell'Asta ^{70a,70b},
 M. Delmastro ⁴, P.A. Delsart ⁶⁰, S. Demers ¹⁷¹, M. Demichev ³⁸, S.P. Denisov ³⁷, L. D'Eramo ¹¹⁴,
 D. Derendarz ⁸⁵, F. Derue ¹²⁶, P. Dervan ⁹¹, K. Desch ²⁴, K. Dette ¹⁵⁴, C. Deutsch ²⁴, P.O. Deviveiros ³⁶,
 F.A. Di Bello ^{57b,57a}, A. Di Ciaccio ^{75a,75b}, L. Di Ciaccio ⁴, A. Di Domenico ^{74a,74b}, C. Di Donato ^{71a,71b},
 A. Di Girolamo ³⁶, G. Di Gregorio ⁵, A. Di Luca ^{77a,77b}, B. Di Micco ^{76a,76b}, R. Di Nardo ^{76a,76b},
 C. Diaconu ¹⁰¹, F.A. Dias ¹¹³, T. Dias Do Vale ¹⁴¹, M.A. Diaz ^{136a,136b}, F.G. Diaz Capriles ²⁴, M. Didenko ¹⁶²,
 E.B. Diehl ¹⁰⁵, L. Diehl ⁵⁴, S. Díez Cornell ⁴⁸, C. Diez Pardos ¹⁴⁰, C. Dimitriadi ^{24,160}, A. Dimitrieva ^{17a},
 W. Ding ^{14b}, J. Dingfelder ²⁴, I.-M. Dinu ^{27b}, S.J. Dittmeier ^{63b}, F. Dittus ³⁶, F. Djama ¹⁰¹, T. Djobava ^{148b},
 J.I. Djuvsland ¹⁶, C. Doglioni ^{100,97}, J. Dolejsi ¹³², Z. Dolezal ¹³², M. Donadelli ^{81c}, B. Dong ^{62c}, J. Donini ⁴⁰,
 A. D'Onofrio ^{14c}, M. D'Onofrio ⁹¹, J. Dopke ¹³³, A. Doria ^{71a}, M.T. Dova ⁸⁹, A.T. Doyle ⁵⁹, M.A. Draguet ¹²⁵,
 E. Drechsler ¹⁴¹, E. Dreyer ¹⁶⁸, I. Drivas-koulouris ¹⁰, A.S. Drobac ¹⁵⁷, M. Drozdova ⁵⁶, D. Du ^{62a},
 T.A. du Pree ¹¹³, F. Dubinin ³⁷, M. Dubovsky ^{28a}, E. Duchovni ¹⁶⁸, G. Duckeck ¹⁰⁸, O.A. Ducu ^{27b},
 D. Duda ¹⁰⁹, A. Dudarev ³⁶, M. D'uffizi ¹⁰⁰, L. Duflot ⁶⁶, M. Dührssen ³⁶, C. Dülsen ¹⁷⁰, A.E. Dumitriu ^{27b},
 M. Dunford ^{63a}, S. Dungs ⁴⁹, K. Dunne ^{47a,47b}, A. Duperrin ¹⁰¹, H. Duran Yildiz ^{3a}, M. Düren ⁵⁸,
 A. Durglishvili ^{148b}, B.L. Dwyer ¹¹⁴, G.I. Dyckes ^{17a}, M. Dyndal ^{84a}, S. Dysch ¹⁰⁰, B.S. Dziedzic ⁸⁵,
 Z.O. Earnshaw ¹⁴⁵, B. Eckerova ^{28a}, M.G. Eggleston ⁵¹, E. Egido Purcino De Souza ^{81b}, L.F. Ehrke ⁵⁶,

- G. Eigen 16, K. Einsweiler 17a, T. Ekelof 160, P.A. Ekman 97, Y. El Ghazali 35b, H. El Jarrari 35e, 147,
 A. El Moussaouy 35a, V. Ellajosyula 160, M. Ellert 160, F. Ellinghaus 170, A.A. Elliot 93, N. Ellis 36,
 J. Elmsheuser 29, M. Elsing 36, D. Emeliyanov 133, A. Emerman 41, Y. Enari 152, I. Ene 17a, S. Epari 13,
 J. Erdmann 49,ab, A. Ereditato 19, P.A. Erland 85, M. Errenst 170, M. Escalier 66, C. Escobar 162, E. Etzion 150,
 G. Evans 129a, H. Evans 67, M.O. Evans 145, A. Ezhilov 37, S. Ezarqtouni 35a, F. Fabbri 59, L. Fabbri 23b,23a,
 G. Facini 95, V. Fadeev 135, R.M. Fakhrutdinov 37, S. Falciano 74a, P.J. Falke 24, S. Falke 36, J. Faltova 132,
 Y. Fan 14a, Y. Fang 14a,14d, G. Fanourakis 46, M. Fanti 70a,70b, M. Faraj 68a,68b, Z. Farazpay 96, A. Farbin 8,
 A. Farilla 76a, T. Farooque 106, S.M. Farrington 52, F. Fassi 35e, D. Fassouliotis 9, M. Faucci Giannelli 75a,75b,
 W.J. Fawcett 32, L. Fayard 66, P. Federicova 130, O.L. Fedin 37,a, G. Fedotov 37, M. Feickert 169, L. Feligioni 101,
 A. Fell 138, D.E. Fellers 122, C. Feng 62b, M. Feng 14b, Z. Feng 113, M.J. Fenton 159, A.B. Fenyuk 37,
 L. Ferencz 48, S.W. Ferguson 45, J. Ferrando 48, A. Ferrari 160, P. Ferrari 113,112, R. Ferrari 72a, D. Ferrere 56,
 C. Ferretti 105, F. Fiedler 99, A. Filipčič 92, E.K. Filmer 1, F. Filthaut 112, M.C.N. Fiolhais 129a,129c,c,
 L. Fiorini 162, F. Fischer 140, W.C. Fisher 106, T. Fitschen 100, I. Fleck 140, P. Fleischmann 105, T. Flick 170,
 L. Flores 127, M. Flores 33d, L.R. Flores Castillo 64a, F.M. Follega 77a,77b, N. Fomin 16, J.H. Foo 154,
 B.C. Forland 67, A. Formica 134, A.C. Forti 100, E. Fortin 101, A.W. Fortman 61, M.G. Foti 17a, L. Fountas 9,
 D. Fournier 66, H. Fox 90, P. Francavilla 73a,73b, S. Francescato 61, S. Franchellucci 56, M. Franchini 23b,23a,
 S. Franchino 63a, D. Francis 36, L. Franco 112, L. Franconi 19, M. Franklin 61, G. Frattari 26, A.C. Freegard 93,
 P.M. Freeman 20, W.S. Freund 81b, N. Fritzsche 50, A. Froch 54, D. Froidevaux 36, J.A. Frost 125, Y. Fu 62a,
 M. Fujimoto 117, E. Fullana Torregrosa 162,* J. Fuster 162, A. Gabrielli 23b,23a, A. Gabrielli 154, P. Gadow 48,
 G. Gagliardi 57b,57a, L.G. Gagnon 17a, G.E. Gallardo 125, E.J. Gallas 125, B.J. Gallop 133, R. Gamboa Goni 93,
 K.K. Gan 118, S. Ganguly 152, J. Gao 62a, Y. Gao 52, F.M. Garay Walls 136a,136b, B. Garcia 29,ag, C. García 162,
 J.E. García Navarro 162, J.A. García Pascual 14a, M. Garcia-Sciveres 17a, R.W. Gardner 39, D. Garg 79,
 R.B. Garg 142, S. Gargiulo 54, C.A. Garner 154, V. Garonne 29, S.J. Gasiorowski 137, P. Gaspar 81b,
 G. Gaudio 72a, V. Gautam 13, P. Gauzzi 74a,74b, I.I. Gavrilenko 37, A. Gavrilyuk 37, C. Gay 163, G. Gaycken 48,
 E.N. Gazis 10, A.A. Geanta 27b,27e, C.M. Gee 135, J. Geisen 97, M. Geisen 99, C. Gemme 57b, M.H. Genest 60,
 S. Gentile 74a,74b, S. George 94, W.F. George 20, T. Geralis 46, L.O. Gerlach 55, P. Gessinger-Befurt 36,
 M. Ghasemi Bostanabad 164, M. Ghneimat 140, K. Ghorbanian 93, A. Ghosal 140, A. Ghosh 159, A. Ghosh 7,
 B. Giacobbe 23b, S. Giagu 74a,74b, N. Giangiacomi 154, P. Giannetti 73a, A. Giannini 62a, S.M. Gibson 94,
 M. Gignac 135, D.T. Gil 84b, A.K. Gilbert 84a, B.J. Gilbert 41, D. Gillberg 34, G. Gilles 113, N.E.K. Gillwald 48,
 L. Ginabat 126, D.M. Gingrich 2,ad, M.P. Giordani 68a,68c, P.F. Giraud 134, G. Giugliarelli 68a,68c, D. Giugni 70a,
 F. Juli 36, I. Gkialas 9,j, L.K. Gladilin 37, C. Glasman 98, G.R. Gledhill 122, M. Glisic 122, I. Gnesi 43b,f,
 Y. Go 29,ag, M. Goblirsch-Kolb 26, B. Gocke 49, D. Godin 107, S. Goldfarb 104, T. Golling 56, M.G.D. Gololo 33g,
 D. Golubkov 37, J.P. Gombas 106, A. Gomes 129a,129b, G. Gomes Da Silva 140, A.J. Gomez Delegido 162,
 R. Goncalves Gama 55, R. Gonçalo 129a,129c, G. Gonella 122, L. Gonella 20, A. Gongadze 38, F. Gonnella 20,
 J.L. Gonski 41, R.Y. González Andana 52, S. González de la Hoz 162, S. Gonzalez Fernandez 13,
 R. Gonzalez Lopez 91, C. Gonzalez Renteria 17a, R. Gonzalez Suarez 160, S. Gonzalez-Sevilla 56,
 G.R. Gonzalvo Rodriguez 162, L. Goossens 36, N.A. Gorasia 20, P.A. Gorbounov 37, B. Gorini 36,
 E. Gorini 69a,69b, A. Gorišek 92, A.T. Goshaw 51, M.I. Gostkin 38, C.A. Gottardo 36, M. Gouighri 35b,
 V. Goumarre 48, A.G. Goussiou 137, N. Govender 33c, C. Goy 4, I. Grabowska-Bold 84a, K. Graham 34,
 E. Gramstad 124, S. Grancagnolo 18, M. Grandi 145, V. Gratchev 37,* P.M. Gravila 27f, F.G. Gravili 69a,69b,
 H.M. Gray 17a, M. Greco 69a,69b, C. Grefe 24, I.M. Gregor 48, P. Grenier 142, C. Grieco 13, A.A. Grillo 135,
 K. Grimm 31,n, S. Grinstein 13,u, J.-F. Grivaz 66, E. Gross 168, J. Grosse-Knetter 55, C. Grud 105,
 A. Grummer 111, J.C. Grundy 125, L. Guan 105, W. Guan 169, C. Gubbels 163, J.G.R. Guerrero Rojas 162,
 G. Guerrieri 68a,68b, F. Guescini 109, R. Gugel 99, J.A.M. Guhit 105, A. Guida 48, T. Guillemin 4,
 E. Guilloton 166,133, S. Guindon 36, F. Guo 14a,14d, J. Guo 62c, L. Guo 66, Y. Guo 105, R. Gupta 48, S. Gurbuz 24,
 S.S. Gurdasani 54, G. Gustavino 36, M. Guth 56, P. Gutierrez 119, L.F. Gutierrez Zagazeta 127, C. Gutschow 95,
 C. Guyot 134, C. Gwenlan 125, C.B. Gwilliam 91, E.S. Haaland 124, A. Haas 116, M. Habedank 48, C. Haber 17a,
 H.K. Hadavand 8, A. Hadef 99, S. Hadzic 109, E.H. Haines 95, M. Haleem 165, J. Haley 120, J.J. Hall 138,
 G.D. Hallewell 101, L. Halser 19, K. Hamano 164, H. Hamdaoui 35e, M. Hamer 24, G.N. Hamity 52, J. Han 62b,
 K. Han 62a, L. Han 14c, L. Han 62a, S. Han 17a, Y.F. Han 154, K. Hanagaki 82, M. Hance 135, D.A. Hangal 41,aa,
 H. Hanif 141, M.D. Hank 39, R. Hankache 100, J.B. Hansen 42, J.D. Hansen 42, P.H. Hansen 42, K. Hara 156,
 D. Harada 56, T. Harenberg 170, S. Harkusha 37, Y.T. Harris 125, N.M. Harrison 118, P.F. Harrison 166,
 N.M. Hartman 142, N.M. Hartmann 108, Y. Hasegawa 139, A. Hasib 52, S. Haug 19, R. Hauser 106,

- M. Havranek ¹³¹, C.M. Hawkes ²⁰, R.J. Hawkings ³⁶, S. Hayashida ¹¹⁰, D. Hayden ¹⁰⁶, C. Hayes ¹⁰⁵, R.L. Hayes ¹⁶³, C.P. Hays ¹²⁵, J.M. Hays ⁹³, H.S. Hayward ⁹¹, F. He ^{62a}, Y. He ¹⁵³, Y. He ¹²⁶, M.P. Heath ⁵², V. Hedberg ⁹⁷, A.L. Heggelund ¹²⁴, N.D. Hehir ⁹³, C. Heidegger ⁵⁴, K.K. Heidegger ⁵⁴, W.D. Heidorn ⁸⁰, J. Heilman ³⁴, S. Heim ⁴⁸, T. Heim ^{17a}, J.G. Heinlein ¹²⁷, J.J. Heinrich ¹²², L. Heinrich ¹⁰⁹, J. Hejbal ¹³⁰, L. Helary ⁴⁸, A. Held ¹⁶⁹, S. Hellesund ¹²⁴, C.M. Helling ¹⁶³, S. Hellman ^{47a,47b}, C. Helsens ³⁶, R.C.W. Henderson ⁹⁰, L. Henkelmann ³², A.M. Henriques Correia ³⁶, H. Herde ⁹⁷, Y. Hernández Jiménez ¹⁴⁴, L.M. Herrmann ²⁴, M.G. Herrmann ¹⁰⁸, T. Herrmann ⁵⁰, G. Herten ⁵⁴, R. Hertenberger ¹⁰⁸, L. Hervas ³⁶, N.P. Hessey ^{155a}, H. Hibi ⁸³, E. Higón-Rodriguez ¹⁶², S.J. Hillier ²⁰, I. Hinchliffe ^{17a}, F. Hinterkeuser ²⁴, M. Hirose ¹²³, S. Hirose ¹⁵⁶, D. Hirschbuehl ¹⁷⁰, T.G. Hitchings ¹⁰⁰, B. Hiti ⁹², J. Hobbs ¹⁴⁴, R. Hobincu ^{27e}, N. Hod ¹⁶⁸, M.C. Hodgkinson ¹³⁸, B.H. Hodkinson ³², A. Hoecker ³⁶, J. Hofer ⁴⁸, D. Hohn ⁵⁴, T. Holm ²⁴, M. Holzbock ¹⁰⁹, L.B.A.H. Hommels ³², B.P. Honan ¹⁰⁰, J. Hong ^{62c}, T.M. Hong ¹²⁸, J.C. Honig ⁵⁴, A. Höngle ¹⁰⁹, B.H. Hooberman ¹⁶¹, W.H. Hopkins ⁶, Y. Horii ¹¹⁰, S. Hou ¹⁴⁷, A.S. Howard ⁹², J. Howarth ⁵⁹, J. Hoya ⁶, M. Hrabovsky ¹²¹, A. Hrynevich ⁴⁸, T. Hrynevich ⁴, P.J. Hsu ⁶⁵, S.-C. Hsu ¹³⁷, Q. Hu ^{41,aa}, Y.F. Hu ^{14a,14d,af}, D.P. Huang ⁹⁵, S. Huang ^{64b}, X. Huang ^{14c}, Y. Huang ^{62a}, Y. Huang ^{14a}, Z. Huang ¹⁰⁰, Z. Hubacek ¹³¹, M. Huebner ²⁴, F. Huegging ²⁴, T.B. Huffman ¹²⁵, M. Huhtinen ³⁶, S.K. Huiberts ¹⁶, R. Hulsken ¹⁰³, N. Huseynov ^{12,a}, J. Huston ¹⁰⁶, J. Huth ⁶¹, R. Hyneman ¹⁴², S. Hyrych ^{28a}, G. Iacobucci ⁵⁶, G. Iakovidis ²⁹, I. Ibragimov ¹⁴⁰, L. Iconomidou-Fayard ⁶⁶, P. Iengo ^{71a,71b}, R. Iguchi ¹⁵², T. Iizawa ⁵⁶, Y. Ikegami ⁸², A. Ilg ¹⁹, N. Ilic ¹⁵⁴, H. Imam ^{35a}, T. Ingebretsen Carlson ^{47a,47b}, G. Introzzi ^{72a,72b}, M. Iodice ^{76a}, V. Ippolito ^{74a,74b}, M. Ishino ¹⁵², W. Islam ¹⁶⁹, C. Issever ^{18,48}, S. Istin ^{21a,ai}, H. Ito ¹⁶⁷, J.M. Iturbe Ponce ^{64a}, R. Iuppa ^{77a,77b}, A. Ivina ¹⁶⁸, J.M. Izen ⁴⁵, V. Izzo ^{71a}, P. Jacka ^{130,131}, P. Jackson ¹, R.M. Jacobs ⁴⁸, B.P. Jaeger ¹⁴¹, C.S. Jagfeld ¹⁰⁸, G. Jäkel ¹⁷⁰, K. Jakobs ⁵⁴, T. Jakoubek ¹⁶⁸, J. Jamieson ⁵⁹, K.W. Janas ^{84a}, G. Jarlskog ⁹⁷, A.E. Jaspan ⁹¹, M. Javurkova ¹⁰², F. Jeanneau ¹³⁴, L. Jeanty ¹²², J. Jejelava ^{148a,z}, P. Jenni ^{54,g}, C.E. Jessiman ³⁴, S. Jézéquel ⁴, J. Jia ¹⁴⁴, X. Jia ⁶¹, X. Jia ^{14a,14d}, Z. Jia ^{14c}, Y. Jiang ^{62a}, S. Jiggins ⁵², J. Jimenez Pena ¹⁰⁹, S. Jin ^{14c}, A. Jinaru ^{27b}, O. Jinnouchi ¹⁵³, P. Johansson ¹³⁸, K.A. Johns ⁷, D.M. Jones ³², E. Jones ¹⁶⁶, P. Jones ³², R.W.L. Jones ⁹⁰, T.J. Jones ⁹¹, R. Joshi ¹¹⁸, J. Jovicevic ¹⁵, X. Ju ^{17a}, J.J. Junggeburth ³⁶, A. Juste Rozas ^{13,u}, S. Kabana ^{136e}, A. Kaczmarska ⁸⁵, M. Kado ^{74a,74b}, H. Kagan ¹¹⁸, M. Kagan ¹⁴², A. Kahn ⁴¹, A. Kahn ¹²⁷, C. Kahra ⁹⁹, T. Kaji ¹⁶⁷, E. Kajomovitz ¹⁴⁹, N. Kakati ¹⁶⁸, C.W. Kalderon ²⁹, A. Kamenshchikov ¹⁵⁴, S. Kanayama ¹⁵³, N.J. Kang ¹³⁵, Y. Kano ¹¹⁰, D. Kar ^{33g}, K. Karava ¹²⁵, M.J. Kareem ^{155b}, E. Karentzos ⁵⁴, I. Karkalias ¹⁵¹, S.N. Karpov ³⁸, Z.M. Karpova ³⁸, V. Kartvelishvili ⁹⁰, A.N. Karyukhin ³⁷, E. Kasimi ¹⁵¹, C. Kato ^{62d}, J. Katzy ⁴⁸, S. Kaur ³⁴, K. Kawade ¹³⁹, K. Kawagoe ⁸⁸, T. Kawamoto ¹³⁴, G. Kawamura ⁵⁵, E.F. Kay ¹⁶⁴, F.I. Kaya ¹⁵⁷, S. Kazakos ¹³, V.F. Kazanin ³⁷, Y. Ke ¹⁴⁴, J.M. Keaveney ^{33a}, R. Keeler ¹⁶⁴, G.V. Kehris ⁶¹, J.S. Keller ³⁴, A.S. Kelly ⁹⁵, D. Kelsey ¹⁴⁵, J.J. Kempster ²⁰, K.E. Kennedy ⁴¹, P.D. Kennedy ⁹⁹, O. Kepka ¹³⁰, B.P. Kerridge ¹⁶⁶, S. Kersten ¹⁷⁰, B.P. Kerševan ⁹², S. Keshri ⁶⁶, L. Keszeghova ^{28a}, S. Ketabchi Haghighat ¹⁵⁴, M. Khandoga ¹²⁶, A. Khanov ¹²⁰, A.G. Kharlamov ³⁷, T. Kharlamova ³⁷, E.E. Khoda ¹³⁷, T.J. Khoo ¹⁸, G. Khoriauli ¹⁶⁵, J. Khubua ^{148b}, Y.A.R. Khwaira ⁶⁶, M. Kiehn ³⁶, A. Kilgallon ¹²², D.W. Kim ^{47a,47b}, E. Kim ¹⁵³, Y.K. Kim ³⁹, N. Kimura ⁹⁵, A. Kirchhoff ⁵⁵, D. Kirchmeier ⁵⁰, C. Kirfel ²⁴, J. Kirk ¹³³, A.E. Kiryunin ¹⁰⁹, T. Kishimoto ¹⁵², D.P. Kisliuk ¹⁵⁴, C. Kitsaki ¹⁰, O. Kivernyk ²⁴, M. Klassen ^{63a}, C. Klein ³⁴, L. Klein ¹⁶⁵, M.H. Klein ¹⁰⁵, M. Klein ⁹¹, S.B. Klein ⁵⁶, U. Klein ⁹¹, P. Klimek ³⁶, A. Klimentov ²⁹, F. Klimentov ¹⁰⁹, T. Klingl ²⁴, T. Klioutchnikova ³⁶, F.F. Klitzner ¹⁰⁸, P. Kluit ¹¹³, S. Kluth ¹⁰⁹, E. Kneringer ⁷⁸, T.M. Knight ¹⁵⁴, A. Knue ⁵⁴, D. Kobayashi ⁸⁸, R. Kobayashi ⁸⁶, M. Kocian ¹⁴², P. Kodyš ¹³², D.M. Koeck ¹⁴⁵, P.T. Koenig ²⁴, T. Koffas ³⁴, M. Kolb ¹³⁴, I. Koletsou ⁴, T. Komarek ¹²¹, K. Köneke ⁵⁴, A.X.Y. Kong ¹, T. Kono ¹¹⁷, N. Konstantinidis ⁹⁵, B. Konya ⁹⁷, R. Kopeliansky ⁶⁷, S. Koperny ^{84a}, K. Korcyl ⁸⁵, K. Kordas ¹⁵¹, G. Koren ¹⁵⁰, A. Korn ⁹⁵, S. Korn ⁵⁵, I. Korolkov ¹³, N. Korotkova ³⁷, B. Kortman ¹¹³, O. Kortner ¹⁰⁹, S. Kortner ¹⁰⁹, W.H. Kostecka ¹¹⁴, V.V. Kostyukhin ¹⁴⁰, A. Kotsokechagia ¹³⁴, A. Kotwal ⁵¹, A. Koulouris ³⁶, A. Kourkoumelis-Charalampidi ^{72a,72b}, C. Kourkoumelis ⁹, E. Kourlitis ⁶, O. Kovanda ¹⁴⁵, R. Kowalewski ¹⁶⁴, W. Kozanecki ¹³⁴, A.S. Kozhin ³⁷, V.A. Kramarenko ³⁷, G. Kramberger ⁹², P. Kramer ⁹⁹, M.W. Krasny ¹²⁶, A. Krasznahorkay ³⁶, J.A. Kremer ⁹⁹, T. Kresse ⁵⁰, J. Kretzschmar ⁹¹, K. Kreul ¹⁸, P. Krieger ¹⁵⁴, F. Krieter ¹⁰⁸, S. Krishnamurthy ¹⁰², A. Krishnan ^{63b}, M. Krivos ¹³², K. Krizka ^{17a}, K. Kroeninger ⁴⁹, H. Kroha ¹⁰⁹, J. Kroll ¹³⁰, J. Kroll ¹²⁷, K.S. Krowpman ¹⁰⁶, U. Kruchonak ³⁸, H. Krüger ²⁴, N. Krumnack ⁸⁰, M.C. Kruse ⁵¹, J.A. Krzysiak ⁸⁵, O. Kuchinskaia ³⁷, S. Kuday ^{3a}, D. Kuechler ⁴⁸, J.T. Kuechler ⁴⁸, S. Kuehn ³⁶, T. Kuhl ⁴⁸, V. Kukhtin ³⁸, Y. Kulchitsky ^{37,a}, S. Kuleshov ^{136d,136b}, M. Kumar ^{33g}, N. Kumari ¹⁰¹, A. Kupco ¹³⁰, T. Kupfer ⁴⁹, A. Kupich ³⁷, O. Kuprash ⁵⁴, H. Kurashige ⁸³, L.L. Kurchaninov ^{155a}, Y.A. Kurochkin ³⁷,

- A. Kurova ³⁷, M. Kuze ¹⁵³, A.K. Kvam ¹⁰², J. Kvita ¹²¹, T. Kwan ¹⁰³, K.W. Kwok ^{64a}, N.G. Kyriacou ¹⁰⁵,
 L.A.O. Laatu ¹⁰¹, C. Lacasta ¹⁶², F. Lacava ^{74a,74b}, H. Lacker ¹⁸, D. Lacour ¹²⁶, N.N. Lad ⁹⁵, E. Ladygin ³⁸,
 B. Laforge ¹²⁶, T. Lagouri ^{136e}, S. Lai ⁵⁵, I.K. Lakomiec ^{84a}, N. Lalloue ⁶⁰, J.E. Lambert ¹¹⁹, S. Lammers ⁶⁷,
 W. Lampl ⁷, C. Lampoudis ¹⁵¹, A.N. Lancaster ¹¹⁴, E. Lançon ²⁹, U. Landgraf ⁵⁴, M.P.J. Landon ⁹³, V.S. Lang ⁵⁴,
 R.J. Langenberg ¹⁰², A.J. Lankford ¹⁵⁹, F. Lanni ³⁶, K. Lantzsch ²⁴, A. Lanza ^{72a}, A. Lapertosa ^{57b,57a},
 J.F. Laporte ¹³⁴, T. Lari ^{70a}, F. Lasagni Manghi ^{23b}, M. Lassnig ³⁶, V. Latonova ¹³⁰, T.S. Lau ^{64a}, A. Laudrain ⁹⁹,
 A. Laurier ³⁴, S.D. Lawlor ⁹⁴, Z. Lawrence ¹⁰⁰, M. Lazzaroni ^{70a,70b}, B. Le ¹⁰⁰, B. Leban ⁹², A. Lebedev ⁸⁰,
 M. LeBlanc ³⁶, T. LeCompte ⁶, F. Ledroit-Guillon ⁶⁰, A.C.A. Lee ⁹⁵, G.R. Lee ¹⁶, L. Lee ⁶¹, S.C. Lee ¹⁴⁷,
 S. Lee ^{47a,47b}, T.F. Lee ⁹¹, L.L. Leeuw ^{33c}, H.P. Lefebvre ⁹⁴, M. Lefebvre ¹⁶⁴, C. Leggett ^{17a}, K. Lehmann ¹⁴¹,
 G. Lehmann Miotti ³⁶, M. Leigh ⁵⁶, W.A. Leight ¹⁰², A. Leisos ^{151,t}, M.A.L. Leite ^{81c}, C.E. Leitgeb ⁴⁸,
 R. Leitner ¹³², K.J.C. Leney ⁴⁴, T. Lenz ²⁴, S. Leone ^{73a}, C. Leonidopoulos ⁵², A. Leopold ¹⁴³, C. Leroy ¹⁰⁷,
 R. Les ¹⁰⁶, C.G. Lester ³², M. Levchenko ³⁷, J. Levêque ⁴, D. Levin ¹⁰⁵, L.J. Levinson ¹⁶⁸, M.P. Lewicki ⁸⁵,
 D.J. Lewis ⁴, A. Li ⁵, B. Li ^{14b}, B. Li ^{62b}, C. Li ^{62a}, C-Q. Li ^{62c}, H. Li ^{62a}, H. Li ^{62b}, H. Li ^{14c}, H. Li ^{62b}, J. Li ^{62c},
 K. Li ¹³⁷, L. Li ^{62c}, M. Li ^{14a,14d}, Q.Y. Li ^{62a}, S. Li ^{14a,14d}, S. Li ^{62d,62c,e}, T. Li ^{62b}, X. Li ¹⁰³, Z. Li ^{62b}, Z. Li ¹²⁵,
 Z. Li ¹⁰³, Z. Li ⁹¹, Z. Li ^{14a,14d}, Z. Liang ^{14a}, M. Liberatore ⁴⁸, B. Liberti ^{75a}, K. Lie ^{64c}, J. Lieber Marin ^{81b},
 K. Lin ¹⁰⁶, R.A. Linck ⁶⁷, R.E. Lindley ⁷, J.H. Lindon ², A. Linss ⁴⁸, E. Lipeles ¹²⁷, A. Lipniacka ¹⁶, A. Lister ¹⁶³,
 J.D. Little ⁴, B. Liu ^{14a}, B.X. Liu ¹⁴¹, D. Liu ^{62d,62c}, J.B. Liu ^{62a}, J.K.K. Liu ³², K. Liu ^{62d,62c}, M. Liu ^{62a},
 M.Y. Liu ^{62a}, P. Liu ^{14a}, Q. Liu ^{62d,137,62c}, X. Liu ^{62a}, Y. Liu ⁴⁸, Y. Liu ^{14c,14d}, Y.L. Liu ¹⁰⁵, Y.W. Liu ^{62a},
 M. Livan ^{72a,72b}, J. Llorente Merino ¹⁴¹, S.L. Lloyd ⁹³, E.M. Lobodzinska ⁴⁸, P. Loch ⁷, S. Loffredo ^{75a,75b},
 T. Lohse ¹⁸, K. Lohwasser ¹³⁸, M. Lokajicek ¹³⁰, J.D. Long ¹⁶¹, I. Longarini ^{74a,74b}, L. Longo ^{69a,69b},
 R. Longo ¹⁶¹, I. Lopez Paz ³⁶, A. Lopez Solis ⁴⁸, J. Lorenz ¹⁰⁸, N. Lorenzo Martinez ⁴, A.M. Lory ¹⁰⁸,
 A. Lösle ⁵⁴, X. Lou ^{47a,47b}, X. Lou ^{14a,14d}, A. Lounis ⁶⁶, J. Love ⁶, P.A. Love ⁹⁰, J.J. Lozano Bahilo ¹⁶²,
 G. Lu ^{14a,14d}, M. Lu ⁷⁹, S. Lu ¹²⁷, Y.J. Lu ⁶⁵, H.J. Lubatti ¹³⁷, C. Luci ^{74a,74b}, F.L. Lucio Alves ^{14c}, A. Lucotte ⁶⁰,
 F. Luehring ⁶⁷, I. Luise ¹⁴⁴, O. Lukianchuk ⁶⁶, O. Lundberg ¹⁴³, B. Lund-Jensen ¹⁴³, N.A. Luongo ¹²²,
 M.S. Lutz ¹⁵⁰, D. Lynn ²⁹, H. Lyons ⁹¹, R. Lysak ¹³⁰, E. Lytken ⁹⁷, F. Lyu ^{14a}, V. Lyubushkin ³⁸,
 T. Lyubushkina ³⁸, H. Ma ²⁹, L.L. Ma ^{62b}, Y. Ma ⁹⁵, D.M. Mac Donell ¹⁶⁴, G. Maccarrone ⁵³,
 J.C. MacDonald ¹³⁸, R. Madar ⁴⁰, W.F. Mader ⁵⁰, J. Maeda ⁸³, T. Maeno ²⁹, M. Maerker ⁵⁰, V. Magerl ⁵⁴,
 H. Maguire ¹³⁸, D.J. Mahon ⁴¹, C. Maidantchik ^{81b}, A. Maio ^{129a,129b,129d}, K. Maj ^{84a}, O. Majersky ^{28a},
 S. Majewski ¹²², N. Makovec ⁶⁶, V. Maksimovic ¹⁵, B. Malaescu ¹²⁶, Pa. Malecki ⁸⁵, V.P. Maleev ³⁷,
 F. Malek ⁶⁰, D. Malito ^{43b,43a}, U. Mallik ⁷⁹, C. Malone ³², S. Maltezos ¹⁰, S. Malyukov ³⁸, J. Mamuzic ¹³,
 G. Mancini ⁵³, G. Manco ^{72a,72b}, J.P. Mandalia ⁹³, I. Mandić ⁹², L. Manhaes de Andrade Filho ^{81a},
 I.M. Maniatis ¹⁵¹, M. Manisha ¹³⁴, J. Manjarres Ramos ⁵⁰, D.C. Mankad ¹⁶⁸, A. Mann ¹⁰⁸, B. Mansoulie ¹³⁴,
 S. Manzoni ³⁶, A. Marantis ¹⁵¹, G. Marchiori ⁵, M. Marcisovsky ¹³⁰, L. Marcoccia ^{75a,75b}, C. Marcon ^{70a,70b},
 M. Marinescu ²⁰, M. Marjanovic ¹¹⁹, E.J. Marshall ⁹⁰, Z. Marshall ^{17a}, S. Marti-Garcia ¹⁶², T.A. Martin ¹⁶⁶,
 V.J. Martin ⁵², B. Martin dit Latour ¹⁶, L. Martinelli ^{74a,74b}, M. Martinez ^{13,u}, P. Martinez Agullo ¹⁶²,
 V.I. Martinez Outschoorn ¹⁰², P. Martinez Suarez ¹³, S. Martin-Haugh ¹³³, V.S. Martoiu ^{27b},
 A.C. Martyniuk ⁹⁵, A. Marzin ³⁶, S.R. Maschek ¹⁰⁹, L. Masetti ⁹⁹, T. Mashimo ¹⁵², J. Masik ¹⁰⁰,
 A.L. Maslennikov ³⁷, L. Massa ^{23b}, P. Massarotti ^{71a,71b}, P. Mastrandrea ^{73a,73b}, A. Mastroberardino ^{43b,43a},
 T. Masubuchi ¹⁵², T. Mathisen ¹⁶⁰, N. Matsuzawa ¹⁵², J. Maurer ^{27b}, B. Maček ⁹², D.A. Maximov ³⁷,
 R. Mazini ¹⁴⁷, I. Maznas ¹⁵¹, M. Mazza ¹⁰⁶, S.M. Mazza ¹³⁵, C. Mc Ginn ^{29,ag}, J.P. Mc Gowan ¹⁰³,
 S.P. Mc Kee ¹⁰⁵, W.P. McCormack ^{17a}, E.F. McDonald ¹⁰⁴, A.E. McDougall ¹¹³, J.A. McFayden ¹⁴⁵,
 G. Mchedlidze ^{148b}, R.P. Mckenzie ^{33g}, T.C. McLachlan ⁴⁸, D.J. McLaughlin ⁹⁵, K.D. McLean ¹⁶⁴,
 S.J. McMahon ¹³³, P.C. McNamara ¹⁰⁴, C.M. Mcpartland ⁹¹, R.A. McPherson ^{164,w}, T. Megy ⁴⁰,
 S. Mehlhase ¹⁰⁸, A. Mehta ⁹¹, B. Meirose ⁴⁵, D. Melini ¹⁴⁹, B.R. Mellado Garcia ^{33g}, A.H. Melo ⁵⁵,
 F. Meloni ⁴⁸, E.D. Mendes Gouveia ^{129a}, A.M. Mendes Jacques Da Costa ²⁰, H.Y. Meng ¹⁵⁴, L. Meng ⁹⁰,
 S. Menke ¹⁰⁹, M. Mentink ³⁶, E. Meoni ^{43b,43a}, C. Merlassino ¹²⁵, L. Merola ^{71a,71b}, C. Meroni ^{70a},
 G. Merz ¹⁰⁵, O. Meshkov ³⁷, J.K.R. Meshreki ¹⁴⁰, J. Metcalfe ⁶, A.S. Mete ⁶, C. Meyer ⁶⁷, J.-P. Meyer ¹³⁴,
 M. Michetti ¹⁸, R.P. Middleton ¹³³, L. Mijović ⁵², G. Mikenberg ¹⁶⁸, M. Mikestikova ¹³⁰, M. Mikuž ⁹²,
 H. Mildner ¹³⁸, A. Milic ³⁶, C.D. Milke ⁴⁴, D.W. Miller ³⁹, L.S. Miller ³⁴, A. Milov ¹⁶⁸, D.A. Milstead ^{47a,47b},
 T. Min ^{14c}, A.A. Minaenko ³⁷, I.A. Minashvili ^{148b}, L. Mince ⁵⁹, A.I. Mincer ¹¹⁶, B. Mindur ^{84a}, M. Mineev ³⁸,
 Y. Mino ⁸⁶, L.M. Mir ¹³, M. Miralles Lopez ¹⁶², M. Mironova ¹²⁵, M.C. Missio ¹¹², T. Mitani ¹⁶⁷, A. Mitra ¹⁶⁶,
 V.A. Mitsou ¹⁶², O. Miu ¹⁵⁴, P.S. Miyagawa ⁹³, Y. Miyazaki ⁸⁸, A. Mizukami ⁸², J.U. Mjörnmark ⁹⁷,
 T. Mkrtchyan ^{63a}, T. Mlinarevic ⁹⁵, M. Mlynarikova ³⁶, T. Moa ^{47a,47b}, S. Mobius ⁵⁵, K. Mochizuki ¹⁰⁷,

- P. Moder⁴⁸, P. Mogg¹⁰⁸, A.F. Mohammed^{14a,14d}, S. Mohapatra⁴¹, G. Mokgatitswane^{33g}, B. Mondal¹⁴⁰, S. Mondal¹³¹, K. Mönig⁴⁸, E. Monnier¹⁰¹, L. Monsonis Romero¹⁶², J. Montejo Berlingen³⁶, M. Montella¹¹⁸, F. Monticelli⁸⁹, N. Morange⁶⁶, A.L. Moreira De Carvalho^{129a}, M. Moreno Llácer¹⁶², C. Moreno Martinez⁵⁶, P. Morettini^{57b}, S. Morgenstern¹⁶⁶, M. Morii⁶¹, M. Morinaga¹⁵², A.K. Morley³⁶, F. Morodei^{74a,74b}, L. Morvaj³⁶, P. Moschovakos³⁶, B. Moser³⁶, M. Mosidze^{148b}, T. Moskalets⁵⁴, P. Moskvitina¹¹², J. Moss^{31,0}, E.J.W. Moyse¹⁰², S. Muanza¹⁰¹, J. Mueller¹²⁸, D. Muenstermann⁹⁰, R. Müller¹⁹, G.A. Mullier⁹⁷, J.J. Mullin¹²⁷, D.P. Mungo¹⁵⁴, J.L. Munoz Martinez¹³, D. Munoz Perez¹⁶², F.J. Munoz Sanchez¹⁰⁰, M. Murin¹⁰⁰, W.J. Murray^{166,133}, A. Murrone^{70a,70b}, J.M. Muse¹¹⁹, M. Muškinja^{17a}, C. Mwewa²⁹, A.G. Myagkov^{37,a}, A.J. Myers⁸, A.A. Myers¹²⁸, G. Myers⁶⁷, M. Myska¹³¹, B.P. Nachman^{17a}, O. Nackenhorst⁴⁹, A. Nag⁵⁰, K. Nagai¹²⁵, K. Nagano⁸², J.L. Nagle^{29,ag}, E. Nagy¹⁰¹, A.M. Nairz³⁶, Y. Nakahama⁸², K. Nakamura⁸², H. Nanjo¹²³, R. Narayan⁴⁴, E.A. Narayanan¹¹¹, I. Naryshkin³⁷, M. Naseri³⁴, C. Nass²⁴, G. Navarro^{22a}, J. Navarro-Gonzalez¹⁶², R. Nayak¹⁵⁰, A. Nayaz¹⁸, P.Y. Nechaeva³⁷, F. Nechansky⁴⁸, L. Nedic¹²⁵, T.J. Neep²⁰, A. Negri^{72a,72b}, M. Negrini^{23b}, C. Nellist¹¹², C. Nelson¹⁰³, K. Nelson¹⁰⁵, S. Nemecek¹³⁰, M. Nessi^{36,h}, M.S. Neubauer¹⁶¹, F. Neuhaus⁹⁹, J. Neundorf⁴⁸, R. Newhouse¹⁶³, P.R. Newman²⁰, C.W. Ng¹²⁸, Y.S. Ng¹⁸, Y.W.Y. Ng⁴⁸, B. Ngair^{35e}, H.D.N. Nguyen¹⁰⁷, R.B. Nickerson¹²⁵, R. Nicolaïdou¹³⁴, J. Nielsen¹³⁵, M. Niemeyer⁵⁵, N. Nikiforou³⁶, V. Nikolaenko^{37,a}, I. Nikolic-Audit¹²⁶, K. Nikolopoulos²⁰, P. Nilsson²⁹, H.R. Nindhito⁵⁶, A. Nisati^{74a}, N. Nishu², R. Nisius¹⁰⁹, J-E. Nitschke⁵⁰, E.K. Nkadameng^{33g}, S.J. Noacco Rosende⁸⁹, T. Nobe¹⁵², D.L. Noel³², Y. Noguchi⁸⁶, T. Nommensen¹⁴⁶, M.A. Nomura²⁹, M.B. Norfolk¹³⁸, R.R.B. Norisam⁹⁵, B.J. Norman³⁴, J. Novak⁹², T. Novak⁴⁸, O. Novgorodova⁵⁰, L. Novotny¹³¹, R. Novotny¹¹¹, L. Nozka¹²¹, K. Ntekas¹⁵⁹, N.M.J. Nunes De Moura Junior^{81b}, E. Nurse⁹⁵, F.G. Oakham^{34,ad}, J. Ocariz¹²⁶, A. Ochi⁸³, I. Ochoa^{129a}, S. Oerdekk¹⁶⁰, A. Ogrodnik^{84a}, A. Oh¹⁰⁰, C.C. Ohm¹⁴³, H. Oide⁸², R. Oishi¹⁵², M.L. Ojeda⁴⁸, Y. Okazaki⁸⁶, M.W. O'Keefe⁹¹, Y. Okumura¹⁵², A. Olariu^{27b}, L.F. Oleiro Seabra^{129a}, S.A. Olivares Pino^{136e}, D. Oliveira Damazio²⁹, D. Oliveira Goncalves^{81a}, J.L. Oliver¹⁵⁹, M.J.R. Olsson¹⁵⁹, A. Olszewski⁸⁵, J. Olszowska^{85,*}, Ö.O. Öncel⁵⁴, D.C. O'Neil¹⁴¹, A.P. O'Neill¹⁹, A. Onofre^{129a,129e}, P.U.E. Onyisi¹¹, M.J. Oreglia³⁹, G.E. Orellana⁸⁹, D. Orestano^{76a,76b}, N. Orlando¹³, R.S. Orr¹⁵⁴, V. O'Shea⁵⁹, R. Ospanov^{62a}, G. Otero y Garzon³⁰, H. Otono⁸⁸, P.S. Ott^{63a}, G.J. Ottino^{17a}, M. Ouchrif^{35d}, J. Ouellette^{29,ag}, F. Ould-Saada¹²⁴, M. Owen⁵⁹, R.E. Owen¹³³, K.Y. Oyulmaz^{21a}, V.E. Ozcan^{21a}, N. Ozturk⁸, S. Ozturk^{21d}, J. Pacalt¹²¹, H.A. Pacey³², K. Pachal⁵¹, A. Pacheco Pages¹³, C. Padilla Aranda¹³, G. Padovano^{74a,74b}, S. Pagan Griso^{17a}, G. Palacino⁶⁷, A. Palazzo^{69a,69b}, S. Palazzo⁵², S. Palestini³⁶, M. Palka^{84b}, J. Pan¹⁷¹, T. Pan^{64a}, D.K. Panchal¹¹, C.E. Pandini¹¹³, J.G. Panduro Vazquez⁹⁴, H. Pang^{14b}, P. Pani⁴⁸, G. Panizzo^{68a,68c}, L. Paolozzi⁵⁶, C. Papadatos¹⁰⁷, S. Parajuli⁴⁴, A. Paramonov⁶, C. Paraskevopoulos¹⁰, D. Paredes Hernandez^{64b}, T.H. Park¹⁵⁴, M.A. Parker³², F. Parodi^{57b,57a}, E.W. Parrish¹¹⁴, V.A. Parrish⁵², J.A. Parsons⁴¹, U. Parzefall⁵⁴, B. Pascual Dias¹⁰⁷, L. Pascual Dominguez¹⁵⁰, V.R. Pascuzzi^{17a}, F. Pasquali¹¹³, E. Pasqualucci^{74a}, S. Passaggio^{57b}, F. Pastore⁹⁴, P. Pasuwan^{47a,47b}, P. Patel⁸⁵, J.R. Pater¹⁰⁰, T. Pauly³⁶, J. Pearkes¹⁴², M. Pedersen¹²⁴, R. Pedro^{129a}, S.V. Peleganchuk³⁷, O. Penc³⁶, E.A. Pender⁵², C. Peng^{64b}, H. Peng^{62a}, K.E. Penski¹⁰⁸, M. Penzin³⁷, B.S. Peralva^{81d}, A.P. Pereira Peixoto⁶⁰, L. Pereira Sanchez^{47a,47b}, D.V. Perepelitsa^{29,ag}, E. Perez Codina^{155a}, M. Perganti¹⁰, L. Perini^{70a,70b,*}, H. Pernegger³⁶, S. Perrella³⁶, A. Perrevoort¹¹², O. Perrin⁴⁰, K. Peters⁴⁸, R.F.Y. Peters¹⁰⁰, B.A. Petersen³⁶, T.C. Petersen⁴², E. Petit¹⁰¹, V. Petousis¹³¹, C. Petridou¹⁵¹, A. Petrukhin¹⁴⁰, M. Pettee^{17a}, N.E. Pettersson³⁶, A. Petukhov³⁷, K. Petukhova¹³², A. Peyaud¹³⁴, R. Pezoa^{136f}, L. Pezzotti³⁶, G. Pezzullo¹⁷¹, T.M. Pham¹⁶⁹, T. Pham¹⁰⁴, P.W. Phillips¹³³, M.W. Phipps¹⁶¹, G. Piacquadio¹⁴⁴, E. Pianori^{17a}, F. Piazza^{70a,70b}, R. Piegaia³⁰, D. Pietreanu^{27b}, A.D. Pilkington¹⁰⁰, M. Pinamonti^{68a,68c}, J.L. Pinfold², B.C. Pinheiro Pereira^{129a}, C. Pitman Donaldson⁹⁵, D.A. Pizzi³⁴, L. Pizzimento^{75a,75b}, A. Pizzini¹¹³, M.-A. Pleier²⁹, V. Plesanovs⁵⁴, V. Pleskot¹³², E. Plotnikova³⁸, G. Poddar⁴, R. Poettgen⁹⁷, L. Poggioli¹²⁶, I. Pogrebnyak¹⁰⁶, D. Pohl²⁴, I. Pokharel⁵⁵, S. Polacek¹³², G. Polesello^{72a}, A. Poley^{141,155a}, R. Polifka¹³¹, A. Polini^{23b}, C.S. Pollard¹²⁵, Z.B. Pollock¹¹⁸, V. Polychronakos²⁹, E. Pompa Pacchi^{74a,74b}, D. Ponomarenko³⁷, L. Pontecorvo³⁶, S. Popa^{27a}, G.A. Popeneiciu^{27d}, D.M. Portillo Quintero^{155a}, S. Pospisil¹³¹, P. Postolache^{27c}, K. Potamianos¹²⁵, I.N. Potrap³⁸, C.J. Potter³², H. Potti¹, T. Poulsen⁴⁸, J. Poveda¹⁶², M.E. Pozo Astigarraga³⁶, A. Prades Ibanez¹⁶², M.M. Prapa⁴⁶, J. Pretel⁵⁴, D. Price¹⁰⁰, M. Primavera^{69a}, M.A. Principe Martin⁹⁸, R. Privara¹²¹, M.L. Proffitt¹³⁷, N. Proklova¹²⁷, K. Prokofiev^{64c}, G. Proto^{75a,75b}, S. Protopopescu²⁹, J. Proudfoot⁶, M. Przybycien^{84a}, J.E. Puddefoot¹³⁸, D. Pudzha³⁷, P. Puzo⁶⁶, D. Pyatiizbyantseva³⁷, J. Qian¹⁰⁵, D. Qichen¹⁰⁰, Y. Qin¹⁰⁰, T. Qiu⁹³, A. Quadrat⁵⁵,

- M. Queitsch-Maitland ¹⁰⁰, G. Quetant ⁵⁶, G. Rabanal Bolanos ⁶¹, D. Rafanoharana ⁵⁴, F. Ragusa ^{70a,70b}, J.L. Rainbolt ³⁹, J.A. Raine ⁵⁶, S. Rajagopalan ²⁹, E. Ramakoti ³⁷, K. Ran ^{48,14d}, N.P. Rapheeha ^{33g}, V. Raskina ¹²⁶, D.F. Rassloff ^{63a}, S. Rave ⁹⁹, B. Ravina ⁵⁵, I. Ravinovich ¹⁶⁸, M. Raymond ³⁶, A.L. Read ¹²⁴, N.P. Readioff ¹³⁸, D.M. Rebuzzi ^{72a,72b}, G. Redlinger ²⁹, K. Reeves ⁴⁵, J.A. Reidelsturz ¹⁷⁰, D. Reikher ¹⁵⁰, A. Reiss ⁹⁹, A. Rej ¹⁴⁰, C. Rembser ³⁶, A. Renardi ⁴⁸, M. Renda ^{27b}, M.B. Rendel ¹⁰⁹, F. Renner ⁴⁸, A.G. Rennie ⁵⁹, S. Resconi ^{70a}, M. Ressegotti ^{57b,57a}, E.D. Ressegueie ^{17a}, S. Rettie ³⁶, B. Reynolds ¹¹⁸, E. Reynolds ^{17a}, M. Rezaei Estabragh ¹⁷⁰, O.L. Rezanova ³⁷, P. Reznicek ¹³², E. Ricci ^{77a,77b}, R. Richter ¹⁰⁹, S. Richter ^{47a,47b}, E. Richter-Was ^{84b}, M. Ridel ¹²⁶, P. Rieck ¹¹⁶, P. Riedler ³⁶, M. Rijssenbeek ¹⁴⁴, A. Rimoldi ^{72a,72b}, M. Rimoldi ⁴⁸, L. Rinaldi ^{23b,23a}, T.T. Rinn ²⁹, M.P. Rinnagel ¹⁰⁸, G. Ripellino ¹⁴³, I. Riu ¹³, P. Rivadeneira ⁴⁸, J.C. Rivera Vergara ¹⁶⁴, F. Rizatdinova ¹²⁰, E. Rizvi ⁹³, C. Rizzi ⁵⁶, B.A. Roberts ¹⁶⁶, B.R. Roberts ^{17a}, S.H. Robertson ^{103,w}, M. Robin ⁴⁸, D. Robinson ³², C.M. Robles Gajardo ^{136f}, M. Robles Manzano ⁹⁹, A. Robson ⁵⁹, A. Rocchi ^{75a,75b}, C. Roda ^{73a,73b}, S. Rodriguez Bosca ^{63a}, Y. Rodriguez Garcia ^{22a}, A. Rodriguez Rodriguez ⁵⁴, A.M. Rodríguez Vera ^{155b}, S. Roe ³⁶, J.T. Roemer ¹⁵⁹, A.R. Roepe-Gier ¹¹⁹, J. Roggel ¹⁷⁰, O. Røhne ¹²⁴, R.A. Rojas ¹⁶⁴, B. Roland ⁵⁴, C.P.A. Roland ⁶⁷, J. Roloff ²⁹, A. Romanouk ³⁷, E. Romano ^{72a,72b}, M. Romano ^{23b}, A.C. Romero Hernandez ¹⁶¹, N. Rompotis ⁹¹, L. Roos ¹²⁶, S. Rosati ^{74a}, B.J. Rosser ³⁹, E. Rossi ⁴, E. Rossi ^{71a,71b}, L.P. Rossi ^{57b}, L. Rossini ⁴⁸, R. Rosten ¹¹⁸, M. Rotaru ^{27b}, B. Rottler ⁵⁴, D. Rousseau ⁶⁶, D. Rousso ³², G. Rovelli ^{72a,72b}, A. Roy ¹⁶¹, A. Rozanov ¹⁰¹, Y. Rozen ¹⁴⁹, X. Ruan ^{33g}, A. Rubio Jimenez ¹⁶², A.J. Ruby ⁹¹, V.H. Ruelas Rivera ¹⁸, T.A. Ruggeri ¹, F. Rühr ⁵⁴, A. Ruiz-Martinez ¹⁶², A. Rummler ³⁶, Z. Rurikova ⁵⁴, N.A. Rusakovich ³⁸, H.L. Russell ¹⁶⁴, J.P. Rutherford ⁷, K. Rybacki ⁹⁰, M. Rybar ¹³², E.B. Rye ¹²⁴, A. Ryzhov ³⁷, J.A. Sabater Iglesias ⁵⁶, P. Sabatini ¹⁶², L. Sabetta ^{74a,74b}, H.F-W. Sadrozinski ¹³⁵, F. Safai Tehrani ^{74a}, B. Safarzadeh Samani ¹⁴⁵, M. Safdari ¹⁴², S. Saha ¹⁰³, M. Sahinsoy ¹⁰⁹, M. Saimpert ¹³⁴, M. Saito ¹⁵², T. Saito ¹⁵², D. Salamani ³⁶, G. Salamanna ^{76a,76b}, A. Salnikov ¹⁴², J. Salt ¹⁶², A. Salvador Salas ¹³, D. Salvatore ^{43b,43a}, F. Salvatore ¹⁴⁵, A. Salzburger ³⁶, D. Sammel ⁵⁴, D. Sampsonidis ¹⁵¹, D. Sampsonidou ^{62d,62c}, J. Sánchez ¹⁶², A. Sanchez Pineda ⁴, V. Sanchez Sebastian ¹⁶², H. Sandaker ¹²⁴, C.O. Sander ⁴⁸, J.A. Sandesara ¹⁰², M. Sandhoff ¹⁷⁰, C. Sandoval ^{22b}, D.P.C. Sankey ¹³³, A. Sansoni ⁵³, L. Santi ^{74a,74b}, C. Santoni ⁴⁰, H. Santos ^{129a,129b}, S.N. Santpur ^{17a}, A. Santra ¹⁶⁸, K.A. Saoucha ¹³⁸, J.G. Saraiva ^{129a,129d}, J. Sardain ⁷, O. Sasaki ⁸², K. Sato ¹⁵⁶, C. Sauer ^{63b}, F. Sauerburger ⁵⁴, E. Sauvan ⁴, P. Savard ^{154,ad}, R. Sawada ¹⁵², C. Sawyer ¹³³, L. Sawyer ⁹⁶, I. Sayago Galvan ¹⁶², C. Sbarra ^{23b}, A. Sbrizzi ^{23b,23a}, T. Scanlon ⁹⁵, J. Schaarschmidt ¹³⁷, P. Schacht ¹⁰⁹, D. Schaefer ³⁹, U. Schäfer ⁹⁹, A.C. Schaffer ⁶⁶, D. Schaile ¹⁰⁸, R.D. Schamberger ¹⁴⁴, E. Schanet ¹⁰⁸, C. Scharf ¹⁸, M.M. Schefer ¹⁹, V.A. Schegelsky ³⁷, D. Scheirich ¹³², F. Schenck ¹⁸, M. Schernau ¹⁵⁹, C. Scheulen ⁵⁵, C. Schiavi ^{57b,57a}, Z.M. Schillaci ²⁶, E.J. Schioppa ^{69a,69b}, M. Schioppa ^{43b,43a}, B. Schlag ⁹⁹, K.E. Schleicher ⁵⁴, S. Schlenker ³⁶, J. Schmeing ¹⁷⁰, M.A. Schmidt ¹⁷⁰, K. Schmieden ⁹⁹, C. Schmitt ⁹⁹, S. Schmitt ⁴⁸, L. Schoeffel ¹³⁴, A. Schoening ^{63b}, P.G. Scholer ⁵⁴, E. Schopf ¹²⁵, M. Schott ⁹⁹, J. Schovancova ³⁶, S. Schramm ⁵⁶, F. Schroeder ¹⁷⁰, H-C. Schultz-Coulon ^{63a}, M. Schumacher ⁵⁴, B.A. Schumm ¹³⁵, Ph. Schune ¹³⁴, A. Schwartzman ¹⁴², T.A. Schwarz ¹⁰⁵, Ph. Schwemling ¹³⁴, R. Schwienhorst ¹⁰⁶, A. Sciandra ¹³⁵, G. Sciolla ²⁶, F. Scuri ^{73a}, F. Scutti ¹⁰⁴, C.D. Sebastiani ⁹¹, K. Sedlaczek ⁴⁹, P. Seema ¹⁸, S.C. Seidel ¹¹¹, A. Seiden ¹³⁵, B.D. Seidlitz ⁴¹, T. Seiss ³⁹, C. Seitz ⁴⁸, J.M. Seixas ^{81b}, G. Sekhniaidze ^{71a}, S.J. Sekula ⁴⁴, L. Selem ⁴, N. Semprini-Cesari ^{23b,23a}, S. Sen ⁵¹, D. Sengupta ⁵⁶, V. Senthilkumar ¹⁶², L. Serin ⁶⁶, L. Serkin ^{68a,68b}, M. Sessa ^{76a,76b}, H. Severini ¹¹⁹, S. Sevova ¹⁴², F. Sforza ^{57b,57a}, A. Sfyrla ⁵⁶, E. Shabalina ⁵⁵, R. Shaheen ¹⁴³, J.D. Shahinian ¹²⁷, D. Shaked Renous ¹⁶⁸, L.Y. Shan ^{14a}, M. Shapiro ^{17a}, A. Sharma ³⁶, A.S. Sharma ¹⁶³, P. Sharma ⁷⁹, S. Sharma ⁴⁸, P.B. Shatalov ³⁷, K. Shaw ¹⁴⁵, S.M. Shaw ¹⁰⁰, Q. Shen ^{62c,5}, P. Sherwood ⁹⁵, L. Shi ⁹⁵, C.O. Shimmin ¹⁷¹, Y. Shimogama ¹⁶⁷, J.D. Shinner ⁹⁴, I.P.J. Shipsey ¹²⁵, S. Shirabe ⁶⁰, M. Shiyakova ³⁸, J. Shlomi ¹⁶⁸, M.J. Shochet ³⁹, J. Shojaii ¹⁰⁴, D.R. Shope ¹²⁴, S. Shrestha ^{118,ah}, E.M. Shrif ^{33g}, M.J. Shroff ¹⁶⁴, P. Sicho ¹³⁰, A.M. Sickles ¹⁶¹, E. Sideras Haddad ^{33g}, A. Sidoti ^{23b}, F. Siegert ⁵⁰, Dj. Sijacki ¹⁵, R. Sikora ^{84a}, F. Sili ⁸⁹, J.M. Silva ²⁰, M.V. Silva Oliveira ³⁶, S.B. Silverstein ^{47a}, S. Simion ⁶⁶, R. Simoniello ³⁶, E.L. Simpson ⁵⁹, N.D. Simpson ⁹⁷, S. Simsek ^{21d}, S. Sindhu ⁵⁵, P. Sinervo ¹⁵⁴, V. Sinetckii ³⁷, S. Singh ¹⁴¹, S. Singh ¹⁵⁴, S. Sinha ⁴⁸, S. Sinha ^{33g}, M. Sioli ^{23b,23a}, I. Siral ³⁶, S.Yu. Sivoklokov ^{37,*}, J. Sjölin ^{47a,47b}, A. Skaf ⁵⁵, E. Skorda ⁹⁷, P. Skubic ¹¹⁹, M. Slawinska ⁸⁵, V. Smakhtin ¹⁶⁸, B.H. Smart ¹³³, J. Smiesko ³⁶, S.Yu. Smirnov ³⁷, Y. Smirnov ³⁷, L.N. Smirnova ^{37,a}, O. Smirnova ⁹⁷, A.C. Smith ⁴¹, E.A. Smith ³⁹, H.A. Smith ¹²⁵, J.L. Smith ⁹¹, R. Smith ¹⁴², M. Smizanska ⁹⁰, K. Smolek ¹³¹, A. Smykiewicz ⁸⁵, A.A. Snesarev ³⁷, H.L. Snoek ¹¹³, S. Snyder ²⁹, R. Sobie ^{164,w}, A. Soffer ¹⁵⁰, C.A. Solans Sanchez ³⁶,

- E.Yu. Soldatov ³⁷, U. Soldevila ¹⁶², A.A. Solodkov ³⁷, S. Solomon ⁵⁴, A. Soloshenko ³⁸, K. Solovieva ⁵⁴, O.V. Solovyanov ³⁷, V. Solovyev ³⁷, P. Sommer ³⁶, A. Sonay ¹³, W.Y. Song ^{155b}, A. Sopczak ¹³¹, A.L. Sopio ⁹⁵, F. Sopkova ^{28b}, V. Sothilingam ^{63a}, S. Sottocornola ^{72a,72b}, R. Soualah ^{115b}, Z. Soumaimi ^{35e}, D. South ⁴⁸, S. Spagnolo ^{69a,69b}, M. Spalla ¹⁰⁹, F. Spanò ⁹⁴, D. Sperlich ⁵⁴, G. Spigo ³⁶, M. Spina ¹⁴⁵, S. Spinali ⁹⁰, D.P. Spiteri ⁵⁹, M. Spousta ¹³², E.J. Staats ³⁴, A. Stabile ^{70a,70b}, R. Stamen ^{63a}, M. Stamenkovic ¹¹³, A. Stampeki ²⁰, M. Standke ²⁴, E. Stanecka ⁸⁵, M.V. Stange ⁵⁰, B. Stanislaus ^{17a}, M.M. Stanitzki ⁴⁸, M. Stankaityte ¹²⁵, B. Stapf ⁴⁸, E.A. Starchenko ³⁷, G.H. Stark ¹³⁵, J. Stark ¹⁰¹, D.M. Starko ^{155b}, P. Staroba ¹³⁰, P. Starovoitov ^{63a}, S. Stärz ¹⁰³, R. Staszewski ⁸⁵, G. Stavropoulos ⁴⁶, J. Steentoft ¹⁶⁰, P. Steinberg ²⁹, A.L. Steinhebel ¹²², B. Stelzer ^{141,155a}, H.J. Stelzer ¹²⁸, O. Stelzer-Chilton ^{155a}, H. Stenzel ⁵⁸, T.J. Stevenson ¹⁴⁵, G.A. Stewart ³⁶, M.C. Stockton ³⁶, G. Stoicea ^{27b}, M. Stolarski ^{129a}, S. Stonjek ¹⁰⁹, A. Straessner ⁵⁰, J. Strandberg ¹⁴³, S. Strandberg ^{47a,47b}, M. Strauss ¹¹⁹, T. Strebler ¹⁰¹, P. Strizenec ^{28b}, R. Ströhmer ¹⁶⁵, D.M. Strom ¹²², L.R. Strom ⁴⁸, R. Stroynowski ⁴⁴, A. Strubig ^{47a,47b}, S.A. Stucci ²⁹, B. Stugu ¹⁶, J. Stupak ¹¹⁹, N.A. Styles ⁴⁸, D. Su ¹⁴², S. Su ^{62a}, W. Su ^{62d,137,62c}, X. Su ^{62a,66}, K. Sugizaki ¹⁵², V.V. Sulin ³⁷, M.J. Sullivan ⁹¹, D.M.S. Sultan ^{77a,77b}, L. Sultanaliyeva ³⁷, S. Sultansoy ^{3b}, T. Sumida ⁸⁶, S. Sun ¹⁰⁵, S. Sun ¹⁶⁹, O. Sunneborn Gudnadottir ¹⁶⁰, M.R. Sutton ¹⁴⁵, M. Svatos ¹³⁰, M. Swiatlowski ^{155a}, T. Swirski ¹⁶⁵, I. Sykora ^{28a}, M. Sykora ¹³², T. Sykora ¹³², D. Ta ⁹⁹, K. Tackmann ^{48,v}, A. Taffard ¹⁵⁹, R. Tafirout ^{155a}, J.S. Tafoya Vargas ⁶⁶, R.H.M. Taibah ¹²⁶, R. Takashima ⁸⁷, K. Takeda ⁸³, E.P. Takeva ⁵², Y. Takubo ⁸², M. Talby ¹⁰¹, A.A. Talyshев ³⁷, K.C. Tam ^{64b}, N.M. Tamir ¹⁵⁰, A. Tanaka ¹⁵², J. Tanaka ¹⁵², R. Tanaka ⁶⁶, M. Tanasini ^{57b,57a}, J. Tang ^{62c}, Z. Tao ¹⁶³, S. Tapia Araya ⁸⁰, S. Tapprogge ⁹⁹, A. Tarek Abouelfadl Mohamed ¹⁰⁶, S. Tarem ¹⁴⁹, K. Tariq ^{62b}, G. Tarna ^{101,27b}, G.F. Tartarelli ^{70a}, P. Tas ¹³², M. Tasevsky ¹³⁰, E. Tassi ^{43b,43a}, A.C. Tate ¹⁶¹, G. Tateno ¹⁵², Y. Tayalati ^{35e}, G.N. Taylor ¹⁰⁴, W. Taylor ^{155b}, H. Teagle ⁹¹, A.S. Tee ¹⁶⁹, R. Teixeira De Lima ¹⁴², P. Teixeira-Dias ⁹⁴, J.J. Teoh ¹⁵⁴, K. Terashi ¹⁵², J. Terron ⁹⁸, S. Terzo ¹³, M. Testa ⁵³, R.J. Teuscher ^{154,w}, A. Thaler ⁷⁸, O. Theiner ⁵⁶, N. Themistokleous ⁵², T. Theveneaux-Pelzer ¹⁸, O. Thielmann ¹⁷⁰, D.W. Thomas ⁹⁴, J.P. Thomas ²⁰, E.A. Thompson ⁴⁸, P.D. Thompson ²⁰, E. Thomson ¹²⁷, E.J. Thorpe ⁹³, Y. Tian ⁵⁵, V. Tikhomirov ^{37,a}, Yu.A. Tikhonov ³⁷, S. Timoshenko ³⁷, E.X.L. Ting ¹, P. Tipton ¹⁷¹, S. Tisserant ¹⁰¹, S.H. Tlou ^{33g}, A. Tnourji ⁴⁰, K. Todome ^{23b,23a}, S. Todorova-Nova ¹³², S. Todt ⁵⁰, M. Togawa ⁸², J. Tojo ⁸⁸, S. Tokár ^{28a}, K. Tokushuku ⁸², R. Tombs ³², M. Tomoto ^{82,110}, L. Tompkins ¹⁴², K.W. Topolnicki ^{84b}, P. Tornambe ¹⁰², E. Torrence ¹²², H. Torres ⁵⁰, E. Torró Pastor ¹⁶², M. Toscani ³⁰, C. Tosciri ³⁹, M. Tost ¹¹, D.R. Tovey ¹³⁸, A. Traeet ¹⁶, I.S. Trandafir ^{27b}, T. Trefzger ¹⁶⁵, A. Tricoli ²⁹, I.M. Trigger ^{155a}, S. Trincaz-Duvoud ¹²⁶, D.A. Trischuk ²⁶, B. Trocmé ⁶⁰, A. Trofymov ⁶⁶, C. Troncon ^{70a}, L. Truong ^{33c}, M. Trzebinski ⁸⁵, A. Trzupek ⁸⁵, F. Tsai ¹⁴⁴, M. Tsai ¹⁰⁵, A. Tsiamis ¹⁵¹, P.V. Tsiareshka ³⁷, S. Tsigaridas ^{155a}, A. Tsirigotis ^{151,t}, V. Tsiskaridze ¹⁴⁴, E.G. Tskhadadze ^{148a}, M. Tsopoulou ¹⁵¹, Y. Tsujikawa ⁸⁶, I.I. Tsukerman ³⁷, V. Tsulaia ^{17a}, S. Tsuno ⁸², O. Tsur ¹⁴⁹, D. Tsybychev ¹⁴⁴, Y. Tu ^{64b}, A. Tudorache ^{27b}, V. Tudorache ^{27b}, A.N. Tuna ³⁶, S. Turchikhin ³⁸, I. Turk Cakir ^{3a}, R. Turra ^{70a}, T. Turtuvshin ^{38,x}, P.M. Tuts ⁴¹, S. Tzamarias ¹⁵¹, P. Tzanis ¹⁰, E. Tzovara ⁹⁹, K. Uchida ¹⁵², F. Ukegawa ¹⁵⁶, P.A. Ulloa Poblete ^{136c}, G. Unal ³⁶, M. Unal ¹¹, A. Undrus ²⁹, G. Unel ¹⁵⁹, J. Urban ^{28b}, P. Urquijo ¹⁰⁴, G. Usai ⁸, R. Ushioda ¹⁵³, M. Usman ¹⁰⁷, Z. Uysal ^{21b}, L. Vacavant ¹⁰¹, V. Vacek ¹³¹, B. Vachon ¹⁰³, K.O.H. Vadla ¹²⁴, T. Vafeiadis ³⁶, A. Vaitkus ⁹⁵, C. Valderanis ¹⁰⁸, E. Valdes Santurio ^{47a,47b}, M. Valente ^{155a}, S. Valentinetti ^{23b,23a}, A. Valero ¹⁶², A. Vallier ¹⁰¹, J.A. Valls Ferrer ¹⁶², T.R. Van Daalen ¹³⁷, P. Van Gemmeren ⁶, M. Van Rijnbach ^{124,36}, S. Van Stroud ⁹⁵, I. Van Vulpen ¹¹³, M. Vanadia ^{75a,75b}, W. Vandelli ³⁶, M. Vandebroucke ¹³⁴, E.R. Vandewall ¹²⁰, D. Vannicola ¹⁵⁰, L. Vannoli ^{57b,57a}, R. Vari ^{74a}, E.W. Varnes ⁷, C. Varni ^{17a}, T. Varol ¹⁴⁷, D. Varouchas ⁶⁶, L. Varriale ¹⁶², K.E. Varvell ¹⁴⁶, M.E. Vasile ^{27b}, L. Vaslin ⁴⁰, G.A. Vasquez ¹⁶⁴, F. Vazeille ⁴⁰, T. Vazquez Schroeder ³⁶, J. Veatch ³¹, V. Vecchio ¹⁰⁰, M.J. Veen ¹⁰², I. Velisek ¹²⁵, L.M. Veloce ¹⁵⁴, F. Veloso ^{129a,129c}, S. Veneziano ^{74a}, A. Ventura ^{69a,69b}, A. Verbytskyi ¹⁰⁹, M. Verducci ^{73a,73b}, C. Vergis ²⁴, M. Verissimo De Araujo ^{81b}, W. Verkerke ¹¹³, J.C. Vermeulen ¹¹³, C. Vernieri ¹⁴², P.J. Verschueren ⁹⁴, M. Vessella ¹⁰², M.C. Vetterli ^{141,ad}, A. Vgenopoulos ¹⁵¹, N. Viaux Maira ^{136f}, T. Vickey ¹³⁸, O.E. Vickey Boeriu ¹³⁸, G.H.A. Viehhauser ¹²⁵, L. Vigani ^{63b}, M. Villa ^{23b,23a}, M. Villaplana Perez ¹⁶², E.M. Villhauer ⁵², E. Vilucchi ⁵³, M.G. Vincter ³⁴, G.S. Virdee ²⁰, A. Vishwakarma ⁵², C. Vittori ^{23b,23a}, I. Vivarelli ¹⁴⁵, V. Vladimirov ¹⁶⁶, E. Voevodina ¹⁰⁹, F. Vogel ¹⁰⁸, P. Vokac ¹³¹, J. Von Ahnen ⁴⁸, E. Von Toerne ²⁴, B. Vormwald ³⁶, V. Vorobel ¹³², K. Vorobev ³⁷, M. Vos ¹⁶², J.H. Vossebeld ⁹¹, M. Vozak ¹¹³, L. Vozdecky ⁹³, N. Vranjes ¹⁵, M. Vranjes Milosavljevic ¹⁵, M. Vreeswijk ¹¹³, R. Vuillermet ³⁶, O. Vujinovic ⁹⁹, I. Vukotic ³⁹, S. Wada ¹⁵⁶, C. Wagner ¹⁰², W. Wagner ¹⁷⁰, S. Wahdan ¹⁷⁰, H. Wahlberg ⁸⁹,

R. Wakasa¹⁵⁶, M. Wakida¹¹⁰, V.M. Walbrecht¹⁰⁹, J. Walder¹³³, R. Walker¹⁰⁸, W. Walkowiak¹⁴⁰, A.M. Wang⁶¹, A.Z. Wang¹⁶⁹, C. Wang^{62a}, C. Wang^{62c}, H. Wang^{17a}, J. Wang^{64a}, R.-J. Wang⁹⁹, R. Wang⁶¹, R. Wang⁶, S.M. Wang¹⁴⁷, S. Wang^{62b}, T. Wang^{62a}, W.T. Wang⁷⁹, X. Wang^{14c}, X. Wang¹⁶¹, X. Wang^{62c}, Y. Wang^{62d}, Y. Wang^{14c}, Z. Wang¹⁰⁵, Z. Wang^{62d,51,62c}, Z. Wang¹⁰⁵, A. Warburton¹⁰³, R.J. Ward²⁰, N. Warrack⁵⁹, A.T. Watson²⁰, H. Watson⁵⁹, M.F. Watson²⁰, G. Watts¹³⁷, B.M. Waugh⁹⁵, A.F. Webb¹¹, C. Weber²⁹, H.A. Weber¹⁸, M.S. Weber¹⁹, S.M. Weber^{63a}, C. Wei^{62a}, Y. Wei¹²⁵, A.R. Weidberg¹²⁵, J. Weingarten⁴⁹, M. Weirich⁹⁹, C. Weiser⁵⁴, C.J. Wells⁴⁸, T. Wenaus²⁹, B. Wendland⁴⁹, T. Wengler³⁶, N.S. Wenke¹⁰⁹, N. Wermes²⁴, M. Wessels^{63a}, K. Whalen¹²², A.M. Wharton⁹⁰, A.S. White⁶¹, A. White⁸, M.J. White¹, D. Whiteson¹⁵⁹, L. Wickremasinghe¹²³, W. Wiedenmann¹⁶⁹, C. Wiel⁵⁰, M. Wieters¹³³, N. Wieseotte⁹⁹, C. Wiglesworth⁴², L.A.M. Wiik-Fuchs⁵⁴, D.J. Wilbern¹¹⁹, H.G. Wilkens³⁶, D.M. Williams⁴¹, H.H. Williams¹²⁷, S. Williams³², S. Willocq¹⁰², P.J. Windischhofer¹²⁵, F. Winklmeier¹²², B.T. Winter⁵⁴, J.K. Winter¹⁰⁰, M. Wittgen¹⁴², M. Wobisch⁹⁶, R. Wölker¹²⁵, J. Wollrath¹⁵⁹, M.W. Wolter⁸⁵, H. Wolters^{129a,129c}, V.W.S. Wong¹⁶³, A.F. Wongel⁴⁸, S.D. Worm⁴⁸, B.K. Wosiek⁸⁵, K.W. Woźniak⁸⁵, K. Wraight⁵⁹, J. Wu^{14a,14d}, M. Wu^{64a}, M. Wu¹¹², S.L. Wu¹⁶⁹, X. Wu⁵⁶, Y. Wu^{62a}, Z. Wu^{134,62a}, J. Wuerzinger¹²⁵, T.R. Wyatt¹⁰⁰, B.M. Wynne⁵², S. Xella⁴², L. Xia^{14c}, M. Xia^{14b}, J. Xiang^{64c}, X. Xiao¹⁰⁵, M. Xie^{62a}, X. Xie^{62a}, S. Xin^{14a,14d}, J. Xiong^{17a}, I. Xiotidis¹⁴⁵, D. Xu^{14a}, H. Xu^{62a}, H. Xu^{62a}, L. Xu^{62a}, R. Xu¹²⁷, T. Xu¹⁰⁵, W. Xu¹⁰⁵, Y. Xu^{14b}, Z. Xu^{62b}, Z. Xu^{14a}, B. Yabsley¹⁴⁶, S. Yacoob^{33a}, N. Yamaguchi⁸⁸, Y. Yamaguchi¹⁵³, H. Yamauchi¹⁵⁶, T. Yamazaki^{17a}, Y. Yamazaki⁸³, J. Yan^{62c}, S. Yan¹²⁵, Z. Yan²⁵, H.J. Yang^{62c,62d}, H.T. Yang^{62a}, S. Yang^{62a}, T. Yang^{64c}, X. Yang^{62a}, X. Yang^{14a}, Y. Yang⁴⁴, Z. Yang^{62a,105}, W.-M. Yao^{17a}, Y.C. Yap⁴⁸, H. Ye^{14c}, H. Ye⁵⁵, J. Ye⁴⁴, S. Ye²⁹, X. Ye^{62a}, Y. Yeh⁹⁵, I. Yeletskikh³⁸, B.K. Yeo^{17a}, M.R. Yexley⁹⁰, P. Yin⁴¹, K. Yorita¹⁶⁷, S. Younas^{27b}, C.J.S. Young⁵⁴, C. Young¹⁴², M. Yuan¹⁰⁵, R. Yuan^{62b,k}, L. Yue⁹⁵, X. Yue^{63a}, M. Zaazoua^{35e}, B. Zabinski⁸⁵, E. Zaid⁵², T. Zakareishvili^{148b}, N. Zakharchuk³⁴, S. Zambito⁵⁶, J.A. Zamora Saa^{136d,136b}, J. Zang¹⁵², D. Zanzi⁵⁴, O. Zaplatilek¹³¹, S.V. Zeißner⁴⁹, C. Zeitnitz¹⁷⁰, J.C. Zeng¹⁶¹, D.T. Zenger Jr²⁶, O. Zenin³⁷, T. Ženiš^{28a}, S. Zenz⁹³, S. Zerradi^{35a}, D. Zerwas⁶⁶, B. Zhang^{14c}, D.F. Zhang¹³⁸, G. Zhang^{14b}, J. Zhang^{62b}, J. Zhang⁶, K. Zhang^{14a,14d}, L. Zhang^{14c}, P. Zhang^{14a,14d}, R. Zhang¹⁶⁹, S. Zhang¹⁰⁵, T. Zhang¹⁵², X. Zhang^{62c}, X. Zhang^{62b}, Y. Zhang^{62c,5}, Z. Zhang^{17a}, Z. Zhang⁶⁶, H. Zhao¹³⁷, P. Zhao⁵¹, T. Zhao^{62b}, Y. Zhao¹³⁵, Z. Zhao^{62a}, A. Zhemchugov³⁸, X. Zheng^{62a}, Z. Zheng¹⁴², D. Zhong¹⁶¹, B. Zhou¹⁰⁵, C. Zhou¹⁶⁹, H. Zhou⁷, N. Zhou^{62c}, Y. Zhou⁷, C.G. Zhu^{62b}, C. Zhu^{14a,14d}, H.L. Zhu^{62a}, H. Zhu^{14a}, J. Zhu¹⁰⁵, Y. Zhu^{62c}, Y. Zhu^{62a}, X. Zhuang^{14a}, K. Zhukov³⁷, V. Zhulanov³⁷, N.I. Zimine³⁸, J. Zinsser^{63b}, M. Ziolkowski¹⁴⁰, L. Živković¹⁵, A. Zoccoli^{23b,23a}, K. Zoch⁵⁶, T.G. Zorbas¹³⁸, O. Zormpa⁴⁶, W. Zou⁴¹, L. Zwalski³⁶

¹ Department of Physics, University of Adelaide, Adelaide; Australia² Department of Physics, University of Alberta, Edmonton AB; Canada³ (a) Department of Physics, Ankara University, Ankara; (b) Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye⁴ LAPP, Univ. Savoie Mont Blanc, CNRS/IN2P3, Annecy; France⁵ APC, Université Paris Cité, CNRS/IN2P3, Paris; France⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America⁷ Department of Physics, University of Arizona, Tucson AZ; United States of America⁸ Department of Physics, University of Texas at Arlington, Arlington TX; United States of America⁹ Physics Department, National and Kapodistrian University of Athens, Athens; Greece¹⁰ Physics Department, National Technical University of Athens, Zografou; Greece¹¹ Department of Physics, University of Texas at Austin, Austin TX; United States of America¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain¹⁴ (a) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; (b) Physics Department, Tsinghua University, Beijing; (c) Department of Physics, Nanjing University, Nanjing;¹⁵ University of Chinese Academy of Science (UCAS), Beijing; China¹⁶ Institute of Physics, University of Belgrade, Belgrade; Serbia¹⁷ Department for Physics and Technology, University of Bergen, Bergen; Norway¹⁸ (a) Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; (b) University of California, Berkeley CA; United States of America¹⁹ Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany²⁰ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland²¹ School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom²² (a) Department of Physics, Bogazici University, Istanbul; (b) Department of Physics Engineering, Gaziantep University, Gaziantep; (c) Department of Physics, Istanbul University, Istanbul; (d) İstinye University, Sarıyer, İstanbul; Türkiye²³ (a) Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; (b) Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia²⁴ Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; (b) INFN Sezione di Bologna; Italy²⁵ Physikalisch-Technische Bundesanstalt, Braunschweig; Germany²⁶ Department of Physics, Boston University, Boston MA; United States of America²⁷ (a) Transilvania University of Brasov, Brasov; (b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; (c) Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; (d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; (e) University Politehnica Bucharest, Bucharest; (f) West University in Timisoara, Timisoara; (g) Faculty of Physics, University of Bucharest, Bucharest; Romania

- 28 ^(a) Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; ^(b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic
- 29 Physics Department, Brookhaven National Laboratory, Upton NY; United States of America
- 30 Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina
- 31 California State University, CA; United States of America
- 32 Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom
- 33 ^(a) Department of Physics, University of Cape Town, Cape Town; ^(b) iThemba Labs, Western Cape; ^(c) Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg; ^(d) National Institute of Physics, University of the Philippines Diliman (Philippines); ^(e) University of South Africa, Department of Physics, Pretoria; ^(f) University of Zululand, KwaDlangezwa; ^(g) School of Physics, University of the Witwatersrand, Johannesburg; South Africa
- 34 Department of Physics, Carleton University, Ottawa ON; Canada
- 35 ^(a) Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies - Université Hassan II, Casablanca; ^(b) Faculté des Sciences, Université Ibn-Tofail, Kénitra; ^(c) Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA, Marrakech; ^(d) LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda; ^(e) Faculté des sciences, Université Mohammed V, Rabat; ^(f) Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- 36 CERN, Geneva; Switzerland
- 37 Affiliated with an institute covered by a cooperation agreement with CERN
- 38 Affiliated with an international laboratory covered by a cooperation agreement with CERN
- 39 Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America
- 40 LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France
- 41 Nevis Laboratory, Columbia University, Irvington NY; United States of America
- 42 Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark
- 43 ^(a) Dipartimento di Fisica, Università della Calabria, Rende; ^(b) INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy
- 44 Physics Department, Southern Methodist University, Dallas TX; United States of America
- 45 Physics Department, University of Texas at Dallas, Richardson TX; United States of America
- 46 National Centre for Scientific Research "Demokritos", Agia Paraskevi; Greece
- 47 ^(a) Department of Physics, Stockholm University; ^(b) Oskar Klein Centre, Stockholm; Sweden
- 48 Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany
- 49 Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany
- 50 Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany
- 51 Department of Physics, Duke University, Durham NC; United States of America
- 52 SUPA - School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom
- 53 INFN e Laboratori Nazionali di Frascati, Frascati; Italy
- 54 Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany
- 55 II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany
- 56 Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland
- 57 ^(a) Dipartimento di Fisica, Università di Genova, Genova; ^(b) INFN Sezione di Genova; Italy
- 58 II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany
- 59 SUPA - School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom
- 60 LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France
- 61 Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America
- 62 ^(a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; ^(b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; ^(c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(d) Tsung-Dao Lee Institute, Shanghai; China
- 63 ^(a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; ^(b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany
- 64 ^(a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; ^(b) Department of Physics, University of Hong Kong, Hong Kong; ^(c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China
- 65 Department of Physics, National Tsing Hua University, Hsinchu; Taiwan
- 66 IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France
- 67 Department of Physics, Indiana University, Bloomington IN; United States of America
- 68 ^(a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; ^(b) ICTP, Trieste; ^(c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy
- 69 ^(a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy
- 70 ^(a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano; Italy
- 71 ^(a) INFN Sezione di Napoli; ^(b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy
- 72 ^(a) INFN Sezione di Pavia; ^(b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy
- 73 ^(a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy
- 74 ^(a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy
- 75 ^(a) INFN Sezione di Roma Tor Vergata; ^(b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy
- 76 ^(a) INFN Sezione di Roma Tre; ^(b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy
- 77 ^(a) INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy
- 78 University Innsbruck, Department of Astroparticle Physics, Innsbruck; Austria
- 79 University of Iowa, Iowa City IA; United States of America
- 80 Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America
- 81 ^(a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; Brazil
- 82 KEK, High Energy Accelerator Research Organization, Tsukuba; Japan
- 83 Graduate School of Science, Kobe University, Kobe; Japan
- 84 ^(a) AGH University of Science and Technology, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland
- 85 Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland
- 86 Faculty of Science, Kyoto University, Kyoto; Japan
- 87 Kyoto University of Education, Kyoto; Japan
- 88 Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka; Japan
- 89 Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina
- 90 Physics Department, Lancaster University, Lancaster; United Kingdom
- 91 Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom
- 92 Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia
- 93 School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom
- 94 Department of Physics, Royal Holloway University of London, Egham; United Kingdom
- 95 Department of Physics and Astronomy, University College London, London; United Kingdom
- 96 Louisiana Tech University, Ruston LA; United States of America
- 97 Fysiska institutionen, Lunds universitet, Lund; Sweden

- 98 Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain
 99 Institut für Physik, Universität Mainz, Mainz; Germany
 100 School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom
 101 CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France
 102 Department of Physics, University of Massachusetts, Amherst MA; United States of America
 103 Department of Physics, McGill University, Montreal QC; Canada
 104 School of Physics, University of Melbourne, Victoria; Australia
 105 Department of Physics, University of Michigan, Ann Arbor MI; United States of America
 106 Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
 107 Group of Particle Physics, University of Montreal, Montreal QC; Canada
 108 Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany
 109 Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany
 110 Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan
 111 Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America
 112 Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands
 113 Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands
 114 Department of Physics, Northern Illinois University, DeKalb IL; United States of America
 115 ^(a) New York University Abu Dhabi, Abu Dhabi; ^(b) University of Sharjah, Sharjah; United Arab Emirates
 116 Department of Physics, New York University, New York NY; United States of America
 117 Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan
 118 Ohio State University, Columbus OH; United States of America
 119 Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America
 120 Department of Physics, Oklahoma State University, Stillwater OK; United States of America
 121 Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic
 122 Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America
 123 Graduate School of Science, Osaka University, Osaka; Japan
 124 Department of Physics, University of Oslo, Oslo; Norway
 125 Department of Physics, Oxford University, Oxford; United Kingdom
 126 LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France
 127 Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America
 128 Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America
 129 ^(a) Laboratório de Instrumentação e Física Experimental de Partículas - LIP, Lisboa; ^(b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; ^(c) Departamento de Física, Universidade de Coimbra, Coimbra; ^(d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; ^(e) Departamento de Física, Universidade do Minho, Braga; ^(f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); ^(g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal
 130 Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic
 131 Czech Technical University in Prague, Prague; Czech Republic
 132 Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic
 133 Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom
 134 IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France
 135 Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America
 136 ^(a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b) Millennium Institute for Subatomic physics at high energy frontier (SAPIR), Santiago; ^(c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; ^(d) Universidad Andres Bello, Department of Physics, Santiago; ^(e) Instituto de Alta Investigación, Universidad de Tarapacá, Arica; ^(f) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile
 137 Department of Physics, University of Washington, Seattle WA; United States of America
 138 Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom
 139 Department of Physics, Shinshu University, Nagano; Japan
 140 Department Physik, Universität Siegen, Siegen; Germany
 141 Department of Physics, Simon Fraser University, Burnaby BC; Canada
 142 SLAC National Accelerator Laboratory, Stanford CA; United States of America
 143 Department of Physics, Royal Institute of Technology, Stockholm; Sweden
 144 Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America
 145 Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom
 146 School of Physics, University of Sydney, Sydney; Australia
 147 Institute of Physics, Academia Sinica, Taipei; Taiwan
 148 ^(a) E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi; ^(b) High Energy Physics Institute, Tbilisi State University, Tbilisi; ^(c) University of Georgia, Tbilisi; Georgia
 149 Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel
 150 Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel
 151 Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece
 152 International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan
 153 Department of Physics, Tokyo Institute of Technology, Tokyo; Japan
 154 Department of Physics, University of Toronto, Toronto ON; Canada
 155 ^(a) TRIUMF, Vancouver BC; ^(b) Department of Physics and Astronomy, York University, Toronto ON; Canada
 156 Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan
 157 Department of Physics and Astronomy, Tufts University, Medford MA; United States of America
 158 United Arab Emirates University, Al Ain; United Arab Emirates
 159 Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America
 160 Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden
 161 Department of Physics, University of Illinois, Urbana IL; United States of America
 162 Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia - CSIC, Valencia; Spain
 163 Department of Physics, University of British Columbia, Vancouver BC; Canada
 164 Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada
 165 Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany
 166 Department of Physics, University of Warwick, Coventry; United Kingdom
 167 Waseda University, Tokyo; Japan
 168 Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel
 169 Department of Physics, University of Wisconsin, Madison WI; United States of America
 170 Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany
 171 Department of Physics, Yale University, New Haven CT; United States of America

- ^a Also Affiliated with an institute covered by a cooperation agreement with CERN.
- ^b Also at An-Najah National University, Nablus; Palestine.
- ^c Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America.
- ^d Also at Bruno Kessler Foundation, Trento; Italy.
- ^e Also at Center for High Energy Physics, Peking University; China.
- ^f Also at Centro Studi e Ricerche Enrico Fermi; Italy.
- ^g Also at CERN, Geneva; Switzerland.
- ^h Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland.
- ⁱ Also at Departament de Fisica de la Universitat Autònoma de Barcelona, Barcelona; Spain.
- ^j Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece.
- ^k Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America.
- ^l Also at Department of Physics and Astronomy, University of Louisville, Louisville, KY; United States of America.
- ^m Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel.
- ⁿ Also at Department of Physics, California State University, East Bay; United States of America.
- ^o Also at Department of Physics, California State University, Sacramento; United States of America.
- ^p Also at Department of Physics, King's College London, London; United Kingdom.
- ^q Also at Department of Physics, University of Fribourg, Fribourg; Switzerland.
- ^r Also at Department of Physics, University of Thessaly; Greece.
- ^s Also at Department of Physics, Westmont College, Santa Barbara; United States of America.
- ^t Also at Hellenic Open University, Patras; Greece.
- ^u Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain.
- ^v Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany.
- ^w Also at Institute of Particle Physics (IPP); Canada.
- ^x Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia.
- ^y Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan.
- ^z Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia.
- ^{aa} Also at Lawrence Livermore National Laboratory, Livermore; United States of America.
- ^{ab} Also at RWTH Aachen University, III. Physikalischs Institut A, Aachen; Germany.
- ^{ac} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China.
- ^{ad} Also at TRIUMF, Vancouver BC; Canada.
- ^{ae} Also at Università di Napoli Parthenope, Napoli; Italy.
- ^{af} Also at University of Chinese Academy of Sciences (UCAS), Beijing; China.
- ^{ag} Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America.
- ^{ah} Also at Washington College, Maryland; United States of America.
- ^{ai} Also at Yeditepe University, Physics Department, İstanbul; Türkiye.

* Deceased.