

# Vaccination protects endangered albatross chicks against avian cholera

Vincent Bourret<sup>1</sup> | Amandine Gamble<sup>1</sup> | Jérémy Tornos<sup>1</sup> | Audrey Jaeger<sup>2</sup> | Karine Delord<sup>3</sup> |  
Christophe Barbraud<sup>3</sup> | Pablo Tortosa<sup>4</sup> | Sarah Kada<sup>1</sup> | Jean-Baptiste Thiebot<sup>5,6</sup> |  
Eric Thibault<sup>7</sup> | Hubert Gantelet<sup>7</sup> | Henri Weimerskirch<sup>3</sup> | Romain Garnier<sup>8</sup> |  
Thierry Boulinier<sup>1</sup>

<sup>1</sup>UMR 5175 CEFE, CNRS-Université de Montpellier, Montpellier, France

<sup>2</sup>Université de La Réunion, UMR ENTROPIE, UR-IRD-CNRS, Saint Denis, La Réunion, France

<sup>3</sup>UMR 7372 CEBC, CNRS-Université de La Rochelle, Villiers-en-Bois, France

<sup>4</sup>Université de La Réunion, UMR PIMIT CNRS 9192-INSERM 1187-IRD 249, GIP CYROI, Saint Denis, La Réunion, France

<sup>5</sup>Réserve Naturelle Nationale des Terres Australes Françaises, TAAF, Saint Pierre, La Réunion, France

<sup>6</sup>National Institute of Polar Research, 10–3 Midori-cho, Tachikawa, Tokyo, Japan

<sup>7</sup>Ceva Biovac Santé Animale, Beaucouzé, France

<sup>8</sup>Department of Veterinary Medicine, Disease Dynamics Unit, University of Cambridge, Cambridge, United Kingdom

## Correspondence

Vincent Bourret, UMR 5175 CEFE, CNRS-  
Université de Montpellier, Montpellier,  
France.

Email: vincent.bourret@cantab.net

## Funding information

Réserve Naturelle Nationale des Terres Aus-  
trales Françaises; Institut Polaire Français  
Paul Emile Victor, Grant/Award Numbers:  
prg. no. 109, 1151; Labex CeMEB; Min-  
istère de l'Enseignement Supérieur et de la  
Recherche; Agence Nationale de la Recherche,  
Grant/Award Number: 'EVEMATA' 11 BSV7  
003; Zone Atelier Antarctique

## Abstract

Global change is contributing to unprecedented expansions of infectious diseases in wildlife. Recurrent avian cholera outbreaks are causing dramatic chick mortality and population decline in endangered albatross colonies on Amsterdam Island, a critical seabird breeding ground in the Southern Indian Ocean. We manufactured a killed vaccine using a *Pasteurella multocida* strain isolated from a dead albatross in the field. We used this same bacterial strain to establish a serological assay allowing the monitoring of antibody levels following bird vaccination. Using this vaccine on chicks 2 weeks posthatching caused 100% seroconversion and reduced the death risk by a factor exceeding 2.5, raising fledging probability from 14% to 46%. These results suggest that using a specifically tailored vaccine could be a key tool to effectively protect endangered seabirds from disease outbreaks threatening them with extinction.

## KEYWORDS

Amsterdam albatross, antibody dynamics, autogenous vaccine, avian cholera, dark-mantled sooty albatross, emerging infectious diseases, Indian yellow-nosed albatross, *Pasteurella multocida*, seabird conservation

## 1 | INTRODUCTION

Emerging infectious diseases are a growing concern for both public health and wildlife conservation (Daszak,

Cunningham, & Hyatt, 2000; Smith, Acevedo-Whitehouse, & Pedersen, 2009). The globally connected marine environment is especially sensitive to environmental change, and oceanic ecosystems are increasingly suspected to be “incubators and

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. Conservation Letters published by Wiley Periodicals, Inc.



**FIGURE 1** Adult Indian yellow-nosed albatross over dead chick at the Entrecasteaux cliffs on Amsterdam Island (37.8°S, 77.6°E) where recurrent massive chick mortality events have been occurring. Photo by Thierry Boulonier

conveyors” of diseases (Harvell et al., 1999, 2002). Understanding and managing infectious diseases in marine ecosystems is thus a current priority (Lafferty & Hofmann, 2016), especially at higher latitudes which include some of the places most dramatically impacted by climate change (Van Hemert, Pearce, & Handel, 2014).

Colonial seabirds are particularly sensitive to changes in their local, regional, and global environments (Croxall, Trathan, & Murphy, 2002) and infectious disease outbreaks. Given the high mortalities it provokes and its recurrent appearances in affected populations, avian cholera, caused by the bacterium *Pasteurella multocida*, has the potential to threaten several wild bird species (Botzler, 1991; Samuel, Botzler, & Wobeser, 2007). This is notably true of birds aggregating at high densities at a time of their life cycle, such as Arctic eider ducks (Descamps, Jenouvrier, Gilchrist, & Forbes, 2012), penguins (Cooper et al., 2009), Cape cormorants (Waller & Underhill, 2007), and albatrosses (Weimerskirch, 2004). Currently, because of the high mortality this disease may cause in these species, attention to albatrosses and large petrels affected by avian cholera is considered an “utmost priority” (reviewed in Uhart, Gallo, & Quintana, 2017).

Amsterdam Island (37.8°S, 77.6°E) is a ~55 km<sup>2</sup> isolated volcanic island that hosts two-thirds of the global population of the endangered Indian yellow-nosed albatross (*Thalassarche carteri*; Figure 1) and a small subpopulation of dark-mantled sooty albatrosses (*Phoebastria fusca*). Since the 1980s, chicks of these two species experience recurrent and dramatic mortality episodes attributed to avian cholera (Weimerskirch, 2004). By contrast, the disease does not seem

to affect the survival of adults (Rolland, Barbraud, & Weimerskirch, 2009). Although the exact case-fatality rate in chicks is not known, acute infection in chicks results in rapid death, as in poultry exposed to virulent strains of *P. multocida*. These intense chick mortality episodes lead to near-zero chick productivity in some years, a major cause of overall population decline in Indian yellow-nosed albatrosses (Rolland et al., 2009). Importantly, Amsterdam Island is also home to the critically endangered and endemic Amsterdam albatross (*Diomedea amsterdamensis*; IUCN, 2017). Although its population has increased over the past decades, the Amsterdam albatross has also suffered from marked chick mortality in 2000 and 2001 (Weimerskirch, 2004). For these reasons, avian cholera is considered the most severe global infectious threat to albatrosses (Phillips et al., 2016; Uhart et al., 2017).

One approach to limit the impact of deadly epizootics in wildlife is to promote the resilience of host populations through vaccination (Groner et al., 2016). However, the availability of vaccines combining safety and efficacy against wildlife pathogens is limited. Here, we manufactured an auto-genous vaccine based on a field strain of *P. multocida* and used it in the wild to test its ability to protect albatross chicks from avian cholera mortality.

## 2 | METHODS

### 2.1 | Vaccine preparation

*P. multocida* strain D2C was isolated from a dead dark-mantled sooty albatross chick collected at an Amsterdam Island colony during the 2011–2012 breeding season, and typed using lipopolysaccharide (LPS, somatic) antibodies as Heddlestone serotype 1 and Namioka serotype 7. The Heddlestone test is widely used to differentiate *P. multocida* strains based on LPS type, although it should be noted that this test may yield ambiguous results, especially compared with a recently developed PCR typing test (Harper et al., 2015). Strain D2C was grown on solid soybean-casein digest agar with yeast extract in a 500 cm<sup>2</sup> tissue culture flask at 37°C for approximately 22 hours. The bacteria were then collected in 20 mL of phosphate buffered saline (PBS), their concentration was estimated using optical density at 580 nm, and they were inactivated (killed) with a 34–38% formaldehyde solution at room temperature (20°C ± 5°C). After centrifugation at 2,934 × g for 30 minutes, the bacterial pellets were resuspended in PBS. Mineral oil adjuvants were used to obtain water-in-oil emulsions containing the equivalent of 3 × 10<sup>9</sup> colony-forming units (CFU) per milliliter (2013 and 2014 vaccination campaigns with 50% v/v adjuvant) or 1 × 10<sup>9</sup> CFU/mL (2015 campaign with 60% v/v adjuvant). Water-in-oil emulsions are extensively used for the formulation of inactivated avian vaccines, where they have proven to be strong immunity adjuvants (Hilgers, Nicolas, Lejeune,

**TABLE 1** Number of albatross chicks that were seropositive/sampled for serum during the 2015–2016 reproductive season according to vaccination status

Treatment group	December 2015	January 2016	February 2016	March 2016
Vaccinated	0/29 <sup>a</sup>	8/28	16/16	11/12
Unvaccinated, sampled	0/11 <sup>a</sup>	0/8	0/3	0/0

<sup>a</sup>In December, one chick in each group could not be sampled.

Dewil, & Boon, 1998; Jansen, Hofmans, Theelen, & Schijns, 2005; Jansen, Hofmans, Theelen, Manders, & Schijns, 2006). Several innocuity and efficacy tests conducted on a series of avian species (chickens, turkeys, and ducks) had indicated that these were appropriate antigen doses to use with these killed vaccine formulations. Endotoxin content of the 2015 vaccine was less than 5,000 EU/mL, as assessed using the kinetic chromogenic method of the European Pharmacopoeia.

The inactivated bacteria were also used to set up an agglutination assay (detailed in Supplementary Online Material, SOM) to detect antibodies directed against this or potential cross-reactive strains in bird plasma samples.

## 2.2 | Field procedures

Rigorous biosecurity measures were observed in the field to avoid spreading the disease between bird colonies and individual birds within the colonies. These involved the use of dedicated clothing consisting of a different set of waterproof leggings, jackets, and rubber boots for each bird colony. Waterproof overalls, any other equipment (e.g., caliper, ruler) and hands were disinfected between contacts with any two birds, respectively using hydrogen peroxide, commercial disinfectant wipes, and hydroalcoholic gel. All pieces of clothing were thoroughly cleaned and disinfected again at the end of each day.

Indian yellow-nosed albatrosses lay only one egg and thus raise at most one chick per nest each year. Seronegative chicks from 30 nests were injected subcutaneously with 0.5 mL of autogenous vaccine on December 18, 2015 (an estimated  $14 \pm 3$  days after hatching, mean  $\pm$  SD). Vaccination date was optimized based on the results from previous vaccination trials in 2013 and 2014 (detailed in SOM.) Two control groups were used to compare their survival with that of the *vaccinated* group. In the first control group, chicks were not vaccinated but were *sampled* to monitor potential seroconversion ( $n = 12$ ). In the second control group, chicks were left entirely *unmanipulated* to assess baseline reproductive success and any potential adverse effect of vaccination or handling on survival ( $n = 24$ ). The nests from the three groups were mixed and dispersed over the same area.

Following injection, chick survival was assessed on three occasions in January, February, and March 2016, fledging occurring in late March. Survival of chicks was analyzed using the Cox proportional hazards model as detailed in SOM.

Blood samples were also taken monthly to investigate seroconversion. Plasma samples were anonymized and randomized before serological testing.


## 3 | RESULTS

Serological analyses showed seroconversion of all vaccinated chicks by February 8, 2016 (Table 1), with mean titres increasing until March 2016 (Figure 2). By contrast, none of the sampled unvaccinated chick seroconverted over that period. Importantly, chick vaccination provided a strong protection against mortality (Figure 2). End survival probability was 46% for the vaccinated chicks versus 14% for the unmanipulated controls, the death risk at a given time being divided by 2.6 (95% confidence intervals or CI: 1.31, 5.07;  $P = 0.006$ ); Cox proportional hazards model. The death risk was also divided by 3.1 (CI: 1.44, 6.72;  $P = 0.004$ ) when comparing vaccinated to unvaccinated sampled chicks. No survival difference was detected between the sampled (and unvaccinated) versus unmanipulated control groups (death risk divided by 0.82 (CI: 0.40, 1.69;  $P = 0.602$ )).

## 4 | DISCUSSION

To our knowledge, this is the first immunization trial in albatrosses, and the first evaluation of an autogenous vaccine in a threatened animal population. Vaccination programs targeting endangered species in the wild generally endeavor to maximize protection of the target population (Livingston et al., 2013; López et al., 2009; Malakoff, 2016) and thus do not include negative control groups. This study directly demonstrates the efficacy of an autogenous vaccine in protecting endangered albatrosses against lethal infection in a controlled field trial.

Vaccine production was inspired from a process initially developed for the poultry industry, with analogies to the production of a bacterin that had shown promising results in giant Canada geese (*Branta canadensis maxima*; Price, 1985) and was later used to investigate the effect of avian cholera on adult survival in lesser snow geese (*Anser caerulescens caerulescens*; Samuel, Takekawa, Baranyuk, & Orthmeyer, 1999). Multilocus sequence typing (MLST) using seven housekeeping genes (*adk*, *est*, *gdh*, *mdh*, *pgi*, *pmi*, *zwf*;



**FIGURE 2** (a) Seroconversion and (b) survival curves for Indian yellow-nosed albatross chicks that were vaccinated (red,  $n = 30$ ), or unvaccinated (blue) and either left entirely unmanipulated (solid line,  $n = 24$ ) or sampled (dotted line,  $n = 12$ ) in the 2015–2016 reproductive season. Error bars indicate standard error of the mean antibody titres (expressed as natural log[serial serum dilutions + 1])

Subaaharan, Blackall, & Blackall, 2010) showed no variation on *Pasteurella* isolated on Amsterdam Island from three albatross individuals sampled in 2011–2012 and three individuals sampled in 2012–2013. This analysis (carried out by AJ and colleagues) showed that the isolates all belonged to multilocus sequence type 61. Only one strain was thus included in vaccine preparation. However, *P. multocida* strains can mutate in antigenic LPS regions that are not examined using MLST, and the potential for heterologous protection across different serotypes with this killed vaccine was not examined. Live mutant *P. multocida* vaccines have been shown to offer better protection against strains differing in their LPS structure compared to killed vaccine (Harper

et al., 2016) and cross-serotype protection (Scott, Markham, & Whithear, 1999). However, the concern that live vaccines might revert to virulence, or be detrimental in immunocompromised hosts, precludes their use in endangered albatrosses and other protected wild animal populations including seabirds on Amsterdam Island. In the future, it will be feasible and might be advisable to manufacture multivalent killed vaccines including different strains of *P. multocida*, notably with different LPS structures to prevent the apparition of vaccine escape mutants with, for instance, shorter LPS. Other bacterial species could also be included if deemed relevant. For the time being, the spectacular improvement in chick survival following immunization with a monovalent (anti-*Pasteurella* only) vaccine is one further confirmation of the role of *P. multocida* in chick die-offs.

Choosing an optimal date for chick vaccination is important. If vaccinating too early, the chick immune system may still be immature and unable to mount a protective response (Mast & Goddeeris, 1999), as suggested by the negative results of the 2013 and 2014 early vaccination protocols (SOM). On the other hand, excessive delay in vaccination may result in an epidemic occurring before chicks are effectively protected. Vaccinating when most chicks were 10–15 days old (mid-December) proved an effective compromise. Complete seroconversion (in the sense of plateauing antibody titres) took well over a month following vaccination (Figure 2A), but differential survival could be noted as early as January, when antibody titres in vaccinated chicks were still low. Thus, even low antibody titres appear associated with protection, and the vaccine may also stimulate other arms of the immune system before antibody titres peak.

The worrying conservation status of numerous albatross species has prompted the Agreement on the Conservation of Albatrosses and Petrels. Given the dramatic consequences of these infections in terms of chick mortality, which can severely impact population viability (Finkelstein, Doak, Nakagawa, Sievert, & Klavitter, 2010), increase in chick survival through vaccination may be a key element in albatross conservation. Successfully vaccinated birds fledged in March 2016 have not yet returned to the colony to breed. Therefore, it is still unclear how long the benefits of vaccination of a given individual are likely to last, and whether there may be long-term benefits of chick vaccination beyond increasing the fledging probability. If the effect of vaccination is limited to improving the fledging rate, models developed in Rolland et al. (2009) suggest that the improvement observed in this study might not be sufficient to stabilize the population, although slight differences in adult mortality or emigration can strongly affect the outcome. Also, in this trial, we vaccinated only about a 10th of the local study colony, itself accounting for about 1% of the Amsterdam Island population of yellow-nosed albatrosses (Rolland et al., 2009). Therefore, the trial likely stands well below the population



immunity threshold above which one infectious individual would not cause an outbreak (Feng, Hill, Smith, & Glasser, 2015). By contrast, vaccination programs involving a high coverage of local albatross populations may benefit from substantial herd immunity effects, proportionally providing even greater protection to the chicks. The possible protection afforded to chicks via the vaccination of mothers and the recurrent transfer of antibodies via the egg yolk is another perspective to explore further (Garnier et al., 2012).

The suite of issues involved in the decision of vaccinating a wild threatened population is quite complex. Scaling up vaccination as a management tool in this system will also necessitate a refined understanding of factors affecting the frequency and timing of epizootics. Modeling studies comparing different vaccination scenarios (Garnier et al., 2012; Haydon et al., 2006) may then help anticipate the outcome of vaccinating yellow-nosed albatrosses or other species in the system. For instance, vaccination may directly benefit the minute population of critically endangered Amsterdam albatross. The vaccination of brown skuas (*Stercorarius antarcticus*), whose predatory behavior, substantial infection rates and high mobility could make them key epidemiological actors (Boulinier et al., 2016), might also prove helpful in controlling the negative effects of avian cholera on Amsterdam Island. Of note, without a much clearer understanding of the epidemiological reservoirs on the island, local extirpation of the pathogen through vaccination only is probably not a realistic perspective at this stage, and further investigations into disease dynamics are certainly warranted.

Finally, avian cholera affects wild bird populations in other very remote areas (Iverson et al., 2016; Leotta, Chinen, Vigo, Pecoraro, & Rivas, 2006). The presence of such infections in these isolated systems raises the question of their origin. On Amsterdam Island, some hypotheses involve human activities, such as a poultry pen that was active until 2006 (Weimerskirch, 2004) or the presence of carrier rats introduced involuntarily about a century ago. In the past, other domesticated animals potentially acting as reservoirs (reviewed in Wilson & Ho, 2013), such as cattle and dogs, were present on the island; a small number of feral cats still remain. Beyond the academic interest, understanding the origin of the diseases may also have ethical implications on the appropriateness of bold human intervention such as vaccination in wild ecosystems (Cleaveland, 2009), where infectious diseases may be a natural component. However, the high density of birds on Amsterdam Island, representing a significant proportion of the global population of the endangered Indian yellow-nosed albatross, and the seemingly strong potential for this pathogen to drive some of these species of conservation concern closer to extinction (Phillips et al., 2016; Uhart et al., 2017) seems to provide an unusually compelling case for intervention with vaccination.

In conclusion, this report identifies a way of promoting resistance of endangered albatross populations to a major infectious threat. This tool will be relevant to the applied conservation of these and other species suffering high levels of avian cholera (and potentially other bacterial disease) mortality. As such, it should be of interest to governments, nongovernmental organizations, and other sustainable development policy actors (such as the International Commission for the Conservation of Antarctic Marine Living Resources) involved in the conservation of wildlife threatened by emerging infectious diseases.

## ACKNOWLEDGMENTS

We are grateful to Nicolas Giraud, David Hémery, Hélène Le Berre, and Romain Bazire for help in the field, Raul Ramos, Nicolas Courbin, and Sylvain Gandon for discussion and advice on analyses and interpretations. This work was supported by the French Polar Institute (IPEV programs ECOPATH-1151 and ORNITHOECO-109), ANR “EVE-MATA” (11 BSV7 003), Réserve Nationale des Terres Australes Françaises (Cédric Marteau), and Zone Atelier Antarctique et Subantarctique. VB was funded through a CeMEB LabEx postdoctoral fellowship and AG is supported via a PhD fellowship from the French Ministry of Research.

The experimental design was approved by the Comité de l'Environnement Polaire and the French Ministry of Research (04939.03).

## REFERENCES

- Botzler, R. G. (1991). Epizootiology of avian cholera in wildfowl. *Journal of Wildlife Diseases*, 27, 367–395.
- Boulinier, T., Kada, S., Ponchon, A., Dupraz, M., Dietrich, M., Gamble, A., ... McCoy, K. D. (2016). Migration, prospecting, dispersal? What host movement matters for infectious agent circulation? *Integrative and Comparative Biology*, 56, 330–342.
- Cleaveland, S. (2009). Viral threats and vaccination: Disease management of endangered species. *Animal Conservation*, 12, 187–189.
- Cooper, J., Crawford, R. J. M., Villiers, M. S. D., Dyer, B. M., Hofmeyr, G. J. G., & Jonker, A. (2009). Disease outbreaks among penguins at sub-Antarctic Marion Island: A conservation concern. *Marine Ornithology*, 37, 193–196.
- Croxall, J. P., Trathan, P. N., & Murphy, E. J. (2002). Environmental change and Antarctic seabird populations. *Science*, 297, 1510–1514.
- Daszak, P., Cunningham, A. A., & Hyatt, A. D. (2000). Emerging infectious diseases of wildlife – Threats to biodiversity and human health. *Science*, 287, 443–449.
- Descamps, S., Jenouvrier, S., Gilchrist, H. G., & Forbes, M. R. (2012). Avian cholera, a threat to the viability of an Arctic seabird colony? *Plos One*, 7(2), e29659.
- Feng, Z., Hill, A. N., Smith, P. J., & Glasser, J. W. (2015). An elaboration of theory about preventing outbreaks in homogeneous populations to include heterogeneity or preferential mixing. *Journal of Theoretical Biology*, 386, 177–187.

- Finkelstein, M. E., Doak, D. F., Nakagawa, M., Sievert, P. R., & Klavitter, J. (2010). Assessment of demographic risk factors and management priorities: Impacts on juveniles substantially affect population viability of a long-lived seabird. *Animal Conservation*, 13, 148–156.
- Garnier, R., Ramos, R., Staszewski, V., Militão, T., Lobato, E., González-Solís, J., & Boulinier, T. (2012). Maternal antibody persistence: A neglected life-history trait with implications from albatross conservation to comparative immunology. *Proceedings of the Royal Society of London Series B, Biological Sciences*, 279, 2033–2041.
- Groner, M. L., Maynard, J., Breyta, R., Carnegie, R. B., Dobson, A., Friedman, C. S., ... Harvell, C. D. (2016). Managing marine disease emergencies in an era of rapid change. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences*, 371, 20150364.
- Harper, M., John, M., Turni, C., Edmunds, M., St. Michael, F., Adler, B., ... Boyce, J. D. (2015). Development of a rapid multiplex PCR assay to genotype *Pasteurella multocida* strains by use of the lipopolysaccharide outer core biosynthesis locus. *Journal of Clinical Microbiology*, 53, 477–485.
- Harper, M., John, M., Edmunds, M., Wright, A., Ford, M., Turni, C., ... Boyce, J. D. (2016). Protective efficacy afforded by live *Pasteurella multocida* vaccines in chickens is independent of lipopolysaccharide outer core structure. *Vaccine*, 34, 1696–1703.
- Harvell, C. D., Kim, K., Burkholder, J. M., Colwell, R. R., Epstein, P. R., Grimes, D. J., ... Vasta, G. R. (1999). Emerging marine diseases – Climate links and anthropogenic factors. *Science*, 285, 1505–1510.
- Harvell, C. D., Mitchell, C. E., Ward, J. R., Altizer, S., Dobson, A. P., Ostfeld, R. S., & Samuel, M. D. (2002). Climate warming and disease risks for terrestrial and marine biota. *Science*, 296, 2158–2162.
- Haydon, D. T., Randall, D. A., Matthews, L., Knobel, D. L., Tallents, L. A., Gravenor, M. B., ... Laurenson, M. K. (2006). Low-coverage vaccination strategies for the conservation of endangered species. *Nature*, 443, 692–695.
- Hilgers, L. A., Nicolas, I., Lejeune, G., Dewil, E., & Boon, B. (1998). Effect of various adjuvants on secondary immune response in chickens. *Veterinary Immunology and Immunopathology*, 66, 159–171.
- IUCN–BirdLife International. 2017. *Diomedea amsterdamensis* (amended version of 2016 assessment). *The IUCN Red List of Threatened Species 2017*: e.T22698310A110677305. <https://doi.org/10.2305/IUCN.UK.2017-1.RLTS.T22698310A110677305.en>.
- Iverson, S. A., Gilchrist, H. G., Soos, C., Buttler, I. I., Harms, N. J., & Forbes, M. R. (2016). Injecting epidemiology into population viability analysis: Avian cholera transmission dynamics at an arctic seabird colony. *Journal of Animal Ecology*, 85, 1481–1490.
- Jansen, T., Hofmans, M. P. M., Theelen, M. J. G., & Schijns, V. E. J. C. (2005). Structure-activity relations of water-in-oil vaccine formulations and induced antigen-specific antibody responses. *Vaccine*, 23, 1053–1060.
- Jansen, T., Hofmans, M. P. M., Theelen, M. J. G., Manders, F., & Schijns, V. E. J. C. (2006). Structure- and oil type-based efficacy of emulsion adjuvants. *Vaccine*, 24, 5400–5405.
- Lafferty, K. D., & Hofmann, E. E. (2016). Marine disease impacts, diagnosis, forecasting, management and policy. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 371: 20150200.
- Leotta, G. A., Chinen, I., Vigo, G. B., Pecoraro, M., & Rivas, M. (2006). Outbreaks of avian cholera in Hope Bay, Antarctica. *Journal of Wildlife Diseases*, 42, 259–270.
- Livingston, M., Fidler, A., Mellor, D., de Kloet, S., Eason, D., Elliott, G., & Moorhouse, R. (2013). Prevalence of IgY antibodies against *Erysipelothrix rhusiopathiae* in a critically endangered parrot (kakapo, *Strigops habroptilus*) and associated responses to vaccination. *Avian Pathol. J. WVPA*, 42, 502–507.
- López, G., López-Parra, M., Fernández, L., Martínez-Granados, C., Martínez, F., Meli, M. L., ... Simón, M. A. (2009). Management measures to control a feline leukemia virus outbreak in the endangered Iberian lynx. *Animal Conservation*, 12, 173–182.
- Malakoff, D. (2016). A race to vaccinate rare seals. *Science*, 352, 1265–1265.
- Mast, J., & Goddeeris, B. M. (1999). Development of immunocompetence of broiler chickens. *Veterinary Immunology and Immunopathology*, 70, 245–256.
- Phillips, R. A., Gales, R., Baker, G. B., Double, M. C., Favero, M., Quintana, F., ... Wolfaardt, A. (2016). The conservation status and priorities for albatrosses and large petrels. *Biological Conservation*, 201, 169–183.
- Price, J. I. (1985). Immunizing Canada geese against avian cholera. *Wildlife Society Bulletin 1973–2006*, 13, 508–515.
- Rolland, V., Barbraud, C., & Weimerskirch, H. (2009). Assessing the impact of fisheries, climate and disease on the dynamics of the Indian yellow-nosed Albatross. *Biological Conservation*, 142, 1084–1095.
- Samuel, M. D., Takekawa, J. Y., Baranyuk, V. V., & Orthmeyer, D. L. (1999). Effects of avian cholera on survival of Lesser Snow Geese *Anser caerulescens*: An experimental approach. *Bird Study*, 46, S239–S247.
- Samuel, M. D., Botzler, R. G., & Wobeser, G. A. (2007). Avian cholera. In N. J. Thomas, D. B. Hunter, & C. T. Atkinson (Eds.), *Infectious diseases of wild birds* (pp. 239–269). Ames, Iowa, USA: Blackwell Publishing Professional.
- Scott, P. C., Markham, J. F., & Whithear, K. G. (1999). Safety and efficacy of two live *Pasteurella multocida* aro-A mutant vaccines in chickens. *Avian Diseases*, 43, 83–88.
- Smith, K. F., Acevedo-Whitehouse, K., & Pedersen, A. B. (2009). The role of infectious diseases in biological conservation. *Animal Conservation*, 12, 1–12.
- Subaaharan, S., Blackall, L. L., & Blackall, P. J. (2010). Development of a multi-locus sequence typing scheme for avian isolates of *Pasteurella multocida*. *Veterinary Microbiology*, 141, 354–361.
- Uhart, M. M., Gallo, L., & Quintana, F. (2017). Review of diseases (pathogen isolation, direct recovery and antibodies) in albatrosses and large petrels worldwide. *Bird Conservation International*, 0, 1–28.
- Van Hemert, C. R., Pearce, J. M., & Handel, C. M. (2014). Wildlife health in a rapidly changing north: Focus on avian disease. *Frontiers in Ecology and the Environment*, 12, 548–556.

- Waller, L. J., & Underhill, L. G. (2007). Management of avian cholera *Pasteurella multocida* outbreaks on Dyer Island, South Africa, 2002–2005. *African Journal of Marine Science*, 29, 105–111.
- Weimerskirch, H. (2004). Diseases threaten Southern Ocean albatrosses. *Polar Biology*, 27, 374–379.
- Wilson, B. A., & Ho, M. (2013). *Pasteurella multocida*: From zoonosis to cellular microbiology. *Clinical Microbiology Reviews*, 26, 631–655.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

**How to cite this article:** Bourret V, Gamble A, Tornos J, et al. Vaccination protects endangered albatross chicks against avian cholera. *Conservation Letters*. 2018;11:e12443. <https://doi.org/10.1111/conl.12443>