



University  
of Glasgow

Mukhopadhyay, A. and Chen, C.Y. and Doerig, C. and Henriquez, F.L. and Roberts, C.W. and Barrett, M.P. (2009) *The Toxoplasma gondii Plastid replication and Repair Enzyme complex, PREX*. Parasitology, 136 (7). pp. 747-755. ISSN 0031-1820

<http://eprints.gla.ac.uk/6078/>

Deposited on: 29 June 2009

# The *Toxoplasma gondii* Plastid replication and Repair Enzyme Complex, PREX

A. MUKHOPADHYAY<sup>1</sup>, C-Y. CHEN<sup>2</sup>, C. DOERIG<sup>3</sup>, F. L. HENRIQUEZ<sup>4</sup>, C. W. ROBERTS<sup>4</sup> and M. P. BARRETT<sup>1\*</sup>

<sup>1</sup>Institute of Biomedical and Life Sciences, Division of Infection and Immunity, Glasgow Biomedical Research Centre, University of Glasgow, Glasgow G12 8TA, UK

<sup>2</sup>Joseph Gottstein Memorial Cancer Research Laboratory, Department of Pathology, University of Washington, Seattle, WA 98195-7705, USA

<sup>3</sup>INSERM U609, Wellcome Centre for Molecular Parasitology, University of Glasgow, 120 University Place, Glasgow, G12 8TA, Scotland, UK. INSERM U609, Global Health Institute, Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

<sup>4</sup>Strathclyde Institute of Pharmacy and Biomedical Sciences, 27 Taylor Street, University of Strathclyde, Glasgow G4 0NR, UK

(Received 4 December 2008; revised 13 February and 4 March 2009; accepted 4 March 2009; first published online 30 April 2009)

## SUMMARY

A plastid-like organelle, the apicoplast, is essential to the majority of medically and veterinary important apicomplexan protozoa including *Toxoplasma gondii* and *Plasmodium*. The apicoplast contains multiple copies of a 35 kb genome, the replication of which is dependent upon nuclear-encoded proteins that are imported into the organelle. In *P. falciparum* an unusual multi-functional gene, *pfprex*, was previously identified and inferred to encode a protein with DNA primase, DNA helicase and DNA polymerase activities. Herein, we report the presence of a *prex* orthologue in *T. gondii*. The protein is predicted to have a bi-partite apicoplast targeting sequence similar to that demonstrated on the PfPREX polypeptide, capable of delivering marker proteins to the apicoplast. Unlike the *P. falciparum* gene that is devoid of introns, the *T. gondii* *prex* gene carries 19 introns, which are spliced to produce a contiguous mRNA. Bacterial expression of the polymerase domain reveals the protein to be active. Consistent with the reported absence of a plastid in *Cryptosporidium* species, *in silico* analysis of their genomes failed to demonstrate an orthologue of *prex*. These studies indicate that *prex* is conserved across the plastid-bearing apicomplexans and may play an important role in the replication of the plastid genome.

Key words: Apicomplexa, DNA replication, *Plasmodium falciparum*, *Toxoplasma gondii*, exon, intron, DNA polymerase, DNA primase, DNA helicase.

## INTRODUCTION

The parasite *Toxoplasma gondii* is a member of the eukaryotic phylum Apicomplexa. The phylum, which includes *Plasmodium falciparum*, a causative agent of malaria, as well as multiple other parasitic genera, is of immense medical and veterinary importance. Members of this phylum share a number of structural attributes. With the likely exception of *Cryptosporidium*, one such shared characteristic is the apicoplast, a relict plastid believed to have been acquired through secondary endosymbiosis (Wilson *et al.* 1994; McFadden *et al.* 1996). The *T. gondii* apicoplast contains its own ~35 kb genome that is replicated and segregated with the organelle during parasite proliferation. With the exception of the microgamete stage, all *T. gondii* life-cycle stages

have been demonstrated to possess a single apicoplast, which is faithfully segregated into the daughter cells (Striepen *et al.* 2000; Ferguson *et al.* 2007). Like many endosymbiont-derived organelles, the apicoplast has a reduced genome containing genes involved principally in its own replication. A variety of the genes originally associated with the prokaryotic endosymbiont have been incorporated into the nuclear genome and their products are targeted back to the apicoplast using a bi-partite plastid targeting peptide. Consequently, many nuclear-encoded enzymes associated with the apicoplast are of prokaryotic ancestry. The organelle is essential as it is the site for numerous biochemical pathways including heme and fatty acid biosynthesis and steroid production (Roos, 1999; Soldati, 1999; McLeod *et al.* 2001). The apicoplast has therefore proven to be a valuable target for drugs active against apicomplexan parasites (Fichera and Roos, 1997).

Genes encoding proteins involved in DNA replication or repair are absent from the apicoplast genome. Certain genes encoding plastid DNA replication enzymes are, however, present in the nuclear

\* Corresponding author: Institute of Biomedical and Life Sciences, Division of Infection and Immunity, Glasgow Biomedical Research Centre, University of Glasgow, Glasgow G12 8TA, UK.

genome, with the protein being imported into the plastid post-translationally. Inhibition of components of the apicoplast DNA replication machinery (e.g. DNA gyrase) by ciprofloxacin (Dar *et al.* 2007) can cause cessation of apicoplast replication and subsequent parasite death (Dar *et al.* 2007; Raghu Ram *et al.* 2007). Thus, the nuclear encoded apicoplast replication machinery is an established drug target in apicomplexan parasites (Ralph *et al.* 2001).

The various enzymatic activities required for replication of the apicoplast genome in *P. falciparum* may be derived in large part from a single gene, *pfprex*, which encodes a multifunctional protein (Seow *et al.* 2005) comprising a T7 bacteriophage homologous primase-helicase domain and a prokaryotic DNA polymerase domain separated by a spacer domain. An N-terminal sequence of PfPREX targeted Green Fluorescent Protein (GFP) to the apicoplast, strongly suggesting that this protein may be involved in DNA replication or repair of the organelle genome. Thus, this single gene provides multiple functional properties required for apicoplast replication and/or repair (Seow *et al.* 2005). Herein, we demonstrate the presence of a single *prex* gene (*Tgprex*) in *T. gondii*. Unlike *prex* in *P. falciparum*, the gene in *T. gondii* is interrupted by multiple introns. The polymerase domain expressed as a recombinant protein in *E. coli* displays DNA polymerase activity. The conservation of unusual prokaryotic-like DNA replication enzymes across the Apicomplexa indicates that PREX could be a potential target for chemotherapy against this class of organisms.

#### MATERIALS AND METHODS

##### *T. gondii* tachyzoite preparation and preparation of cDNA

*T. gondii* tachyzoites (RH strain) were grown in the peritoneum of BALB/c mice as previously described (Roberts and Alexander, 1992). Tachyzoites were harvested and washed in sterile phosphate-buffered saline (PBS; 0.1 M NaCl, 2.7 mM KCl, 10 mM Na<sub>2</sub>HPO<sub>4</sub>, 1.76 mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4) by centrifugation at 500 g for 5 min at 4 °C and stored as pellets at -70 °C. RNA isolation from the tachyzoites was performed using Trizol<sup>®</sup> (Invitrogen, UK) following a protocol, based on a single-step acid guanidinium thiocyanate-phenol-chloroform for RNA isolation (Chomczynski and Sacchi, 1987).

Complementary DNA (cDNA) was produced from total RNA as previously described (Campbell *et al.* 2004). In a 13 µl volume, 1 µg of RNA was added to 1 µl of random hexamer mix (Promega, Southampton, UK) and 1 µl of a 10 mM deoxy-nucleoside triphosphate mixture, incubated at 65 °C for 5 min, and chilled on ice for 1 min. Then 80 units

of RNasin ribonuclease inhibitor (Promega, UK), 4 µl of 5 × first strand buffer (250 mM Tris-HCl, pH 8.3, 375 mM KCl, 15 mM MgCl<sub>2</sub>), 1 µl of 0.1 M DTT and 200 units of Superscript III reverse transcriptase RNase H<sup>-</sup> (Invitrogen, UK) were added to the reaction and incubated for 5 min at 25 °C followed by 55 °C for 60 min. Inactivation of the reverse transcriptase was performed by heating at 70 °C for 10 min.

##### Identification and sequencing of the *prex* gene in *T. gondii*

Initially the PfPREX homologues were identified in the *T. gondii* genome project by BLASTP analysis. As the *P. falciparum* gene was AT rich, the protein sequence was used as a query and the protein match was extrapolated to its cognate gene in the genome. The two protein coding genes are situated on chromosome VIIb in the *T. gondii* genome project (www.toxodb.org release 4) (Gajria *et al.* 2008). Oligonucleotide primers were designed to confirm the *prex* gene structure by PCR. The primer locations are listed in relation to the actual gene model (Fig. 1A) determined in this study and are listed in order of their positions on the gene (Table 1). Primer pairs for each PCR are designated in Fig. 1A. The primers situated within the consensus exons predicted by different gene prediction algorithms (GLEAN, GlimmerHMM, TwinScan, TwinScan/Eimeria and TigrScan) are in bold font (Table 1). After confirmation of predicted sequences these primers were coupled with primers from other predicted exons to confirm the coding sequence.

PCR reactions were performed on both genomic and cDNA from *T. gondii* (RH strain) tachyzoites for comparison. One µl of the cDNA or 500 ng of the genomic DNA template was amplified by Expand Hi-Fidelity polymerase enzyme (Roche) using 0.2 mM dNTP mix, 0.4 µM of sequence specific forward and reverse primers with 1.5 µM/2.0 µM of Mg<sup>2+</sup> in enzyme-specific buffer. Amplified PCR products from the cDNA were cloned into the pGEM-T Easy vector (Promega) and sequenced to verify the ORF. The cDNA contig was assembled using the Contig Express programme of the Vector NTI suite 9 and 10.

To clarify the 5' end of the coding sequence in the *prex* cDNA, Rapid Amplification of the cDNA end (RACE) was performed on *T. gondii* total RNA. The cDNA was amplified (Superscript III, Invitrogen) using a gene-specific reverse primer (RACE 3) situated in the first exon sequenced (Table 1). A poly-cytosine tail was added to the transcribed cDNA using recombinant terminal transferase (Roche). Nested PCR was performed using a poly-guanosine forward primer and 2 gene-specific reverse primers (RACE 2 and RACE 1 respectively) as described in Table 1.




Fig. 1. The exon–intron structure (Guo *et al.* 2007) and a diagrammatic representation of the domain structure of the *Toxoplasma gondii* ORF 7740 present on chromosome VIIb of *Toxoplasma gondii* as elucidated by PCR. The structure of the *T. gondii* ORF 7740 was confirmed by the sequencing of cDNA from the RH strain of the organism, using primers listed in Table 1 (A). The 7740 bp long ORF was composed of 20 exons separated by 19 introns and different reading frames for the introns are designated. The exon intron structure of the gene was created using GSDS (<http://gsds.cbi.pku.edu.cn/>) (Guo *et al.* 2007). Primer pairs used for PCR are delineated in relation to the exon structure of the *Tgprex* gene. The primers designed on the consensus exon sequences are in bold font. (B) 5'RACE was performed to confirm the initiation codon where RT+ and RT– refers to reactions with or without reverse transcriptase, respectively. (C) The 5' end of the ORF was confirmed by comparing PCR on cDNA (c) and genomic DNA (g) using primers F.I and R3.1 (Table 1) generating 2345 bp and 3178 bp long PCR products, respectively. (D) The continuity of the single gene was confirmed by PCR using primers F11.2 and R12.3 (Table 1) showing a 1766 bp long PCR product from cDNA (c) compared to a 3334 bp long product from genomic DNA (g). The DNA marker is designated by M. (E) The domain structure of the TgPREX putative protein is diagrammatically represented.

#### TgPREX polymerase domain cloning and expression

From the alignment of the predicted PREX proteins in different apicomplexan organisms (data not shown), the exonuclease-polymerase domain of the *T. gondii* PREX was identified and a 3060 bp region encompassing the entire domain was amplified from *T. gondii* tachyzoite cDNA using primers Tg\_polymeraseF and R20.2 (Table 1). The fragment was cloned into the pGEM-T Easy vector and digested with *NheI* (shown in the Tg\_polymeraseF primer in underlined italics in Table 1) and *SacI* (restriction enzyme site in the pGEM-T Easy plasmid) enzymes, and subsequently ligated into the pET28a<sup>+</sup> vector. In the resulting construct, TgPREX is fused to an

N-terminal hexa-histidine tag and expressed under the control of the T7 promoter.

The pET28a-TgPREX polymerase plasmid was used for transformation of the *E. coli* BL21-AI (Invitrogen) and cells were grown at 37 °C in 3 L of LB medium containing 0.1% glucose and 50 µg/ml carbenicillin; L-arabinose (0.2% final concentration) was added at OD<sub>600</sub> = ~0.6. After further incubation for 4 h at 37 °C, cells were harvested by centrifugation and resuspended in 60 ml of buffer A [50 mM Tris-HCl (pH 8.0), 50 mM NaCl, 1 mM EDTA, 1 mM PMSF, 2 mM B-mercaptoethanol, 2 mM benzamidine and 10% (w/v) glycerol], and adjusted to 500 µg/ml lysozyme prior to storage at –80 °C.

Table 1. List of primers used for confirmation of the *Tgprex* gene coding sequence on chromosome VIIIb of *Toxoplasma gondii* and expression of recombinant polymerase domain of TgPREX protein

(The table shows the primer name, nucleotide sequence and the position of each primer according to the sequenced gene model presented in Fig. 1A. Primer pairs used for PCR amplification are delineated in Fig. 1A. The reverse primers are highlighted in the grey background. Primers situated in consensus exons predicted by ToxoDB are in bold font. The 5'RACE primers were used for the clarification of the initiation codon of the gene that was confirmed by PCR amplification of the 5' region using the 5'UTR primer. The cDNA primers were used to establish the coding sequence of the gene. The 3'UTR primer was used to ensure the 3' end of the *Tgprex* gene. Polymerase domain of the TgPREX protein was cloned using forward primer Tg\_polymeraseF in conjunction with R20.2 reverse primer. The *Nhe I* restriction site on Tg\_polymeraseF primer is shown in italicized font.)

Primer Name	5'→3' Nucleotide sequence	Location
5'UTR primers		
FI	GCCTCTCCTTGTTCCCTGCTTC	175 bp 5' start codon
5'RACE primers		
RACE.1	CCAGAGGAGAAAGAACTGTAC	Exon 1
RACE.2	AACGAAGAAAACCGCAGGG	Exon 1
RACE.3	CGAGAAGGAAAAGACGACG	Exon 1
cDNA primers		
F1.1	ATGCGTCCGGTTGAGTACCGG	5'ATG
F1.2	TACTGCCACCGCTGCGGGTGG	Exon 1
R3.1	CAGCGAGTTCTCTGTCTTAC	Exon 3
F4.1	GAAACTCGGAATCGGGAGAT	Exon 4
R4.2	TGTCGGGTGTCGGCTCTTCT	Exon 4
F4.3	AAGGACGCGAACGAGGCGCTC	Exon 4
R4.4	CTCAGATCCCTGAAGGTGAGGATTTG	Exon 4
R8.1	CAACATGAACTGCAAGTTGTC	Exon 8
F8.1	GACGTAGGCCATGTTGTTCTCGAC	Exon 8
F8.2	GTTCAGGCGATTGCGCCACAGTC	Exon 8
R9	GAAGTCCGCGACGAGGCAGAG	Exon 9
R10.1	CCGCTGGTGGGTGCAGCGAGGC	Exon 10
F10.1	GCCTCGCTGCACCCACAGCG	Exon 10
R11.1	CGAAGTGACCTGGAACGCGG	Exon 11
F11.2	TTTCGTCTCCTCAGCA	Exon 11
F11.3	ATGCCTCTCCTGCGAGAACTG	Exon 11
R12.1	GGTCGAATTGTCCGTTATGG	Exon 12
R12.2	ATCGCTCTGCAAGTCGCCGA	Exon 12
F12.2	TCGGCGACTTGCAGAGCGAT	Exon 12
R12.3	CATGACTCCAGCTTCCACGAC	Exon 12
R13.1	GTTTCTGTTGAAGGCGTTGC	Exon 13
F13.1	GCAACGCCTTCAACAGAAAC	Exon 13
R14.1	CTCTTTCCATCGTTCGTTGGTTCG	Exon 14
F14.2	CTCCACCCGATCGTCCCTGAA	Exon 14
F16	CAGCTGCGATTCTCCGAA	Exon 16
R17	CCGCACTCATTCCATAAATC	Exon 17
F17	GATTTATGGAATGAGTGCGG	Exon 17
R18	AAAGTGCTCTCCGTCTTGGC	Exon 18
F19	CGGCGGTGCGGCTGGTTCATGTG	Exon 19
R20.1	CAACACACGGGACAAAGCGAAG	Exon 20
R20.2	CTACGGCTTGTTCTGCCAGCTGTCCGC	Exon 20
3'UTR primer		
R.I	C GACTACGCCGCGCCG	32 bp 3' stop codon
TgPREX polymerase cloning		
Tg_polymeraseF	GGGGGCTAGCATGCCTCTCCTGCGGAGAACTG	Exon 11

The cell suspension was thawed, immediately supplemented with 0.1× protease inhibitor cocktail (EMD Biosciences) and sonicated on ice. The following procedures were carried out at 4 °C. After centrifugation at 20 000 *g* for 30 min, the supernatant was preserved, and the cell pellet resuspended in 15 ml of buffer A; the sonification and centrifugation

steps were repeated and the supernatant fractions were combined (Fraction I). Fraction I was loaded onto a DEAE cellulose (Sigma) column (4.9 cm<sup>2</sup> × 4 cm) equilibrated in buffer A and the column was washed with 100 ml of buffer A. Proteins were eluted by using a 120 ml linear gradient of 50–500 mM NaCl in buffer A. Fractions (5 ml) containing DNA



polymerase activity were pooled (Fraction II) and dialysed against buffer B [50 mM Tris-HCl (pH 8.0), 50 mM NaCl, 1 mM EDTA, 0.1 mM PMSF, 2 mM  $\beta$ -mercaptoethanol, 2 mM benzamidine and 10% (w/v) glycerol]. Following dialysis, Fraction II was applied to a P-11 (GE Healthcare, Piscataway, NJ) column (1.8 cm<sup>2</sup> × 7 cm) equilibrated in buffer B. The column was washed with 60 ml of buffer B, followed by elution with a 60 ml linear gradient of 50–500 mM NaCl in buffer B. The fractions (2.5 ml) were analysed in 10% SDS-polyacrylamide gels, and fractions containing DNA polymerase activity were pooled (Fraction III) and dialysed against buffer C [50 mM sodium phosphate (pH 8.0), 250 mM NaCl, 10 mM imidazole, 1 mM  $\beta$ -mercaptoethanol, 2 mM benzamidine and 5% (w/v) glycerol]. Dialysed Fraction III was loaded onto a Qiagen nickel-nitrilotriacetic acid (Ni-NTA) affinity column (1.77 cm × 3 cm) equilibrated in buffer C. The column was washed with 25 ml of buffer C and then with a 30 ml linear gradient of 10–300 mM imidazole (pH 8.0) in buffer C. Fractions (2.5 ml) containing TgPREX polymerase proteins, identified by electrophoresis in 10% SDS-polyacrylamide gels, were pooled (fraction IV). After dialysis against storage buffer [50 mM Tris-HCl (pH 7.4), 1 mM DTT, 30 mM NaCl and 10% (w/v) glycerol], the preparation was concentrated using an Amicon filter unit (MW cut-off 50 000). Aliquots were made and stored at –80 °C. The purified TgPREX polymerase protein (Fraction IV) appeared homogenous, as assessed in a 10% SDS-polyacrylamide gel followed by the Commassie Blue G-250 staining. Protein concentration was determined by the Bradford reaction (Bradford, 1976) using the Bio-Rad Protein Assay with bovine serum albumin as a standard.

#### Mass spectrometric analysis of the expressed protein

The purified protein was used for trypsin digestion (Bridges *et al.* 2008). Tryptic peptide samples were separated on an LC system (Famos/Switchos/Ultimate, LC Packings) before being analysed by electrospray ionisation (ESI) Mass Spectrometry (MS) on a Q-STAR<sup>®</sup> Pulsar i hybrid LC/MS/MS System. Peptide separation was performed on a Pepmap C18 reversed phase column (LC Packings), using a 5–85% v/v acetonitrile gradient (in 0.5% v/v formic acid) run over 45 min. The flow rate was maintained at 0.2  $\mu$ l/min. Mass spectrometric analysis was performed using a 3-sec survey MS scan followed by up to 4 MS/MS analyses of the most abundant peptides (3 sec per peak) in Information Dependent Acquisition (IDA) mode, choosing 2+ to 4+ ions above threshold of 30 counts, with dynamic exclusion for 120 sec.

Data generated from the Q-STAR<sup>®</sup> Pulsar i hybrid mass spectrometer was analysed using Applied Biosystems Analyst QS (v1.1) software and the

automated Matrix Science Mascot Daemon server (v2.1.06). Protein identifications were assigned using the Mascot search engine, which gives each protein a probability-based MOWSE score. In all cases variable methionine oxidation was allowed in searches. An MS tolerance of 1.2 Da for MS and 0.4 Da for MS/MS analysis was used.

#### DNA polymerase activity assay

DNA polymerase activity was assayed as described (Glick *et al.* 2002). Briefly, the activity was measured at 37 °C for 15 min in 10  $\mu$ l reaction mixtures containing 1  $\mu$ g of activated calf thymus DNA, 25  $\mu$ M each dNTP, 1  $\mu$ Ci  $\alpha$ [<sup>32</sup>P] dTTP (3000 Ci/mmol) (PerkinElmer, Waltham, MA, USA), and 1  $\mu$ l of enzyme in 30 mM Tris-HCl, pH 7.0, 10 mM MgCl<sub>2</sub>, 1 mM DTT, 250  $\mu$ g BSA, and 2.5% glycerol. The reaction was terminated by the addition of 100  $\mu$ l of 0.1 M sodium pyrophosphate/0.05 M EDTA. An aliquot (100  $\mu$ l) of the reaction mixture was transferred to a well in a 96-microwell<sup>®</sup> plate (Biodyne<sup>®</sup> B; NUNC<sup>™</sup>) mounted on a 96-vacuum manifold (Beckman Coulter, Fullerton, CA, USA). The plates were washed 3 times with 250  $\mu$ l of 0.1 M sodium pyrophosphate. The filter was removed from the plate, dried, and the amount of radioactivity associated with the filter was quantified by phosphorimager analysis using ImageQuant<sup>®</sup> software (GE Healthcare, Piscataway, NJ).

#### Software

Vector NTI Advance<sup>™</sup> suite 9 and 10 (Infomax 2003 from Invitrogen) was used for sequence analysis. The exon-intron structure of the *Tgprex* gene was created using GSDS, a gene structure display software (<http://gsds.cbi.pku.edu.cn/>) (Guo *et al.* 2007). SignalP 3.0 Server (<http://www.cbs.dtu.dk/services/SignalP/>) (Bendtsen *et al.* 2004; Nielsen *et al.* 1997; Nielsen and Krogh, 1998), SIG-Pred ([http://www.bioinformatics.leeds.ac.uk/prot\\_analysis/Signal.html](http://www.bioinformatics.leeds.ac.uk/prot_analysis/Signal.html)), SIGFIND (<http://139.91.72.10/sigfind/sigfind.html>), PrediSi (Hiller *et al.* 2004) and ChloroP 1.1 Server (<http://www.cbs.dtu.dk/services/ChloroP/#submission>) (Emanuelsson *et al.* 1999) were used for signal and transit peptide identification of TgPREX protein.

## RESULTS

#### Identification of a PREX homologue in ToxoDB

Two genes (55.m04962/TGME49\_061920 and 55.m04960/TGME49\_061800) were identified in ToxoDB (release 4 and 5 respectively) on chromosome VIIb of *T. gondii* genomic sequence on contigs 994 270 and 994 314 in the antisense strand (Kissinger *et al.* 2003). The coding sequence of this region




Fig. 2. SDS-PAGE and functional analysis of the purified TgPREX polymerase protein. The SDS-PAGE of purified TgPREX revealed a 114 kDa protein band in purified (2) followed by concentrated (1) protein samples. Functional properties of the purified TgPREX protein. (B) For TgPREX DNA polymerase activity  $Mg^{2+}$  ions are required. In the absence of  $Mg^{2+}$  ions, the amount of incorporated  $[\alpha]^{32}P$ -dTTP by the TgPREX to the activated calf thymus DNA is similar to that without the enzyme addition (Lanes 3 and 2, respectively). The amounts of incorporated  $[\alpha]^{32}P$ -dTTP to the activated calf thymus DNA increases as the concentrations (3·1, 6·3, 12·5, 25, 50 and 100 nm, respectively) of TgPREX increases in the reaction (Lanes 4–9, respectively). The DNA polymerase activity of *E. coli* Klenow (1 unit) is also shown as a control (Lane 1). (C) TgPREX polymerase activity measured as a function of  $MgCl_2$  concentration; (D) TgPREX polymerase activity measured as a function of pH; (E) TgPREX polymerase activity measured as a function of temperature. Error bars represent the standard deviation of 3 independent experimental determinations.

(1 488 133 to 1 515 551 bp) on chromosome VIIb was clarified by PCR amplification from *T. gondii* cDNA using primer pairs delineated in Fig. 1A.

Following sequencing, it appeared that a single gene homologous to *Pfprex* is present in *T. gondii* cDNA; thereafter, we call this gene *Tgprex*. The *Tgprex* gene comprises of 20 exons as shown in Fig. 1A. The 5' end of the ORF was elucidated by 5' RACE (Fig. 1B). The initiation codon of the ORF (Fig. 1C) was confirmed by comparing the 2345 bp long sequenced PCR product from the cDNA with that of genomic DNA (Fig. 1C). Sequencing of the 1766 bp long cDNA PCR product in comparison to genomic DNA PCR (Fig. 1D) confirmed the connection between the putative primase-helicase and the polymerase region of the *Tgprex* gene. The putative domain structure of the translated TgPREX protein is represented in Fig. 1E.

In summary, a 7740 bp long ORF, (Tgondii-ORF7740/*Tgprex*, GenBank Accession no. FJ665392) was identified from the sequenced region of chromosome VIIb (between 1,491,400 and 1,512,118). A sequencing gap between contigs 994 270 and 994 314 in the database was sequenced from both the genomic DNA and the cDNA of the RH strain of the organism and a 146 bp long sequence was subsequently deposited in the database.

#### Prediction of localization of TgPREX in silico

The TgPREX translated protein sequence possessed a primase, helicase and polymerase homologous domain structure (Fig. 1E) similar to that of the PREX protein of *Plasmodium*. The peptide region of the TgPREX protein that separated the primase-helicase and the polymerase domains has no similarity to the spacer region occupying the same region of the PfPREX predicted peptide sequence.

*T. gondii* uses the same general mechanism to mediate transport of proteins into the apicoplast as *Plasmodium* (He *et al.* 2001). A bipartite leader peptide, including a primary secretory domain followed by a secondary plastid transit domain, can be predicted by specialized tools including PlasmoAP (Foth *et al.* 2003) and PATS (Waller *et al.* 1998; Zuegge *et al.* 2001) in PlasmoDB. These programmes, however, failed in systemically identifying the apicoplast targeted proteins in *T. gondii* (Harb *et al.* 2004). For example, known *T. gondii* nuclear encoded apicoplast proteins (e.g. acyl carrier protein [AAC63956], beta-hydroxyacyl-ACP dehydratase [AAC72191], small ribosomal protein S9 [AAC63957], large ribosomal protein L28 [AAC63958] (Waller *et al.* 1998) and a putative ferredoxin NADP<sup>+</sup> oxidoreductase [CAC15394] (Vollmer *et al.* 2001) were all

Table 2. PREX homologous genes in other apicomplexa

Organism	Gene/Chromosome/ Contig/Protein	Probability
<i>Toxoplasma gondii</i>		
<i>Plasmodium yoelii</i>	Protein XP_729420	2e-136
<i>Plasmodium berghei</i>	Protein XP_679621	7e-135
<i>Plasmodium vivax</i>	Protein XP_001616839	2e-131
<i>Plasmodium knowlesi</i>	Protein XP_002260883	1e-129
<i>Plasmodium falciparum</i>	Protein XP_001348285	8e-106
<i>Plasmodium chabaudi</i>	Protein XP_746033	4e-135
<i>Theileria annulata</i>	Protein XP_954352	9e-131
<i>Theileria parva</i>	Protein XP_765913	3e-134
<i>Babesia bovis</i>	Protein XP_001610510	4e-148

unrecognized. For each of these proteins, however, the signal peptide component was recognized by SignalP 3.0 and the plastid targeting peptide following the signal peptide was recognized by ChloroP 1.1.

The N-terminal end of the TgPREX protein was analysed sequentially for the presence of a bipartite leader peptide. A 70 amino acid long N-terminal sequence of TgPREX was used as a query in PrediSi and SIGFIND software. Amino acid residues from 30 to 70 were used as a query in SignalP 3.0 as the maximum length of sequence input is restricted to 30 in this programme. All the analyses predicted TgPREX as an apicoplast-localized protein similar to its homologous counterpart PfPREX whose N-terminus was shown to direct GFP to the apicoplast (Seow *et al.* 2005). TgPREX appears to possess a putative, comparatively long N-terminal signal peptide, 60 to 61 amino acids in length. The cleavage sequence has been identified within the multiple serine residues (VLS-SS or VLSS-SS). The signal peptide was followed by a 61-amino acid long, putative plastid transit peptide as identified by the ChloroP1.1 programme.

#### Analysis of the PREX polymerase

The recombinant polymerase domain of the TgPREX, with a predicted mass of 114 kDa was expressed in *E. coli*. The histidine-tagged protein was purified by Ni<sup>2+</sup> affinity chromatography. The protein was analysed by SDS-PAGE and the identity was confirmed by mass spectrometry. The identified peptides are detailed in the Supplementary Data 1 (Online version only).

The purified TgPREX polymerase recombinant protein (Fraction IV) (Fig. 2A) was tested for its DNA polymerase activity. Activity was optimal at a temperature of 50 °C (Fig. 1C) and at pH 7.0 (Fig. 1D). The enzyme required Mg<sup>2+</sup> ions for activity (Fig. 1B), which reached the optimum at 5 mM MgCl<sub>2</sub> (Fig. 1E).

#### DISCUSSION

PREX, the plastid replication and repair enzyme complex of apicomplexan parasites was first identified in *Plasmodium*. Homology to DNA primase, helicase and polymerase in other systems indicated that the gene codes for all of these key functions. The presence of a predicted plastid targeting sequence and the fact that a similar sequence in *P. falciparum* guides marker proteins to the apicoplast suggest that PREX is instrumental in the replication of apicoplast DNA.

Among the annotated genes related to *Tgprex* in ToxoDB, TGME49\_061800 expression is linked to one EST (Expressed Sequence Tag) and 7.6 normalized SAGE (serial analysis of gene expression) tag counts (Gajria *et al.* 2008). There are no such data related to the TGME\_061920 gene. Mass spectrometry analysis has identified 2 peptide sequences (ARPLSPEHSALNESAGCAR and LFLESATPVPHAQILTFR) (Xia *et al.* 2008) which correspond to peptides in exon 4 and 5 of the sequenced *Tgprex* gene respectively. The lack of good EST coverage for this gene probably corresponds to a low level expression as documented in ToxoDB at the 32.2 and 25.2 percentile for TGME49\_061800 and TGME\_061920, respectively. A similarly low level of expression was also observed in *P. falciparum* (Le Roch *et al.* 2003).

BLAST searches identified full length PREX orthologues in other *Plasmodium* species, in *Theileria parva* [XP\_765913] and *Theileria annulata* [XP\_954352] and in *Babesia bovis* (XP\_001610510) (Table 2) (Alignment in Supplementary Data 2 – Online version only). Homologues representing fragments of the total predicted protein were also identified in *Babesia bigemina*, *Eimeria tenella*, *Neospora caninum* and *Sarcocystis neurona* where respective genome sequencing is incomplete. Interestingly, no homologous protein was found in the *Cryptosporidium* species. Genome sequencing also suggests that *Cryptosporidium hominis* (Xu *et al.* 2004) and *Cryptosporidium parvum* (Abrahamsen *et al.* 2004) lack an apicoplast.



Interestingly, the *prex* gene of *T. gondii* differs from that of *P. falciparum* in that it is interrupted by 20 introns. The intron footprint of eukaryotic, nuclear-encoded genes with possible prokaryotic cyanobacterial ancestry has made a significant contribution to our understanding of gene evolution. The presence of introns in such genes suggests the insertion of introns into pre-assembled genes of eukaryotes late in evolution ('intron late') as the ancestral cyanobacterial genes were devoid of introns. Similar comparison of ribosomal protein S9, L28 and acyl carrier protein genes between *P. falciparum* and *T. gondii* suggests a process of continuous intron insertion during evolution even after the divergence of *Plasmodium* and *T. gondii* from their common ancestor (Schaap *et al.* 2001).

The polymerase function of this putative protein, identified only in apicoplast-bearing apicomplexans, was confirmed in *T. gondii*. The nuclear encoded PREX appears to serve as a replicative enzyme for the 35 kb genome of the apicoplast. The participation of nuclear-encoded proteins in apicoplast replication has also been suggested for DNA gyrase subunits A and B (Dar *et al.* 2007; Raghu Ram *et al.* 2007), a DNA ligase and 2 hypothetical proteins bearing similarities with DNA-repair proteins (Dahl and Rosenthal, 2008).

Given that inhibitors of other enzymes involved in plastid replication (e.g. inhibitors of DNA gyrase) are toxic to apicomplexan parasites the PREX protein complex may be considered a target for chemotherapy. This may apply particularly to the primase and polymerase domains since the helicase domain has substantial homology to the Twinkle helicase that is active in mitochondria of most eukaryotic species.

We thank Dr Richard J. S. Burchmore of Sir Henry Wellcome Functional Genomics Facility, Institute of Biomedical and Life Sciences, University of Glasgow, for performing the MS analysis of protein in our study. A.M. was funded by the Wellcome Trust as part of their "Molecular Functions in Disease" programme.

## REFERENCES

- Abrahamsen, M. S., Templeton, T. J., Enomoto, S., Abrahante, J. E., Zhu, G., Lancto, C. A., Deng, M., Liu, C., Widmer, G., Tzipori, S., Buck, G. A., Xu, P., Bankier, A. T., Dear, P. H., Konfortov, B. A., Spriggs, H. F., Iyer, L., Anantharaman, V., Aravind, L. and Kapur, V. (2004). Complete genome sequence of the apicomplexan, *Cryptosporidium parvum*. *Science* **304**, 441–445.
- Bendtsen, J. D., Nielsen, H., von Heijne, G. and Brunak, S. (2004). Improved prediction of signal peptides: SignalP 3.0. *Journal of Molecular Biology* **340**, 783–795.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**, 248–254.
- Bridges, D. J., Pitt, A. R., Hanrahan, O., Brennan, K., Voorheis, H. P., Herzyk, P., de Koning, H. P. and Burchmore, R. J. (2008). Characterisation of the plasma membrane subproteome of bloodstream form *Trypanosoma brucei*. *Proteomics* **8**, 83–99.
- Campbell, S. A., Richards, T. A., Mui, E. J., Samuel, B. U., Coggins, J. R., McLeod, R. and Roberts, C. W. (2004). A complete shikimate pathway in *Toxoplasma gondii*: an ancient eukaryotic innovation. *International Journal for Parasitology* **34**, 5–13.
- Chomczynski, P. and Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry* **162**, 156–159.
- Dahl, E. L. and Rosenthal, P. J. (2008). Apicoplast translation, transcription and genome replication: targets for antimalarial antibiotics. *Trends in Parasitology* **24**, 279–284.
- Dar, M. A., Sharma, A., Mondal, N. and Dhar, S. K. (2007). Molecular cloning of apicoplast-targeted *Plasmodium falciparum* DNA gyrase genes: unique intrinsic ATPase activity and ATP-independent dimerization of PfGyrB subunit. *Eukaryotic Cell* **6**, 398–412.
- Emanuelsson, O., Nielsen, H. and von Heijne, G. (1999). ChloroP, a neural network-based method for predicting chloroplast transit peptides and their cleavage sites. *Protein Science* **8**, 978–984.
- Ferguson, D. J., Campbell, S. A., Henriquez, F. L., Phan, L., Mui, E., Richards, T. A., Muench, S. P., Allary, M., Lu, J. Z., Prigge, S. T., Tomley, F., Shirley, M. W., Rice, D. W., McLeod, R. and Roberts, C. W. (2007). Enzymes of type II fatty acid synthesis and apicoplast differentiation and division in *Eimeria tenella*. *International Journal for Parasitology* **37**, 33–51.
- Fichera, M. E. and Roos, D. S. (1997). A plastid organelle as a drug target in apicomplexan parasites. *Nature, London* **390**, 407–409.
- Foth, B. J., Ralph, S. A., Tonkin, C. J., Struck, N. S., Fraunholz, M., Roos, D. S., Cowman, A. F. and McFadden, G. I. (2003). Dissecting apicoplast targeting in the malaria parasite *Plasmodium falciparum*. *Science* **299**, 705–708.
- Gajria, B., Bahl, A., Brestelli, J., Dommer, J., Fischer, S., Gao, X., Heiges, M., Iodice, J., Kissinger, J. C., Mackey, A. J., Pinney, D. F., Roos, D. S., Stoeckert, C. J., Jr., Wang, H. and Brunk, B. P. (2008). ToxoDB: an integrated *Toxoplasma gondii* database resource. *Nucleic Acids Research*, **36**, D553–D556.
- Glick, E., Anderson, J. P. and Loeb, L. A. (2002). In vitro production and screening of DNA polymerase eta mutants for catalytic diversity. *Biotechniques* **33**, 1136–1142, 1144.
- Guo, A. Y., Zhu, Q. H., Chen, X. and Luo, J. C. (2007). [GSDS: a gene structure display server]. *Yi. Chuan* **29**, 1023–1026.
- Harb, O. S., Chatterjee, B., Fraunholz, M. J., Crawford, M. J., Nishi, M. and Roos, D. S. (2004). Multiple functionally redundant signals mediate

- targeting to the apicoplast in the apicomplexan parasite *Toxoplasma gondii*. *Eukaryotic Cell* **3**, 663–674.
- He, C. Y., Striepen, B., Pletcher, C. H., Murray, J. M. and Roos, D. S.** (2001). Targeting and processing of nuclear-encoded apicoplast proteins in plastid segregation mutants of *Toxoplasma gondii*. *Journal of Biological Chemistry* **276**, 28436–28442.
- Hiller, K., Grote, A., Scheer, M., Munch, R. and Jahn, D.** (2004). PrediSi: prediction of signal peptides and their cleavage positions. *Nucleic Acids Research* **32**, W375–W379.
- Kissinger, J. C., Gajria, B., Li, L., Paulsen, I. T. and Roos, D. S.** (2003). ToxoDB: accessing the *Toxoplasma gondii* genome. *Nucleic Acids Research* **31**, 234–236.
- Le Roch, K. G., Zhou, Y., Blair, P. L., Grainger, M., Moch, J. K., Haynes, J. D., De, L. V., Holder, A. A., Batalov, S., Carucci, D. J. and Winzeler, E. A.** (2003). Discovery of gene function by expression profiling of the malaria parasite life cycle. *Science* **301**, 1503–1508.
- McFadden, G. I., Reith, M. E., Munholland, J. and Lang-Unnasch, N.** (1996). Plastid in human parasites. *Nature, London* **381**, 482.
- McLeod, R., Muench, S. P., Rafferty, J. B., Kyle, D. E., Mui, E. J., Kirisits, M. J., Mack, D. G., Roberts, C. W., Samuel, B. U., Lyons, R. E., Dorris, M., Milhous, W. K. and Rice, D. W.** (2001). Triclosan inhibits the growth of *Plasmodium falciparum* and *Toxoplasma gondii* by inhibition of apicomplexan Fab I. *International Journal for Parasitology* **31**, 109–113.
- Nielsen, H., Engelbrecht, J., Brunak, S. and von Heijne, G.** (1997). Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Engineering Design and Selection* **10**, 1–6.
- Nielsen, H. and Krogh, A.** (1998). Prediction of signal peptides and signal anchors by a hidden Markov model. *Proceedings of the International Conference on Intelligent Systems for Molecular Biology* **6**, 122–130.
- Raghu Ram, E. V., Kumar, A., Biswas, S., Kumar, A., Chaubey, S., Siddiqi, M. I. and Habib, S.** (2007). Nuclear gyrB encodes a functional subunit of the *Plasmodium falciparum* gyrase that is involved in apicoplast DNA replication. *Molecular and Biochemical Parasitology* **154**, 30–39.
- Ralph, S. A., D’Ombrain, M. C. and McFadden, G. I.** (2001). The apicoplast as an antimalarial drug target. *Drug Resistance Updates* **4**, 145–151.
- Roberts, C. W. and Alexander, J.** (1992). Studies on a murine model of congenital toxoplasmosis: vertical disease transmission only occurs in BALB/c mice infected for the first time during pregnancy. *Parasitology* **104**, 19–23.
- Roos, D. S.** (1999). The apicoplast as a potential therapeutic target in *Toxoplasma* and other apicomplexan parasites: some additional thoughts. *Parasitology Today* **15**, 41.
- Schaap, D., van Poppel, N. F. and Vermeulen, A. N.** (2001). Intron invasion in protozoal nuclear encoded plastid genes. *Molecular and Biochemical Parasitology* **115**, 119–121.
- Seow, F., Sato, S., Janssen, C. S., Riehle, M. O., Mukhopadhyay, A., Phillips, R. S., Wilson, R. J. and Barrett, M. P.** (2005). The plastidic DNA replication enzyme complex of *Plasmodium falciparum*. *Molecular and Biochemical Parasitology* **141**, 145–153.
- Soldati, D.** (1999). The apicoplast as a potential therapeutic target in and other apicomplexan parasites. *Parasitology Today* **15**, 5–7.
- Striepen, B., Crawford, M. J., Shaw, M. K., Tilney, L. G., Seeber, F. and Roos, D. S.** (2000). The plastid of *Toxoplasma gondii* is divided by association with the centrosomes. *Journal of Cell Biology* **151**, 1423–1434.
- Vollmer, M., Thomsen, N., Wiek, S. and Seeber, F.** (2001). Apicomplexan parasites possess distinct nuclear-encoded, but apicoplast-localized, plant-type ferredoxin-NADP<sup>+</sup> reductase and ferredoxin. *Journal of Biological Chemistry* **276**, 5483–5490.
- Waller, R. F., Keeling, P. J., Donald, R. G., Striepen, B., Handman, E., Lang-Unnasch, N., Cowman, A. F., Besra, G. S., Roos, D. S. and McFadden, G. I.** (1998). Nuclear-encoded proteins target to the plastid in *Toxoplasma gondii* and *Plasmodium falciparum*. *Proceedings of the National Academy of Sciences, USA* **95**, 12352–12357.
- Wilson, R. J., Williamson, D. H. and Preiser, P.** (1994). Malaria and other Apicomplexans: the “plant” connection. *Infectious Agents and Disease* **3**, 29–37.
- Xia, D., Sanderson, S. J., Jones, A. R., Prieto, J. H., Yates, J. R., Bromley, E., Tomley, F. M., Lal, K., Sinden, R. E., Brunk, B. P., Roos, D. S. and Wastling, J. M.** (2008). The proteome of *Toxoplasma gondii*: integration with the genome provides novel insights into gene expression and annotation. *Genome Biology* **9**, R116.
- Xu, P., Widmer, G., Wang, Y., Ozaki, L. S., Alves, J. M., Serrano, M. G., Puiu, D., Manque, P., Akiyoshi, D., Mackey, A. J., Pearson, W. R., Dear, P. H., Bankier, A. T., Peterson, D. L., Abrahamsen, M. S., Kapur, V., Tzipori, S. and Buck, G. A.** (2004). The genome of *Cryptosporidium hominis*. *Nature, London* **431**, 1107–1112.
- Zuegge, J., Ralph, S., Schmuker, M., McFadden, G. I. and Schneider, G.** (2001). Deciphering apicoplast targeting signals—feature extraction from nuclear-encoded precursors of *Plasmodium falciparum* apicoplast proteins. *Gene* **280**, 19–26.