



Hinks, A. et al. (2012) *Investigation of rheumatoid arthritis susceptibility loci in juvenile idiopathic arthritis confirms high degree of overlap*. *Annals of the Rheumatic Diseases*, 71 (7). pp. 1117-1121. ISSN 0003-4967

Copyright © 2012 BMJ

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

The content must not be changed in any way or reproduced in any format or medium without the formal permission of the copyright holder(s)

When referring to this work, full bibliographic details must be given

<http://eprints.gla.ac.uk/76454>

Deposited on: 15 March 2013

EXTENDED REPORT

Investigation of rheumatoid arthritis susceptibility loci in juvenile idiopathic arthritis confirms high degree of overlap

Anne Hinks,¹ Joanna Cobb,¹ Marc Sudman,² Stephen Eyre,¹ Paul Martin,¹ Edward Flynn,¹ Jonathon Packham,³ Childhood Arthritis Prospective Study (CAPS),* UK RA Genetics (UKRAG) Consortium,* British Society of Paediatric and Adolescent Rheumatology (BSPAR) Study Group,* Anne Barton,¹ Jane Worthington,¹ Carl D Langefeld,⁴ David N Glass,² Susan D Thompson,² Wendy Thomson¹

► Additional supplementary tables are published online only. To view these files, please visit the journal online (<http://ard.bmj.com/content/71/7.toc>).

¹Arthritis Research UK Epidemiology Unit, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK

²Division of Rheumatology, Cincinnati Children's Hospital Medical Centre, Cincinnati, Ohio, USA

³Keele University, Stoke on Trent, UK

⁴Department of Biostatistical Sciences, Wake Forest University School of Medicine, Winston-Salem, North Carolina, USA

*Members of the Childhood Arthritis Prospective Study (CAPS), UKRAG Consortium and the British Society of Paediatric and Adolescent Rheumatology (BSPAR) Study Group are shown at the end of the paper

Correspondence to

Anne Hinks, University of Manchester, Arthritis Research UK Epidemiology Unit, Manchester M13 9PT, UK; anne.hinks@manchester.ac.uk

Received 13 December 2011
Accepted 20 December 2011
Published Online First
31 January 2012

ABSTRACT

Objectives Rheumatoid arthritis (RA) shares some similar clinical and pathological features with juvenile idiopathic arthritis (JIA); indeed, the strategy of investigating whether RA susceptibility loci also confer susceptibility to JIA has already proved highly successful in identifying novel JIA loci. A plethora of newly validated RA loci has been reported in the past year. Therefore, the aim of this study was to investigate these single nucleotide polymorphisms (SNP) to determine if they were also associated with JIA.

Methods Thirty-four SNP that showed validated association with RA and had not been investigated previously in the UK JIA cohort were genotyped in JIA cases (n=1242), healthy controls (n=4281), and data were extracted for approximately 5380 UK Caucasian controls from the Wellcome Trust Case-Control Consortium 2. Genotype and allele frequencies were compared between cases with JIA and controls using PLINK. A replication cohort of 813 JIA cases and 3058 controls from the USA was available for validation of any significant findings.

Results Thirteen SNP showed significant association (p<0.05) with JIA and for all but one the direction of association was the same as in RA. Of the eight loci that were tested, three showed significant association in the US cohort.

Conclusions A novel JIA susceptibility locus was identified, *CD247*, which represents another JIA susceptibility gene whose protein product is important in T-cell activation and signalling. The authors have also confirmed association of the *PTPN2* and *IL2RA* genes with JIA, both reaching genome-wide significance in the combined analysis.

share genetic susceptibility risk factors.^{3,4} JIA is a relatively rare disease and as such it has taken longer to collect the large and appropriately powered sample sizes required for genome-wide association studies (GWAS). International collaborations have been established and GWAS for JIA have been performed or are in progress.⁵ However, in the meantime, there are other approaches that can be used to identify genetic risk factors for JIA. We and other investigators have successfully exploited the fact that many complex autoimmune diseases share common genetic risk factors; for example, protein tyrosine phosphatase non-receptor 22 (*PTPN22*) and interleukin 2 receptor α (*IL2RA*) both confer susceptibility to multiple autoimmune diseases, such as type 1 diabetes (T1DM), rheumatoid arthritis (RA) and autoimmune thyroid disease, and markers in both these genes have been associated with JIA.^{4,6} Of the autoimmune diseases, RA shares the most similar clinical and pathological features with JIA, and we have previously reported considerable overlap in genetic susceptibility loci for the two diseases, finding evidence of association for four loci (*STAT4*, *TRAF1/C5*, *TNFAIP3* and *PRKCCQ*) with JIA.⁷ The first three of which have further evidence of association with JIA in independent cohorts.^{8,9} In the past few years many additional RA loci have been identified¹⁰⁻¹² and these represent excellent candidate JIA loci. Therefore the aim of this study was to investigate whether newly identified RA susceptibility loci also confer susceptibility to JIA.

MATERIALS AND METHODS

SNP selection

Single-nucleotide polymorphisms (SNP) that showed validated association with RA and had not been investigated previously in the UK JIA cohort were selected for genotyping. These were identified from three recent RA publications.¹⁰⁻¹² The first was a GWAS in cases and controls from North America and from that study SNP in *REL* (rs13031237), *BLK* (rs2736340) and *CTLA4* (rs231735) were selected for genotyping.¹⁰ The second was a meta-analysis of published GWAS in which 13 novel RA SNP were identified.¹¹ Finally, a second larger meta-

Juvenile idiopathic arthritis (JIA) is a collective term that encompasses all forms of arthritis, with an unknown cause, that have an onset before the age of 16 years and that persist for more than 6 weeks.¹ There are seven disease categories as defined by the International League of Associations for Rheumatology (ILAR) classification criteria² and, while there is heterogeneity between the subtypes in terms of disease presentation, clinical symptoms and prognosis, they do appear to



This paper is freely available online under the BMJ Journals unlocked scheme, see <http://ard.bmj.com/info/unlocked.dtl>

Clinical and epidemiological research

analysis of RA cases and controls identified a further 21 SNP with replicated association with RA. Of these, 10 had achieved genome-wide significance in the combined analysis and 11 had highly suggestive evidence of association although not exceeding the threshold for claims of genome-wide significance.¹² Results for three of these SNP (in *IL2RA*, *AFF3* and *c12orf30*) had previously been investigated in the UK JIA cohort,^{4 13 14} and so were not included in this study. In total, therefore, 34 SNP were selected for genotyping.

Subjects

UK cohort

DNA was available for 1242 UK Caucasian JIA patients from three sources: The British Society for Paediatric and Adolescent Rheumatology national repository of JIA (n=654); a group of UK Caucasian patients with long-standing JIA (n=201), described previously,¹⁵ and a third cohort collected as part of the Childhood Arthritis Prospective study, a prospective inception cohort study of JIA cases from five centres across the UK (n=387).¹⁶ JIA cases were classified according to ILAR criteria² (see supplementary table 1, available online only). Healthy Caucasian control DNA samples (n=4281) were available from five centres in the UK, collected as part of the UK RA genetics (UKRAG) consortium, as described previously.¹⁷ All individuals were recruited with ethics approval and provided informed consent (North-West Multi-Centre Research Ethics Committee (MREC 99/8/84) and the University of Manchester Committee on the Ethics of Research on Human Beings). Additional control genotyping data were extracted from the Wellcome Trust Case-Control Consortium 2 (WTCCC2) European Genome-Phenome Archive website (<http://www.ebi.ac.uk/ega/>). SNP had been genotyped on either the Illumina (n=5200) or Affymetrix (n=5380) platform. If a SNP had been genotyped on both platforms then the results from the analysis with the larger control set was presented.

All SNP were genotyped in the UK JIA cases using the Sequenom iPLEX MassARRAY platform according to manufacturer's instructions (Sequenom, San Diego, California, USA; <http://www.sequenom.com/>). A 90% sample quality control rate and 90% genotyping success rate was imposed.

US cohort

For validation of the findings in the UK cohort, genotype information was available from a cohort of 813 JIA cases and 3058 controls from the USA. These data are from a GWAS, which has previously been investigated for autoimmune overlap.⁵ For the JIA children, with polyarticular (rheumatoid factor negative) and oligoarticular JIA, approximately 95% of the cases were recruited at the Cincinnati Children's Hospital Medical Center (CCHMC) or as part of a National Institute of Arthritis and Musculoskeletal and Skin Diseases-supported JIA affected sibpair registry. The remaining cases were contributed by collaborating centres that included the Children's Hospital of Wisconsin, Schneider Children's Hospital and Children's Hospital of Philadelphia. JIA cases were classified according to ILAR criteria² and comprised cases from three subtypes of JIA (see supplementary table 1, available online only). The control cohort (n=3058) comprised 658 healthy children aged between 3 and 18 years, recruited from the general population to represent the geographical region served by CCHMC. In addition, 2400 healthy controls of European ancestry were genotyped at the Broad Institute on the Affymetrix SNP array 6.0, which have been collected as part of the Molecular Genetics of Schizophrenia GWAS. Both cases and controls were subjected to principal component analysis, and any outliers were excluded

from analysis. SNP were genotyped using the Affymetrix genome-wide human SNP array 6.0. The study had full ethics committee approval (CCHMC Institutional Review Board) and was fully compliant with the Declaration of Helsinki.⁵

Statistical analysis

Power calculations were performed using QUANTO to calculate the prior probability of detecting association in the current UK sample size at the allele frequency and effect sizes reported previously in RA.^{10–12} Calculations assumed a log-additive model and an α value of 0.05. Power calculations were also performed to determine the power of the US cohort to validate the findings of the UK cohort. Genotype and allele frequencies were compared between cases with JIA and controls using the Cochran–Armitage trend test implemented in PLINK¹⁸ and allelic OR and their 95% CI were calculated.

Analysis was performed in all JIA cases for the UK cohort initially but then the analysis was restricted to subtypes equivalent to those in the US cohort (see supplementary table 1, available online only) to allow for comparison of OR and for combining the data in a meta-analysis.

RESULTS

The JIA case number after quality control was 1229. The power calculation for each SNP is shown in supplementary table 3 (available online only).

Of the 34 SNP selected for genotyping, four failed to genotype (rs892188 (*ICAM1*), rs934734 (*SPRED2*), rs7155603 (*BATF*) and rs840016 (*CD247*), although a SNP in the *CD247* gene identified in the Raychaudhuri study was genotyped,¹¹ rs1773560, which has $r^2=0.68$ with rs840016. Therefore, there were 30 SNP for analysis. The SNP in *IRF5* (rs10488631) failed genotyping in the UKRAG control dataset, therefore the comparison was limited to the WTCCC2 control data while conversely the SNP in *TAGAP* (rs394581) failed to genotype in the WTCCC2 control cohort so the comparison was limited to the UKRAG control dataset.

Thirteen SNP showed nominal evidence of association with JIA ($p<0.05$) (table 1). Results for all SNP are provided in supplementary table 2 (available online only). Figure 1 shows a comparison between the association analysis results in RA and JIA. For all but one (*NHLH2*) of the 13 loci the direction of association was the same as that observed in the RA studies, with similar OR and sometimes stronger effect sizes in JIA (figure 1).

Of the 13 loci that showed nominal association with JIA in our UK cohort, eight had genotype data available in the US validation cohort (table 2). There were no data for the *IL2*, *ANKRD55*, *C5orf30* or the *IRF5* SNP. Of the eight SNP, three (rs1773560 in *CD247*, rs706778 in *IL2RA* and rs7234029 in *PTPN2*) showed association with JIA in the US cohort, and meta-analysis of the two cohorts strengthened the association.

DISCUSSION

The now well-established overlap of genetic susceptibility risk factors across quite clinically and phenotypically different autoimmune diseases has been an exciting discovery that has emerged from the GWAS era. It suggests there will be shared pathogenic pathways and the potential for shared therapies for these diseases. Furthermore, it can be exploited as a strategy for the identification of novel risk factors for related autoimmune diseases, including JIA. There is already compelling evidence that many susceptibility loci are shared between RA and JIA (*PTPN22*,^{5 6} *IL2RA*,^{4 19} *STAT4*,^{5 7 20} *IL2*,^{5 13 21} *TRAF1/C5*^{7 20}

Table 1 RA-associated SNP nominally associated with JIA (all JIA subtypes) ($p \leq 0.05$)

SNP	CHR	Position	Gene	Minor allele	Major allele	MAF cases	MAF controls	HWE controls	Case 11 (%)	Case 12 (%)	Case 22 (%)	Control 11 (%)	Control 12 (%)	Control 22 (%)	pTREND	OR 95% CI
rs4272626	1	116149950	NHLH2	T	C	0.34	0.36	0.79	517 (44.2)	511 (43.7)	142 (12.1)	3264 (40.8)	3702 (46.3)	1035 (12.9)	0.05	0.91 (0.83 to 1.00)
rs11586238	1	117064661	CD2	G	C	0.26	0.24	0.48	636 (54.4)	455 (38.9)	79 (6.8)	4586 (57.3)	2957 (37.0)	456 (5.7)	0.03	1.11 (1.01 to 1.23)
rs1773560	1	16568387	CD247	G	A	0.38	0.42	0.48	456 (39.2)	520 (44.7)	187 (16.1)	2652 (33.4)	3849 (48.5)	1443 (18.2)	0.0003	0.85 (0.78 to 0.93)
rs10919563	1	196967065	PTPRC	G	C	0.09	0.13	0.92	917 (83.0)	171 (15.5)	17 (1.5)	5994 (75.3)	1824 (22.9)	137 (1.7)	2.57×10^{-7}	0.67 (0.58 to 0.78)
rs874040	4	25717295	RBPJ	G	C	0.33	0.30	0.62	445 (44.4)	451 (45.0)	107 (10.7)	3938 (48.7)	3401 (42.0)	754 (9.3)	0.01	1.14 (1.03 to 1.26)
rs13119723	4	123437763	IL2, IL21	G	A	0.13	0.16	0.24	721 (76.0)	204 (21.5)	24 (2.5)	5631 (71.0)	2090 (26.3)	214 (2.7)	0.004	0.81 (0.71 to 0.93)
rs10040327*	5	55470977	ANKRD55	A	C	0.09	0.11	0.20	825 (82.4)	167 (16.7)	9 (0.9)	6417 (79.1)	1583 (19.5)	112 (1.4)	0.01	0.81 (0.69 to 0.95)
rs26232	5	102624619	C5orf30	T	C	0.29	0.33	0.35	506 (50.4)	417 (41.6)	80 (8.0)	3579 (45.0)	3536 (44.5)	831 (10.5)	0.0003	0.83 (0.75 to 0.92)
rs10488631	7	128381419	IRF5	C	T	0.13	0.11	0.47	902 (76.5)	257 (21.8)	20 (1.7)	4119 (79.6)	1003 (19.4)	54 (1.0)	0.009	1.20 (1.05 to 1.37)
rs2736340	8	11381382	BLK	T	C	0.27	0.24	0.88	522 (52.4)	406 (40.8)	68 (6.8)	4631 (57.1)	3002 (37.0)	481 (5.9)	0.006	1.16 (1.04 to 1.28)
rs706778	10	6138955	IL2RA	T	C	0.45	0.40	0.80	314 (31.4)	479 (47.9)	208 (20.8)	2926 (36.1)	3883 (47.9)	1304 (16.1)	5.61×10^{-5}	1.21 (1.10 to 1.33)
rs2872507	17	35294289	ORMDL3	A	G	0.52	0.48	0.75	221 (23.2)	474 (49.8)	256 (26.9)	2199 (27.3)	4001 (49.7)	1846 (22.9)	0.0009	1.18 (1.07 to 1.29)
rs7234029	18	12867060	PTPN2	G	A	0.19	0.16	0.33	766 (65.5)	362 (30.9)	42 (3.6)	5682 (71.1)	2127 (26.6)	182 (2.3)	1.25×10^{-5}	1.28 (1.15 to 1.43)

* $r^2=0.86$ with rs6859219 the single-nucleotide polymorphism (SNP) genotyped in the study by Stahl *et al.*¹²
Case 11, case genotype counts and percentage of major allele homozygotes; Case 12, case genotype counts and percentage of heterozygotes; Case 22, case genotype counts and percentage of minor allele homozygotes; Control 11, control genotype counts and percentage of major allele homozygotes; Control 12, control genotype counts and percentage of heterozygotes; Control 22, control genotype counts and percentage of minor allele homozygotes, p value for deviation from HWE; CHR, chromosome; HWE, Hardy-Weinberg equilibrium; JIA, juvenile idiopathic arthritis; MAF, minor allele frequency; pTREND, p value for the Armitage test for trend as generated in PLINK; RA, rheumatoid arthritis.




Figure 1 Plot of OR for minor allele for 13 loci showing nominal evidence of association with juvenile idiopathic arthritis (JIA), comparison with rheumatoid arthritis (RA). Odds ratios obtained from Raychaudhuri *et al.*¹¹ Stahl *et al.*¹² and Gregersen *et al.*¹⁰

amongst others). In this study we have identified and validated association with three novel loci, originally associated with RA.

One of these identifies a new JIA susceptibility gene, *CD247*, a good candidate for JIA as it encodes the zeta chain of the T-cell receptor-CD3 complex and adds another gene to the list of JIA susceptibility loci that play a role in T-cell activation and signalling (*PTPN22*, *IL2RA*, *PTPN2*, *SH2B3* and *STAT4*). In addition to the reported associations with RA, SNP within *CD247* have also been associated with coeliac disease²² and with systemic sclerosis.²³ The JIA-associated SNP is in complete linkage disequilibrium (LD) ($r^2=1$) with the SNP associated with systemic sclerosis (rs2056626) and in strong LD ($r^2=0.86$) with the SNP most significantly associated with coeliac disease (rs864537). The risk variant of the SNP has also been correlated with *cis* gene expression of *CD247*.^{22 24}

We found strong evidence for association of a SNP, rs7234029, in the protein tyrosine phosphatase non-receptor 2 (*PTPN2*) gene with JIA in our UK cohort. There is already replicated evidence for association of SNP within this gene with JIA in US and German cohorts.⁵ In the US study, the SNP with the strongest association was also rs7234029, and it reached genome-wide levels of significance; in addition there was association with four additional SNP. Meta-analysis of the UK JIA cohort study with the US data further established *PTPN2* as an important JIA susceptibility locus (combined p value 8.1×10^{-13}). The *PTPN2* gene has been associated with multiple autoimmune diseases, including T1DM,^{25 26} Graves' disease,²⁵ Crohn's disease²⁷ and coeliac disease.²⁶ In T1DM, fine-mapping suggests there are two independently associated SNP, rs45450798 and rs478582.²⁶ These SNP are only in modest LD with rs7234029. Further fine-mapping of the gene in RA and JIA will be required to identify the causal variant, whether there are multiple independent effects and whether the associations are the same across the different autoimmune diseases. *PTPN2* encodes a protein tyrosine phosphatase, similar to *PTPN22*, which also plays a role in the activation and regulation of T and B cells.

SNP within *IL2RA* have previously been shown to be associated with JIA using these same UK and US cohorts⁴ as well as many other autoimmune diseases such as RA,¹⁹ T1DM²⁸ and multiple sclerosis,²⁹ but the association is complex with multiple independent effects and different associations across the different diseases. The novel SNP associated here is in only

Table 2 Analysis of all JIA nominally associated SNP in oligoarthritis and polyarthritis subtypes in UK discovery cohort, the US validation cohort and combined meta-analysis

SNP	Chr	Position	Gene	Minor allele	Major allele	UK JIA oligoarthritis and RF-negative polyarthritis cases (n=798)			US JIA oligoarthritis and RF-negative polyarthritis cases (n=813)			Combined meta-analysis				
						MAF cases	MAF controls	pTREND	OR (95% CI)	MAF cases	MAF controls	pTREND	OR (95% CI)	p value		
rs4272626	1	116149950	NHLH2	T	C	0.34	0.36	0.07	0.90 (0.81 to 1.01)	0.36	0.37	0.33	0.95 (0.84 to 1.06)	0.05	0.92 (0.85 to 1.0)	0.55
rs11586238	1	117064661	CD2	G	C	0.26	0.24	0.10	1.11 (0.98 to 1.25)	0.22	0.24	0.39	0.94 (0.83 to 1.08)	0.56	1.03 (0.94 to 1.12)	0.08
rs1773560	1	165688387	CD247	A	G	0.39	0.42	0.008	0.86 (0.77 to 0.96)	0.38	0.41	0.01	0.87 (0.78 to 0.97)	2.8×10^{-4}	0.86 (0.8 to 0.92)	0.93
rs10919563	1	196967065	PTPRC	A	G	0.09	0.13	4.2×10^{-6}	0.65 (0.54 to 0.78)	0.14	0.12	0.08	1.16 (0.98 to 1.36)	0.04	0.88 (0.78 to 0.99)	3.3×10^{-6}
rs874040	4	25717295	ABR1	G	C	0.33	0.30	0.04	1.14 (1.01 to 1.28)	0.31	0.30	0.59	1.03 (0.92 to 1.16)	0.07	1.08 (0.99 to 1.18)	0.27
rs2736340	8	11381382	BLK	T	C	0.28	0.24	0.007	1.19 (1.05 to 1.36)	0.26	0.26	0.99	1.00 (0.88 to 1.13)	0.06	1.09 (1.0 to 1.19)	0.05
rs706778	10	6138955	IL2RA	T	C	0.45	0.40	0.002	1.20 (1.07 to 1.35)	0.46	0.40	0.0002	1.24 (1.11 to 1.38)	9.5×10^{-7}	1.22 (1.13 to 1.32)	0.74
rs2872507	17	35294289	ORMDL3	A	G	0.53	0.48	0.001	1.22 (1.08 to 1.37)	0.46	0.46	0.93	0.99 (0.89 to 1.11)	0.03	1.09 (1.01 to 1.18)	0.01
rs7234029	18	12867060	PTPN2	G	A	0.20	0.16	9.3×10^{-6}	1.35 (1.18 to 1.54)	0.23	0.16	1.3×10^{-8}	1.48 (1.29 to 1.69)	8.1×10^{-13}	1.41 (1.28 to 1.55)	0.33

Breslow-Day p value is a measure of whether there is significant difference in the OR in the two cohorts.

Chr, chromosome; JIA, juvenile idiopathic arthritis; MAF, minor allele frequency; pTREND, p value for the Armitage test for trend as generated in PLINK; SNP, single-nucleotide polymorphism.

weak LD with the most associated SNP from the previous study, rs2104286, $r^2=0.24$. However, conditional logistic regression analysis in the UK dataset suggests that rs2104286 is driving the association with JIA. After conditioning on rs2104286 there was no significant association at rs706778, whereas after conditioning on rs706778, there was still significant association at rs2104286 (data not shown). Meta-analysis of the UK and US studies found highly significant evidence for association with JIA (combined p value 3.3×10^{-8}), which now brings this locus to a genome-wide significance level.

In the initial analysis of the UK cohort only the most significant association was for a SNP in the protein-tyrosine phosphatase receptor type C (*PTPRC*) gene, which again is a promising candidate gene. This gene encodes the common leucocyte antigen, CD45, which is a haemopoietic cell-specific tyrosine phosphatase. CD45 is essential for the activation of T and B cells by mediating cell-to-cell contacts and regulating protein-tyrosine kinases involved in signal transduction.³⁰ However, this was not validated in the US cohort; in fact, there was a trend towards an association but in the opposite direction. There are a number of possible reasons for this, including a false positive in the initial UK dataset, a false negative in the US dataset, unrecognised population stratification, phenotypic heterogeneity and different environmental exposures. Analysis of this SNP in additional larger cohorts is required to confirm or refute association with this locus.

The JIA association findings were not validated in the US cohort for *CD2*, *RBPJ*, *BLK* and *ORMDL3*, and only *ORMDL3* remained significant in the combined analysis. The US cohort was underpowered to detect an effect with all of these loci apart from *ORMDL3* (see supplementary table 3, available online only), therefore further validation is needed.

In addition it should be noted that for the 16 SNP that were not significantly associated with JIA in this study, we had between 29% and 99% power to detect an association; therefore, for some of these SNP, larger datasets and meta-analyses will be required to exclude association with JIA completely.

JIA is a heterogeneous disease and it may be expected that there are genetic differences across the JIA subtypes and some subtype-specific effects have been identified.^{31 32} In this study our strategy was to look for shared autoimmune or inflammatory arthritis susceptibility loci, and it would be interesting to investigate whether these associations are common to all JIA subtypes or are restricted to some subtypes. However, stratified analysis leads to small sample sizes for many of the subtypes and further issues with multiple testing. Larger cohorts of the ILAR subtypes are required to improve the power to detect any subtype-specific effects. We have stratified our analysis just to investigate the oligoarthritis and rheumatoid factor-negative polyarthritis subtypes as these were the subtypes comprising the validation cohort.

For all these loci identified for JIA we have only tested one SNP, the SNP most significantly associated with RA from GWAS. In most cases for the genes identified to date for RA and other autoimmune diseases, the actual causal variant has yet to be identified, which may reduce the power of this study even further, so further fine-mapping of the genes/regions is now required.

In conclusion, the strategy of testing loci identified in other autoimmune diseases, such as RA, for association with JIA is proving successful in the identification of novel JIA loci and is a complementary approach to the hypothesis-free methods of GWAS. In the past few years we have investigated 45 loci that confer susceptibility to RA for association with JIA.^{7 13 33}

Of these, 22 show association ($p < 0.05$) with JIA in our UK cohort and 11 of these show validated association with JIA in independent cohorts.^{5 8 9 34 35} Clearly, there is a selection bias given the strategy used to select SNP to test; however, there still remains quite considerable overlap between two distinct clinical entities. Many of these loci also confer susceptibility to other autoimmune diseases and may thus represent genes controlling general immune function, the defects of which may lead to an enhanced autoimmune response. Despite the success of this approach in identifying novel JIA loci, large well-powered GWAS will be required to identify disease-specific loci for JIA and its subtypes.

Childhood Arthritis Prospective Study (CAPS): Eileen Baildam, Lynsey Brown, Joanne Buckley, Alice Chieng, Joyce Davidson, Michael Eltringham, Helen Foster, Mark Friswell, Janet Gardner-Medwin, Paul Gilbert, Kimmehyrich, Julie Jones, Sham Lal, Mark Lay, Carol Lydon, Alexandra Meijer, Vicki Price, Jane Sim, Maureen Todd, Peter Ward, Lucy Wedderburn.

UKRAG Consortium: Arthritis Research UK Epidemiology Unit, University of Manchester: Stephen Eyre, Anne Hinks, John Bowes, Edward Flynn, Paul Martin, Wendy Thomson, Anne Barton, Jane Worthington; Leeds Institute of Molecular Medicine, Section of Musculoskeletal Disease, University of Leeds: Stephen Martin, James I Robinson, Ann W Morgan, Paul Emery; School of Medicine and Biomedical Sciences, University of Sheffield: Anthony G. Wilson; Clinical and Academic Rheumatology, Kings College Hospital NHS Foundation Trust, University of London: Sophia Steer; Bone Research Group, Department of Medicine and Therapeutics, University of Aberdeen: Lynne Hocking, David M Reid; University of Oxford Institute of Musculoskeletal Sciences, University of Oxford: Pille Harrison, Paul Wordsworth

British Society of Paediatric and Adolescent Rheumatology (BSPAR) Study Group: M Abinum, M Becker, A Bell, A Craft, E Crawley, J David, H Foster, J Gardener-Medwin, J Griffin, A Hall, M Hall, A Herrick, P Hollingworth, L Holt, S Jones, G Pountain, C Ryder, T Southwood, I Stewart, H Venning, L Wedderburn, P Woo, S Wyatt

Acknowledgements This study makes use of data generated by the WTCCC2. A full list of the investigators who contributed to the generation of the data is available from <http://www.wtccc.org.uk>. Funding for the project was provided by the Wellcome Trust under award 085475. The authors gratefully acknowledge contributions from physicians at CCHMC, the Medical College of Wisconsin, Schneider Children's Hospital and the Children's Hospital of Philadelphia for the collection of patient samples, and the assistance of Sandy Kramer for patient recruitment at CCHMC and coordination of clinical information. The normal control DNA collection including all genotypes was supported and made available by CCHMC.

Funding This work was supported by Arthritis Research UK grant reference no. 17552 and the National Institutes of Health (RC1-AR-058587, P30-AR-473639 to SDT, K23-AR-50177 to SP, R01-AR-057106 to CDL and SDT, P01-AR048929 and N01-AR-42272 to DNG); the Federal Ministry of Education and Research Germany (BMBF) (grants 01GM0907 and 01 ZZ 0403 to JPH); the Val A. Browning Foundation; the Texas Scottish Rite Hospital for Children (grant 0305756 to CW and MP); and the Arthritis Foundation.

Ethics approval Ethics approval was obtained from the North-West Multi-Centre Research Ethics Committee (MREC 99/8/84) and the University of Manchester Committee on the Ethics of Research on Human Beings.

Patient consent Obtained.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Ravelli A, Martini A. Juvenile idiopathic arthritis. *Lancet* 2007;**369**:767–78.
- Petty RE, Southwood TR, Manners P, et al. International League of Associations for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001. *J Rheumatol* 2004;**31**:390–2.
- Hinks A, Barton A, John S, et al. Association between the *PTPN22* gene and rheumatoid arthritis and juvenile idiopathic arthritis in a UK population: further support that *PTPN22* is an autoimmunity gene. *Arthritis Rheum* 2005;**52**:1694–9.
- Hinks A, Ke X, Barton A, et al. Association of the *IL2RA/CD25* gene with juvenile idiopathic arthritis. *Arthritis Rheum* 2009;**60**:251–7.
- Thompson SD, Sudman M, Ramos PS, et al. The susceptibility loci juvenile idiopathic arthritis shares with other autoimmune diseases extend to *PTPN22*, *COG6*, and *ANGPT1*. *Arthritis Rheum* 2010;**62**:3265–76.
- Hinks A, Worthington J, Thomson W. The association of *PTPN22* with rheumatoid arthritis and juvenile idiopathic arthritis. *Rheumatology (Oxford)* 2006;**45**:365–8.
- Hinks A, Eyre S, Ke X, et al. Overlap of disease susceptibility loci for rheumatoid arthritis and juvenile idiopathic arthritis. *Ann Rheum Dis* 2010;**69**:1049–53.
- Prahalad S, Hansen S, Whiting A, et al. Variants in *TNFAIP3*, *STAT4*, and *C12orf30* loci associated with multiple autoimmune diseases are also associated with juvenile idiopathic arthritis. *Arthritis Rheum* 2009;**60**:2124–30.
- Albers HM, Kurreeman FA, Houwing-Duistermaat JJ, et al. The *TRAF1/C5* region is a risk factor for polyarthritis in juvenile idiopathic arthritis. *Ann Rheum Dis* 2008;**67**:1578–80.
- Gregersen PK, Amos CI, Lee AT, et al. REL, encoding a member of the NF-kappaB family of transcription factors, is a newly defined risk locus for rheumatoid arthritis. *Nat Genet* 2009;**41**:820–3.
- Raychaudhuri S, Thomson BP, Remmers EF, et al. Genetic variants at CD28, PRDM1 and *CD2/CD58* are associated with rheumatoid arthritis risk. *Nat Genet* 2009;**41**:1313–18.
- Stahl EA, Raychaudhuri S, Remmers EF, et al. Genome-wide association study meta-analysis identifies seven new rheumatoid arthritis risk loci. *Nat Genet* 2010;**42**:508–14.
- Hinks A, Eyre S, Ke X, et al. Association of the *AFF3* gene and *IL2/IL21* gene region with juvenile idiopathic arthritis. *Genes Immun* 2010;**11**:194–8.
- Hinks A, Martin P, Flynn E, et al. Investigation of type 1 diabetes and coeliac disease susceptibility loci for association with juvenile idiopathic arthritis. *Ann Rheum Dis* 2010;**69**:2169–72.
- Packham JC, Hall MA. Long-term follow-up of 246 adults with juvenile idiopathic arthritis: functional outcome. *Rheumatology (Oxford)* 2002;**41**:1428–35.
- Adib N, Hyrich K, Thornton J, et al. Association between duration of symptoms and severity of disease at first presentation to paediatric rheumatology: results from the Childhood Arthritis Prospective Study. *Rheumatology (Oxford)* 2008;**47**:991–5.
- Thomson W, Barton A, Ke X, et al. Rheumatoid arthritis association at 6q23. *Nat Genet* 2007;**39**:1431–3.
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;**81**:559–75.
- Barton A, Thomson W, Ke X, et al. Rheumatoid arthritis susceptibility loci at chromosomes 10p15, 12q13 and 22q13. *Nat Genet* 2008;**40**:1156–9.
- Barton A, Thomson W, Ke X, et al. Re-evaluation of putative rheumatoid arthritis susceptibility genes in the post-genome wide association study era and hypothesis of a key pathway underlying susceptibility. *Hum Mol Genet* 2008;**17**:2274–9.
- Barton A, Eyre S, Ke X, et al. Identification of AF4/FMR2 family, member 3 (*AFF3*) as a novel rheumatoid arthritis susceptibility locus and confirmation of two further pan-autoimmune susceptibility genes. *Hum Mol Genet* 2009;**18**:2518–22.
- Dubois PC, Trynka G, Franke L, et al. Multiple common variants for celiac disease influencing immune gene expression. *Nat Genet* 2010;**42**:295–302.
- Radstake TR, Gorlova O, Rueda B, et al. Genome-wide association study of systemic sclerosis identifies *CD247* as a new susceptibility locus. *Nat Genet* 2010;**42**:426–9.
- Zhernakova A, Stahl EA, Trynka G, et al. Meta-analysis of genome-wide association studies in celiac disease and rheumatoid arthritis identifies fourteen non-HLA shared loci. *PLoS Genet* 2011;**7**:e1002004.
- Todd JA, Walker NM, Cooper JD, et al. Robust associations of four new chromosome regions from genome-wide analyses of type 1 diabetes. *Nat Genet* 2007;**39**:857–64.
- Smyth DJ, Plagnol V, Walker NM, et al. Shared and distinct genetic variants in type 1 diabetes and celiac disease. *N Engl J Med* 2008;**359**:2767–77.
- Barrett JC, Hansoul S, Nicolae DL, et al. Genome-wide association defines more than 30 distinct susceptibility loci for Crohn's disease. *Nat Genet* 2008;**40**:955–62.
- Lowe CE, Cooper JD, Brusko T, et al. Large-scale genetic fine mapping and genotype-phenotype associations implicate polymorphism in the *IL2RA* region in type 1 diabetes. *Nat Genet* 2007;**39**:1074–82.
- Maier LM, Lowe CE, Cooper J, et al. *IL2RA* genetic heterogeneity in multiple sclerosis and type 1 diabetes susceptibility and soluble interleukin-2 receptor production. *PLoS Genet* 2009;**5**:e1000322.
- Tchilian EZ, Beverley PC. Altered CD45 expression and disease. *Trends Immunol* 2006;**27**:146–53.
- Hinks A, Martin P, Flynn E, et al. Subtype specific genetic associations for juvenile idiopathic arthritis: ERAP1 with the enthesitis related arthritis subtype and *IL23R* with juvenile psoriatic arthritis. *Arthritis Res Ther* 2011;**13**:R12.
- Stock CJ, Ogilvie EM, Samuel JM, et al. Comprehensive association study of genetic variants in the *IL-1* gene family in systemic juvenile idiopathic arthritis. *Genes Immun* 2008;**9**:349–57.
- Hinks A, Martin P, Flynn E, et al. Association of the *CCR5* gene with juvenile idiopathic arthritis. *Genes Immun* 2010;**11**:584–9.
- Albers HM, Kurreeman FA, Stoeken-Rijsbergen G, et al. Association of the autoimmunity locus 4q27 with juvenile idiopathic arthritis. *Arthritis Rheum* 2009;**60**:901–4.
- Prahalad S, Bohnsack JF, Jorde LB, et al. Association of two functional polymorphisms in the *CCR5* gene with juvenile rheumatoid arthritis. *Genes Immun* 2006;**7**:468–75.



Investigation of rheumatoid arthritis susceptibility loci in juvenile idiopathic arthritis confirms high degree of overlap

Anne Hinks, Joanna Cobb, Marc Sudman, et al.

Ann Rheum Dis 2012 71: 1117-1121 originally published online January 31, 2012

doi: 10.1136/annrheumdis-2011-200814

Updated information and services can be found at:

<http://ard.bmj.com/content/71/7/1117.full.html>

	<i>These include:</i>
Data Supplement	"Web Only Data" http://ard.bmj.com/content/suppl/2012/01/31/annrheumdis-2011-200814.DC1.html
References	This article cites 33 articles, 9 of which can be accessed free at: http://ard.bmj.com/content/71/7/1117.full.html#ref-list-1
Open Access	This paper is freely available online under the BMJ Journals unlocked scheme, see http://ard.bmj.com/info/unlocked.dtl
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (275 articles)
[Connective tissue disease](#) (3093 articles)
[Degenerative joint disease](#) (3346 articles)
[Immunology \(including allergy\)](#) (3659 articles)
[Musculoskeletal syndromes](#) (3592 articles)
[Rheumatoid arthritis](#) (2338 articles)
[Genetics](#) (699 articles)

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>